

# REGULARIZATION MATRICES FOR DISCRETE ILL-POSED PROBLEMS IN SEVERAL SPACE-DIMENSIONS

Laura Dykes\*, Guangxin Huang†, Silvia Noschese‡, and Lothar Reichel§

**Abstract.** Many applications in science and engineering require the solution of large linear discrete ill-posed problems that are obtained by the discretization of a Fredholm integral equation of the first kind in several space-dimensions. The matrix that defines these problems is very ill-conditioned and generally numerically singular, and the right-hand side, which represents measured data, typically is contaminated by measurement error. Straightforward solution of these problems generally is not meaningful due to severe error propagation. Tikhonov regularization seeks to alleviate this difficulty by replacing the given linear discrete ill-posed problem by a penalized least-squares problem, whose solution is less sensitive to the error in the right-hand side and to round-off errors introduced during the computations. This paper discusses the construction of penalty terms that are determined by solving a matrix-nearness problem. These penalty terms allow partial transformation to standard form of Tikhonov regularization problems that stem from the discretization of integral equations on a cube in several space-dimensions.

**Key words.** Discrete ill-posed problems; Tikhonov regularization; standard form problems; matrix nearness problems; Krylov subspace iterative methods

**AMS subject classifications.** 65F10; 65F22; 65R30

**1. Introduction.** We consider the solution of linear discrete ill-posed problems that arise from the discretization of a Fredholm integral equation of the first kind on a cube in two or more space-dimensions. Discretization of the integral operator yields a matrix  $K \in \mathbb{R}^{M \times N}$ , which we assume to be large. A vector  $\mathbf{b} \in \mathbb{R}^M$  that represents measured data, and therefore is error-contaminated, is available and we would like to compute an approximate solution of the least-square problem

$$\min_{\mathbf{x} \in \mathbb{R}^N} \|K\mathbf{x} - \mathbf{b}\|. \quad (1.1)$$

The matrix  $K$  has many “tiny” singular values of different orders of magnitude. This makes  $K$  severely ill-conditioned; in fact,  $K$  may be numerically singular. Least-squares problems (1.1) with a matrix of this kind are commonly referred to as linear discrete ill-posed problems.

Let  $\mathbf{e} \in \mathbb{R}^M$  denote the (unknown) error in  $\mathbf{b}$ . Thus,

$$\mathbf{b} = \widehat{\mathbf{b}} + \mathbf{e},$$

where  $\widehat{\mathbf{b}} \in \mathbb{R}^M$  stands for the unknown error-free vector associated with  $\mathbf{b}$ . We will assume the unavailable linear system of equations

$$K\mathbf{x} = \widehat{\mathbf{b}} \quad (1.2)$$

to be consistent.

---

\*University School, Hunting Valley, OH 44022, USA. E-mail: [ldykes@math.kent.edu](mailto:ldykes@math.kent.edu).

†Geomathematics Key Laboratory of Sichuan, Department of Information and Computing Science, Chengdu University of Technology, Chengdu, 610059, P. R. China. E-mail: [huangx@cdut.edu.cn](mailto:huangx@cdut.edu.cn).

‡Department of Mathematics, SAPIENZA Università di Roma, P.le A. Moro, 2, I-00185 Roma, Italy. E-mail: [noschese@mat.uniroma1.it](mailto:noschese@mat.uniroma1.it).

§Department of Mathematical Sciences, Kent State University, Kent, OH 44242, USA. E-mail: [reichel@math.kent.edu](mailto:reichel@math.kent.edu).

Let  $K^\dagger$  denote the Moore–Penrose pseudoinverse of the matrix  $K$ . We are interested in determining the solution  $\hat{\mathbf{x}}$  of (1.2) of minimal Euclidean norm; it is given by  $K^\dagger \hat{\mathbf{b}}$ . We note that the solution of minimal Euclidean norm of (1.1),

$$K^\dagger \mathbf{b} = K^\dagger \hat{\mathbf{b}} + K^\dagger \mathbf{e} = \hat{\mathbf{x}} + K^\dagger \mathbf{e},$$

generally is not a meaningful approximation of  $\hat{\mathbf{x}}$  due to a large propagated error  $K^\dagger \mathbf{e}$ . This difficulty stems from the fact that  $\|K^\dagger\|$  is large. Throughout this paper  $\|\cdot\|$  denotes the Euclidean vector norm or spectral matrix norm. We also will use the Frobenius norm of a matrix  $A$  and the associated inner product between two commensurate matrices  $A_1$  and  $A_2$ , defined by

$$\|A\|_F = \sqrt{\text{trace}(A^T A)}, \quad \langle A_1, A_2 \rangle = \text{trace}(A_1^T A_2),$$

respectively. The superscript  $T$  stands for transposition.

To avoid severe error propagation, one replaces the least-squares problem (1.1) by a nearby problem, whose solution is less sensitive to the error  $\mathbf{e}$  in  $\mathbf{b}$ . This replacement is known as regularization. Tikhonov regularization, which is one of the most popular regularization methods, replaces (1.1) by a penalized least-squares problem of the form

$$\min_{\mathbf{x} \in \mathbb{R}^N} \{ \|K\mathbf{x} - \mathbf{b}\|^2 + \mu \|L\mathbf{x}\|^2 \}, \quad (1.3)$$

where  $L \in \mathbb{R}^{J \times N}$  is referred to as the regularization matrix and the scalar  $\mu > 0$  as the regularization parameter; see, e.g., [2, 8, 10]. We assume the matrix  $L$  to be chosen so that

$$\mathcal{N}(K) \cap \mathcal{N}(L) = \{\mathbf{0}\},$$

where  $\mathcal{N}(M)$  denotes the null space of the matrix  $M$ . Then the minimization problem (1.3) has a unique solution

$$\mathbf{x}_\mu = (K^T K + \mu L^T L)^{-1} K^T \mathbf{b}$$

for any  $\mu > 0$ .

Common choices of  $L$  include the identity matrix and discretizations of differential operators. The Tikhonov minimization problem (1.3) is said to be in *standard form* when  $L = I$ ; otherwise it is in *general form*. Numerous computed examples in the literature, see, e.g., [4, 5, 9, 26], illustrate that the choice of  $L$  may be important for the quality of the computed approximation  $\mathbf{x}_\mu$  of  $\hat{\mathbf{x}}$ . The regularization matrix  $L$  should be chosen so that known important features of the desired solution  $\hat{\mathbf{x}}$  of (1.2) are not damped. This can be achieved by choosing  $L$  so that  $\mathcal{N}(L)$  contains vectors that represent these features, because vectors in  $\mathcal{N}(L)$  are not damped by  $L$ .

Several approaches to construct regularization matrices with desirable properties are described in the literature; see, e.g., [1, 4, 5, 6, 12, 19, 23, 24, 26, 28]. Huang et al. [16] proposed the construction of square regularization matrices with a user-specified null space by solving a matrix nearness problem in the Frobenius norm. The regularization matrices so obtained are designed for linear discrete ill-posed problems in one space-dimension. This paper extends this approach to problems in higher space-dimensions. The regularization matrices of this paper generalize those applied by Bouhamidi and Jbilou [1] by allowing a user-specified null space.



Following Bouhamidi and Jbilou [1], we consider square matrices  $K$  with a tensor product structure, i.e.,

$$K = K^{(2)} \otimes K^{(1)}. \quad (1.10)$$

We assume for simplicity that  $K^{(1)}, K^{(2)} \in \mathbb{R}^{n \times n}$  with  $N = n^2$ . However, we note that the regularization matrices described in this paper can be applied also when the matrix  $K$  in (1.1) does not have a tensor product structure.

Bouhamidi and Jbilou [1] are concerned with applications to image restoration and achieve restorations of high quality. However, for linear discrete ill-posed problems in one space-dimension, analysis presented in [4, 12] indicates that the regularization matrix (1.6), with a non-trivial null space, may give approximate solutions of higher quality than the matrix (1.9), which has a trivial null space. This depends on that the latter matrix may introduce artifacts close to the boundary; see also [5, 6, 26] for related discussions and illustrations. It is the aim of the present paper to develop a generalization of the regularization matrix (1.8) that has a non-trivial null space. The null spaces in the regularization matrices described in this paper allow an additive constant or linear growth in some or all space-dimensions without damping. It is also easy to construct regularization matrices with null spaces that are chosen not to damp certain shapes in specified positions by choosing a null space that contains a representation of these shapes.

Our approach to define regularization matrices generalizes the technique proposed in [16] from one to several space-dimensions. Specifically, the regularization matrix is defined by solving a matrix nearness problem in the Frobenius norm. The regularization matrix so obtained allows a partial transformation of the Tikhonov regularization problem (1.3) to standard form. When the matrix  $K$  is square, Arnoldi-type iterative solution methods can be used. Arnoldi-type iterative solution methods often require fewer matrix-vector product evaluations than iterative solution methods based on Golub–Kahan bidiagonalization, because they do not require matrix-vector product evaluations with  $K^T$ ; see, e.g., [22] for illustrations. A nice recent survey of iterative solution methods for linear discrete ill-posed problems is provided by Gazzola et al. [9].

This paper is organized as follows. Section 2 describes our construction of new regularization matrices for problems in two space-dimensions. The section also discusses iterative methods for the solution of the Tikhonov minimization problems obtained. We consider both the situation when  $K$  is a general matrix and when  $K$  has tensor product structure. Section 3 generalizes the results of Section 2 to more than two space-dimensions. Computed examples can be found in Section 4, and Section 5 contains concluding remarks.

We conclude this section by noting that while this paper focuses on iterative solution methods for large-scale Tikhonov minimization problems (1.3), the regularization matrices described also can be applied in direct solution methods for small to medium-sized problems that are based on the generalized singular value decomposition (GSVD); see, e.g., [7, 10] for discussions.

**2. Regularization matrices for problems in two space-dimensions.** Many image restoration problems as well as problems from certain other applications (1.1) have a matrix  $K \in \mathbb{R}^{N \times N}$  that is the Kronecker product of two matrices  $K^{(1)}, K^{(2)} \in \mathbb{R}^{n \times n}$  with  $N = n^2$ , cf. (1.10). We will consider this situation in most of this section; the case when  $K$  is a general square matrix without Kronecker product structure is

commented on at the end of the section. Extension to rectangular matrices  $K$ ,  $K^{(1)}$ , and  $K^{(2)}$  is straightforward.

We will use regularization matrices with a Kronecker product structure,

$$L = L^{(2)} \otimes L^{(1)} \quad (2.1)$$

and will discuss the choice of square regularization matrices  $L^{(1)}, L^{(2)} \in \mathbb{R}^{n \times n}$ . The following result is an extension of [16, Proposition 2.1] to problems with a Kronecker product structure. Let  $\mathcal{R}(M)$  denote the range of the matrix  $M$ . Throughout this section  $N = n^2$ .

**PROPOSITION 2.1.** *Let the matrices  $V^{(1)} \in \mathbb{R}^{n \times \ell_1}$  and  $V^{(2)} \in \mathbb{R}^{n \times \ell_2}$  have orthonormal columns, and let  $\mathcal{B}$  denote the subspace of matrices of the form  $B = B^{(2)} \otimes B^{(1)}$ , where the null space of  $B^{(i)} \in \mathbb{R}^{n \times n}$  contains  $\mathcal{R}(V^{(i)})$  for  $i = 1, 2$ . Introduce for  $i = 1, 2$  the orthogonal projectors  $P^{(i)} = I - V^{(i)}V^{(i)T}$  with null space  $\mathcal{R}(V^{(i)})$ , and define  $P = P^{(2)} \otimes P^{(1)}$ . Let  $A = A^{(2)} \otimes A^{(1)}$  with  $A^{(i)} \in \mathbb{R}^{n \times n}$ ,  $i = 1, 2$ . Then  $\hat{A} = AP$  is a closest matrix to  $A = A^{(2)} \otimes A^{(1)}$  in  $\mathcal{B}$  in the Frobenius norm.*

*Proof.* The matrix  $\hat{A}$  satisfies the following conditions:

1.  $\hat{A} \in \mathcal{B}$ ;
2. if  $A \in \mathcal{B}$ , then  $\hat{A} \equiv A$ ;
3. if  $B \in \mathcal{B}$ , then  $\langle B, A - \hat{A} \rangle = 0$ .

In fact,

$$\hat{A}(V^{(2)} \otimes V^{(1)}) = A(P^{(2)}V^{(2)} \otimes P^{(1)}V^{(1)}) = 0,$$

which shows the first property. The fact that  $A \in \mathcal{B}$  implies that

$$A^{(2)}V^{(2)} = 0, \quad A^{(1)}V^{(1)} = 0,$$

from which it follows that

$$\hat{A} = (A^{(2)} - A^{(2)}V^{(2)}V^{(2)T}) \otimes (A^{(1)} - A^{(1)}V^{(1)}V^{(1)T}) = A^{(2)} \otimes A^{(1)} = A.$$

Finally, for any  $B \in \mathcal{B}$  of the form  $B = B^{(2)} \otimes B^{(1)}$ , one has that

$$B^{(2)}V^{(2)} = V^{(2)T}B^{(2)T} = 0, \quad B^{(1)}V^{(1)} = V^{(1)T}B^{(1)T} = 0,$$

so that

$$\begin{aligned} \langle B, A - \hat{A} \rangle &= \text{trace}(B^T A - B^T \hat{A}) \\ &= \text{trace}(B^{(2)T} A^{(2)}) \text{trace}(B^{(1)T} A^{(1)} V^{(1)} V^{(1)T}) \\ &\quad + \text{trace}(B^{(2)T} A^{(2)} V^{(2)} V^{(2)T}) \text{trace}(B^{(1)T} A^{(1)}) \\ &\quad - \text{trace}(B^{(2)T} A^{(2)} V^{(2)} V^{(2)T}) \text{trace}(B^{(1)T} A^{(1)} V^{(1)} V^{(1)T}) = 0, \end{aligned}$$

where the last equality follows from the cyclic property of the trace.  $\square$

**Example 2.1.** Let  $L_2$  and  $\tilde{L}_2$  be defined by (1.6) and (1.9), respectively. Proposition 2.1 shows that a closest matrix to  $\tilde{L} = \tilde{L}_2 \otimes \tilde{L}_2$  in the Frobenius norm with null space  $\mathcal{N}(L_2 \otimes L_2)$  is

$$L = \tilde{L}_2 P_2 \otimes \tilde{L}_2 P_2,$$

where  $P_2$  is the orthogonal projector onto  $\mathcal{N}(L_2)^\perp$ .  $\square$

Example 2.2. Define the nonsingular square bidiagonal regularization matrix

$$\tilde{L}_1 = \frac{1}{2} \begin{bmatrix} 1 & -1 & & & 0 \\ & 1 & -1 & & \\ & & 1 & -1 & \\ & & & \ddots & \ddots \\ 0 & & & & -1 \\ & & & & & 1 \end{bmatrix} \in \mathbb{R}^{n \times n}. \quad (2.2)$$

A closest matrix to  $\tilde{L} = \tilde{L}_1 \otimes \tilde{L}_1$  in the Frobenius norm with null space  $\mathcal{N}(L_1 \otimes L_1)$  is given by

$$L = \tilde{L}_1 P_1 \otimes \tilde{L}_1 P_1,$$

where  $P_1$  is the orthogonal projector onto  $\mathcal{N}(L_1)^\perp$ ; see, e.g., [16].  $\square$

The following result is concerned with the situation when the order of the nonsingular matrices  $\tilde{L}_i$  and projectors  $P_i$  in Examples 2.1 and 2.2 is reversed. We first consider the case when  $\tilde{L}$  is a square matrix without Kronecker product structure, since this situation is of independent interest.

**PROPOSITION 2.2.** *Let  $\tilde{L} \in \mathbb{R}^{n \times n}$  and let  $\mathcal{V}$  be a subspace of  $\mathbb{R}^n$ . Define the orthogonal projector  $P_{\mathcal{V}^\perp}$  onto  $\mathcal{V}^\perp$ . Then the closest matrix  $\hat{L} \in \mathbb{R}^{n \times n}$  to  $\tilde{L}$  in the Frobenius norm, such that  $\mathcal{R}(\hat{L}) \subset \mathcal{V}^\perp$ , is given by  $\hat{L} = P_{\mathcal{V}^\perp} \tilde{L}$ .*

*Proof.* Consider the problem of determining a closest matrix  $\hat{L}^T \in \mathbb{R}^{n \times n}$  to  $\tilde{L}^T$  in the Frobenius norm whose null space contains  $\mathcal{V}$ . It is shown in [16, Proposition 2.3] that  $\hat{L}^T = \tilde{L}^T P_{\mathcal{V}^\perp}$  is such a matrix. The Frobenius norm is invariant under transposition and orthogonal projectors are symmetric. Therefore,

$$\|\tilde{L}^T P_{\mathcal{V}^\perp} - \tilde{L}^T\|_F = \|P_{\mathcal{V}^\perp} \tilde{L} - \tilde{L}\|_F.$$

Moreover,  $\mathcal{R}(P_{\mathcal{V}^\perp}) = \mathcal{V}^\perp$ . It follows that a closest matrix to  $\tilde{L}$  in the Frobenius norm whose range is a subset of  $\mathcal{V}^\perp$  is given by  $P_{\mathcal{V}^\perp} \tilde{L}$ .  $\square$

The following result extends Proposition 2.2 to matrices with a tensor product structure. We formulate the result similarly as Proposition 2.1.

**COROLLARY 2.3.** *Let the matrices  $V^{(1)} \in \mathbb{R}^{n \times \ell_1}$  and  $V^{(2)} \in \mathbb{R}^{n \times \ell_2}$  have orthonormal columns, and let  $\mathcal{B}$  denote the subspace of matrices of the form  $B = B^{(2)} \otimes B^{(1)}$ , where the range of  $B^{(i)} \in \mathbb{R}^{n \times n}$  is orthogonal to  $\mathcal{R}(V^{(i)})$  for  $i = 1, 2$ . Introduce for  $i = 1, 2$  the orthogonal projectors  $P^{(i)} = I - V^{(i)} V^{(i)T}$  and define  $P = P^{(2)} \otimes P^{(1)}$ . Let  $A = A^{(2)} \otimes A^{(1)}$  with  $A^{(i)} \in \mathbb{R}^{n \times n}$ ,  $i = 1, 2$ . Then  $\hat{A} = PA$  is a closest matrix to  $A$  in  $\mathcal{B}$  in the Frobenius norm.*

*Proof.* The result can be shown by applying Propositions 2.1 or 2.2.  $\square$

Example 2.3. Let  $L_2$  be defined by (1.6) and  $\tilde{L}_2$  by (1.9). Corollary 2.3 shows that a closest matrix to  $\tilde{L} = \tilde{L}_2 \otimes \tilde{L}_2$  with range in  $\mathcal{R}(L_2 \otimes L_2)$  is

$$L = P_2 \tilde{L}_2 \otimes P_2 \tilde{L}_2,$$

where  $P_2 = \text{diag}[0, 1, 1, \dots, 1, 0] \in \mathbb{R}^{n \times n}$ .  $\square$

Example 2.4. Let the matrices  $L_1$  and  $\tilde{L}_1$  be given by (1.5) and (2.2). It follows from Corollary 2.3 that a closest matrix to  $\tilde{L} = \tilde{L}_1 \otimes \tilde{L}_1$  with range in  $\mathcal{R}(L_1 \otimes L_1)$  is given by

$$L = P_1 \tilde{L}_1 \otimes P_1 \tilde{L}_1,$$

where  $P_1 = \text{diag}[1, 1, \dots, 1, 0] \in \mathbb{R}^{n \times n}$ .  $\square$

Using (1.10) and (2.1), the Tikhonov regularization problem (1.3) can be expressed as

$$\min_{\mathbf{x} \in \mathbb{R}^N} \left\{ \|(K^{(2)} \otimes K^{(1)})\mathbf{x} - \mathbf{b}\|^2 + \mu \|(L^{(2)} \otimes L^{(1)})\mathbf{x}\|^2 \right\}. \quad (2.3)$$

It is convenient to introduce the operator  $\text{vec}$ , which transforms a matrix  $Y \in \mathbb{R}^{n \times n}$  to a vector of size  $n^2$  by stacking the columns of  $Y$ . Let  $A$ ,  $B$ , and  $Y$ , be matrices of commensurate sizes. Then

$$\text{vec}(AYB) = (B^T \otimes A)\text{vec}(Y);$$

see, e.g., [15] for operations on matrices with Kronecker product structure. We can apply this identity to express (2.3) in the form

$$\min_{X \in \mathbb{R}^{n \times n}} \left\{ \|K^{(1)}XK^{(2)T} - B\|_F^2 + \mu \|L^{(1)}XL^{(2)T}\|_F^2 \right\}, \quad (2.4)$$

where the matrix  $B \in \mathbb{R}^{n \times n}$  satisfies  $\mathbf{b} = \text{vec}(B)$ .

Let the regularization matrices in (2.4) be of the forms

$$L^{(1)} = P^{(1)}\tilde{L}^{(1)}, \quad L^{(2)} = P^{(2)}\tilde{L}^{(2)},$$

where the matrices  $\tilde{L}^{(i)} \in \mathbb{R}^{n \times n}$  are nonsingular and the  $P^{(i)}$  are orthogonal projectors. We easily can transform (2.4) to a form with an orthogonal projector regularization matrix,

$$\min_{Y \in \mathbb{R}^{n \times n}} \left\{ \|K_1^{(1)}YK_1^{(2)T} - B\|_F^2 + \mu \|P^{(1)}YP^{(2)}\|_F^2 \right\}, \quad (2.5)$$

where

$$K_1^{(i)} = K^{(i)}(\tilde{L}^{(i)})^{-1}, \quad i = 1, 2.$$

We will solve (2.5) by an iterative method. The structure of the minimization problem makes it convenient to apply an iterative method based on the global Arnoldi process, which was introduced and first analyzed by Jbilou et al. [17, 18]. We refer to matrices with many more rows than columns as ‘‘block vectors’’. The block vectors  $U, W \in \mathbb{R}^{N \times n}$  are said to be *F-orthogonal* if  $\langle U, W \rangle = 0$ ; they are *F-orthonormal* if in addition  $\|U\|_F = \|W\|_F = 1$ .

The application of  $k$  steps of the global Arnoldi method to the solution of (2.5) yields an *F-orthonormal* basis  $\{V_1, V_2, \dots, V_k\}$  of block vectors  $V_j$  for the block Krylov subspace

$$\mathcal{K}_k = \text{span}\{B, K_1^{(1)}BK_1^{(2)T}, \dots, (K_1^{(1)})^{k-1}B(K_1^{(2)T})^{k-1}\}. \quad (2.6)$$

In particular  $V_1 = B/\|B\|_F$ . The use of the global Arnoldi method to the solution of (2.5) is mathematically equivalent to applying a standard Arnoldi method to (2.3). An advantage of the global Arnoldi method is that it can be implemented by using matrix-matrix operations, while the standard Arnoldi method applies matrix-vector and vector-vector operations. This can lead to faster execution of the global Arnoldi method on many modern computers. Algorithm 1 outlines the global Arnoldi method; see [17, 18] for further discussions of this and other block methods.

---

**Algorithm 1:** Global Arnoldi for computing an  $F$ -orthonormal basis for (2.6)

---

- 1 compute  $V_1 = B/\|B\|_F$
  - 2 for  $j = 1, 2, \dots, k$  compute
    - 3  $V = K_1^{(1)}V_j$
    - 4  $V = VK_1^{(2)T}$
    - 5 for  $i = 1, 2, \dots, j$ 
      - 6  $h_{i,j} = \langle V, V_i \rangle$
      - 7  $V = V - h_{i,j}V_i$
    - 8 end
    - 9  $h_{j+1,j} = \|V\|_F$
    - 10 if  $h_{j+1,j} = 0$  stop
    - 11  $V_{j+1} = V/h_{j+1,j}$
    - 12 end
  - 13 construct the  $n \times kn$  matrix  $\widehat{V}_k = [V_1, \dots, V_k]$  with  $F$ -orthonormal block columns  $V_j$ . The block columns span the space (2.6)
  - 14 construct the  $(k+1) \times k$  Hessenberg matrix  $\widetilde{H}_k = [h_{i,j}]_{i=1,2,\dots,k+1, j=1,2,\dots,k}$
- 

We determine an approximate solution of (2.5) in the global Arnoldi subspace (2.6). This is described by Algorithm 2 for a given  $\mu > 0$ . The solution subspace (2.6) is independent of the orthogonal projectors that determine the regularization term in (2.5). This approach to generate a solution subspace for the solution of Tikhonov minimization problems in general form was first discussed in [13]; see also [9] for examples.

---

**Algorithm 2:** Tikhonov regularization based on the global Arnoldi process

---

- 1 construct  $\widehat{V}_k = [V_1, V_2, \dots, V_k]$  and  $\widetilde{H}_k$  using Algorithm 1
- 2 solve for a given  $\mu > 0$ ,

$$\min_{\mathbf{y} \in \mathbb{R}^k} \left\{ \left\| \widetilde{H}_k \mathbf{y} - \|B\|_F \mathbf{e}_1 \right\|^2 + \mu \left\| \sum_{i=1}^k y_i P^{(1)} V_i P^{(2)} \right\|_F^2 \right\}, \quad (2.7)$$

where  $\mathbf{e}_1 = [1, 0, \dots, 0]^T \in \mathbb{R}^{k+1}$  and  $\mathbf{y} = [y_1, y_2, \dots, y_k]^T$

- 3 compute  $Y_{\mu,k} = \sum_{i=1}^k V_i y_i$
- 

We briefly comment on the evaluation of the penalty term in (2.7). Let  $M_i = P^{(1)} V_i P^{(2)}$ ,  $1 \leq i \leq k$ . Then the penalty term can be expressed as

$$\left\| \sum_{i=1}^k y_i M_i \right\|_F^2 = \text{trace} \left( \left( \sum_{i=1}^k y_i M_i^T \right) \left( \sum_{j=1}^k y_j M_j \right) \right) = \sum_{i,j=1}^k y_i y_j \text{trace} (M_i^T M_j).$$

Introduce the matrix  $N = [n_{i,j}] \in \mathbb{R}^{k \times k}$  with elements  $n_{i,j} = \text{trace} (M_i^T M_j)$ . Then (2.7) can be expressed as

$$\min_{\mathbf{y} \in \mathbb{R}^k} \left\{ \left\| \widetilde{H}_k \mathbf{y} - \|B\|_F \mathbf{e}_1 \right\|^2 + \mu \mathbf{y}^T N \mathbf{y} \right\}. \quad (2.8)$$



There are many techniques for determining a suitable value of the regularization parameter  $\mu$  in (2.8) including the discrepancy principle and the generalized cross validation (GCV) criterion; see, e.g., [10, 20, 25]. The discrepancy principle is a popular approach to determine the regularization parameter when a bound  $\varepsilon$  for the norm of the error  $\mathbf{e}$  in  $\mathbf{b}$  is known, i.e.,  $\|\mathbf{e}\| \leq \varepsilon$ . It prescribes that  $\mu > 0$  be chosen so that the computed approximate solution  $Y_{\mu,k}$  of (2.5) satisfies

$$\|K_2^{(1)} Y_{\mu,k} K_2^{(2)T} - B\|_F = \eta \varepsilon, \quad (2.9)$$

where  $\eta \geq 1$  is a user-chosen constant independent of  $\varepsilon$ . The nonlinear equation (2.9) for  $\mu$  can be solved by a variety of methods such as Newton's method; see, e.g., [13] for a discussion.

We note that the regularization matrices of this section also can be applied when the matrix  $K$  in (1.3) does not have a Kronecker product structure (1.10). Let  $\mathbf{x} = \text{vec}(X)$ . Then the matrix expression in the penalty term of (2.5) can be written as

$$\text{vec}(P^{(1)} \tilde{L}^{(1)} X \tilde{L}^{(2)T} P^{(2)}) = ((P^{(2)} \tilde{L}^{(2)}) \otimes (P^{(1)} \tilde{L}^{(1)})) \mathbf{x} = (P^{(2)} \otimes P^{(1)}) (\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}) \mathbf{x}.$$

The analogue of the minimization problem (2.5) therefore can be expressed as

$$\min_{\mathbf{x} \in \mathbb{R}^N} \left\{ \|K \mathbf{x} - \mathbf{b}\|^2 + \mu \|(P^{(2)} \otimes P^{(1)}) (\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}) \mathbf{x}\|^2 \right\}. \quad (2.10)$$

The matrix  $\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$  is invertible; we have  $(\tilde{L}^{(2)} \otimes \tilde{L}^{(1)})^{-1} = (\tilde{L}^{(2)})^{-1} \otimes (\tilde{L}^{(1)})^{-1}$ . It follows that the problem (2.10) can be transformed to

$$\min_{\mathbf{y} \in \mathbb{R}^N} \left\{ \|K ((\tilde{L}^{(2)})^{-1} \otimes (\tilde{L}^{(1)})^{-1}) \mathbf{y} - \mathbf{b}\|^2 + \mu \|(P^{(2)} \otimes P^{(1)}) \mathbf{y}\|^2 \right\}. \quad (2.11)$$

The matrix  $P^{(2)} \otimes P^{(1)}$  is an orthogonal projector. It is described in [23] how Tikhonov regularization problems with a regularization term that is determined by an orthogonal projector with a low-dimensional null space easily can be transformed to standard form. However,  $\dim(\mathcal{N}(P^{(2)} \otimes P^{(1)})) \geq n$ , which generally is quite large in problems of interest to us. It is therefore impractical to transform the Tikhonov minimization problem (2.11) to standard form. We can solve (2.11), e.g., by generating a (standard) Krylov subspace determined by the matrix  $\hat{K} = K((\tilde{L}^{(2)})^{-1} \otimes (\tilde{L}^{(1)})^{-1})$  and vector  $\mathbf{b}$ , similarly as described in [13]. When the matrix  $K$  is square, then also the matrix  $\hat{K}$  is square, and the Arnoldi process can be applied to  $\hat{K}$  to generate a solution subspace; when  $K$  is rectangular, partial Golub–Kahan bidiagonalization of  $\hat{K}$  can be used. The latter approach requires matrix-vector product evaluations with both  $\hat{K}$  and  $\hat{K}^T$ ; see [13] for further details. The matrices  $(\tilde{L}^{(i)})^{-1}$ ,  $i = 1, 2$ , of course, do not have to be explicitly formed.

### 3. Regularization matrices for problems in higher space-dimensions.

Proposition 2.1 can be extended to higher space-dimensions. In addition to allowing  $d \geq 2$  space-dimensions, we remove the requirement that all blocks be square and of equal size.

**PROPOSITION 3.1.** *Let  $V_{\ell_i}^{(i)} \in \mathbb{R}^{n_i \times \ell_i}$  have  $1 \leq \ell_i < n_i$  orthonormal columns for  $i = 1, 2, \dots, d$ , and let  $\mathcal{B}$  denote the subspace of matrices of the form*

$$B = B^{(d)} \otimes B^{(d-1)} \otimes \dots \otimes B^{(1)},$$

where the null space of  $B^{(i)} \in \mathbb{R}^{p_i \times n_i}$  contains  $\mathcal{R}(V_{\ell_i}^{(i)})$  for all  $i$ . Let  $I_k$  denote the identity matrix of order  $k$  and define the orthogonal projectors

$$P = P^{(d)} \otimes P^{(d-1)} \otimes \dots \otimes P^{(1)}, \quad P^{(i)} = I_{n_i} - V_{\ell_i}^{(i)} V_{\ell_i}^{(i)T}, \quad i = 1, 2, \dots, d. \quad (3.1)$$

Let  $A = A^{(d)} \otimes A^{(d-1)} \otimes \dots \otimes A^{(1)}$ , where  $A^{(i)} \in \mathbb{R}^{p_i \times n_i}$ ,  $i = 1, 2, \dots, d$ . Then  $\hat{A} = AP$  is a closest matrix to  $A$  in  $\mathcal{B}$  in the Frobenius norm.

*Proof.* The proof is a straightforward modification of the proof of Proposition 2.1.

□

Let  $\tilde{L}^{(1)}, \tilde{L}^{(2)}, \dots, \tilde{L}^{(d)}$  be a sequence of square nonsingular matrices, and let  $L^{(1)}, L^{(2)}, \dots, L^{(d)}$  be regularization matrices with desirable null spaces. It follows from Proposition 3.1 that a closest matrix to

$$\tilde{L} = \tilde{L}^{(d)} \otimes \tilde{L}^{(d-1)} \otimes \dots \otimes \tilde{L}^{(1)}$$

with null space  $\mathcal{N}(L^{(d)} \otimes L^{(d-1)} \otimes \dots \otimes L^{(1)})$  is

$$L = \tilde{L}^{(d)} P^{(d)} \otimes \tilde{L}^{(d-1)} P^{(d-1)} \otimes \dots \otimes \tilde{L}^{(1)} P^{(1)},$$

where the orthogonal projectors  $P^{(i)}$  are defined by (3.1) and the matrix  $V_{\ell_i}^{(i)} \in \mathbb{R}^{n_i \times \ell_i}$  has  $1 \leq \ell_i < n_i$  orthonormal columns that span  $\mathcal{N}(L^{(i)})$  for  $i = 1, 2, \dots, d$ .

The following result generalizes Corollary 2.3 to higher space-dimensions and to rectangular blocks of different sizes.

**PROPOSITION 3.2.** *Let  $V_{\ell_i}^{(i)} \in \mathbb{R}^{n_i \times \ell_i}$  have  $1 \leq \ell_i < n_i$  orthonormal columns for  $i = 1, 2, \dots, d$ , and let  $\mathcal{B}$  denote the subspace of matrices of the form*

$$B = B^{(d)} \otimes B^{(d-1)} \otimes \dots \otimes B^{(1)},$$

where the range of  $B^{(i)} \in \mathbb{R}^{p_i \times n_i}$  is orthogonal to  $\mathcal{R}(V_{\ell_i}^{(i)})$  for all  $i$ . Let  $P$  be defined by (3.1) and let  $A = A^{(d)} \otimes A^{(d-1)} \otimes \dots \otimes A^{(1)}$ , where  $A^{(i)} \in \mathbb{R}^{p_i \times n_i}$ ,  $i = 1, 2, \dots, d$ . Then  $\hat{A} = PA$  is a closest matrix to  $A$  in  $\mathcal{B}$  in the Frobenius norm.

*Proof.* The result can be shown by modifying the proof of Propositions 2.1 or 2.2.

□

Let  $L^{(1)}, L^{(2)}, \dots, L^{(d)}$  be a sequence of regularization matrices with desirable ranges, and let  $\tilde{L}^{(1)}, \tilde{L}^{(2)}, \dots, \tilde{L}^{(d)}$  be full rank matrices. It follows from Proposition 3.2 that a closest matrix to

$$\tilde{L} = \tilde{L}^{(d)} \otimes \tilde{L}^{(d-1)} \otimes \dots \otimes \tilde{L}^{(1)}$$

with range in  $\mathcal{R}(L^{(d)} \otimes L^{(d-1)} \otimes \dots \otimes L^{(1)})$  is

$$L = P^{(d)} \tilde{L}^{(d)} \otimes P^{(d-1)} \tilde{L}^{(d-1)} \otimes \dots \otimes P^{(1)} \tilde{L}^{(1)},$$

where the orthogonal projectors  $P^{(i)}$  are defined by (3.1) and the matrix  $V_{\ell_i}^{(i)} \in \mathbb{R}^{n_i \times \ell_i}$  has  $1 \leq \ell_i < n_i$  orthonormal columns that span  $\mathcal{N}(L^{(i)})$  for  $i = 1, 2, \dots, d$ .

We conclude this section with an extension of (2.5) to higher space-dimensions and assume that the problem has nested tensor structure, i.e.,

$$K^{(i)} = K^{(i,2)} \otimes K^{(i,1)},$$

where  $K^{(1,i)} \in \mathbb{R}^{n_i \times n_i}$ ,  $K^{(2,i)} \in \mathbb{R}^{s_i \times s_i}$ ,  $i = 1, 2$ , and that

$$B = B^{(2)} \otimes B^{(1)},$$

where  $B^{(i)} \in \mathbb{R}^{n_i \times s_i}$  for  $i = 1, 2$ . The minimization problem (1.1) with

$$K = K^{(2,2)} \otimes K^{(2,1)} \otimes K^{(1,2)} \otimes K^{(1,1)}$$

and  $\mathbf{b} = \text{vec}(B)$  reads

$$\min_{X \in \mathbb{R}^{n \times s}} \left\{ \|(K^{(1,2)} \otimes K^{(1,1)})X(K^{(2,2)T} \otimes K^{(2,1)T}) - B^{(2)} \otimes B^{(1)}\|_F^2 \right\}.$$

Let the regularization matrices have a nested tensor structure

$$L^{(i)} = L^{(i,2)} \otimes L^{(i,1)}, \quad i = 1, 2.$$

The penalized least-squares problem to be solved is of the form

$$\min_{X \in \mathbb{R}^{n \times s}} \left\{ \|(K^{(1,2)} \otimes K^{(1,1)})X(K^{(2,2)T} \otimes K^{(2,1)T}) - B^{(2)} \otimes B^{(1)}\|_F^2 + \mu \|(L^{(1,2)} \otimes L^{(1,1)})X(L^{(2,2)T} \otimes L^{(2,1)T})\|_F^2 \right\}. \quad (3.2)$$

If, moreover, the solution is separable of the form  $X = X^{(2)} \otimes X^{(1)}$ , where  $X^{(i)} \in \mathbb{R}^{n_i \times s_i}$  for  $i = 1, 2$ , then we obtain the minimization problem

$$\min_{\substack{X^{(1)} \in \mathbb{R}^{n_1 \times s_1} \\ X^{(2)} \in \mathbb{R}^{n_2 \times s_2}}} \left\{ \|(K^{(1,2)} X^{(2)} K^{(2,2)T}) \otimes (K^{(1,1)} X^{(1)} K^{(2,1)T}) - B^{(2)} \otimes B^{(1)}\|_F^2 + \mu \|(L^{(1,2)} X^{(2)} L^{(2,2)T}) \otimes (L^{(1,1)} X^{(1)} L^{(2,1)T})\|_F^2 \right\}. \quad (3.3)$$

When the regularization matrices are of the form  $L^{(i,j)} = P^{(i,j)} \tilde{L}^{(i,j)}$ ,  $1 \leq i, j \leq 2$ , where the  $P^{(i,j)}$  are orthogonal projectors and the  $\tilde{L}^{(i,j)}$  are square and invertible, the minimization problems (3.2) and (3.3) can be transformed similarly as equation (2.4) was transformed into (2.5).

**4. Computed examples.** We illustrate the performance of regularization matrices of the form  $L = L^{(2)} \otimes L^{(1)}$  with  $L^{(i)} = P^{(i)} \tilde{L}^{(i)}$  or  $L^{(i)} = \tilde{L}^{(i)} P^{(i)}$  for  $i = 1, 2$ , and compare with the regularization matrices  $L^{(i)} = \tilde{L}^{(i)}$  for  $i = 1, 2$ . The noise level is given by

$$\nu := \frac{\|E\|_F}{\|\hat{B}\|_F}.$$

Here  $E = B - \hat{B}$  is the error matrix,  $B$  is the available error-contaminated matrix in (2.4), and  $\hat{B}$  is the associated unknown error-free matrix, i.e.,  $\hat{\mathbf{b}} = \text{vec}(\hat{B})$  in (1.2). In all examples, the entries of the matrix  $E$  are normally distributed with zero mean and are scaled to correspond to a specified noise level. We let  $\eta = 1.01$  in (2.9) in all examples. The quality of computed approximate solutions  $X_{\mu,k}$  of (2.4) is measured with the relative error norm

$$e_k := \frac{\|X_{\mu,k} - \hat{X}\|_F}{\|\hat{X}\|_F},$$

where  $\hat{X}$  is the desired solution of the unknown error-free problem, i.e.,  $\hat{\mathbf{x}} = \text{vec}(\hat{X})$ .

The number of (outer) iterations,  $k$ , in Algorithm 2 is determined by the following stopping criteria:

1. The relative change of the computed approximate solution  $Y_{\mu_k, k}$  drops below a user-specified threshold  $\tau > 0$ , i.e., we terminate the iterations as soon as

$$\|Y_{\mu_k, k} - Y_{\mu_{k-1}, k-1}\|_F / \|Y_{\mu_{k-1}, k-1}\|_F < \tau.$$

2. The number of (outer) iterations,  $k$ , is bounded by  $k_{max}$ .

Thus, the computations are terminated as soon as one of these criteria is satisfied.

We set  $k_{max} = 24$  and  $\tau = 5 \cdot 10^{-4}$  in Example 4.1, and  $k_{max} = 30$  and  $\tau = 1 \cdot 10^{-4}$  for Examples 4.2-4.4. The choices of  $k_{max}$  and  $\tau$  are such that the computed solution does not change much with the iteration number when the iterations are terminated. In particular, nearby choices of these parameters yield computed solutions of about the same quality. All computations were carried out in MATLAB R2017a with about 15 significant decimal digits on a laptop computer with an Intel Core i7-6700HQ CPU @ 2.60GHz processor and 16GB RAM.

regularization matrix	$k$	relative error $e_k$
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	24	$8.30 \cdot 10^{-2}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	24	$8.23 \cdot 10^{-2}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	24	$8.23 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	24	$9.34 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	24	$8.15 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	19	$9.13 \cdot 10^{-2}$

TABLE 4.1

Example 4.1: Number of iterations  $k$  and relative error  $e_k$  in computed approximate solutions  $X_{\mu_k, k}$  determined by Tikhonov regularization based on the global Arnoldi process and several regularization matrices for noise level  $\nu = 1 \cdot 10^{-3}$ .




FIG. 4.1. Example 4.1: Computed approximate solution  $X_{\mu_k}$  for noise level  $\nu = 1 \cdot 10^{-3}$  and regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  using the discrepancy principle.

Example 4.1. Consider the Fredholm integral equation of the first kind in two space-dimensions,

$$\int \int_{\Omega} \kappa(\tau, \sigma; x, y) f(x, y) dx dy = g(\tau, \sigma), \quad (\tau, \sigma) \in \Omega, \quad (4.1)$$

where  $\Omega = [-\pi/2, \pi/2] \times [-\pi/2, \pi/2]$ . The kernel is given by

$$\kappa(\tau, \sigma; x, y) = \kappa_1(\tau, x)\kappa_1(\sigma, y), \quad (\tau, \sigma), (x, y) \in \Omega,$$

where

$$\kappa_1(\tau, \sigma) = (\cos(\sigma) + \sin(\tau))^2 \left( \frac{\sin(\xi)}{\xi} \right)^2, \quad \xi = \pi(\sin(\sigma) + \cos(\tau)).$$

The right-hand side function is of the form

$$g(\tau, \sigma) = h(\tau)h(\sigma),$$

where  $h(\sigma)$  is chosen so that the solution is the sum of two Gaussian functions and a constant. We use the MATLAB code `shaw` from [11] to discretize (4.1) by a Galerkin method with  $1000 \times 1000$  orthonormal box functions as test and trial functions. This code produces the matrix  $K \in \mathbb{R}^{1000 \times 1000}$  that approximates the analogue of the integral operator (4.1) in one space-dimension, and a discrete approximate solution  $\mathbf{x}_1$  in one space-dimension. Adding the vector  $\mathbf{n}_1 = [1, 1, \dots, 1]^T$  yields the vector  $\hat{\mathbf{x}}_1 \in \mathbb{R}^{1000}$ , from which we construct the scaled discrete approximation  $\hat{X} = \hat{\mathbf{x}}_1 \hat{\mathbf{x}}_1^T$  of the solution of (4.1). The error-free right-hand side is computed by  $\hat{B} = K \hat{X} K^T$ . The error matrix  $E \in \mathbb{R}^{1000 \times 1000}$  models white Gaussian noise with noise levels  $\nu = 1 \cdot 10^{-3}$ . The data matrix  $B$  in (2.4) is computed as  $B = \hat{B} + E$ . The regularization matrices  $L$  used are constructed like in Examples 2.1-2.4. We compare the performance of these regularization matrices to the performance of the nonsingular regularization matrices  $L = \tilde{L}^{(i)} \otimes \tilde{L}^{(i)}$ ,  $i = 1, 2$ . We note that the regularization matrix for  $i = 1$  does not damp an additive constant in the computed solution, while the regularization matrix for  $i = 2$  allows linear growth of the computed solution in both the horizontal and vertical directions without damping. We also apply the regularization matrix  $L = \tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$ , which allows an arbitrary additive constant in the computed solution without damping, as well as linear growth of the computed solution in the horizontal direction.

The regularization parameter  $\mu$  is essentially determined by the discrepancy principle. More precisely, we first determine  $\mu$  so that the discrepancy principle holds and then multiply  $\mu$  by 0.9. We observed ‘‘under regularization’’ in this manner to improve the quality of the computed approximate solutions in all examples of this section. We therefore applied this technique in all examples.

Table 4.1 displays results obtained for the different regularization matrices. The table shows the regularization matrix  $P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$  to yield the smallest relative errors. Figure 4.1 shows the computed approximate solution for the noise level  $\nu = 1 \cdot 10^{-3}$  when the regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  is used. The computed approximation cannot be visually distinguished from the desired exact solution  $\hat{X}$ . We therefore do not show the latter.  $\square$

Example 4.2. We consider the restoration of the test image `satellite`, which is represented by an array of  $256 \times 256$  pixels. The available image, represented by the matrix  $B \in \mathbb{R}^{256 \times 256}$ , is corrupted by Gaussian blur and additive zero-mean white Gaussian noise; it is shown in Figure 4.2(a). Figure 4.2(b) displays the desired blur- and noise-free image. It is represented by the matrix  $\hat{X} \in \mathbb{R}^{256 \times 256}$ , and is assumed not to be known. The blurring matrices  $K^{(i)} \in \mathbb{R}^{256 \times 256}$ ,  $i = 1, 2$ , are Toeplitz matrices. We let  $K^{(1)} = K^{(2)} = K$ , where  $K$  is analogous to the matrix generated by the MATLAB function `blur` from [11] using the parameter values `band = 5` and

regularization matrix	$k$	$\mu_k$	relative error $e_k$
noise level $\nu = 1 \cdot 10^{-2}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	17	$7.84 \cdot 10^{-2}$	$8.60 \cdot 10^{-2}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	9	$6.84 \cdot 10^{-2}$	$8.38 \cdot 10^{-2}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	28	$7.53 \cdot 10^{-2}$	$8.55 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	13	$1.22 \cdot 10^{-1}$	$8.75 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	21	$1.09 \cdot 10^{-1}$	$8.62 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	13	$1.16 \cdot 10^{-1}$	$8.69 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	17	$2.25 \cdot 10^{-1}$	$8.90 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	13	$1.94 \cdot 10^{-1}$	$8.73 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	17	$2.17 \cdot 10^{-1}$	$8.86 \cdot 10^{-2}$
noise level $\nu = 1 \cdot 10^{-3}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	11	$1.29 \cdot 10^{-3}$	$1.95 \cdot 10^{-2}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	12	$1.11 \cdot 10^{-3}$	$1.81 \cdot 10^{-2}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	11	$1.26 \cdot 10^{-3}$	$1.92 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	20	$1.42 \cdot 10^{-3}$	$2.08 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	20	$1.23 \cdot 10^{-3}$	$1.93 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	11	$1.41 \cdot 10^{-3}$	$1.94 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	20	$1.64 \cdot 10^{-3}$	$2.16 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	20	$1.42 \cdot 10^{-3}$	$2.00 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	20	$1.57 \cdot 10^{-3}$	$2.11 \cdot 10^{-2}$
noise level $\nu = 1 \cdot 10^{-4}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	19	$9.04 \cdot 10^{-5}$	$2.40 \cdot 10^{-3}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	20	$7.66 \cdot 10^{-5}$	$2.08 \cdot 10^{-3}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	19	$8.66 \cdot 10^{-5}$	$2.32 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	19	$1.25 \cdot 10^{-4}$	$3.03 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	19	$1.06 \cdot 10^{-4}$	$2.64 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	19	$9.68 \cdot 10^{-5}$	$2.45 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	30	$3.24 \cdot 10^{-4}$	$6.54 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	30	$2.82 \cdot 10^{-4}$	$5.82 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	19	$1.23 \cdot 10^{-4}$	$2.88 \cdot 10^{-3}$

TABLE 4.2

Example 4.2: Number of iterations  $k$ , regularization parameter  $\mu_k$ , and relative error  $e_k$  in computed approximate solutions  $X_{\mu_k, k}$  determined by Tikhonov regularization based on the global Arnoldi process for three noise levels and several regularization matrices.

$\sigma = 1.5$ . We show results for the noise levels  $\nu = 1 \cdot 10^{-j}$ ,  $j = 2, 3, 4$ . The data matrix  $B$  in (2.4) is determined similarly as in Example 4.1.

Table 4.2 shows the regularization parameters  $\mu_k$  and the relative errors  $e_k$  in the computed approximate solutions  $X_{\mu_k, k}$  determined by the global Arnoldi process with data matrices contaminated by noise of levels  $\nu = 1 \cdot 10^{-j}$ ,  $j = 2, 3, 4$ , for several regularization matrices. The iterations are terminated as soon as the discrepancy principle can be satisfied and the regularization parameter then is chosen so that (2.9) holds. Table 4.2 shows the global Arnoldi process with the regularization




FIG. 4.2. *Example 4.2: (a) Available blur- and noise-contaminated satellite image represented by the matrix  $B$ , (b) desired image, (c) restored image for the noise level  $\nu = 1 \cdot 10^{-3}$  and regularization matrix  $P^{(1)}\tilde{L}^{(1)} \otimes P^{(1)}\tilde{L}^{(1)}$ , and (d) restored image for the same noise level and regularization matrix  $P^{(2)}\tilde{L}^{(2)} \otimes P^{(2)}\tilde{L}^{(2)}$ .*

matrix  $P^{(1)}\tilde{L}^{(1)} \otimes P^{(1)}\tilde{L}^{(1)}$  to determine the best approximations of  $\hat{X}$  for all noise levels. Figures 4.2(c) and 4.2(d) show the computed approximate solutions determined by the global Arnoldi process with  $\nu = 1 \cdot 10^{-3}$  and the regularization matrices  $P^{(1)}\tilde{L}^{(1)} \otimes P^{(1)}\tilde{L}^{(1)}$  and  $P^{(2)}\tilde{L}^{(2)} \otimes P^{(2)}\tilde{L}^{(2)}$ , respectively. The quality of the computed restorations is visually indistinguishable.  $\square$

**Example 4.3.** This example is similar to the previous one; only the image to be restored differs. Here we consider the restoration of the test image `QRcode`, which is represented by an array of  $256 \times 256$  pixels corrupted by Gaussian blur and additive zero-mean white Gaussian noise. Figure 4.3(a) shows the corrupted image that we would like to restore. It is represented by the matrix  $B \in \mathbb{R}^{256 \times 256}$ . The desired blur- and noise-free image is depicted in Figure 4.3(b). The blurring matrices  $K^{(i)} \in \mathbb{R}^{256 \times 256}$ ,  $i = 1, 2$ , are Toeplitz matrices. They are generated like in Example

regularization matrix	$k$	$\mu_k$	relative error $e_k$
noise level $\nu = 1 \cdot 10^{-2}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	30	$4.86 \cdot 10^{-1}$	$4.96 \cdot 10^{-2}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	20	$4.13 \cdot 10^{-1}$	$4.76 \cdot 10^{-2}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	30	$4.30 \cdot 10^{-1}$	$4.82 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	30	$9.42 \cdot 10^{-1}$	$5.05 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	30	$8.12 \cdot 10^{-1}$	$4.84 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	30	$8.42 \cdot 10^{-1}$	$4.90 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	27	$2.77 \cdot 10^0$	$5.06 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	27	$2.45 \cdot 10^0$	$4.90 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	30	$2.50 \cdot 10^0$	$4.95 \cdot 10^{-2}$
noise level $\nu = 1 \cdot 10^{-3}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	16	$1.00 \cdot 10^{-2}$	$1.32 \cdot 10^{-2}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	16	$8.33 \cdot 10^{-3}$	$1.25 \cdot 10^{-2}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	16	$9.56 \cdot 10^{-2}$	$1.30 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	20	$1.32 \cdot 10^{-2}$	$1.35 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	17	$1.11 \cdot 10^{-2}$	$1.28 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	20	$1.25 \cdot 10^{-2}$	$1.33 \cdot 10^{-2}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	17	$1.90 \cdot 10^{-2}$	$1.40 \cdot 10^{-2}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	13	$1.84 \cdot 10^{-2}$	$1.37 \cdot 10^{-2}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	17	$1.80 \cdot 10^{-2}$	$1.38 \cdot 10^{-2}$
noise level $\nu = 1 \cdot 10^{-4}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	15	$4.11 \cdot 10^{-4}$	$2.10 \cdot 10^{-3}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	16	$3.40 \cdot 10^{-4}$	$1.83 \cdot 10^{-3}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	16	$3.89 \cdot 10^{-4}$	$2.00 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	15	$5.21 \cdot 10^{-4}$	$2.34 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	15	$4.27 \cdot 10^{-4}$	$2.06 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	15	$4.35 \cdot 10^{-4}$	$2.09 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	13	$8.27 \cdot 10^{-4}$	$2.98 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	13	$6.78 \cdot 10^{-4}$	$2.68 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	14	$4.88 \cdot 10^{-4}$	$2.18 \cdot 10^{-3}$

TABLE 4.3

Example 4.3: Number of iterations  $k$ , regularization parameter  $\mu_k$ , and relative error  $e_k$  in computed approximate solutions  $X_{\mu_k, k}$  determined by Tikhonov regularization based on the global Arnoldi process for three noise levels and several regularization matrices.

4.2. The regularization matrices  $L$  are the same as in Example 4.2.

Table 4.3 is analogous to Table 4.2. The table shows the regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  to give the most accurate approximations of  $\hat{X}$ . Figures 4.3(c) and 4.3(d) show the restorations determined for  $\nu = 1 \cdot 10^{-3}$  with the regularization matrices  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  and  $P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$ , respectively. One cannot visually distinguish the quality of these restorations.  $\square$

**Example 4.4.** This example is similar to the previous one; only the image to be restored is larger. Here we consider the restoration of the test image **Barbara**,






FIG. 4.3. *Example 4.3: (a) Available blur- and noise-contaminated QR code image represented by the matrix  $B$ , (b) desired image, (c) restored image for the noise level  $\nu = 1 \cdot 10^{-3}$  and regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$ , and (d) restored image for the same noise level and regularization matrix  $P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$ .*

which is represented by an array of  $510 \times 510$  pixels corrupted by Gaussian blur and additive zero-mean white Gaussian noise. Figure 4.4(a) shows the corrupted image that we would like to restore. It is represented by the matrix  $B \in \mathbb{R}^{510 \times 510}$ . The desired blur- and noise-free image is depicted in Figure 4.4(b). The blurring matrices  $K^{(i)} \in \mathbb{R}^{510 \times 510}$ ,  $i = 1, 2$ , are Toeplitz matrices. They are generated like in Example 4.2. The regularization matrices  $L$  are the same as in Example 4.2.

Table 4.4 is analogous to Table 4.2. The table shows the regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  to yield the most accurate approximations of  $\hat{X}$ . Figures 4.3(c) and 4.3(d) show the restorations determined for  $\nu = 1 \cdot 10^{-3}$  with the regularization matrices  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$  and  $P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$ , respectively.  $\square$

**5. Concluding remarks.** This paper presents a novel method to determine regularization matrices for discrete ill-posed problems in several space-dimensions by

regularization matrix	$k$	$\mu_k$	relative error $e_k$
noise level $\nu = 1 \cdot 10^{-3}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	12	$1.35 \cdot 10^{-2}$	$8.44 \cdot 10^{-3}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	13	$1.14 \cdot 10^{-2}$	$7.93 \cdot 10^{-3}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	13	$1.28 \cdot 10^{-2}$	$8.30 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	12	$1.58 \cdot 10^{-2}$	$8.57 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	13	$1.36 \cdot 10^{-2}$	$8.08 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	12	$1.52 \cdot 10^{-2}$	$8.44 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	17	$4.26 \cdot 10^{-2}$	$9.32 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	12	$3.85 \cdot 10^{-2}$	$8.68 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	17	$4.11 \cdot 10^{-2}$	$9.21 \cdot 10^{-3}$
noise level $\nu = 1 \cdot 10^{-4}$			
$\tilde{L}^{(1)} \otimes \tilde{L}^{(1)}$	12	$9.20 \cdot 10^{-4}$	$1.52 \cdot 10^{-3}$
$P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$	12	$7.76 \cdot 10^{-4}$	$1.41 \cdot 10^{-3}$
$\tilde{L}^{(1)} P^{(1)} \otimes \tilde{L}^{(1)} P^{(1)}$	12	$8.70 \cdot 10^{-4}$	$1.49 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(1)}$	12	$1.16 \cdot 10^{-3}$	$1.61 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(1)} \tilde{L}^{(1)}$	12	$9.88 \cdot 10^{-4}$	$1.50 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(1)} P^{(1)}$	12	$9.96 \cdot 10^{-3}$	$1.51 \cdot 10^{-3}$
$\tilde{L}^{(2)} \otimes \tilde{L}^{(2)}$	11	$2.68 \cdot 10^{-3}$	$2.02 \cdot 10^{-3}$
$P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$	11	$2.33 \cdot 10^{-3}$	$1.87 \cdot 10^{-3}$
$\tilde{L}^{(2)} P^{(2)} \otimes \tilde{L}^{(2)} P^{(2)}$	11	$1.72 \cdot 10^{-3}$	$1.66 \cdot 10^{-3}$

TABLE 4.4

Example 4.4: Number of iterations  $k$ , regularization parameter  $\mu_k$ , and relative error  $e_k$  in computed approximate solutions  $X_{\mu_k, k}$  determined by Tikhonov regularization based on the global Arnoldi process for two noise levels and several regularization matrices.

solving a matrix nearness problem. Numerical examples illustrate the effectiveness of the regularization matrices determined in this manner. While all examples use the discrepancy principle to determine a suitable value of the regularization parameter, other parameter choice rules also can be applied; see, e.g., [10, 20, 25] for discussions on alternate techniques.

**Acknowledgment.** The authors would like to thank the referees for comments. Research by GH was supported in part by the Fund of Application Foundation of Science and Technology Department of the Sichuan Province (2016JY0249) and NNSF (11501392, 2017YFC0601505, 41672325), research by SN was partially supported by SAPIENZA Università di Roma and INdAM-GNCS, and research by LR was partially supported by NSF grants DMS-1729509 and DMS-1720259.

## REFERENCES

- [1] A. Bouhamidi and K. Jbilou, *Sylvester Tikhonov-regularization methods in image restoration*, J. Comput. Appl. Math., 206 (2007), pp. 86–98.
- [2] C. Brezinski, M. Redivo-Zaglia, G. Rodriguez, and S. Seatzu, *Extrapolation techniques for ill-conditioned linear systems*, Numer. Math., 81 (1998), pp. 1–29.
- [3] D. Calvetti, B. Lewis, and L. Reichel, *A hybrid GMRES and TV-norm based method for image restoration*, in Advanced Signal Processing Algorithms, Architectures, and Implementa-



FIG. 4.4. *Example 4.4:* (a) Available blur- and noise-contaminated *Barbara* image represented by the matrix  $B$ , (b) desired image, (c) restored image for the noise level  $\nu = 1 \cdot 10^{-3}$  and regularization matrix  $P^{(1)} \tilde{L}^{(1)} \otimes P^{(1)} \tilde{L}^{(1)}$ , and (d) restored image for the same noise level and regularization matrix  $P^{(2)} \tilde{L}^{(2)} \otimes P^{(2)} \tilde{L}^{(2)}$ .

- tions XII, ed. F. T. Luk, Proceedings of the Society of Photo-Optical Instrumentation Engineers (SPIE), vol. 4791, The International Society for Optical Engineering, Bellingham, WA, 2002, pp. 192–200.
- [4] D. Calvetti, L. Reichel, and A. Shuibi, *Invertible smoothing preconditioners for linear discrete ill-posed problems*, Appl. Numer. Math., 54 (2005), pp. 135–149.
- [5] M. Donatelli, A. Neuman, and L. Reichel, *Square regularization matrices for large linear discrete ill-posed problems*, Numer. Linear Algebra Appl., 19 (2012), pp. 896–913.
- [6] M. Donatelli and L. Reichel, *Square smoothing regularization matrices with accurate boundary conditions*, J. Comput. Appl. Math., 272 (2014), pp. 334–349.
- [7] L. Dykes, S. Noschese, and L. Reichel, *Rescaling the GSVD with application to ill-posed problems*, Numer. Algorithms, 68 (2015), pp. 531–545.
- [8] H. W. Engl, M. Hanke, and A. Neubauer, *Regularization of Inverse Problems*, Kluwer, Dordrecht, 1996.
- [9] S. Gazzola, P. Novati, and M. R. Russo, *On Krylov projection methods and Tikhonov regularization*, Electron. Trans. Numer. Anal., 44 (2015), pp. 83–123.
- [10] P. C. Hansen, *Rank-Deficient and Discrete Ill-Posed Problems*, SIAM, Philadelphia, 1998.

- [11] P. C. Hansen, *Regularization tools version 4.0 for MATLAB 7.3*, Numer. Algorithms, 46 (2007), pp. 189–194.
- [12] P. C. Hansen and T. K. Jensen, *Smoothing norm preconditioning for regularizing minimum residual methods*, SIAM J. Matrix Anal. Appl., 29 (2006), pp. 1–14.
- [13] M. E. Hochstenbach and L. Reichel, *An iterative method for Tikhonov regularization with a general linear regularization operator*, J. Integral Equations Appl., 22 (2010), pp. 463–480.
- [14] M. E. Hochstenbach, L. Reichel, and X. Yu, *A Golub–Kahan-type reduction method for matrix pairs*, J. Sci. Comput., 65 (2015), pp. 767–789.
- [15] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
- [16] G. Huang, S. Noschese, and L. Reichel, *Regularization matrices determined by matrix nearness problems*, Linear Algebra Appl., 502 (2016), pp. 41–57.
- [17] K. Jbilou, A. Messaoudi, and H. Sadok, *Global FOM and GMRES algorithms for matrix equations*, Appl. Numer. Math., 31 (1999), pp. 49–63.
- [18] K. Jbilou, H. Sadok, and A. Tinzette, *Oblique projection methods for linear systems with multiple right-hand sides*, Electron. Trans. Numer. Anal., 20 (2005), pp. 119–138.
- [19] M. E. Kilmer, P. C. Hansen, and M. I. Español, *A projection-based approach to general-form Tikhonov regularization*, SIAM J. Sci. Comput., 29 (2007), pp. 315–330.
- [20] S. Kindermann, *Convergence analysis of minimization-based noise level-free parameter choice rules for linear ill-posed problems*, Electron. Trans. Numer. Anal., 38 (2011), pp. 233–257.
- [21] A. Lanza, S. Morigi, L. Reichel, and F. Sgallari, *A generalized Krylov subspace method for  $\ell_p$ - $\ell_q$  minimization*, SIAM J. Sci. Comput., 37 (2015), pp. S30–S50.
- [22] B. Lewis and L. Reichel, *Arnoldi–Tikhonov regularization methods*, J. Comput. Appl. Math., 226 (2009), pp. 92–102.
- [23] S. Morigi, L. Reichel, and F. Sgallari, *Orthogonal projection regularization operators*, Numer. Algorithms, 44 (2007), pp. 99–114.
- [24] S. Noschese and L. Reichel, *Inverse problems for regularization matrices*, Numer. Algorithms, 60 (2012), pp. 531–544.
- [25] L. Reichel and G. Rodriguez, *Old and new parameter choice rules for discrete ill-posed problems*, Numer. Algorithms, 63 (2013), pp. 65–87.
- [26] L. Reichel and Q. Ye, *Simple square smoothing regularization operators*, Electron. Trans. Numer. Anal., 33 (2009), pp. 63–83.
- [27] L. Reichel and X. Yu, *Matrix decompositions for Tikhonov regularization*, Electron. Trans. Numer. Anal., 43 (2015), pp. 223–243.
- [28] L. Reichel and X. Yu, *Tikhonov regularization via flexible Arnoldi reduction*, BIT, 55 (2015), pp. 1145–1168.