

Article

When functionalisation comes in useful: ionic liquids with a “sweet” appended moiety demonstrate drastically reduced toxicological effects

Floriana Billeci, Francesca D'Anna, Marta Feroci, Patrizia Cancemi, Salvatore Feo, Antonella Forlino, Francesca Tonnelli, Kenneth Richard Seddon, H. Q. Nimal Gunaratne, and Natalia V Plechkova

ACS Sustainable Chem. Eng., Just Accepted Manuscript • DOI: 10.1021/acsuschemeng.9b05507 • Publication Date (Web): 09 Dec 2019

Downloaded from pubs.acs.org on December 12, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

When functionalisation comes in useful: ionic liquids with a “sweet” appended moiety demonstrate drastically reduced toxicological effects

Floriana Billeci,^{1,2} Francesca D’Anna,^{2} Marta Feroci,³ Patrizia Cancemi,⁴ Salvatore Feo,⁴*

Antonella Forlino,⁵ Francesca Tonnelli,⁵ Kenneth R. Seddon, H. Q. Nimal Gunaratne^{1,6,}*

Natalia V. Plechkova^{1,7}

¹The QUILL Research Centre, School of Chemistry and Chemical Engineering Queen's University of Belfast, Stranmillis Road, Belfast, Northern Ireland BT9 5AG, UK.

E-mail: n.gunaratne@qub.ac.uk

²Università degli Studi di Palermo, Dipartimento STEBICEF, Viale delle Scienze, Ed. 17 90128 Palermo (Italy). E-mail: francesca.danna@unipa.it.

³Dept. SBAI, Sapienza University of Rome, via Castro Laurenziano, 7, 00161 Rome, Italy.

⁴Università degli Studi di Palermo, Dipartimento STEBICEF, Viale delle Scienze, Ed. 16 90128
Palermo (Italy).

⁵Dept Molecular Medicine, Unit of Biochemistry, University of Pavia, 27100 Pavia (Italy).

⁶School of Chemistry and Chemical Engineering Queen's University of Belfast, Stranmillis
Road, Belfast, Northern Ireland BT9 5AG, UK

⁷Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and
Biomedical Sciences, The Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL, UK

KEYWORDS. ionic liquids, gluconic acid derivatives, ecotoxicity, cytotoxicity.

ABSTRACT. The growing number of applications of ionic liquids (ILs) in industry have brought attention to the green credentials of synthesis, as well as their cyto- and ecotoxicities both for their use and accidental leakage into the environment.

With the above in mind, we designed a class of ILs with either cations bearing gluconamide motif and aliphatic side chains or the anion incorporating gluconic acid (derived from food waste) moiety. An ionic liquid with imidazolium cation with an appended gluconic amide (bearing 5 hydroxyl groups) moiety was also synthesised for a useful comparison.

Different structural features were considered, placing emphasis on the nature and length of the alkyl chain and the nature of the anion. For comparison, two ILs with one hydroxyl group and another without any hydroxyl-groups were prepared.

Apart from the typical characterisation of ILs, such as differential scanning calorimetry (DSC), thermal gravimetric analysis (TGA), conductivity and viscosity measurements, the main focus was placed on the evaluation of their cyto- and ecotoxicities, performing tests with three cell cancer lines (HeLa, HTC-116 and MCF-7) and fish embryos (Zebrafish).

Data obtained have shed light on the relationship working between the physico-chemical properties and the structural features of the ILs. Interestingly, these ILs are able to inhibit cell growth only at very high concentrations ($IC_{50} \sim 10^{-3}$ M) and they do not affect the vitality of fish embryos, allowing them to be classified as harmless solvents.

To our surprise, the data collected show that derivatisation with the “sweet” residue on the imidazolium cation completely removes the well known toxicity effects of imidazolium ILs.

KEYWORDS: ionic liquids, gluconic acid, cytotoxicity, ecotoxicity, physico-chemical properties

INTRODUCTION

Ionic liquids (ILs), organic salts with melting temperatures lower than 100 °C,¹ have been considered for a long time as an advantageous alternative to using conventional organic solvents. Their high thermal stability, low vapor pressure and flammability favoured their application as solvents,² catalysts,³ and reagents.⁴ As more applications of ILs appear, there is also a growing attention to their biodegradability and toxicity.⁵⁻⁹

Since a large number of ILs are water soluble, they can persist in the environment and interact with living organisms.¹⁰⁻¹¹ Furthermore, their degradation and persistence in the environment are heavily dependent both on the cation and anion structure.¹⁰ As a consequence, over the past decade, there has been an increasing emphasis on the design and synthesis of ILs (focussing on the greener aspects).¹²⁻¹⁵

Among the factors considered are: the nature of the cation, the length of the alkyl chain attached to it and the presence of functional groups on the chain. Each of these plays a significant role in determining the toxicity and biodegradability of the ILs.^{6,16}


In particular, biodegradation and toxicology investigations have highlighted that chemical functionalities like hydroxyl, carboxyl and ester groups could, if inserted in the IL structure, make them less harmful towards the ecosystem.^{17,18}

Aliphatic cations are usually more biodegradable and less toxic than the corresponding aromatic ones¹⁹, and phosphonium ILs may show toxicity similar to that of imidazolium ILs.²⁰

Imidazolium ILs are probably the most widely used ILs, from an industrial point of view, they are applied in, for example, BASILTM and CELLIONICTM processes.^{21,22} They are also used in lithium batteries, catalytic and separation processes,²¹⁻²⁵ notwithstanding the issues concerning toxicity. Indeed, it is well known that the degradation of imidazolium cations leads to the persistence of some cation fragments in the environment.²⁶ To decrease and eventually eliminate the toxicity of aromatic imidazolium ILs is a present day challenge. In this respect, Gathergood *et al.* have demonstrated that ether and ester groups on the side chain(s) of imidazolium-based ILs can reduce their bacterial toxicity,²⁷ as the ester bond represents a site of possible enzymatic cleavage.¹² Furthermore, in certain cases, etherase and sulfatase enzymes can also take part in the degradation

processes.¹² Greener ILs have also been obtained using reagents derived from natural amino acids and organic acids.²⁸⁻²⁹

With this in mind, we synthesised new eco-friendly ILs bearing a motif derived from gluconic acid (Scheme 1). In all cases, the cation is indicated as $[N_{xyzGAw}]^+$ where x and y represent the length of alkyl groups depending on the diamine used, z is the number of the carbon atoms of the alkyl spacer of diamine, GA stands for gluconic amide and w represents the length of the alkyl chain used in the quaternization step.



Scheme 1. Structure of a) gluconic acid in equilibrium with glucono- δ -lactone; b) amide-based ILs; c) gluconate-based ILs.

Gluconic acid is a natural organic acid obtained from food waste.³⁰⁻³¹ Gluconic acid is obtained from the oxidation of glucose (Scheme 1a), and in nature, this process is carried out by several bacteria and fungi.³²⁻³⁴ This organic acid can be produced by chemical, electrochemical and biochemical methods.³⁵ Moreover, the use of gluconate in several medicaments and medical research as an anion for salts like potassium, calcium, and iron led to the safe use of this molecule

1
2
3 in preserving human health.³⁶⁻³⁸ Different studies have demonstrated that this organic acid is safe
4
5 for the human metabolism.^{39,40}
6
7

8
9 All the above considerations led us to supposing that the presence of such a residue on the cation
10
11 or anion structure of ILs could allow obtaining salts of both low cyto- and ecotoxicity.
12
13 Furthermore, the origin of gluconic acid and its wide availability make it inexpensive and
14
15 economically sustainable.
16
17

18
19 From a structural point of view, “sweet” ILs based on this molecule take advantage of the presence
20
21 of the oligo-hydroxylated chain. The presence of several OH groups could play a dual function.
22
23 Indeed, as stated above, it could contribute to decreasing the IL toxicity, but it also endows the
24
25 structure with high coordinating ability. By incorporating this feature, some of these ILs have been
26
27 obtained⁴¹⁻⁴³ and in some cases, they have been applied to create ionic liquid gels and
28
29 thermochromic systems aimed at environmental preservation.⁴⁴⁻⁴⁵
30
31

32
33 With respect to previous reports in literature,⁴¹⁻⁴³ our synthetic approach allowed inserting a
34
35 gluconic-like residue on the cation structure. Consequently, ILs in this work can be separated into
36
37 two different classes, discerning gluconate- and gluconamide-based ILs.
38
39
40

41
42 The ILs structures were designed to evaluate the roles played by the alkyl chain length and
43
44 branching, as well as the length of the spacer on the cation and the anion nature. Some previous
45
46 reports have highlighted that branching on the alkyl chain could induce a significant decrease in
47
48 the ILs toxicity with respect to the corresponding linear ones.⁴⁶
49
50

51
52 ILs obtained were fully characterised by determining the physico-chemical properties such as the
53
54 TGA, DSC, viscosity and conductivity.
55
56
57
58
59
60

Apart from considerations of a “greener” ionic liquid synthesis, their cyto- and ecotoxicities were assessed (with regard to potential leakage of ILs).

In particular, three human cell lines, namely HeLa, HTC-116 and MCF-7 from cervical, colon and breast carcinoma were chosen as prototypes of human cells. Furthermore, haemolysis investigations were performed by using ovine blood cells.

As far as the aquatic toxicity is concerned, Zebrafish are considered as an ideal model for assessing the developmental toxicity of exposure to toxicants during the early stages of life. Several studies have been widely applied to test IL toxicity⁴⁷ as a consequence of its transparency and rapid development of embryos. Moreover, the Zebrafish shares 70% of its genes with humans and, if orthologous genes are considered, 47% of human genes have a one-to-one relationship with Zebrafish genes.^{48,49}

In light of all the above considerations, the main aim of this work was to obtain a new collection of ILs based on their structure, moieties (derived from renewable sources) and ensuring both low cyto- and ecotoxicities. This should contribute to making their whole life cycle more sustainable. By focussing on synthetic procedures, we were able to identify a protocol that is able to produce ILs in good yields with a minimum impact on the environment.

EXPERIMENTAL SECTION

Materials. Glucono- δ -lactone (Sigma Aldrich, ≥ 99.0 %), *N,N*-dimethylethylenediamine (TCI, > 98.0 %), *N,N*-dimethyl-1,3-propanediamine (Sigma Aldrich, 99.0 %), 1-(3-aminopropyl)imidazole (Sigma Aldrich, ≥ 97 %), 2-ethylhexyl bromide (Sigma Aldrich, 95.0 %),

1
2
3 1-bromobutane (Sigma Aldrich, 99.0 %), 1-bromooctane (Sigma Aldrich, 99 %), 1-
4 bromododecane (Sigma Aldrich, 97 %), 1-iodobutane (Sigma Aldrich, 99 %), ethyl glycolate
5
6 (Sigma Aldrich, 98 %), hexanoyl chloride (Sigma Aldrich, 97 %), *N,N,N',N'*-tetramethylguanidine
7
8 (Sigma Aldrich, 99 %), D-gluconic acid aqueous solution (Sigma Aldrich, 49-53 wt. % in H₂O)
9
10 and lithium bis{(trifluoromethyl)sulfonyl}imide salt (TCI, > 98.0 %) were purchased and used
11
12 without further purification. Dichloromethane (Merck, > 99.8 %), 1-butanol (Merck, > 99.8 %),
13
14 methanol (Merck, HPLC grade), ethanol absolute (Merck, for analysis EMPARTA® ACS),
15
16 acetonitrile (Merck, HPLC grade) and ethyl acetate (Merck, 98.8 %) were purchased and used
17
18 without further purification.
19
20
21
22
23

24
25 **Cell cultures.** HeLa (human cervix adenocarcinoma), HCT-116 (human Colorectal
26
27 Adenocarcinoma) and MCF-7 (human breast adenocarcinoma) cell lines were cultured in
28
29 Dulbecco's modified Eagle's medium (DMEM) supplemented with 10 % fetal bovine serum,
30
31 100 U mL⁻¹ penicillin and 100 mg mL⁻¹ streptomycin. Cells were maintained at 37 °C in a
32
33 humidified atmosphere of 5 % CO₂, as previously described.⁵⁰⁻⁵² Cells having a narrow range of
34
35 passage number were used for all experiments.
36
37
38
39

40 **Cell viability assay.** To test the cytotoxic effects of the ILs, the cells were plated at 5000 cells
41
42 per well in 96-well plates and incubated for 24 h at 37 °C in a CO₂ incubator, as previously
43
44 described.^{52,53} ILs, diluted to the desiderated concentrations in the culture medium, starting from
45
46 10⁻² M solutions in water, were added to the wells for 24 h with respective vehicle control (H₂O).
47
48 At the end of treatment, 20 μL of the Cell titer 96®AQueous reagent was added to each well and
49
50 incubated for 1–4 h at 37 °C in a CO₂ incubator. The absorbance was recorded at 490 nm using a
51
52 96-well plate reader (Sparks 20 M, Tecan Trading AG, Switzerland). The CellTiter 96® AQueous
53
54 One Solution Reagent contains a tetrazolium compound [3-(4,5-dimethylthiazol-2-yl)-5-(3-
55
56
57
58
59
60

carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium, inner salt; MTS] and an electron coupling reagent (phenazine ethosulfate; PES). PES has enhanced chemical stability, which allows it to be combined with MTS to form a stable solution. The quantity of formazan product as measured by the amount of 490 nm absorbance is directly proportional to the number of living cells in culture. The percentage of cell viability was calculated with respect to the untreated control cells for each compound concentration after the subtraction of the blank. The concentration necessary for 50 % of growth inhibition (IC_{50}) for each IL was calculated using a dose–response model, obtained from the sigmoidal fitting of response curves of percent inhibition versus logarithmic concentration of ILs using Graph Pad Prism software. Each result was the mean value obtained from experiments performed in triplicate.

Hemolytic activity assay. Hemolytic activity was evaluated with a blood agar medium supplemented with 5 % sheep blood. The blood agar was poured into 100 mm plastic plates. A tip was used to dispense 10 μ L of each IL (10^{-2} M concentrated). Plates were incubated for seven days at 37 °C. The test results were monitored and the presence of a distinctive translucent halo around the inoculum site was measured macroscopically.

Animals. Wild-type AB Zebrafish were obtained by European Zebrafish Research Center (Germany) and housed in ZebTEC semi-closed recirculation housing systems (Techniplast) at 28 °C, pH = 7.5 and conductivity 500 μ S on a 14/10 light/dark cycle. Fish were fed three times a day alternating dry food and brine shrimps. For the experiments, Zebrafish fertilized eggs were collected in Petri dishes in fish water (1.2 mM $NaHCO_3$, 0.1 g/L instant ocean, 1.4 mM $CaSO_4$, methylene blue 0.00002 % w/v) at 28 °C and soon after transferred to a 96 well plate, one embryo per well in 0.2 mL of embryo water. All the experiments were performed in agreement with EU

Directive 2010/63/EU. The experimental protocol was approved by the Italian Ministry of Health (Approval Animal Protocol No.1191/2016-PR).

Fish embryo toxicity test. Fertilized Zebrafish eggs were exposed to different concentrations of test chemicals compound for 96 h. Every 24 h, four observations were noted as indicators of lethality: the coagulation of fertilized eggs, the lack of somite formation, the lack of detachment of the tail-bud from the yolk sac, and the lack of heartbeat. At the end of the exposure period, toxicity was determined based on a positive outcome in any of the four apical observations recorded. Each experiment was done in triplicate using 20 embryos for a group. Data are normalized to control lethality and expressed as mean \pm standard error mean (SEM).

Exposure experiments. The chemical compound stocks were dissolved in a physiological solution ($pH = 7.2$) and kept at 4 °C. A working solution of 2 μ M and 0.2 mM was prepared fresh daily in the embryo water. The water for the Embryos, without (control) or with the chemicals, was changed daily.

Freshly fertilized zebrafish eggs were added to 0.2 mL of exposure medium per egg and incubated at 28 ± 1 °C for 24 h, 48 h, 72 and 96 h. For control, freshly fertilized zebrafish eggs were added to 0.2 mL of exposure medium without toxic compounds and incubated at 28 ± 1 °C for 24 h, 48 h, 72 h and 96 h. The exposure experiments were initiated as soon as possible after the fertilization of the eggs. Daily observation was performed under a stereomicroscope (Leica M165 FC).

NMR measurements. ^1H NMR and ^{13}C NMR spectra were recorded using Bruker AV-300, Bruker ultrashield 400 plus, Bruker-spectroscopin 400 ultrashield and Ascend™ 600 Bruker nuclear magnetic resonance spectrometers. Chemical shifts were reported relative to SiMe_4 .

Mass spectrometric measurements. ESI-MS mass spectrometric measurements were carried out on a Waters ICI Premier instrument with an Adylon Triversa NanoMate injection system (cone voltage 50 V, source 120 °C). Both positive and negative ions were detected. For each sample, a methanol solution was prepared.

Thermogravimetric analysis (TGA). Thermogravimetric analysis was performed using a TGA/DSC thermogravimetric analyser from Mettler-Toledo, Inc. The samples were measured in alumina crucibles, at a heating rate of 5 K min⁻¹, under a dinitrogen atmosphere. The onset of the weight loss in each thermogram was used as a measure of the decomposition temperature (the point at 5 % wt. loss of the sample).

Differential Scanning Calorimetry (DSC). DSC measurements were carried out using TA Instruments Modulated DSC Q 2000 V24.4 Build 116 with a refrigerated cooling system RCS 90, capable of controlling the temperature down to 220 K. Samples were weighed and hermetically sealed in aluminium pans. Heating and cooling rates were 10 °C min⁻¹ for the ionic liquid sample. The maximum heating temperature was chosen depending on the thermal stability of the sample. The minimum cooling temperature was set at -50 °C under dinitrogen atmosphere. Two heating-cooling cycles were performed.

Viscosity measurements. Viscosity measurements were carried out using Marveln instrument Bohlin Gemini HR nano rheometer. Each sample was analysed in the range of temperature from 290 up to 375 K, with a rate of 2 K min⁻¹. The viscosity trends were fitted using the VFT model.


Conductivity measurements. The electrical conductivity σ (S/m) of the liquid salts was measured with a Keithley 2000 Multimeter in a two-points configuration on pure salts, at T = 297.15 K. Due to the small amount of sample to be analysed, a home-made cell with two

microelectrodes was used; the cell constant was calculated using a commercial IL, $[\text{C}_8\text{C}_1\text{Im}][\text{BF}_4]$, whose conductivity is already reported in literature.⁵⁴ The conductivity values are the average of three measurements.

RESULTS AND DISCUSSION

Synthesis. As previously stated, two classes of “sweet” ILs were synthesized with the sugar moiety either present in the cation or the anion.


In order to introduce the sugar moiety to the cation, the 1-(3-aminopropyl)imidazole or aliphatic diamine was heated to reflux with glucono- δ -lactone, in a MeOH solution for 24 h. In the second step, the amide intermediate was quaternised using different alkyl halides (Scheme 2).



Scheme 2. Synthetic pathways for glucono-based ILs.


All the bis{(trifluoromethyl)sulfonyl}amide salts were prepared by the metathesis reaction with LiNTf₂, using 1-butanol as a solvent. The obtained ILs differ in terms of the length of the spacer between the amide and ammonium group, the alkyl chain length or branching and also the anion nature (Scheme 1).

As far as glycolic and hexanoyl derivatives ([N₁₁₂Gly₈]**Br** and [N₁₁₂HexA₈]**Br**) are concerned, the *N,N*-dimethylethylamine was reacted with ethylglycolate or hexanoyl chloride to obtain amide intermediates. These were successively quaternised by reaction with the 2-ethylhexyl bromide to obtain the corresponding ammonium salts (Scheme 3).



Scheme 3. Synthetic pathways for the glycolic (a) and hexanoyl-based ILs (b).

The synthesis of ionic liquids with the gluconate anion (Scheme 4) was described in our previous publication.⁴⁴ The ILs that we prepared differ in terms of the core of the cation nature (*i.e.* ammonium, phosphonium or imidazolium) and, in some cases, the length of the alkyl chains (see Scheme 1).



Scheme 4. Synthetic pathways for the gluconate-based ILs.

It is worth noting the fact that synthetic protocols used to obtain “sweet” ILs adhere to the Principles of Green Chemistry to a good extent, according to the metric approach proposed by Clark *et al.* (Table S1).⁵⁵ Indeed, with only two exceptions related to the synthesis of [N_{112HexA8}]**Br** and [N_{113GA8}][NTf₂], for which yields were lower than 70%, in all other cases they were higher than 89% (15 cases out of 22) or included in the range 70-89% (5 cases out of 22).

Atom economy (AE), with the only exception of the production of *N*-2-(dimethylamino)ethyl glucosamide and *N*-2-(dimethylamino)ethyl hexylamide, is always equal to 100% (Table S1). On the other hand, in most cases, reaction mass efficiency (RME) and optimum efficiency (OE) reach good levels as they change in the range 69.0-95.9%. Values falling in the above range are the ones corresponding to the synthesis of [NTf₂]-ILs, [N_{112HexA8}]**Br** and for the synthesis of the above intermediates.

Mass Intensity (MI) values, as referred to single step processes, fall in the range 1.19 - 49.85. The only unfavourable case is represented by the *N*-2-(dimethylamino)ethyl hexylamide. The lowest MI values were calculated for the synthesis of gluconate ILs.

The analysis of MIs calculated for gluconamide ILs shows that the workup of the product, consisting of washing with ethyl acetate (30 mL), contributes to a large extent in the determining of the value (crf columns 4 and 5 of Table S1). However, this contribution becomes less significant taking into consideration that a recommended and recyclable solvent was used.

As for the solvent nature, with only the exception of *N*-2-(dimethylamino)ethyl hexylamide, most of the reactions were performed in MeOH. *n*-BuOH was used for the anion exchange to obtain [NTf₂]⁻ ILs and in the case of gluconate-based ILs, water was used to carry out the anion exchange. Furthermore, if necessary, products were washed using benign ethyl acetate.

The energy considerations demonstrate how the reactions were performed under relatively mild conditions, as the highest temperature used was 65 °C (boiling point of MeOH).

In Vitro Cytotoxicity. All ILs were tested for their cytotoxicity. As above stated, three different human cancer cell lines were chosen as the prototype of human cells, namely HeLa, HCT-116 and MCF-7, derived from cervical, colon and breast cancer lines respectively and previously used to test ILs cytotoxicity.⁵⁶⁻⁵⁷⁻⁵⁸ Cancer cells are generally more susceptible to toxicity than normal cells since they have a highly unstable genetic load and undergo rapid cell divisions.⁵⁹

The cytotoxic effect of ILs was investigated using the MTT assay.⁶⁰ The results, expressed as IC₅₀ values, were calculated from the dose-response curves.




Figure 1. IC₅₀ values as a function of the nature of ILs.

In the first set of experiments carried out using stock solutions in water at 10^{-6} M, the toxicity effect was recorded only for **[P₆₆₆₁₄][Glu]**, which proved to be the most toxic.

For this reason, the ILs toxicity was tested again by using more concentrated solutions (10^{-3} M) (Figure 1 and Table S2). Interestingly, despite the use of highly concentrated solutions, some ILs did not affect cell viability at all (see later).

In other cases, IC₅₀ values range from 0.06 mM up to 12.6 mM. They proved to be quite high, with respect to the values generally reported in literature.^{57,61}

1
2
3 In general, taking into consideration salts bearing the gluconate anion, IC_{50} values were
4 significantly higher in the presence of ammonium or guanidinium cations than the phosphonium
5 one (Figure 1).
6
7
8
9

10
11 It is worth noting the increase in IC_{50} values is due to the decrease of the alkyl chain length on
12 the phosphonium cation, going from $[P_{66614}][Glu]$ to $[P_{4444}][Glu]$. Indeed, the later exhibited a
13 toxicity effect only when a highly concentrated solution (10^{-3} M) was used. The above trend is in
14 accordance with other findings in literature.^{62,63}
15
16
17
18
19

20
21 Conversely, all iodide-based ILs bearing a butyl chain and the gluconamide residue on the cation
22 were found to be safe, regardless of the spacer length ($[N_{112GA4}][I]$ and $[N_{113GA4}][I]$) and cation nature
23 ($[Im_{3GA4}][I]$ and $[N_{113GA4}][I]$). The most surprising observation was the ultra-low toxicity of the
24 imidazolium derivative, for which the presence of gluconamide moiety on the side chain appears
25 to have removed the toxicity effects completely. Indeed, toxicity effects were not detected even
26 when cells were in contact with a 10^{-2} M solution of the salt. This result represents a significant
27 improvement with respect to the corresponding 1-hexyl-3-methylimidazolium halides.⁶⁴
28
29
30
31
32
33
34
35
36
37

38
39 The results obtained by the biological investigation on this component of the new imidazolium
40 class of compounds are quite interesting. The low toxicity, due to the insertion of the gluconic
41 moiety on the cation, is in accordance with other findings in literature, demonstrating the relevant
42 role of the hydroxylated chain in decreasing IL toxicity.¹⁶
43
44
45
46
47
48

49 According to literature,¹⁸ ammonium salts bearing hydroxyl groups on the side chain proved less
50 toxic than the corresponding alkyl derivatives, as accounted for by the comparison between
51 $[N_{112Gly8}][Br]$ and $[N_{112Hex8}][Br]$. However, the IC_{50} value does not significantly change for glycolate
52
53
54
55
56
57
58
59
60

and gluconamide salts, evidencing that the obtained result is not affected by the number of hydroxyl groups on the side chain.

Cytotoxicity also depends on the nature of the anion. Indeed, dissimilar to $[\text{N}_{112\text{GA}4}]\text{I}$, the toxicity effects were detected for $[\text{N}_{112\text{GA}4}]\text{Br}$ (Figure 1). Furthermore, the analysis of the data collected for 2-ethylhexyl derivatives sheds light on the role played by the length of the spacer. Indeed, both Br^- and $[\text{NTf}_2]^-$ ILs show a decrease in cytotoxicity from $[\text{N}_{112\text{GA}8}]^+$ to $[\text{N}_{113\text{GA}8}]^+$. Probably, the longer the spacer, the higher is the conformational freedom of the cation that hampers the interaction with the cell membrane. This hypothesis is well supported by the data previously reported in literature about the increased biological activity of some bicyclic nucleosides with reduced conformational freedom in side chain.^{65,66}

On the other hand, with the spacer being the same $[\text{N}_{112\text{GA}n}]^+$, the cytotoxicity is also affected by the nature of the alkyl chain, linear or branched. In particular, linear alkyl chain results in higher toxicity, as accounted for by the decrease in the IC_{50} value from $[\text{N}_{112\text{GA}8}]\text{Br}$ to $[\text{N}_{112\text{GA}8\text{L}}]\text{Br}$ (Figure 1). However, the above effect depends on the spacer length being less significant in the case of the propyl spacer. Moreover, the comparison between the IC_{50} values collected for $[\text{N}_{112\text{GA}8\text{L}}]\text{Br}$ and $[\text{N}_{112\text{GA}12\text{L}}]\text{Br}$, evidences that the further lengthening of the alkyl chain, from octyl to dodecyl derivative, differently affects cytotoxicity, exhibiting a dependence on the nature of the cell line (Figure 1; Table S2).

Hemolytic Activity. All ILs were also tested for their hemolytic activity. In this respect, it is well known that erythrocytes represent good systems to assess the cytotoxicity of different molecules.⁶⁷ We used a blood agar test, in which 5% of sheep's blood was included in an agar

medium. ILs were tested at 10^{-2} M, and in the majority of cases they did not show significant hemolytic activity (Figure 2).




Figure 2. Blood agar hemolysis of ILs at 10^{-2} M, after 24h of incubation (1: $[N_{112GA8}]Br$; 2: $[N_{113GA8}]Br$; 3: $[N_{113GA4}]I$; 4: $[Im_{3GA4}]I$; 5: $[N_{112GA4}]I$; 6: $[N_{112GA8}][NTf_2]$; 7: $[N_{113GA8}][NTf_2]$; 8: $[N_{112GA8L}]Br$; 9: $[N_{113GA8}]Br$; 10: $[N_{112GA12}]Br$; 11: $[N_{112Gly8}]Br$; 12: $[N_{112GA4}]Br$; 13: $[N_{112Hex8}]Br$; 14: $[P_{66614}][Glu]$; 15: $[P_{4444}][Glu]$; 16: $[N_{4444}][Glu]$; 17: $[tmgH][Glu]$). + and – represent positive and negative control, respectively.

A slight activity was detected in the case of $[P_{66614}]Glu$, according to the detected cytotoxicity and indicating that the antiproliferative effects are related to unspecific membrane damage. Less significant toxicity effects were detected for $[N_{112GA12}]Br$ and $[N_{113GA8L}]Br$.

Fish embryo toxicity test. The toxicity of ILs was evaluated in Zebrafish from 24 to 96 hours post fertilization (hfp).⁶⁸ The coagulation of fertilized eggs, the lack of heartbeat, the lack of detachment of the tail-bud from the yolk sac and the lack of somite formation were checked daily. In this case, based on the results collected through cytotoxicity tests, we selected only some gluconamide-based ILs to assess the effect of the spacer, anion and alkyl chain length.

Furthermore, we tested the ILs bearing the gluconate anion and the hexylamide derivative, $[N_{112Hex8}]Br$, to evaluate the relevance of the presence of hydroxyl groups on the side chain. All the ILs were tested at $2\ \mu M$ and $0.2\ mM$. The above concentrations were chosen on the basis of previous reports in the literature.^{63,69} No significant toxicity was detected at both concentrations, as we did not observe significant differences in the treated and untreated fish (Figure 3 and Table S3).




Figure 3. Results of toxicity test performed on a Fish embryo (lethality at 24/48 h). For each ionic liquid, the survival of the treated fish was calculated by normalizing the value to the untreated value that was set equal to one.

We did not detect a toxicity effect also for the imidazolium derivative, indicating that the presence of the oligo-hydroxyl motif is able to compensate for the negative effect of the imidazolium cation.

Physical properties of ‘sweet appended’ ionic liquids

TGA investigation. The thermal stability of ILs was determined using TGA measurements (Figure S1 and Table S4 of SI). In all cases, the temperature corresponding to a 5% weight loss was considered (T_d).

In the case of gluconoamide-based ILs, T_d ranges from 127.8 up to 210.3 °C. For the sake of clarity, the data collected will be discussed as a function of the changing structural factors. (Figure 4).




Figure 4. Trend of T_d (°C) values as a function of IL nature.

As far as the length of the alkyl spacer is considered, the trend of T_d values depends on the anion nature and the length of the alkyl chain on the ammonium head. Indeed, for the 2-ammonium derivatives ([N112GA8]Br, [N113GA8]Br, [N112GA4]I, [N113GA4]I, [N112GA8][NTf2] and [N113GA8][NTf2]), the lengthening of the alkyl spacer induces an increase in T_d values for Br[−] – ILs and a decrease in

the case of I^- ILs and $[\text{NTf}_2]^-$ ILs. More significant changes in ΔT were observed for halide anions compared to the $[\text{NTf}_2]^-$ anion ($\Delta T = 26.7, 18.6$ and 2.6 °C for Br^- , I^- and $[\text{NTf}_2]^-$ ILs, respectively). The effect of the anion nature on the thermal stability of ILs is well documented in literature.⁷⁰⁻⁷¹ Furthermore, in our case, the comparison between T_d values corresponding to $[\text{N}_{112\text{GA}4}]\text{Br}$ and $[\text{N}_{112\text{GA}4}]\text{I}$ allows stating that the thermal stability increases with a decrease in the anion nucleophilicity, according to the data previously reported for imidazolium salts.⁷²

Regardless of the spacer length, the branching of the alkyl chain induces a decrease in thermal stability ($[\text{N}_{112\text{GA}8}]\text{Br}$ and $[\text{N}_{112\text{GA}8\text{L}}]\text{Br}$, $[\text{N}_{113\text{GA}8}]\text{Br}$ and $[\text{N}_{113\text{GA}8\text{L}}]\text{Br}$; $\Delta T = 32.2$ and 5.5 °C for $[\text{N}_{112\text{GA}n}]^+$ and $[\text{N}_{113\text{GA}n}]^+$ ILs, respectively).

On the other hand, as far as the effect of the alkyl chain length on the ammonium head is concerned, the data collected allow the identification of the linear octyl chain as the superior feature. Indeed, T_d values change along the following trend: $[\text{N}_{112\text{GA}4}]\text{Br} < [\text{N}_{112\text{GA}8\text{L}}]\text{Br} > [\text{N}_{112\text{GA}12}]\text{Br}$.

Finally, the comparison between T_d values collected for $[\text{Im}_{3\text{GA}4}]\text{I}$ and $[\text{N}_{113\text{GA}4}]\text{I}$ indicates a higher thermal stability of the imidazolium salt with respect to the corresponding ammonium one, according to what was previously observed by Afonso *et al.*, when they compared the thermal properties of $[\text{N}_{1888}]^+$ and $[\text{C}_8\text{C}_1\text{im}]^+$ ILs.³²

As stated above, to evaluate the effect of the presence of the oligo-hydroxylated chain, glycolic and hexyl derivatives, $[\text{N}_{112\text{GlyA}8}]\text{Br}$ and $[\text{N}_{112\text{HexA}8}]\text{Br}$, were also synthesized. In both cases, the comparison of T_d values with the one corresponding to $[\text{N}_{112\text{GA}8}]\text{Br}$, evidences how the presence of a “sweet” chain represents a warranty of higher thermal stability.

As for gluconate-based ILs, T_d ranges from 103.1 up to 172.5 °C. Among the analysed salts, [tmgH][Glu] showed the lowest thermal stability, whereas the [P₆₆₆₁₄][Glu] was the most thermally stable ionic liquid. In general, T_d values depended on the cation nature, with ammonium IL exhibiting a higher thermal stability with respect to the corresponding phosphonium IL ([N₄₄₄₄][Glu] and [P₄₄₄₄][Glu]). Finally, for phosphonium ILs, the elongation of the alkyl chain induces a corresponding increase in T_d ([P₄₄₄₄][Glu] and [P₆₆₆₁₄][Glu]), according to the previous reports.⁷⁰

DSC investigation. The solid-liquid transition phase of the new ILs was investigated by performing DSC measurements. In general, they showed glass transitions (T_g) detected in the heating cycle and displayed in Figure 5 (see also Table S4 and Figure S2).

According to the previous reports, this behavior indicates that they have a weak tendency to crystallize.⁷³⁻⁷⁴




Figure 5. (a) Glass transition temperatures (T_g) for amide-based ILs; (b) Glass transition temperatures (T_g) for gluconate-based ILs.

With the only exception of $[\text{N}_{444}][\text{Glu}]$, T_g values lower than 100 °C were detected, allowing us to classify almost all salts as ILs. In the case of gluconamide-based ILs, they range from -55.5 ([$\text{N}_{112\text{GA}8\text{L}}\text{Br}$]) up to -0.1 °C ([$\text{N}_{113\text{GA}8}\text{Br}$]).

For the above ILs, T_g values are affected by both the anion nature and alkyl chain length. In the case of butyl derivatives, they increase from [$\text{N}_{112\text{GA}4}\text{Br}$] to [$\text{N}_{112\text{GA}4}\text{I}$] and, taking into consideration the iodide salts, they also decrease by lengthening the alkyl spacer between the amide and ammonium group ([$\text{N}_{112\text{GA}4}\text{I}$] and [$\text{N}_{113\text{GA}4}\text{I}$]).

Spacer being the same, the lengthening of the alkyl chain on the ammonium head induces firstly a decrease and then an increase in T_g values, as accounted for by comparing the values collected for [$\text{N}_{112\text{GA}4}\text{Br}$], [$\text{N}_{112\text{GA}8\text{L}}\text{Br}$] and [$\text{N}_{112\text{GA}12}\text{Br}$]. This trend perfectly agrees with the one generally detected for alkylimidazolium salts, that shows a decrease in melting temperature until conformational freedom of the alkyl chain prevails over van der Waals interactions.⁷⁵

Among the structural factors, the branching of the alkyl chain also plays a role. Indeed, irrespective of the length of the spacer, salts bearing the linear alkyl chain exhibit T_g values lower than the one bearing the 2-ethylhexyl chain.

A more complex trend was detected as far as the effect of the length of the spacer is concerned. Indeed, in the case of bromide derivatives, T_g increases from [$\text{N}_{112\text{GA}n}\text{Br}$] to [$\text{N}_{113\text{GA}n}\text{Br}$] ILs. No significant changes were detected for [NTf_2] ILs, while a decrease in T_g values was collected from [$\text{N}_{112\text{GA}4}\text{I}$] to [$\text{N}_{113\text{GA}4}\text{I}$].

Comparing the values collected for [$\text{N}_{112\text{GA}8}\text{Br}$], [$\text{N}_{112\text{GlyA}8}\text{Br}$] and [$\text{N}_{112\text{HexA}8}\text{Br}$] accounts for a decrease in T_g in parallel with the decrease in the number of hydroxyl groups present in the amide

chain. Finally, T_g values also decrease moving from imidazolium to ammonium salt (**[Im₃GA₄]**I**** and **[N₁₁GA₄]**I****).

In general, gluconate-based salts show higher T_g values with respect to gluconamide-based salts. Indeed, in this case, they range from -24.1 (for **[tmgH][Glu]**) to 120 °C (for **[N₄₄₄₄][Glu]**).

[P₄₄₄₄][Glu] shows two glass transitions, the first one occurring at -21.8 °C and the second one at 80.2 °C. For these salts, changes in the cation nature significantly affect T_g values, as accounted for by the comparison between **[N₄₄₄₄][Glu]** and **[P₄₄₄₄][Glu]**. On the other hand, for phosphonium salts, a significant increase in T_g was also detected as a consequence of the lengthening of alkyl chains in (**[P₄₄₄₄][Glu]** and **[P₆₆₆₁₄][Glu]**).

Viscosity measurements. Some gluconamide ILs were sufficiently fluid at room temperature to allow their viscosity to be determined using a rheometer equipped with a thermostat. Viscosity values as a function of the IL nature are displayed in Figure 6 (Table S5).




Figure 6. Viscosity (η / mPa s) values for gluconamide-based ILs.

η values range from 106.3 mPa s up to 41700 mPa s, indicating $[\mathbf{N}_{112\text{GA}4}]\mathbf{Br}$ and $[\mathbf{N}_{112\text{GA}4}]\mathbf{I}$ as the ILs showing the lowest and the highest viscosities, respectively. The above observation shows that, in the case of halide based ILs, η depends on the ion size and polarizability, according to previous reports in literature.⁷⁶

As far as the length of the alkyl spacer is concerned, for halide-based ILs ($[\mathbf{N}_{112\text{GA}4}]\mathbf{I}$, $[\mathbf{N}_{113\text{GA}4}]\mathbf{I}$, $[\mathbf{N}_{112\text{GA}8}]\mathbf{Br}$ and $[\mathbf{N}_{113\text{GA}8}]\mathbf{Br}$), η values decrease with the increase in the length of the alkyl spacer. Surprisingly, in the case of $[\text{NTf}_2]^-$ ILs ($[\mathbf{N}_{112\text{GA}8}][\text{NTf}_2]$ and $[\mathbf{N}_{113\text{GA}8}][\text{NTf}_2]$), an opposite trend was observed.

The length of the alkyl chains on the ammonium head also affects viscosity, as accounted for by the decrease in η values on going from $[\mathbf{N}_{113\text{GA}8}]\mathbf{Br}$ to $[\mathbf{N}_{113\text{GA}8\text{L}}]\mathbf{Br}$, indicating that the branching of the alkyl chain gives rise to less viscous solvents.⁷⁷

Finally, alkyl chain length and anion being the same, η values proved lower for imidazolium than for ammonium-based ILs, as accounted for by the data collected for $[\mathbf{N}_{113\text{GA}4}]\mathbf{I}$ and $[\mathbf{Im}_{3\text{GA}4}]\mathbf{I}$, in perfect agreement with the data previously reported in literature.⁷⁸

Viscosity values were also measured as a function of the temperature (from 300 up to 370 K) and the trends observed were analysed with the VFT (Vogel-Fulcher-Tamman) model (Figure S3). The model is represented by the eq. (1)⁷⁹

$$\log \eta = A + \frac{D \cdot T_0}{(T - T_0)} \quad (1)$$

where A stands for the viscosity at the infinite temperature limit, T_0 is the temperature at the infinite viscosity and D represents the fragility that measures the strength for glass forming materials.⁸⁰

In general, in the case of fragile glass materials featured by the occurrence of feeble interactions and hydrogen bonds, D values are lower than 10.

Interestingly, D values are related to the fragility index, m , by the eq. (2):⁸¹

$$m = 16 + \frac{590}{D} \quad (2)$$

This parameter is related to the structural stability of the materials to thermal degradation. For glass-forming materials, m values are higher than 100.




Figure 7. Fragility index (m) for gluconamide-based ILs.

m values obtained for gluconamide-based ILs are displayed in Figure 7 (see also Table S5 and Figure S3). Analysis of both D and m values account for the glass material nature of our ILs.

Indeed, D range from 1.9 ($[\text{Im}_{3\text{GA}4}]\text{I}$) up to 6.8 ($[\text{N}_{112\text{GA}8}]\text{Br}$). Conversely, m values range from 102.3 (for $[\text{N}_{112\text{GA}8}]\text{Br}$) and 319.7 (for $[\text{Im}_{3\text{GA}4}]\text{I}$).

Both parameters significantly change as a function of all the structural features considered so far. In particular, for butyl derivatives, m values decrease as a function of the anion nature; *i.e.* according to the increase in size and polarizability of the anion ($[\text{N}_{112\text{GA}4}]\text{Br} > [\text{N}_{112\text{GA}4}]\text{I}$).

It is worth noting the fact that, as far as the alkyl spacer is concerned, the trend observed is also significantly affected by the length of the alkyl chain on the ammonium head and the anion nature. Indeed, in the case of butyl derivatives ($[\text{N}_{112\text{GA}4}]\text{I}$ and $[\text{N}_{113\text{GA}4}]\text{I}$) and for 2-ethyl-hexyl derivatives associating with the $[\text{NTf}_2]^-$ anion ($[\text{N}_{112\text{GA}8}][\text{NTf}_2]$ and $[\text{N}_{113\text{GA}8}][\text{NTf}_2]$), m values increase on going from the ethylene to the propylene spacer. Conversely, for $[\text{N}_{112\text{GA}8}]\text{Br}$ and $[\text{N}_{113\text{GA}8}]\text{Br}$, an opposite trend was detected.

The nature of the alkyl chain, linear or branched, affects the m values to a lower extent as accounted for by the comparison between $[\text{N}_{113\text{GA}8}]\text{Br}$ and $[\text{N}_{113\text{GA}8\text{L}}]\text{Br}$. Finally, the nature of the cation affects fragility index values as testified by the significant decrease in m values, going from $[\text{N}_{113\text{GA}4}]\text{I}$ to $[\text{Im}_{3\text{GA}4}]\text{I}$.

Conductivity measurements. Conductivity measurements were carried out for most gluconamide ILs, to gain insights into the effects that structural changes may have on their ability to act as charge carriers. σ (S/m) values, measured using $[\text{C}_1\text{C}_8\text{Im}][\text{BF}_4]$ as standard, are displayed in Figure 8 (see also Table S6 of SI).




Figure 8. Conductivity (σ) values for some of ILs obtained.

The collected σ values range from 0.002 up to 0.162 S/m. Analysis of the above values shows that one of the main factors determining the conductivity of our ILs is represented by the length of the alkyl linker, as accounted for by its decrease going from an ethyl to a propyl spacer. Interestingly, the degree of decrease depends on the anion nature, more significantly with the increase in the anion coordination ability ($\text{Br}^- > \text{I}^- > [\text{NTf}_2]^-$).

With anion being the same, conductivity significantly increases from $[\text{N}_{11\text{InGA}8}]^+$ to $[\text{N}_{11\text{InGA}8\text{L}}]^+$ ILs, according to the decrease in the branching of the alkyl chain. On the contrary, the ability of charge transport increases from $[\text{N}_{112\text{GA}8}]\text{Br}$ to $[\text{N}_{112\text{HexA}8}]\text{Br}$, where OH groups on the side chain range from 5 to none. The former, favoring the formation of hydrogen bonds, induces a parallel increase of the viscosity of the ILs together with decrease in the corresponding σ values.

CONCLUSIONS

Taking into consideration the sustainability endorsement and the incessant request for the identification of new solvents potentially applicable in the industry with low environmental impact, we have presented here, a new class of organic salts bearing in the structure a residue derived from gluconic acid. Since gluconic acid can be harnessed from food waste, its utilization to obtain non-toxic solvents from renewable sources is highly desirable from a ‘green chemistry’ point of view.

In particular, we prepared aliphatic organic salts (with ammonium and phosphonium core structure), bearing the “sweet” residue on the cation or anion. The synthetic procedures used comply, to a good extent, with the Green Chemistry metrics approach and principles.

The analysis of solid-liquid transition and thermal stability allowed stating that the majority of these salts fall in the class of ILs that are suitable solvents for applications at high temperatures.

As expected, their physico-chemical properties are significantly affected by changes in the structure. The toxicity of ILs obtained was assessed using three different cancer cell lines and fish embryos. To our surprise, we demonstrated that they are able to exert moderate or low anti-proliferative effects and prove safe towards human erythrocytes. Toxicity effects were detected only when using unusually highly concentrated solutions ($IC_{50} \sim 10^{-3} \text{ M}$), and their change depends upon the presence of the gluconic moiety on the cation or on the anion. In general, lower IC_{50} values were detected for amide based ILs. Furthermore, as far as ecotoxicity studies are concerned, “sweet” ILs proved completely safe towards to Zebrafish embryos, as no lethality events were observed over 24 or 48 hours.

It is noteworthy, that the above considerations can be also drawn for the corresponding imidazolium ILs. The above results testify that the presence of a “sweet” residue, such as the one derived from gluconic acid, could pave the way for the production of safer aromatic ILs, with favorable consequences from an industrial point of view.

ASSOCIATED CONTENT

Supporting Information. Synthetic details; DSC and TGA traces, VFT traces, Table of IC₅₀ values, Table of toxicity test performed in the presence of zebrafish embryos, Table of DSC and TGA data, Table of viscosity data, Table of conductivity data, ¹H and ¹³C NMR spectra.

The following files are available free of charge.

AUTHOR INFORMATION

Corresponding Authors

*Prof. Francesca D’Anna, Università degli Studi di Palermo, Dipartimento STEBICEF, Viale delle Scienze, Ed. 17 90128 Palermo (Italy); email: francesca.danna@unipa.it

*Dr. H. Q. Nimal Gunaratne, The QUILL Research Centre, School of Chemistry and Chemical Engineering Queen's University of Belfast, Stranmillis Road, Belfast, Northern Ireland BT9 5AG, UK; email: n.gunaratne@qub.ac.uk

Author Contributions

The manuscript was written through the contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

University of Palermo FFR 2017

Declaration of interest

The authors declare no competing financial interest

ACKNOWLEDGMENT

We thank the University of Palermo for financial support. FB gratefully acknowledges the support of QUILL. This article is dedicated to the memory of our co-author, Prof. Kenneth R. Seddon, who with his enthusiasm and unwavering support moved the whole project forward.

REFERENCES

- (1) Rogers, R.; Seddon, K.: *In Green Industrial Applications of Ionic Liquids*, 2002; Vol. 818.
- (2) Meindersma, G. W.; Podt, A.; Meseguer, M. G.; de Haan, A. B.: Ionic Liquids as Alternatives to Organic Solvents in Liquid-Liquid Extraction of Aromatics. In *Ionic Liquids IIIB: Fundamentals, Progress, Challenges, and Opportunities*; ACS Symposium Series 902; American Chemical Society, 2005; Vol. 902; pp 57-71.
- (3) Marullo, S.; Rizzo, C.; D'Anna, F. Task-Specific Organic Salts and Ionic Liquids Binary Mixtures: A Combination to Obtain 5-Hydroxymethylfurfural From Carbohydrates. *Frontiers in Chemistry* **2019**, 7, DOI 10.3389/fchem.2019.00134.
- (4) Mohanazadeh, F.; Aghvami, M. Ionic Liquids as Reagent and Reaction Medium: Preparation of Alkyl Aryl Ethers. *Monatshefte für Chemie - Chemical Monthly* **2007**, 138, 47-49, DOI 10.1007/s00706-006-0562-5.
- (5) Gathergood, N.; Garcia, M. T.; Scammells, P. J. Biodegradable ionic liquids: Part I. Concept, preliminary targets and evaluation. *Green Chem.* **2004**, 6, 166-175, DOI 10.1039/B315270G.
- (6) Jordan, A.; Gathergood, N. Biodegradation of ionic liquids – a critical review. *Chem. Soc. Rev.* **2015**, 44, 8200-8237, DOI 10.1039/C5CS00444F.
- (7) Krishnan, S.; Quraishi, K. S.; Aminuddin, N. F.; Mazlan, F. A.; Leveque, J.-M. Biodegradability of imidazolium, pyridinium, piperidinium and pyrrolidinium based ionic liquid in different water source. *AIP Conf. Proc.* **2016**, 1787, 040017, DOI 10.1063/1.4968096.
- (8) Wu, S.; Zeng, L.; Wang, C.; Yang, Y.; Zhou, W.; Li, F.; Tan, Z. Assessment of the cytotoxicity of ionic liquids on *Spodoptera frugiperda* 9 (Sf-9) cell lines via in vitro assays. *J. Hazard. Mater.* **2018**, 348, 1-9, DOI 10.1016/j.jhazmat.2018.01.028.

- (9) Sydow, M.; Owsianiak, M.; Framski, G.; Woźniak-Karczewska, M.; Piotrowska-Cyplik, A.; Ławniczak, Ł.; Szulc, A.; Zgoła-Grześkowiak, A.; Heipieper, H. J.; Chrzanowski, Ł. Biodiversity of soil bacteria exposed to sub-lethal concentrations of phosphonium-based ionic liquids: Effects of toxicity and biodegradation. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 157-164, DOI 10.1016/j.ecoenv.2017.08.026.
- (10) Thuy Pham, T. P.; Cho, C.-W.; Yun, Y.-S. Environmental fate and toxicity of ionic liquids: A review. *Water Res.* **2010**, *44*, 352-372, <https://doi.org/10.1016/j.watres.2009.09.030>.
- (11) Losada-Pérez, P.; Khorshid, M.; Renner, F. U. Interactions of Aqueous Imidazolium-Based Ionic Liquid Mixtures with Solid-Supported Phospholipid Vesicles. *PLoS One* **2016**, *11*, e0163518-e0163518, DOI 10.1371/journal.pone.0163518.
- (12) Coleman, D.; Gathergood, N. Biodegradation studies of ionic liquids. *Chem. Soc. Rev.* **2010**, *39*, 600-637, DOI 10.1039/B817717C.
- (13) Agatemor, C.; Ibsen, K. N.; Tanner, E. E. L.; Mitragotri, S. Ionic liquids for addressing unmet needs in healthcare. *Bioengineering & Translational Medicine* **2018**, *3*, 7-25, DOI 10.1002/btm2.10083.
- (14) Reid, J. E. S. J.; Prydderch, H.; Spulak, M.; Shimizu, S.; Walker, A. J.; Gathergood, N. Green profiling of aprotic versus protic ionic liquids: Synthesis and microbial toxicity of analogous structures. *Sustainable Chemistry and Pharmacy* **2018**, *7*, 17-26, DOI 10.1016/j.scp.2017.11.001.
- (15) Montalbán, M. G.; Villora, G.; Licence, P. Ecotoxicity assessment of dicationic versus monocationic ionic liquids as a more environmentally friendly alternative. *Ecotoxicol. Environ. Saf.* **2018**, *150*, 129-135, DOI 10.1016/j.ecoenv.2017.11.073.
- (16) Vraneš, M.; Tot, A.; Jovanović-Šanta, S.; Karaman, M.; Dožić, S.; Tešanović, K.; Kojić, V.; Gadžurić, S. Toxicity reduction of imidazolium-based ionic liquids by the oxygenation of the alkyl substituent. *RSC Adv.* **2016**, *6*, 96289-96295, DOI 10.1039/C6RA16182K.
- (17) Zhao, D.; Liao, Y.; Zhang, Z. Toxicity of Ionic Liquids. *CLEAN – Soil, Air, Water* **2007**, *35*, 42-48, doi:10.1002/clen.200600015.
- (18) Markiewicz, M.; Maszkowska, J.; Nardello-Rataj, V.; Stolte, S. Readily biodegradable and low-toxic biocompatible ionic liquids for cellulose processing. *RSC Adv.* **2016**, *6*, 87325-87331, DOI 10.1039/C6RA14435G.
- (19) Ventura, S. P. M.; Gonçalves, A. M. M.; Sintra, T.; Pereira, J. L.; Gonçalves, F.; Coutinho, J. A. P. Designing ionic liquids: the chemical structure role in the toxicity. *Ecotoxicology* **2013**, *22*, 1-12, 10.1007/s10646-012-0997-x.
- (20) Ventura, S. P. M.; Marques, C. S.; Rosatella, A. A.; Afonso, C. A. M.; Gonçalves, F.; Coutinho, J. A. P. Toxicity assessment of various ionic liquid families towards *Vibrio fischeri* marine bacteria. *Ecotoxicol. Environ. Saf.* **2012**, *76*, 162-168, <https://doi.org/10.1016/j.ecoenv.2011.10.006>.
- (21) Plechkova, N. V.; Seddon, K. R. Applications of ionic liquids in the chemical industry. *Chem. Soc. Rev.* **2008**, *37*, 123-150, DOI 10.1039/B006677J.
- (22) Swatloski, R. P.; Spear, S. K.; Holbrey, J. D.; Rogers, R. D. Dissolution of Cellulose with Ionic Liquids. *J. Am. Chem. Soc.* **2002**, *124*, 4974-4975, 10.1021/ja025790m.
- (23) Ozokwelu, D.; Zhang, S.; Okafor, O. C.; Cheng, W.; Litombe, N.: Chapter 5 - Separation Science and Technology. In *Novel Catalytic and Separation Processes Based on Ionic Liquids*; Ozokwelu, D., Zhang, S., Okafor, O. C., Cheng, W., Litombe, N., Eds.; Elsevier: Amsterdam, 2017; pp 193-202.

- (24) Ozokwelu, D.; Zhang, S.; Okafor, O. C.; Cheng, W.; Litombe, N.: Chapter 4 - Catalytic Reaction in Ionic Liquids. In *Novel Catalytic and Separation Processes Based on Ionic Liquids*; Ozokwelu, D., Zhang, S., Okafor, O. C., Cheng, W., Litombe, N., Eds.; Elsevier: Amsterdam, 2017; pp 111-191.
- (25) Balducci, A.: Ionic Liquids in Lithium-Ion Batteries. In *Top Curr Chem (Z)* 2017; Vol. 375:20.
- (26) Liwarska-Bizukojc, E.; Maton, C.; Stevens, C. V. Biodegradation of imidazolium ionic liquids by activated sludge microorganisms. *Biodegradation* **2015**, *26*, 453-463, DOI 10.1007/s10532-015-9747-0.
- (27) Morrissey, S.; Pegot, B.; Coleman, D.; Garcia, M. T.; Ferguson, D.; Quilty, B.; Gathergood, N. Biodegradable, non-bactericidal oxygen-functionalised imidazolium esters: A step towards 'greener' ionic liquids. *Green Chem.* **2009**, *11*, 475-483, DOI 10.1039/B812809J.
- (28) Davis, J. H.: Ionic Liquids Derived from Natural Products and other Novel Chemistries. In *Green Industrial Applications of Ionic Liquids*; Rogers, R. D., Seddon, K. R., Volkov, S., Eds.; Springer Netherlands: Dordrecht, 2002; pp 385-401.
- (29) Ohno, H.; Fukumoto, K. Amino Acid Ionic Liquids. *Acc. Chem. Res.* **2007**, *40*, 1122-1129, DOI 10.1021/ar700053z.
- (30) Mato, I.; Huidobro, J. F.; Sánchez, M. P.; Muniategui, S.; Fernández-Muiño, M. A.; Sancho, M. T. Enzymatic determination of total d-Gluconic acid in honey. *J. Agric. Food Chem.* **1997**, *45*, 3550-3553, DOI 10.1021/jf970012c.
- (31) Sanarico, D.; Motta, S.; Bertolini, L.; Antonelli, A. HPLC determination of organic acids in traditional balsamic vinegar of Reggio Emilia. *J. Liq. Chromatogr. Relat. Technol.* **2003**, *26*, 2177-2187, DOI 10.1081/JLC-120022402.
- (32) Kulkarni, P. S.; Branco, L. C.; Crespo, J. G.; Nunes, M. C.; Raymundo, A.; Afonso, C. A. M. Comparison of Physicochemical Properties of New Ionic Liquids Based on Imidazolium, Quaternary Ammonium, and Guanidinium Cations. *Chem. Eur. J.* **2007**, *13*, 8478-8488, DOI 10.1002/chem.200700965.
- (33) Rohr, M.; Kubicek, C.; Kominek, J.: Gluconic Acid. In *Biotechnology: A Comprehensive Treatise in 8 Volumes. Volume 3. Biomass, Microorganisms for Special Applications, Microbial Products I, Energy from Renewable Resources.*; 04/14 ed.; Rehm, H. R., G., Eds., Ed.; Cambridge University Press, 1983; Vol. 3; pp 455-465.
- (34) Ramachandran, S.; Nair, S.; Larroche, C.; Pandey, A.: Gluconic Acid. In *Current Developments in Biotechnology and Bioengineering Production, Isolation and Purification of Industrial Products*; Pandey, A. N., S.; Soccol, C. R., Eds., Ed.; Elsevier, 2017; pp 886.
- (35) Pal, P.; Kumar, R.; Banerjee, S. Manufacture of gluconic acid: A review towards process intensification for green production. *Chemical Engineering and Processing: Process Intensification* **2016**, *104*, 160-171.
- (36) Boley, S. J.; Schultz, L.; Schwartz, S.; Katz, A.; Allen, A. C. Potassium Citrate and Potassium Gluconate Versus Potassium Chloride: Experimental Evaluation of Relative Intestinal Toxicity. *JAMA* **1967**, *199*, 215-217.
- (37) Helms, J. A.; Della-Fera, M. A.; Mott, A. E.; Frank, M. E. Effects of chlorhexidine on human taste perception. *Arch. Oral Biol.* **1995**, *40*, 913-920, DOI 10.1016/0003-9969(95)00062-T.
- (38) Jackson, J. L.; Peterson, C.; Lesho, E. A Meta-analysis of Zinc Salts Lozenges and the Common Cold. *JAMA Internal Medicine* **1997**, *157*, 2373-2376, DOI 10.1001/archinte.1997.00440410105012.

- (39) Stetten, M. R.; Stetten, D., Jr. The metabolism of gluconic acid. *J. Biol. Chem.* **1950**, *187*, 241-252.
- (40) Rohatgi, N.; Nielsen, T. K.; Bjorn, S. P.; Axelsson, I.; Paglia, G.; Voldborg, B. G.; Palsson, B. O.; Rolfsson, O. Biochemical characterization of human gluconokinase and the proposed metabolic impact of gluconic acid as determined by constraint based metabolic network analysis. *PLoS One* **2014**, *9*, e98760, DOI 10.1371/journal.pone.0098760.
- (41) Chen, Y.; Yang, Q.; Chen, J.; Bai, G.; Zhuo, K. Interaction of gluconate-based ionic liquids with common solvents: A study of volumetric, viscosity and conductivity properties. *J. Mol. Liq.* **2016**, *223*, 1013-1020, DOI 10.1016/j.molliq.2016.09.022.
- (42) Costa, A.; Forte, A.; Zalewska, K.; Tiago, G.; Petrovski, Z.; Branco, L. C. Novel biocompatible ionic liquids based on gluconate anion. *Green Chem. Lett. Rev.* **2015**, *8*, 8-12, DOI 10.1080/17518253.2014.951695.
- (43) Javed, F.; Ullah, F.; Akil Hazizan, M.: Synthesis, characterization and cellulose dissolution capabilities of ammonium-based room temperature ionic liquids (RTILs). In *Pure Appl. Chem.*, 2018; Vol. 90; pp 1019.
- (44) Billeci, F.; D'Anna, F.; Gunaratne, H. Q. N.; Plechkova, N. V.; Seddon, K. R. "Sweet" ionic liquid gels: materials for sweetening of fuels. *Green Chem.* **2018**, *20*, 4260-4276, DOI 10.1039/C8GC01615A.
- (45) Billeci, F.; Gunaratne, H. Q. N.; D'Anna, F.; Morgan, G. G.; Seddon, K. R.; Plechkova, N. V. A magnetic self-contained thermochromic system with convenient temperature range. *Green Chem.* **2019**, *21*, 1412-1416, DOI 10.1039/C8GC03906B.
- (46) Petkovic, M.; Ferguson, J. L.; Gunaratne, H. Q. N.; Ferreira, R.; Leitão, M. C.; Seddon, K. R.; Rebelo, L. P. N.; Pereira, C. S. Novel biocompatible cholinium-based ionic liquids—toxicity and biodegradability. *Green Chem.* **2010**, *12*, 643-649, DOI 10.1039/B922247B.
- (47) OECD: *Test No. 236: Fish Embryo Acute Toxicity (FET) Test*, 2013.
- (48) Howe, K.; Clark, M. D.; Torroja, C. F.; Torrance, J.; Berthelot, C.; Muffato, M.; Collins, J. E.; Humphray, S.; McLaren, K.; Matthews, L.; McLaren, S.; Sealy, I.; Caccamo, M.; Churcher, C.; Scott, C.; Barrett, J. C.; Koch, R.; Rauch, G.-J.; White, S.; Chow, W.; Kilian, B.; Quintais, L. T.; Guerra-Assunção, J. A.; Zhou, Y.; Gu, Y.; Yen, J.; Vogel, J.-H.; Eyre, T.; Redmond, S.; Banerjee, R.; Chi, J.; Fu, B.; Langley, E.; Maguire, S. F.; Laird, G. K.; Lloyd, D.; Kenyon, E.; Donaldson, S.; Sehra, H.; Almeida-King, J.; Loveland, J.; Trevanion, S.; Jones, M.; Quail, M.; Willey, D.; Hunt, A.; Burton, J.; Sims, S.; McLay, K.; Plumb, B.; Davis, J.; Clee, C.; Oliver, K.; Clark, R.; Riddle, C.; Elliott, D.; Threadgold, G.; Harden, G.; Ware, D.; Begum, S.; Mortimore, B.; Kerry, G.; Heath, P.; Phillimore, B.; Tracey, A.; Corby, N.; Dunn, M.; Johnson, C.; Wood, J.; Clark, S.; Pelan, S.; Griffiths, G.; Smith, M.; Glithero, R.; Howden, P.; Barker, N.; Lloyd, C.; Stevens, C.; Harley, J.; Holt, K.; Panagiotidis, G.; Lovell, J.; Beasley, H.; Henderson, C.; Gordon, D.; Auger, K.; Wright, D.; Collins, J.; Raisen, C.; Dyer, L.; Leung, K.; Robertson, L.; Ambridge, K.; Leongamornlert, D.; McGuire, S.; Gilderthorp, R.; Griffiths, C.; Manthravadi, D.; Nichol, S.; Barker, G. The zebrafish reference genome sequence and its relationship to the human genome. *Nature* **2013**, *496*, 498, 10.1038/nature12111
<https://www.nature.com/articles/nature12111#supplementary-information>.
- (49) Kong, E. Y.; Cheng, S. H.; Yu, K. N. Zebrafish as an In Vivo Model to Assess Epigenetic Effects of Ionizing Radiation. *International Journal of Molecular Sciences* **2016**, *17*, 2108,

- (50) Cancemi, P.; Albanese, N. N.; DiCara, G.; Marabeti, M. R.; Costantini, F.; Minafra, S.; Pucci-Minafra, I. Multiple changes induced by fibroblasts on breast cancer cells. *Connective tissue research* **2010**, *51*, 88-104, DOI 10.3109/03008200903100651.
- (51) Di Cara, G.; Marengo, G.; Albanese, N. N.; Marabeti, M. R.; Musso, R.; Cancemi, P.; Pucci-Minafra, I. Proteomic Profiling of Trastuzumab (Herceptin®)-sensitive and -resistant SKBR-3 Breast Cancer Cells. *Anticancer Res.* **2013**, *33*, 489-503,
- (52) Pucci-Minafra, I.; Cancemi, P.; Cara, G. D.; Minafra, L.; Feo, S.; Forlino, A.; Tira, M. E.; Tenni, R.; Martini, D.; Ruggeri, A.; Minafra, S. Decorin Transfection Induces Proteomic and Phenotypic Modulation in Breast Cancer Cells 8701-BC. *Connective Tissue Research* **2008**, *49*, 30-41, DOI 10.1080/03008200701820443.
- (53) Buttacavoli, M.; Albanese, N. N.; Di Cara, G.; Alduina, R.; Faleri, C.; Gallo, M.; Pizzolanti, G.; Gallo, G.; Feo, S.; Baldi, F.; Cancemi, P. Anticancer activity of biogenerated silver nanoparticles: an integrated proteomic investigation. *Oncotarget* **2017**, *9*, 9685-9705, DOI 10.18632/oncotarget.23859.
- (54) Stoppa, A.; Zech, O.; Kunz, W.; Buchner, R. The Conductivity of Imidazolium-Based Ionic Liquids from (-35 to 195) °C. A. Variation of Cation's Alkyl Chain. *J. Chem. Eng. Data* **2010**, *55*, 1768-1773, DOI 10.1021/je900789j.
- (55) McElroy, C. R.; Constantinou, A.; Jones, L. C.; Summerton, L.; Clark, J. H. Towards a holistic approach to metrics for the 21st century pharmaceutical industry. *Green Chem.* **2015**, *17*, 3111-3121, DOI 10.1039/C5GC00340G.
- (56) Cancemi, P.; Buttacavoli, M.; D'Anna, F.; Feo, S.; Fontana, R. M.; Noto, R.; Sutura, A.; Vitale, P.; Gallo, G. The effects of structural changes on the anti-microbial and anti-proliferative activities of diimidazolium salts. *New J. Chem.* **2017**, *41*, 3574-3585, DOI 10.1039/C6NJ03904A.
- (57) Malhotra, S. V.; Kumar, V. A profile of the in vitro anti-tumor activity of imidazolium-based ionic liquids. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 581-585, DOI 10.1016/j.bmcl.2009.11.085.
- (58) Xia, X.; Wan, R.; Wang, P.; Huo, W.; Dong, H.; Du, Q. Toxicity of imidazoles ionic liquid [C16mim]Cl to Hela cells. *Ecotoxicol. Environ. Saf.* **2018**, *162*, 408-414, DOI 10.1016/j.ecoenv.2018.07.022.
- (59) Pavel, A. B.; Korolev, K. S. Genetic load makes cancer cells more sensitive to common drugs: evidence from Cancer Cell Line Encyclopedia. *Scientific Reports* **2017**, *7*, 1938, DOI 10.1038/s41598-017-02178-1.
- (60) Chacon, E.; Acosta, D.; Lemasters, J. J.: 9 - Primary Cultures of Cardiac Myocytes as In Vitro Models for Pharmacological and Toxicological Assessments. In *In Vitro Methods in Pharmaceutical Research*; Castell, J. V., Gómez-Lechón, M. J., Eds.; Academic Press: San Diego, 1997; pp 209-223.
- (61) Kumar, V.; Malhotra, S. V. Study on the potential anti-cancer activity of phosphonium and ammonium-based ionic liquids. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 4643-4646, DOI 10.1016/j.bmcl.2009.06.086.
- (62) Frade, R. F. M.; Rosatella, A. A.; Marques, C. S.; Branco, L. C.; Kulkarni, P. S.; Mateus, N. M. M.; Afonso, C. A. M.; Duarte, C. M. M. Toxicological evaluation on human colon carcinoma cell line (CaCo-2) of ionic liquids based on imidazolium, guanidinium, ammonium, phosphonium, pyridinium and pyrrolidinium cations. *Green Chem.* **2009**, *11*, 1660-1665, DOI 10.1039/B914284N.

- (63) Pretti, C.; Renzi, M.; Ettore Focardi, S.; Giovani, A.; Monni, G.; Melai, B.; Rajamani, S.; Chiappe, C. Acute toxicity and biodegradability of N-alkyl-N-methylmorpholinium and N-alkyl-DABCO based ionic liquids. *Ecotoxicol. Environ. Saf.* **2011**, *74*, 748-753, DOI 10.1016/j.ecoenv.2010.10.032.
- (64) Cao, L.; Zhu, P.; Zhao, Y.; Zhao, J. Using machine learning and quantum chemistry descriptors to predict the toxicity of ionic liquids. *J. Hazard. Mater.* **2018**, *352*, 17-26, DOI 10.1016/j.jhazmat.2018.03.025.
- (65) Carangio, A.; Srinivasan, S.; McGuigan, C.; Andrei, G.; Snoeck, R.; De Clercq, E.; Balzarini, J. Bicyclic Nucleoside Inhibitors of Varicella-Zoster Virus: Effect of Terminal Aryl Substitution in the Side-Chain. *Antiviral Chemistry and Chemotherapy* **2002**, *13*, 263-271, DOI 10.1177/095632020201300501.
- (66) McGuigan, C.; Brancale, A.; Barucki, H.; Srinivasan, S.; Jones, G.; Pathirana, R.; Carangio, A.; Blewett, S.; Luoni, G.; Bidet, O.; Jukes, A.; Jarvis, C.; Andrei, G.; Snoeck, R.; De Clercq, E.; Balzarini, J. Furano Pyrimidines as Novel Potent and Selective Anti-VZV Agents. *Antiviral Chemistry and Chemotherapy* **2001**, *12*, 77-89, DOI 10.1177/095632020101200201.
- (67) Pagano, M.; Faggio, C. The use of erythrocyte fragility to assess xenobiotic cytotoxicity. *Cell Biochem. Funct.* **2015**, *33*, 351-355, DOI 10.1002/cbf.3135.
- (68) Gioia, R.; Tonelli, F.; Ceppi, I.; Biggiogera, M.; Leikin, S.; Fisher, S.; Tenedini, E.; Yorgan, T. A.; Schinke, T.; Tian, K.; Schwartz, J.-M.; Forte, F.; Wagener, R.; Villani, S.; Rossi, A.; Forlino, A. The chaperone activity of 4PBA ameliorates the skeletal phenotype of Chihuahua, a zebrafish model for dominant osteogenesis imperfecta. *Hum. Mol. Genet.* **2017**, *26*, 2897-2911, DOI 10.1093/hmg/ddx171.
- (69) Ruokonen, S.-K.; Sanwald, C.; Sundvik, M.; Polnick, S.; Vyavaharkar, K.; Duša, F.; Holding, A. J.; King, A. W. T.; Kilpeläinen, I.; Lämmerhofer, M.; Panula, P.; Wiedmer, S. K. Effect of Ionic Liquids on Zebrafish (*Danio rerio*) Viability, Behavior, and Histology; Correlation between Toxicity and Ionic Liquid Aggregation. *Environ. Sci. Technol.* **2016**, *50*, 7116-7125, DOI 10.1021/acs.est.5b06107.
- (70) Anderson, J. L.; Armstrong, D. W. High-Stability Ionic Liquids. A New Class of Stationary Phases for Gas Chromatography. *Anal. Chem.* **2003**, *75*, 4851-4858, DOI 10.1021/ac0345749.
- (71) Baranyai, K. J.; Deacon, G. B.; MacFarlane, D. R.; Pringle, J. M.; Scott, J. L. Thermal Degradation of Ionic Liquids at Elevated Temperatures. *Aust. J. Chem.* **2004**, *57*, 145-147, DOI 10.1071/CH03221.
- (72) Kroon, M. C.; Buijs, W.; Peters, C. J.; Witkamp, G.-J. Quantum chemical aided prediction of the thermal decomposition mechanisms and temperatures of ionic liquids. *Thermochim. Acta* **2007**, *465*, 40-47, DOI 10.1016/j.tca.2007.09.003.
- (73) Huddleston, J. G.; Visser, A. E.; Reichert, W. M.; Willauer, H. D.; Broker, G. A.; Rogers, R. D. Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation. *Green Chem.* **2001**, *3*, 156-164, DOI 10.1039/B103275P.
- (74) Tokuda, H.; Ishii, K.; Susan, M. A. B. H.; Tsuzuki, S.; Hayamizu, K.; Watanabe, M. Physicochemical Properties and Structures of Room-Temperature Ionic Liquids. 3. Variation of Cationic Structures. *J. Phys. Chem. B* **2006**, *110*, 2833-2839, DOI 10.1021/jp053396f.
- (75) Chauvin, Y.; Gilbert, B.; Guibard, I. Catalytic dimerization of alkenes by nickel complexes in organochloroaluminate molten salts. *J. Chem. Soc., Chem. Commun.* **1990**, 1715-1716, DOI 10.1039/C39900001715.

(76) Branco, L. C.; Carrera, G. V. S. M.; Aires-de-Sousa, J.; Martin, I. L.; Frade, R.; Afonso, C. A. M.: Physico-Chemical Properties of Task-Specific Ionic Liquids. In *Ionic Liquids: Theory, Properties, New Approaches*, 2011.

(77) Ichikawa, T.; Kato, T.; Ohno, H. Dimension control of ionic liquids. *Chem. Commun.* **2019**, 55, 8205-8214, DOI 10.1039/C9CC04280F.


(78) Stolte, S.; Matzke, M.; Arning, J.; Bösch, A.; Pitner, W.-R.; Welz-Biermann, U.; Jastorff, B.; Ranke, J. Effects of different head groups and functionalised side chains on the aquatic toxicity of ionic liquids. *Green Chem.* **2007**, 9, 1170-1179, DOI 10.1039/B711119C.

(79) Mauro, N. A.; Blodgett, M.; Johnson, M. L.; Vogt, A. J.; Kelton, K. F. A structural signature of liquid fragility. *Nature Communications* **2014**, 5, 4616, DOI 10.1038/ncomms5616.

(80) Angell, C. A. Relaxation in liquids, polymers and plastic crystals — strong/fragile patterns and problems. *J. Non-Cryst. Solids* **1991**, 131-133, 13-31, DOI 10.1016/0022-3093(91)90266-9.

(81) Ramos, J. M.; Afonso, C.; Branco, L. Glass transition relaxation and fragility in two room temperature ionic liquids. *J. Therm. Anal. Calorim.* **2003**, 71, 659-666, DOI 10.1023/a:1022884716750.

TOC



SYNOPSIS. "Sweet" ionic liquids incorporating moiety deriving from food waste were obtained. They show good physico-chemical properties and drastically reduced toxicity.