



Uncoupling Protein 2 as a pathogenic determinant and therapeutic target in cardiovascular and metabolic diseases

Journal:	<i>Current Neuropharmacology</i>
Manuscript ID	CN-2021-0047.R1
Manuscript Type:	Invited Review
Date Submitted by the Author:	n/a
Complete List of Authors:	stanzione, rosita; IRCCS Neurological Institute of Southern Italy NEUROMED forte, maurizio; IRCCS Neurological Institute of Southern Italy NEUROMED cotugno, maria; IRCCS Neurological Institute of Southern Italy NEUROMED bianchi, franca; IRCCS Neurological Institute of Southern Italy NEUROMED marchitti, simona; IRCCS Neurological Institute of Southern Italy NEUROMED busceti, carla letizia; IRCCS Neurological Institute of Southern Italy NEUROMED fornai, francesco; IRCCS Neurological Institute of Southern Italy NEUROMED rubattu, speranza; IRCCS Neurological Institute of Southern Italy NEUROMED
Keywords:	UCP2, stroke, cardiovascular disease, diabetes, obesity, therapeutics

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Uncoupling Protein 2 as a pathogenic determinant and therapeutic target in cardiovascular and metabolic diseases

Rosita Stanzione^{1*}, Maurizio Forte¹, Maria Cotugno¹, Franca Bianchi¹, Simona Marchitti¹, Carla Letizia Busceti¹,
Francesco Fornai^{1,3}, Speranza Rubattu^{1,2*}

¹IRCCS Neuromed, Pozzilli (Isernia);

²Department of Clinical and Molecular Medicine, School of Medicine and Psychology, Sapienza University of Rome, Rome;

³Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa; Italy

Running title: UCP2 and cardiovascular-metabolic disorders

*Corresponding addresses:

1) Rosita Stanzione Ph.D. IRCCS Neuromed, Località Camerelle, 86077 Pozzilli, Is, Italy; phone: +390865915224/23; fax: +390865927575. e-mail: stanzione@neuromed.it

2) Speranza Rubattu M.D. Clinical and Molecular Medicine Department, School of Medicine and Psychology, Sapienza University of Rome, Ospedale S.Andrea, 00189 Rome, Italy; IRCCS Neuromed, Località Camerelle, 86077 Pozzilli, Is, Italy; phone: +390865915224/23; fax: +390865927575. e-mail: rubattu.speranza@neuromed.it

Abstract word count: 248; Text word count: 5035 (including figure legends)

Abstract

Uncoupling protein 2 (UCP2) is a mitochondrial protein that acts as an anion carrier. It is involved in the regulation of several processes including mitochondrial membrane potential, generation of reactive oxygen species within the inner mitochondrial membrane and calcium homeostasis. UCP2 expression can be regulated at different levels: genetic (gene variants), transcriptional [by peroxisome proliferator-activated receptors (PPARs) and microRNAs], and post-translational. Experimental evidence indicates that activation of UCP2 expression through the AMPK/PPAR- α axis exerts a protective effect toward renal damage and stroke occurrence in an animal model of ischemic stroke (IS) associated with hypertension. UCP2 plays a key role in heart diseases (myocardial infarction and cardiac hypertrophy) and metabolic disorders (obesity and diabetes). In humans, UCP2 genetic variants (-866G/A and Ala55Val) associate with an increased risk of type 2 diabetes mellitus and of IS development.

Over the last few years, many agents that modulate UCP2 expression have been identified. Some of them are natural compounds of plant origin such as Brassica oleracea, curcumin, berberine and resveratrol. Other molecules, currently used in clinical practice, include anti-diabetic (gliptin) and chemotherapeutic (doxorubicin and taxol) drugs. This evidence highlights the relevant role of UCP2 for the treatment of a wide range of diseases, which affect the national health systems of the Western countries.

We will review current knowledge on the physiological and pathological implications of UCP2 with particular regard to cardiovascular and metabolic disorders and will focus on the available therapeutic approaches affecting UCP2 level for the treatment of human diseases.

Keywords: UCP2, stroke, cardiovascular disease, diabetes, obesity, therapeutics

1. Introduction

The uncoupling proteins (UCPs) family includes four components [1, 2]. UCP1, or thermogenin, was the first member of the family to be discovered in hibernators and small rodents, where UCP1 is highly expressed in the mitochondria of brown adipose tissue (BAT). Herein, UCP1 plays a physiological role in the body temperature regulation by fat storage and mobilization control [3, 4]. UCP1 knock-out mice are unable to regulate the body temperature [5]. The other components of UCPs family (UCP2-UCP5) were characterized later on [6, 7]. UCP2 is ubiquitously expressed. UCP3 is more expressed in the skeletal muscle. UCP4 and UCP5 are present in the brain. The different tissue expression of UCPs reflects different functions. In fact, whereas UCP1 acts as a thermogenesis regulator in the BAT of vertebrates [8], UCP2 uncouples the proton flux and the ATP (adenosine triphosphate) synthesis and converts the energy generated in this process into heat instead of ATP. Moreover, the UCP2-UCP5 members of the family have a physiological role as reactive oxygen species (ROS) scavengers and regulators of ATP-dependent processes, including secretion [7, 9]. Interestingly, UCP2 and UCP3 knockout mice show increased ROS production [10]. This evidence has important implications for human health. In fact, it is known that high concentrations of ROS contribute to the development of several diseases [10]. As a matter of fact, evidence obtained in both animal models and humans demonstrates that UCPs are involved in the pathogenesis of metabolic, neurodegenerative and cardiovascular diseases. High levels of UCP2 exert a protective effect in atherosclerosis [11], heart diseases [myocardial ischemia/reperfusion (I/R) injury] [11] and ischemic stroke (IS) [12]. Consistently, a reduced expression of UCP2 correlates with increased renal and cerebrovascular damage in the stroke-prone spontaneously hypertensive rat (SHRSP) model [13, 14]. On the other hand, increased UCP2 expression inhibits insulin secretion from pancreatic β -cells and favors the development of type 2 diabetes mellitus (T2DM).

In this review, we describe the main physiological and pathological implications of UCP2 with a focus on its involvement in major cardiovascular and metabolic disorders, based on both experimental and human evidences. Furthermore, we discuss the therapeutic implications of UCP2 and the compounds available for the clinical use affecting UCP2 level.

2. Physiological functions of UCP2

The sequence of human UCP2 is very similar to that of rat and mouse (99% and 95% of homology, respectively) [15]. The strong conservation of the gene sequence between the different species indicates that this protein exerts an important physiological role in all organisms. Transcription and translation, as well as UCP2 activity, may be controlled by many stimuli such as free fatty acids (FFA), retinoic acid, lipopolysaccharides, superoxide anion and hormones,

1
2
3 including leptin and thyroid hormones [16, 17]. The main physiological role of UCP2 is related to the control of
4 mitochondrial ROS production [9, 18]. Additional evidence shows that UCP2 plays an important role in the process of
5 oxidative phosphorylation, in the regulation of ATP synthesis, in the maintenance of redox potential membrane, in the
6 control of calcium homeostasis, of cell apoptosis and death [17, 19].
7
8
9

10 11 12 **2.1 Oxidative phosphorylation, ATP synthesis and ROS production**

13
14 Oxidative phosphorylation is a complex process that, by a series of chemical reactions, generates highly energetic
15 molecules of ATP. This process takes place within the inner mitochondrial membrane through the electron transport
16 chain (ETC) made by five enzymatic complexes (labeled I to V) (Figure 1). During this process a flow of electrons is
17 generated by electron transporters starting from NADH (Complex I) or FADH₂ (Complex II). The Complex III
18 (cytochrome oxidoreductase) and the Complex IV (cytochrome c oxidase) also pump protons through the membrane in
19 the intermembranes space. The electrochemical gradient, together with the return of protons across the inner membrane,
20 activates the last enzyme of the chain, the ATP synthase (Complex V). The latter produces ATP through the reaction
21 $ADP + P_i + 3H^+ \rightleftharpoons ATP + H_2O$. The amount of ATP molecules generated from glucose catabolism varies across species
22 and is related to various cellular processes. Under physiological conditions, when the energy demand of the cells is low,
23 UCP2 dissociates the oxidation of different substrates (NADH, coenzyme Q, cytochrome c) from ADP phosphorylation.
24 It consequently reduces the ATP synthesis and the electrochemical membrane potential ($\Delta\Psi$) [20]. UCP2, by decreasing
25 the $\Delta\Psi$, protects the cells from ROS damage and death. The energy generated in this process is dissipated in the form of
26 heat [21]. As a consequence of the reduced ATP synthesis by UCP2, pancreatic β -cells from UCP2 knockout mice show
27 an increased ATP, ATP/ADP ratio and insulin secretion, the latter due to the blockade of ATP-sensitive channels and
28 the opening of the voltage-dependent Ca²⁺ channel involved in insulin secretion [22-24].
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Mitochondria are the major source of ROS (mainly superoxide or hydrogen peroxide) in the aerobic cells [25]. They are
44 generated at several sites of ETC, although the two main sites are Complex I (on the matrix side) and Complex III (on
45 both the intermembrane and the matrix side) (Figure 1) [9]. At physiological concentration, ROS act as signaling factors
46 whereas high levels exert detrimental effects. Superoxide anion (O₂⁻) is a dangerous element that rapidly interacts with
47 the lipid component of the cell membrane causing lipid peroxidation, protein and DNA damage. ROS level is kept
48 under the control of few antioxidant systems such as superoxide dismutase (SOD), catalase (CAT), glutathione
49 peroxidase (GPx) and glutathione reductase (GR). ROS level is also influenced by $\Delta\Psi$ [17] and is modulated by UCP2
50 activity [18]. UCP2, by subtracting electrons from ETC, reduces superoxide formation and limits the cell damage.
51
52
53
54
55
56
57
58
59
60

1
2
3 Animal models lacking UCP2 show higher ROS level. Conversely, overexpression of UCP2 exerts a protective function
4 against the oxidative stress damage [18, 26].
5
6
7

8 9 **2.2 Calcium regulation**

10 The calcium ion (Ca^{2+}) is an important intracellular messenger, which acts as a regulator of cellular processes including
11 the neurotransmitters release from neurons and the contraction of all muscle cell types. In addition, it is a fundamental
12 cofactor of many enzymes. Ca^{2+} is also important for maintaining the difference in potential across membranes of
13 excitable cells. Mitochondria and the endoplasmic reticulum sequester Ca^{2+} to ensure that homeostasis is maintained
14 [27]. Cytosolic Ca^{2+} level is directly related to $\Delta\Psi$. Since UCP2 is involved in the regulation of $\Delta\Psi$, it can influence the
15 calcium dynamic. In normal condition, $\Delta\Psi$ equals 30mV. However, this value can undergo small changes and it can
16 alter important cellular functions in the presence of increased UCP2 level [28]. A drastic reduction of Ca^{2+} can
17 influence cell signaling. On the other hand, a massive flow of Ca^{2+} in the cytosol can increase neuronal excitotoxicity,
18 as it happens following IS or in neurodegenerative diseases [29, 30]. An excess of Ca^{2+} can induce apoptosis and
19 necrosis [31].
20
21
22
23
24
25
26
27
28
29
30

31 **2.3 Regulation of cell apoptosis and death**

32 The mitochondrion and some of the mitochondrial proteins play a key role in the control of apoptosis and cell death.
33 Several studies demonstrated that UCP2 regulates mitochondrial apoptotic signaling. A recent study, performed in a
34 cerebral ischemia/reperfusion (I/R) mouse model, demonstrated that UCP2 deletion exacerbated cell apoptosis [32].
35 Conversely, in the middle cerebral artery occlusion (MCAO) rat model, UCP2 inhibits cell death linked to
36 mitochondrial dysfunction by inducing depolarization of the mitochondrial inner membrane, subsequent Ca^{2+} intake and
37 ROS reduction [17, 33]. On the other hand, overexpression of UCP2 inhibits mitochondrial death pathway in primary
38 cultures of neonatal rat cardiac ventricular myocytes [34].
39
40
41
42
43
44
45
46
47

48 **3. Implications of UCP2 in pre-clinical models and human diseases**

49 UCP2 is implicated in a wide range of physiological and pathological conditions including aging [35] obesity [36],
50 inflammation [26], cerebral and myocardial ischemia [14, 37], hyperinsulinemia [36, 38]. With regard to aging, we
51 reported an age-related decrease of UCP2 expression in the brain, heart and kidneys of the SHRSP model, likely
52 involved into the higher predisposition to the hypertensive organ damage development of this strain [39]. Ren et al. also
53 highlighted the relationship between UCP2, aging and cardiac dysfunction. UCP2 level decreased with advanced aging
54
55
56
57
58
59
60

1
2
3 along with abnormal autophagy, mitophagy, and mitochondrial integrity. UCP2 level was lower in isolated
4 cardiomyocytes from old mice compared to young ones [40, 41]. Moreover, aging was reported to suppress
5 mitochondrial proteins, including UCP2, with a more pronounced effect in the aged mitochondrial aldehyde
6 dehydrogenase (ALDH2) deleted mouse hearts [42]. This evidence clearly implies that increased ROS level, along with
7 decreased UCP2 activity, contribute to the aging process.
8
9
10
11
12
13

14 3.1 UCP2 and IS

15
16 IS is the most common form of stroke and it results from cerebral blood vessel occlusion by an embolus or a thrombus.
17 The initial event is followed by depolarization of neuronal membrane, ion imbalance, such as K^+ and Ca^{2+} , and a
18 massive release of glutamate from the presynaptic vesicles into the extracellular space. Higher Ca^{2+} level increases both
19 ROS production and caspase activation causing degradation of membrane phospholipids, structural proteins and nucleic
20 acids. An increase of glutamate level, instead, contributes to excitotoxic neuronal death through overstimulation of
21 glutamate receptors. All these events contribute to amplify cellular damage [43-45].
22
23
24
25
26
27

28 The upregulation of UCP2 exerted a neuroprotective role in the MCAO mouse and rat models, mainly due to its ability
29 to keep low the depolarization level of the mitochondrial inner membrane (approximately 10 mV). As a consequence,
30 the ROS production, the calcium uptake and the induction of cell death-inducing factors (caspases and other apoptosis
31 mediators) decreased [30]. Conversely, low UCP2 expression induced a membrane hyperpolarization with an increase
32 of $\Delta\Psi$ and consequent ROS accumulation [26, 33].
33
34
35
36
37

38 We previously demonstrated that both cerebral and renal expression of UCP2 were downregulated in the SHRSP model
39 when fed with a stroke permissive high-sodium/low potassium diet (Japanese-style diet, JD), as compared to the stroke-
40 resistant strain (SHRSR). In this experimental condition, we found that UCP2 downregulation was induced by an
41 upregulation of the microRNA-503 [37]. The treatment of JD-fed SHRSP with a Brassica oleracea (BO) sprout extract
42 or with fenofibrate, a drug used to treat abnormal blood lipid levels, significantly decreased brain expression of mir-
43 503, rescued UCP2 expression and consequently reduced ischemic brain damage [37, 46, 47]. Interestingly, the
44 epigenetic control of UCP2 plays a role also in the kidney of JD-fed SHRSP, where renal damage precedes stroke
45 events. In fact, we demonstrated that renal UCP2 expression was regulated by microRNA-34a and microRNA-24 [48]
46
47
48
49
50
51
52
53 (Table 1).

54
55 Several studies showed that the antioxidant activity of UCP2 was mediated by the peroxisome expression of many
56 genes. In particular, PPAR- α activation by fenofibrate, BO juice and curcumin administration exerted a protective role
57
58
59
60

1
2
3 toward stroke and renal damage in JD-fed SHRSP [49, 50]. Activation of PPAR- γ can also modulate UCP2 expression
4 in different tissues [51]. At the brain level, several PPAR- γ agonists, including troglitazone, an antidiabetic drug that
5 improves insulin resistance in peripheral tissues, led to an increase of UCP2 level, improved the *in vitro* motoneurons
6 survival [52] and the outcome following cerebral ischemia [53, 54]. This evidence confirmed that PPARs play a
7 fundamental role in modulating the beneficial effects of UCP2 [46].

8
9
10
11
12 The role of UCP2 has also been investigated in relation to stroke in humans. UCP2 and UCP5 expression was analyzed
13 in brain ischemic lesions where they appeared significantly elevated as compared to the intact area. The upregulation of
14 both UCPs was considered as an adaptive response of neuronal cells to the ischemic insult [55].

15
16
17
18 In recent years, some human studies identified different single nucleotide polymorphisms (SNPs) within *UCP2* in
19 relation to stroke occurrence. The -866G>A (rs659366) is a functional SNP located within the promoter region of
20 *UCP2*, probably at the level of one or more transcriptional factors binding sites [56]. The A allele variant at -866
21 position is associated with increased *UCP2* expression. Many studies demonstrated that the G allele is associated with
22 lower *UCP2* expression and higher ROS accumulation [57]. This gene variant was related to an increased risk of IS in
23 Chinese women with T2DM in a 4-year prospective study [58, 59]. In a recent study, the relationship between this SNP
24 and the functional prognosis was investigated in patients with embolic IS after early recanalization. Patients carrying the
25 AA genotype showed a better functional outcome compared to patients harboring the AG or GG genotypes. This result
26 suggests that the AA genotype acts as an independent marker of favorable prognosis in IS patients after recanalization
27 of proximal MCAO [60] (Table 2).

3.2 UCP2 and heart diseases

38
39
40
41 UCP2 is constitutively expressed in the heart of humans and rodents and its expression level depends on the plasma
42 FFA concentration. The overexpression of UCP2 by adenoviral construct in primary cultures of neonatal rat
43 cardiomyocytes protected them from the damage related to increased oxidative stress [34]. During myocardial infarction
44 (MI), the same detrimental processes occurring in IS, such as increased oxidative stress and cell death, are detected. Wu
45 et al. demonstrated that UCP2 protects the heart from myocardial I/R injury via induction of mitophagy, the selective
46 form of autophagy for mitochondria. In fact, in preconditioning myocardial I/R performed in rats, UCP2 was
47 upregulated in association with an increase of microtubule-associated protein-light chain 3 (LC3) and a decrease of p62,
48 two markers of autophagy, indicating a strong activation of this process. In addition, mitochondrial dysfunction and
49 cardiomyocytes death were reduced. By contrast, the protective effect of UCP2 in cardiomyocytes was abolished by
50
51
52
53
54
55
56
57
58
59
60

1
2
3 mdivi-1, a specific inhibitor of dynamin 1-like, resulting in the inhibition of mitochondrial fission-induced autophagy
4 [61].

5
6 UCP2 expression is differently modulated in cardiac hypertrophy. In this regard, Strøm et al. reported that Wistar rats
7
8 subjected to running on a treadmill developed physiological adaptive hypertrophy with an increase of left and right
9
10 ventricular mass, as compared to sedentary rats. Furthermore, gene expression profiling analysis of total RNA from
11
12 hearts revealed that UCP2 was downregulated in this experimental context, since ATP demand increased during
13
14 exercise. In contrast, UCP2 was upregulated in the pathological or maladaptive hypertrophy resulting from MI induced
15
16 by ligating the left coronary artery [62]. This evidence suggests that the downregulation of UCP2 acts as an adaptive
17
18 response in compensated hypertrophy whereas its upregulation is associated with the detrimental effects in maladaptive
19
20 hypertrophy [62, 63]. Cardiac hypertrophy may also be induced in experimental animal models by exposure to low
21
22 temperatures (4°C) along with increased cellular oxidative stress, apoptosis and autophagy. In this experimental
23
24 condition, UCP2 is upregulated as a likely consequence of its role in thermogenesis regulation in specific conditions
25
26 [64, 65] (Table 1).

27
28 UCP2 is also upregulated in animal models of cardiomyopathy induced by chronic β -adrenergic receptor stimulation. In
29
30 fact, UCP2 upregulation may be partially reverted by β -adrenergic receptor blockade or by angiotensin converting
31
32 enzyme inhibition [66, 67]. In addition, UCP2 deleted mice exposed to pulmonary arterial banding appeared
33
34 significantly protected against pressure overload-induced right heart failure as compared to UCP2 wild type mice [68]
35
36 In humans, the -866 AA and GA genotypes associated with reduced survival and higher levels of myeloperoxidase post-
37
38 MI in diabetic patients [69]. This SNP associated with coronary artery disease also in young South African Indians [70].
39
40 Of interest, the 45 bp insertion/deletion within exon 8 of *UCP2* was related to heart rate variability and high blood
41
42 pressure level in Japanese men [71] (Table 2).

43 44 45 **3.3 UCP2 in thermogenesis, obesity, T2DM and metabolic syndrome**

46
47 Soon after its discovery, UCP2 was mainly interpreted as a thermogenic protein involved in the regulation of energy
48
49 expenditure and obesity [10]. In fact, FFA increase the expression of UCP2 skeletal muscle mRNA, implying that
50
51 UCP2 is somehow involved in fatty acid metabolism. Later on, it was understood that, differently from UCP1, UCP2 is
52
53 not normally thermogenic and can be involved in this process only under specific conditions [10, 64, 65].

54
55 Obesity and T2DM are two strictly related pathological conditions and their prevalence is increasing worldwide [72].
56
57 UCP2 is involved in the development of obesity as well as of diabetes. Zhang et al. demonstrated that UCP2 acts as a

1
2
3 negative regulator of insulin secretion playing a key role in the determination of pancreatic β -cell dysfunction and
4 consequently in the development of obesity and diabetes in experimental conditions [23] (Table 1). Consistently, when
5 UCP2 expression was suppressed by microRNA-15a in mouse pancreatic β -cells, an increase of insulin secretion was
6 observed [73]. UCP2 is downregulated in the streptozotocin (STZ)-induced diabetic mouse model. This model develops
7 cardiomyopathy, a drastic reduction of mitochondrial $\Delta\Psi$ and an increase of cell death. The overexpression of ALDH2
8 in STZ-induced diabetic mice exerted beneficial effects on cardiac structure and function, mitochondrial function and
9 cell survival. UCP2 levels were preserved in STZ-ALDH2 transgenic mice as compared to STZ mice [42, 74].

10
11
12
13
14
15
16 The correlation between the -866G/A SNP and a greater risk of developing diabetes in the presence of obesity has been
17 established in humans [75]. The wild-type G allele of the -866G/A SNP was associated with reduced UCP2 mRNA
18 expression in adipose tissue, reduced transcriptional activity *in vitro* and increased risk of obesity *in vivo*. Kempler et al.
19 showed that the pancreatic transcription factor paired box protein (PAX6) preferentially binds and activates the UCP2
20 promoter in the presence of the wild-type G allele in the insulinoma cells (INS1-E) derived from rat pancreatic β -cells.
21 In the same study, the correlation between UCP2 genotypes and the pancreatic β -cells function was analyzed in 39 non-
22 diabetic obese patients. Subjects carrying the G allele displayed a greater disposition index, the product of insulin
23 sensitivity (SI) and acute insulin response (AIR) to glucose, as compared to subjects carrying the A allele variant. In
24 addition, obese subjects with and without T2DM and the different UCP2 genotypes were compared. This analysis
25 revealed that the G allele was associated with a twofold reduction of T2DM risk in obese subjects compared to carrier
26 of the A allele variant. Additional case/control and meta-analysis studies confirmed the relationship between -866G/A,
27 obesity and T2DM in several human populations such as Finnish [76], Iranian [77], Spanish [78] and Danish [79]
28 populations. The wild type G allele within the UCP2 promoter, although frequently encountered in obese subjects,
29 protected them from T2DM development [80] (Table 2). A more recent meta-analysis confirmed that -866G/A SNP is
30 significantly associated with T2DM especially in Asian populations [81].

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45 Yang et al. observed that -866G/A SNP, in association with the Trp64Arg mutation of the β 3-adrenergic receptor gene,
46 was related with the therapeutic efficacy of rosiglitazone, an anti-diabetic drug, in Chinese T2DM patients [82].

47
48
49 Finally, the Ala55Val SNP (rs660339), a missense variant located within the exon 4 of UCP2 (where there is a change
50 of C to T in the position 164 of the transcript), is commonly associated with higher risk of obesity and higher incidence
51 of T2DM [83] (Table 2). On the other hand, additional studies reported that this SNP associated with an increased
52 insulin secretion in European-American women [84] and with a reduced risk of T2DM in Asian Indians [85].

53
54
55
56
57
58
59 The presence of obesity at the abdominal level is one of the four conditions needed to diagnose metabolic syndrome
60 (MS), along with high blood glucose, high serum triglycerides and low serum high-density lipoprotein (HDL) levels

[86]. Patients with MS show a 2-fold higher risk of developing cardiovascular events, such as MI and IS, and of developing T2DM compared to subjects without MS [87]. In the MS rat model (Mets) UCP2 expression is increased in the liver, where it acts as an adaptive mechanism of protection toward obesity-related oxidative stress [88]. In this model, the administration of both resveratrol and quercetin leads to increased UCP2 and PPAR- α expression in the abdominal white adipose tissue, and it improves metabolic parameters such as dyslipidemia, central adiposity and insulin resistance [89]. Recently, Wei D et al. demonstrated that a treatment with Shexiang Baoxin Pill (SBP) (containing a mixture composed of seven raw medicinal materials including Radix ginseng and Venenum bufonis) in the MS rat model exerted a protective therapeutic effect against this disorder, probably due to the interaction of anti-inflammatory and antioxidant mechanisms which are able to improve lipid metabolism and protect mitochondrial function. In particular, the authors demonstrated that the expression level of UCP2 increased in several tissues of MS rats, especially in the heart, after SBP treatment. Conversely, the expression level of cytochrome b, the main subunit of mitochondrial complex III, was downregulated in the liver, heart, skeletal muscle and adipose tissue of MS rats. The expression of ATPase (mitochondrial Complex V) appeared to be upregulated after treatment with SBP for 12 weeks [90, 91] (Table 1).

Recently, the association of some components of MS with the -866G/A SNP has been observed in patients with non-alcoholic fatty liver disease (NAFLD). In a case/control study involving 75 patients with NAFLD and 76 healthy individuals, subjects carrying the GA genotype had higher values of central obesity indices and metabolic abnormalities compared with healthy individuals [92].

The 45 bp insertion/deletion (I/D) within the 3'-untranslated region (UTR) of *UCP2* was studied in relation to some MS markers such as adiponectin, leptin, tumor necrosis factor- α (TNF α), and interleukin-6 (IL-6) levels. In a study performed in post-menopausal obese women, subjects carrying the DD genotype, but not those carrying the ID genotype, showed an increased level of adiponectin whereas leptin, TNF α , and IL-6 levels were significantly decreased after exercise training [93] (Table 2). The mechanism underlying this phenomenon remains obscure.

4. Pharmacological modulation of UCP2

As discussed above, UCP2 expression can be regulated at multiple levels [94]. In addition, the UCP2 biological effects are tissue specific (Figure 1). Any therapeutic approach targeting UCP2 should take into account the type of target tissue [17]. Much of the therapeutic potential of UCP2 depends on its antioxidant and anti-inflammatory properties [53,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

54] which make UCP2 an attractive therapeutic tool in many diseases, including heart and brain diseases, and inflammatory disorders [17].

Several available drugs affect UCP2 expression levels, although we cannot precisely state if they target directly UCP2.

Among them, the dipeptidyl peptidase-4 (DPP-4) inhibitors or gliptins, a class of oral anti-diabetic drugs, are able to restore endothelial function in hypertension by reducing oxidative stress [95]. In particular, chronic sitagliptin administration attenuated endothelium-dependent contraction (EDC) and reduced ROS level by the upregulation of Glucagon-like Peptide-1 (GLP-1)/5-adenosine monophosphate-activated protein kinase (AMPK)/UCP2 pathway and the downregulation of cyclooxygenase-2 (COX-2) expression in renal arteries from spontaneously hypertensive rats (SHR) and angiotensin II (Ang II)-infused mouse aorta. Conversely, sitagliptin did not affect EDC in Ang II-infused UCP2-KO mice. In addition, the beneficial effects of sitagliptin on endothelial function were inhibited by genipin, a UCP2 inhibitor, in renal arteries of SHR confirming the relevance of UCP2 in this context.

Consistently, UCP2 overexpression, obtained by an adenoviral construct, improved the vascular function in the same experimental rat model. The administration of exendin-4 (Ex-4), a GPL-1 agonist, also improved renal vascular function in renal arteries of both SHR and hypertensive patients through UCP2 level increase with a parallel decrease of COX2 and ROS levels [96, 97].

Other anti-diabetic molecules, such as metformin and berberine, modulate UCP2 level. Metformin, the most common anti-diabetic drug, acts by AMPK upregulation. In fact, AMPK inhibition by compound C prevented the UCP2 increase [98]. Berberine, an isoquinolone alkaloid extracted from plants and used in traditional Chinese medicine, shows anti-diabetic and anti-atherogenic properties, improves dyslipidemia and endothelial function, and reduces the incidence of cardiovascular diseases. Berberine significantly increases UCP2 mRNA and protein level in an AMPK-dependent manner both *in vitro* and in a mouse model of atherosclerosis [99].

Previous studies have shown that PPAR agonists are able to modulate the UCP2 expression level. These drugs are a group of structurally diverse compounds that bind and activate PPAR receptors. Some of them, the thiazolidinediones (TZD), also known as glitazones, are potent PPAR- γ agonists [100], commonly used in the clinical practice for the treatment of diabetes and MS with beneficial effects. These drugs cause a marked reduction of FFA plasmatic level and inhibit lipolysis in T2DM patients. Plasmatic FFA reduction, in turn, leads to fat mobilization from the muscle and the liver and improves insulin sensitivity in these organs [101]. Glitazones also contribute to lower blood pressure level and to improve lipid metabolism by increasing HDL cholesterol level and by reducing triglycerides level [102, 103]. Several studies revealed that the TZD administration, mainly rosiglitazone, induced a rapid increase of UCP2 both in mouse and human adipocyte cell lines [104, 105] as compared to control cells. Its effect appeared to be mediated by PPAR- γ [104,

1
2
3 106]. In addition, administration of rosiglitazone in Sprague-Dawley rats reduced mitochondrial hydrogen peroxide
4
5 level in rostral ventrolateral medulla, a region that contains the sympathetic premotor neurons for the maintenance of
6
7 vasomotor tone and of systemic arterial pressure. Rosiglitazone can exert a protective effect against oxidative stress-
8
9 associated neurogenic hypertension through UCP2 upregulation and ROS reduction [107].

10
11 Additional pre-clinical and *in vitro* studies analyzed the effect of pioglitazone, a well-known PPAR- γ agonist, toward
12
13 oxidative stress in the rat brain [97]. It was demonstrated that this drug, through PPAR- γ activation and UCP2
14
15 upregulation, attenuated oxidative stress in cerebral arteries of aging rats. They also showed that one-month oral
16
17 administration of pioglitazone improved endothelium-dependent relaxation by increasing both eNOS phosphorylation
18
19 and nitric oxide (NO) availability. PPAR- γ appears as a promising target for the treatment of diabetes and of age-related
20
21 cerebrovascular dysfunction.

22
23 No side effects dependent from UCP2 upregulation have been so far reported with the available compounds, although it
24
25 is known that NO, a mediator of UCP2, may contribute to exacerbate neuronal cell dysfunction and diminish cell
26
27 survival after brain injury [108]. Exaggerate NO production is also associated to acute pancreatitis in rats [109] and
28
29 liver damage in mice [110].

30
31 Interestingly, some chemotherapeutic drugs, such as doxorubicin (DOX) and taxol, used for the treatment of different
32
33 types of cancers, downregulate UCP2 expression by activation of c-Jun N-terminal kinase (JNK)/mitogen-activated
34
35 protein kinase (MAPK)/extracellular-signal-regulated kinase (ERK) signaling, with consequent ROS accumulation and
36
37 increase of apoptosis in melanoma cells [111]. UCP2 downregulation and consequent ROS accumulation have also
38
39 been associated with a severe cardiotoxicity in both animal models and cardiomyocytes treated with DOX.
40
41 Interestingly, administration of melatonin counteracted the cardiac toxic effects of DOX by improvement of
42
43 mitochondrial damage and upregulation of UCP2 [112].

44
45 Apart from the above mentioned drugs, UCP2 expression can be modulated by nutraceuticals as documented in the
46
47 SHRSP model [13, 37] and in humans [13]. Pagliaro et al. described the beneficial effects exerted in the cardiovascular
48
49 system by different natural compounds such as BO, curcumin, resveratrol and berberine. These natural compounds of
50
51 plant origin, through the activation of both AMPK and silent mating type information regulation-1 (SIRT-1), upregulate
52
53 PPARs (mainly PPAR- α and PPAR- γ) leading to UCP2 increase, reduction of ROS level and of inflammation,
54
55 improvement of brain ischemic injury and decrease of stroke occurrence [113]. The beneficial effects of BO in humans
56
57 were tested in a small clinical trial in which broccoli extract pills and fresh broccoli were administered simultaneously
58
59 for 7 days to healthy subjects. At the end of the study, a significant reduction of both total and LDL cholesterol, along
60
with a decrease of urinary 8-isoprostanes and of other markers of oxidative stress, was observed [114]. The

1
2
3 administration of broccoli juice for 12 weeks in 32 men with hypercholesterolemia significantly reduced plasma LDL
4 cholesterol and increased both HDL cholesterol and GPx activity, thus lowering cardiovascular disease risk [115].
5
6 Interestingly, treatment with resveratrol, known for its beneficial effects on diabetes progression, oxidative stress and
7 lipid metabolism, exerts a protective effect towards diabetic cardiomyopathy induced by STZ in a UCP2 dependent
8 manner [116]. In fact, resveratrol administration significantly increased UCP2 expression, reduced ROS level and
9 increased the manganese superoxide dismutase level. Conversely, UCP2 inhibition by siRNA hinders the
10 cardioprotective effects of resveratrol on ROS production and apoptosis level in high glucose-treated cardiomyocytes
11 [117].
12
13
14
15
16
17
18
19
20

21 **Conclusions**

22 The physiological role of UCP2 appears complex and it still needs to be fully dissected out. UCP2 expression and
23 functions are tissue specific. Most of the beneficial properties of UCP2 depend on its role as ROS scavenger. UCP2 is
24 also involved in the regulation of $\Delta\Psi$, ATP synthesis, calcium level and cell death.
25

26 UCP2 upregulation attenuates mitochondrial ROS production, consequently the cellular oxidative damage, and exerts
27 protective effects in several organs. On the other hand, UCP2 inhibition protects from hyperglycemia and diabetes
28 through the control of insulin secretion in the pancreatic β -cells.
29

30 Therefore, UCP2 behaves as a critical determinant of many human disorders such as atherosclerosis, ischemic heart
31 disease, neurodegenerative diseases, including IS, and diabetes. Current evidence suggests that UCP2 modulation may
32 represent a suitable tool for the treatment of several human diseases (Figure 2).
33

34 Further experimental studies will allow to improve our understanding of the molecular pathways involved in UCP2
35 activation. Human studies may reveal important new insights on the pathogenic role of UCP2 gene variants. The new
36 knowledge could facilitate the identification of novel pharmacological strategies that, by controlling the expression of
37 UCP2, can reduce the occurrence and progression of diseases in which UCP2 plays a fundamental role. In this latter
38 regard, additional investigations are necessary to identify the exact relationship between molecular mechanisms, cellular
39 effects and clinical consequences, along with potential caveats, upon pharmacological stimulation of UCP2.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **CONSENT FOR PUBLICATION**
4

5 Not applicable.
6

7
8 **CONFLICT OF INTEREST**
9

10 The author declares no conflict of interest, financial or otherwise.
11
12

13 **ACKNOWLEDGEMENTS**
14

15
16 This work was supported by a grant from the Italian Ministry of Health
17

18 **Authors' contributions**
19

20
21 Conceptualization: [RS, SR]; Writing- original draft preparation: [RS]; Writing - review and editing: [RS, MF, SR];
22 Literature search: [RS, MC, FB, SM]; Critical revision: [CB, FF, SR]. All authors read and approved the final version of
23 the manuscript.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Nedergaard J, Ricquier D, Kozak LP. Uncoupling proteins: current status and therapeutic prospects. *EMBO Rep.* (2005);6(10):917-21.
2. Rousset S, Alves-Guerra MC, Mozo J, Miroux B, Cassard-Doulier AM, Bouillaud F, et al. The biology of mitochondrial uncoupling proteins. *Diabetes.* (2004);53 Suppl 1:S130-5.
3. Nicholls DG. A history of UCP1. *Biochem Soc Trans.* (2001);29(Pt 6):751-5.
4. Nicholls DG, Locke RM. Thermogenic mechanisms in brown fat. *Physiol Rev.* (1984);64(1):1-64.
5. Feldmann HM, Golozoubova V, Cannon B, Nedergaard J. UCP1 ablation induces obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by living at thermoneutrality. *Cell Metab.* (2009);9(2):203-9.
6. Bouillaud F, Couplan E, Pecqueur C, Ricquier D. Homologues of the uncoupling protein from brown adipose tissue (UCP1): UCP2, UCP3, BMCP1 and UCP4. *Biochim Biophys Acta.* (2001);1504(1):107-19.
7. Erlanson-Albertsson C. Uncoupling proteins--a new family of proteins with unknown function. *Nutr Neurosci.* (2002);5(1):1-11.
8. Gaudry MJ, Jastroch M. Molecular evolution of uncoupling proteins and implications for brain function. *Neurosci Lett.* (2019);696:140-5.
9. Mailloux RJ, Harper ME. Uncoupling proteins and the control of mitochondrial reactive oxygen species production. *Free Radic Biol Med.* (2011);51(6):1106-15.
10. Brand MD, Esteves TC. Physiological functions of the mitochondrial uncoupling proteins UCP2 and UCP3. *Cell Metab.* (2005);2(2):85-93.
11. Tian XY, Ma S, Tse G, Wong WT, Huang Y. Uncoupling Protein 2 in Cardiovascular Health and Disease. *Front Physiol.* (2018);9:1060.
12. Lan C, Chen X, Zhang Y, Wang W, Wang WE, Liu Y, et al. Curcumin prevents strokes in stroke-prone spontaneously hypertensive rats by improving vascular endothelial function. *BMC Cardiovasc Disord.* (2018);18(1):43.
13. Pierelli G, Stanzione R, Forte M, Migliarino S, Perelli M, Volpe M, et al. Uncoupling Protein 2: A Key Player and a Potential Therapeutic Target in Vascular Diseases. *Oxid Med Cell Longev.* (2017);2017:7348372.
14. Busceti CL, Cotugno M, Bianchi F, Forte M, Stanzione R, Marchitti S, et al. Brain Overexpression of Uncoupling Protein-2 (UCP2) Delays Renal Damage and Stroke Occurrence in Stroke-Prone Spontaneously Hypertensive Rats. *Int J Mol Sci.* (2020);21(12).
15. Hidaka S, Kakuma T, Yoshimatsu H, Yasunaga S, Kurokawa M, Sakata T. Molecular cloning of rat uncoupling protein 2 cDNA and its expression in genetically obese Zucker fatty (fa/fa) rats. *Biochim Biophys Acta.* (1998);1389(3):178-86.
16. Thompson MP, Kim D. Links between fatty acids and expression of UCP2 and UCP3 mRNAs. *FEBS Lett.* (2004);568(1-3):4-9.
17. Mattiasson G, Sullivan PG. The emerging functions of UCP2 in health, disease, and therapeutics. *Antioxid Redox Signal.* (2006);8(1-2):1-38.

18. Cadenas S. Mitochondrial uncoupling, ROS generation and cardioprotection. *Biochim Biophys Acta Bioenerg.* (2018);1859(9):940-50.
19. Brand MD, Chien LF, Ainscow EK, Rolfe DF, Porter RK. The causes and functions of mitochondrial proton leak. *Biochim Biophys Acta.* (1994);1187(2):132-9.
20. Suski J, Lebiezinska M, Bonora M, Pinton P, Duszynski J, Wieckowski MR. Relation Between Mitochondrial Membrane Potential and ROS Formation. *Methods Mol Biol.* (2018);1782:357-81.
21. Zhao RZ, Jiang S, Zhang L, Yu ZB. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int J Mol Med.* (2019);44(1):3-15.
22. Joseph JW, Koshkin V, Saleh MC, Sivitz WI, Zhang CY, Lowell BB, et al. Free fatty acid-induced beta-cell defects are dependent on uncoupling protein 2 expression. *J Biol Chem.* (2004);279(49):51049-56.
23. Zhang CY, Baffy G, Perret P, Krauss S, Peroni O, Grujic D, et al. Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. *Cell.* (2001);105(6):745-55.
24. Pecqueur C, Alves-Guerra C, Ricquier D, Bouillaud F. UCP2, a metabolic sensor coupling glucose oxidation to mitochondrial metabolism? *IUBMB Life.* (2009);61(7):762-7.
25. Cadenas E, Davies KJ. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med.* (2000);29(3-4):222-30.
26. Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, et al. Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. *Nat Genet.* (2000);26(4):435-9.
27. Nicholls DG. Mitochondrial calcium function and dysfunction in the central nervous system. *Biochim Biophys Acta.* (2009);1787(11):1416-24.
28. Waldeck-Weiermair M, Malli R, Naghdi S, Trenker M, Kahn MJ, Graier WF. The contribution of UCP2 and UCP3 to mitochondrial Ca(2+) uptake is differentially determined by the source of supplied Ca(2+). *Cell Calcium.* (2010);47(5):433-40.
29. Takarada T, Fukumori R, Yoneda Y. Mitochondrial uncoupling protein-2 in glutamate neurotoxicity. *Nihon Yakurigaku Zasshi.* (2013);142(1):13-6.
30. Mehta SL, Li PA. Neuroprotective role of mitochondrial uncoupling protein 2 in cerebral stroke. *J Cereb Blood Flow Metab.* (2009);29(6):1069-78.
31. Orrenius S, Gogvadze V, Zhivotovsky B. Calcium and mitochondria in the regulation of cell death. *Biochem Biophys Res Commun.* (2015);460(1):72-81.
32. He M, Zhang T, Fan Y, Ma Y, Zhang J, Jing L, et al. Deletion of mitochondrial uncoupling protein 2 exacerbates mitophagy and cell apoptosis after cerebral ischemia and reperfusion injury in mice. *Int J Med Sci.* (2020);17(17):2869-78.
33. Mattiasson G, Shamloo M, Gido G, Mathi K, Tomasevic G, Yi S, et al. Uncoupling protein-2 prevents neuronal death and diminishes brain dysfunction after stroke and brain trauma. *Nat Med.* (2003);9(8):1062-8.
34. Teshima Y, Akao M, Jones SP, Marban E. Uncoupling protein-2 overexpression inhibits mitochondrial death pathway in cardiomyocytes. *Circ Res.* (2003);93(3):192-200.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

35. Mizuno T, Miura-Suzuki T, Yamashita H, Mori N. Distinct regulation of brain mitochondrial carrier protein-1 and uncoupling protein-2 genes in the rat brain during cold exposure and aging. *Biochem Biophys Res Commun.* (2000);278(3):691-7.
36. Fleury C, Neverova M, Collins S, Raimbault S, Champigny O, Levi-Meyrueis C, et al. Uncoupling protein-2: a novel gene linked to obesity and hyperinsulinemia. *Nat Genet.* (1997);15(3):269-72.
37. Rubattu S, Stanzione R, Bianchi F, Cotugno M, Forte M, Della Ragione F, et al. Reduced brain UCP2 expression mediated by microRNA-503 contributes to increased stroke susceptibility in the high-salt fed stroke-prone spontaneously hypertensive rat. *Cell Death Dis.* (2017);8(6):e2891.
38. Joseph JW, Koshkin V, Zhang CY, Wang J, Lowell BB, Chan CB, et al. Uncoupling protein 2 knockout mice have enhanced insulin secretory capacity after a high-fat diet. *Diabetes.* (2002);51(11):3211-9.
39. Rubattu S, Bianchi F, Busceti CL, Cotugno M, Stanzione R, Marchitti S, et al. Differential modulation of AMPK/PPARalpha/UCP2 axis in relation to hypertension and aging in the brain, kidneys and heart of two closely related spontaneously hypertensive rat strains. *Oncotarget.* (2015);6(22):18800-18.
40. Zhang Y, Mi SL, Hu N, Doser TA, Sun A, Ge J, et al. Mitochondrial aldehyde dehydrogenase 2 accentuates aging-induced cardiac remodeling and contractile dysfunction: role of AMPK, Sirt1, and mitochondrial function. *Free Radic Biol Med.* (2014);71:208-20.
41. Ren J, Yang L, Zhu L, Xu X, Ceylan AF, Guo W, et al. Akt2 ablation prolongs life span and improves myocardial contractile function with adaptive cardiac remodeling: role of Sirt1-mediated autophagy regulation. *Aging Cell.* (2017);16(5):976-87.
42. Zhang Y, Babcock SA, Hu N, Maris JR, Wang H, Ren J. Mitochondrial aldehyde dehydrogenase (ALDH2) protects against streptozotocin-induced diabetic cardiomyopathy: role of GSK3beta and mitochondrial function. *Bmc Med.* (2012);10:40.
43. Bano D, Nicotera P. Ca²⁺ signals and neuronal death in brain ischemia. *Stroke.* (2007);38(2 Suppl):674-6.
44. Sanganalmath SK, Gopal P, Parker JR, Downs RK, Parker JC, Jr., Dawn B. Global cerebral ischemia due to circulatory arrest: insights into cellular pathophysiology and diagnostic modalities. *Mol Cell Biochem.* (2017);426(1-2):111-27.
45. Lai TW, Zhang S, Wang YT. Excitotoxicity and stroke: identifying novel targets for neuroprotection. *Prog Neurobiol.* (2014);115:157-88.
46. Rubattu S, Di Castro S, Cotugno M, Bianchi F, Mattioli R, Baima S, et al. Protective effects of Brassica oleracea sprouts extract toward renal damage in high-salt-fed SHRSP: role of AMPK/PPARalpha/UCP2 axis. *J Hypertens.* (2015);33(7):1465-79.
47. Gallo G, Forte M, Stanzione R, Cotugno M, Bianchi F, Marchitti S, et al. Functional Role of Natriuretic Peptides in Risk Assessment and Prognosis of Patients with Mitral Regurgitation. *J Clin Med.* (2020);9(5).
48. Di Castro S, Scarpino S, Marchitti S, Bianchi F, Stanzione R, Cotugno M, et al. Differential modulation of uncoupling protein 2 in kidneys of stroke-prone spontaneously hypertensive rats under high-salt/low-potassium diet. *Hypertension.* (2013);61(2):534-41.
49. Hou X, Shen YH, Li C, Wang F, Zhang C, Bu P, et al. PPARalpha agonist fenofibrate protects the kidney from hypertensive injury in spontaneously hypertensive rats via inhibition of oxidative stress and MAPK activity. *Biochem Biophys Res Commun.* (2010);394(3):653-9.

- 1
2
3 50. Gelosa P, Banfi C, Gianella A, Brioschi M, Pignieri A, Nobili E, et al. Peroxisome proliferator-activated
4 receptor $\{\alpha\}$ agonism prevents renal damage and the oxidative stress and inflammatory processes affecting the
5 brains of stroke-prone rats. *J Pharmacol Exp Ther.* (2010);335(2):324-31.
6
7 51. Medvedev AV, Snedden SK, Raimbault S, Ricquier D, Collins S. Transcriptional regulation of the mouse
8 uncoupling protein-2 gene. Double E-box motif is required for peroxisome proliferator-activated receptor-gamma-
9 dependent activation. *J Biol Chem.* (2001);276(14):10817-23.
10
11 52. Nishijima C, Kimoto K, Arakawa Y. Survival activity of troglitazone in rat motoneurons. *J Neurochem.*
12 (2001);76(2):383-90.
13
14 53. Shimazu T, Inoue I, Araki N, Asano Y, Sawada M, Furuya D, et al. A peroxisome proliferator-activated
15 receptor-gamma agonist reduces infarct size in transient but not in permanent ischemia. *Stroke.* (2005);36(2):353-9.
16
17 54. Sundararajan S, Gamboa JL, Victor NA, Wanderi EW, Lust WD, Landreth GE. Peroxisome proliferator-
18 activated receptor-gamma ligands reduce inflammation and infarction size in transient focal ischemia. *Neuroscience.*
19 (2005);130(3):685-96.
20
21 55. Nakase T, Yoshida Y, Nagata K. Amplified expression of uncoupling proteins in human brain ischemic
22 lesions. *Neuropathology.* (2007);27(5):442-7.
23
24 56. Dalgaard LT, Andersen G, Larsen LH, Sorensen TI, Andersen T, Drivsholm T, et al. Mutational analysis of the
25 UCP2 core promoter and relationships of variants with obesity. *Obes Res.* (2003);11(11):1420-7.
26
27 57. Lapice E, Pinelli M, Pisu E, Monticelli A, Gambino R, Pagano G, et al. Uncoupling protein 2 G(-866)A
28 polymorphism: a new gene polymorphism associated with C-reactive protein in type 2 diabetic patients. *Cardiovasc*
29 *Diabetol.* (2010);9:68.
30
31 58. Chai Y, Gu B, Qiu JR, Yi HG, Zhu Q, Zhang L, et al. The uncoupling protein 2 -866G > a polymorphism is
32 associated with the risk of ischemic stroke in Chinese type 2 diabetic patients. *CNS Neurosci Ther.* (2012);18(8):636-
33 40.
34
35 59. Chai Y, Gu B, Qiu JR, Yi HG, Zhu Q, Zhang L, et al. Effects of uncoupling protein 2 -866G/A polymorphism
36 on platelet reactivity and prognosis in Chinese patients with type 2 diabetes and ischemic stroke. *Int J Neurosci.*
37 (2013);123(11):752-8.
38
39 60. Diaz-Maroto Cicuendez I, Fernandez-Diaz E, Garcia-Garcia J, Jordan J, Fernandez-Cadenas I, Montaner J, et
40 al. The UCP2-866G/A Polymorphism Could be Considered as a Genetic Marker of Different Functional Prognosis in
41 Ischemic Stroke After Recanalization. *Neuromolecular Med.* (2017);19(4):571-8.
42
43 61. Wu H, Ye M, Liu D, Yang J, Ding JW, Zhang J, et al. UCP2 protect the heart from myocardial
44 ischemia/reperfusion injury via induction of mitochondrial autophagy. *J Cell Biochem.* (2019);120(9):15455-66.
45
46 62. Strom CC, Aplin M, Ploug T, Christoffersen TE, Langfort J, Viese M, et al. Expression profiling reveals
47 differences in metabolic gene expression between exercise-induced cardiac effects and maladaptive cardiac
48 hypertrophy. *FEBS J.* (2005);272(11):2684-95.
49
50 63. Strom CC, Kruhoffer M, Knudsen S, Stensgaard-Hansen F, Jonassen TE, Orntoft TF, et al. Identification of a
51 core set of genes that signifies pathways underlying cardiac hypertrophy. *Comp Funct Genomics.* (2004);5(6-7):459-70.
52
53 64. Zhang Y, Li L, Hua Y, Nunn JM, Dong F, Yanagisawa M, et al. Cardiac-specific knockout of ET(A) receptor
54 mitigates low ambient temperature-induced cardiac hypertrophy and contractile dysfunction. *J Mol Cell Biol.*
55 (2012);4(2):97-107.
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

65. Kong X, Liu H, He X, Sun Y, Ge W. Unraveling the Mystery of Cold Stress-Induced Myocardial Injury. *Front Physiol.* (2020);11:580811.
66. Gaussin V, Tomlinson JE, Depre C, Engelhardt S, Antos CL, Takagi G, et al. Common genomic response in different mouse models of beta-adrenergic-induced cardiomyopathy. *Circulation.* (2003);108(23):2926-33.
67. Murakami K, Mizushige K, Noma T, Tsuji T, Kimura S, Kohno M. Perindopril effect on uncoupling protein and energy metabolism in failing rat hearts. *Hypertension.* (2002);40(3):251-5.
68. Esfandiary A, Kutsche HS, Schreckenber R, Weber M, Pak O, Kojonazarov B, et al. Protection against pressure overload-induced right heart failure by uncoupling protein 2 silencing. *Cardiovasc Res.* (2019);115(7):1217-27.
69. Palmer BR, Devereaux CL, Dhamrait SS, Mocatta TJ, Pilbrow AP, Frampton CM, et al. The common G-866A polymorphism of the UCP2 gene and survival in diabetic patients following myocardial infarction. *Cardiovasc Diabetol.* (2009);8:31.
70. Phulukdaree A, Moodley D, Khan S, Chuturgoon AA. Uncoupling protein 2 -866G/A and uncoupling protein 3 -55C/T polymorphisms in young South African Indian coronary artery disease patients. *Gene.* (2013);524(2):79-83.
71. Matsunaga T, Gu N, Yamazaki H, Tsuda M, Adachi T, Yasuda K, et al. Association of UCP2 and UCP3 polymorphisms with heart rate variability in Japanese men. *J Hypertens.* (2009);27(2):305-13.
72. Kopelman PG. Obesity as a medical problem. *Nature.* (2000);404(6778):635-43.
73. Sun LL, Jiang BG, Li WT, Zou JJ, Shi YQ, Liu ZM. MicroRNA-15a positively regulates insulin synthesis by inhibiting uncoupling protein-2 expression. *Diabetes Res Clin Pract.* (2011);91(1):94-100.
74. Ren J, Pulakat L, Whaley-Connell A, Sowers JR. Mitochondrial biogenesis in the metabolic syndrome and cardiovascular disease. *J Mol Med (Berl).* (2010);88(10):993-1001.
75. Dalgaard LT. Genetic Variance in Uncoupling Protein 2 in Relation to Obesity, Type 2 Diabetes, and Related Metabolic Traits: Focus on the Functional -866G>A Promoter Variant (rs659366). *J Obes.* (2011);2011:340241.
76. Salopuro T, Pulkkinen L, Lindstrom J, Kolehmainen M, Tolppanen AM, Eriksson JG, et al. Variation in the UCP2 and UCP3 genes associates with abdominal obesity and serum lipids: the Finnish Diabetes Prevention Study. *Bmc Med Genet.* (2009);10:94.
77. Heidari J, Akrami SM, Heshmat R, Amiri P, Fakhrzadeh H, Pajouhi M. Association study of the -866G/A UCP2 gene promoter polymorphism with type 2 diabetes and obesity in a Tehran population: a case control study. *Arch Iran Med.* (2010);13(5):384-90.
78. Ochoa MC, Santos JL, Azcona C, Moreno-Aliaga MJ, Martinez-Gonzalez MA, Martinez JA, et al. Association between obesity and insulin resistance with UCP2-UCP3 gene variants in Spanish children and adolescents. *Mol Genet Metab.* (2007);92(4):351-8.
79. Andersen G, Dalgaard LT, Justesen JM, Anthonsen S, Nielsen T, Thorner LW, et al. The frequent UCP2 -866G>A polymorphism protects against insulin resistance and is associated with obesity: a study of obesity and related metabolic traits among 17 636 Danes. *Int J Obes (Lond).* (2013);37(2):175-81.
80. Krempler F, Esterbauer H, Weitgasser R, Ebenbichler C, Patsch JR, Miller K, et al. A functional polymorphism in the promoter of UCP2 enhances obesity risk but reduces type 2 diabetes risk in obese middle-aged humans. *Diabetes.* (2002);51(11):3331-5.

- 1
2
3 81. Xu L, Chen S, Zhan L. Association of uncoupling protein-2 -866G/A and Ala55Val polymorphisms with
4 susceptibility to type 2 diabetes mellitus: A meta-analysis of case-control studies. *Medicine (Baltimore)*.
5 (2021);100(6):e24464.
6
- 7 82. Yang M, Huang Q, Wu J, Yin JY, Sun H, Liu HL, et al. Effects of UCP2 -866 G/A and ADRB3 Trp64Arg on
8 rosiglitazone response in Chinese patients with Type 2 diabetes. *Br J Clin Pharmacol*. (2009);68(1):14-22.
9
- 10 83. Xu K, Zhang M, Cui D, Fu Y, Qian L, Gu R, et al. UCP2 -866G/A and Ala55Val, and UCP3 -55C/T
11 polymorphisms in association with type 2 diabetes susceptibility: a meta-analysis study. *Diabetologia*.
12 (2011);54(9):2315-24.
13
- 14 84. Willig AL, Casazza KR, Divers J, Bigham AW, Gower BA, Hunter GR, et al. Uncoupling protein 2 Ala55Val
15 polymorphism is associated with a higher acute insulin response to glucose. *Metabolism*. (2009);58(6):877-81.
16
- 17 85. Vimaleswaran KS, Radha V, Ghosh S, Majumder PP, Sathyanarayana Rao MR, Mohan V. Uncoupling protein
18 2 and 3 gene polymorphisms and their association with type 2 diabetes in asian indians. *Diabetes Technol Ther*.
19 (2011);13(1):19-25.
20
- 21 86. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract*. (2014);2014:943162.
22
- 23 87. Ford ES, Li C, Sattar N. Metabolic syndrome and incident diabetes: current state of the evidence. *Diabetes*
24 *Care*. (2008);31(9):1898-904.
25
- 26 88. Ruiz-Ramirez A, Chavez-Salgado M, Peneda-Flores JA, Zapata E, Masso F, El-Hafidi M. High-sucrose diet
27 increases ROS generation, FFA accumulation, UCP2 level, and proton leak in liver mitochondria. *Am J Physiol*
28 *Endocrinol Metab*. (2011);301(6):E1198-207.
29
- 30 89. Castrejon-Tellez V, Rodriguez-Perez JM, Perez-Torres I, Perez-Hernandez N, Cruz-Lagunas A, Guarner-Lans
31 V, et al. The Effect of Resveratrol and Quercetin Treatment on PPAR Mediated Uncoupling Protein (UCP-) 1, 2, and 3
32 Expression in Visceral White Adipose Tissue from Metabolic Syndrome Rats. *Int J Mol Sci*. (2016);17(7).
33
- 34 90. Lu L, Sun X, Chen C, Qin Y, Guo X. Shexiang Baoxin Pill, Derived From the Traditional Chinese Medicine,
35 Provides Protective Roles Against Cardiovascular Diseases. *Front Pharmacol*. (2018);9:1161.
36
- 37 91. Wei D, Zheng N, Zheng L, Wang L, Song L, Sun L. Shexiang Baoxin Pill Corrects Metabolic Disorders in a
38 Rat Model of Metabolic Syndrome by Targeting Mitochondria. *Front Pharmacol*. (2018);9:137.
39
- 40 92. Abbasalizad Farhangi M, Mohseni F, Farajnia S, Jafarabadi MA. Major components of metabolic syndrome
41 and nutritional intakes in different genotype of UCP2 -866G/A gene polymorphisms in patients with NAFLD. *J Transl*
42 *Med*. (2016);14(1):177.
43
- 44 93. Lim KI, Shin YA. Impact of UCP2 polymorphism on long-term exercise-mediated changes in adipocytokines
45 and markers of metabolic syndrome. *Aging Clin Exp Res*. (2014);26(5):491-6.
46
- 47 94. Donadelli M, Dando I, Fiorini C, Palmieri M. UCP2, a mitochondrial protein regulated at multiple levels. *Cell*
48 *Mol Life Sci*. (2014);71(7):1171-90.
49
- 50 95. Crepaldi G, Carruba M, Comaschi M, Del Prato S, Frajese G, Paolisso G. Dipeptidyl peptidase 4 (DPP-4)
51 inhibitors and their role in Type 2 diabetes management. *J Endocrinol Invest*. (2007);30(7):610-4.
52
- 53 96. Lemos NE, Dieter C, Carlessi R, Rheinheimer J, Brondani LA, Leitao CB, et al. Renal effects of exendin-4 in
54 an animal model of brain death. *Mol Biol Rep*. (2019);46(2):2197-207.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

97. Liu L, Liu J, Tian XY, Wong WT, Lau CW, Xu A, et al. Uncoupling protein-2 mediates DPP-4 inhibitor-induced restoration of endothelial function in hypertension through reducing oxidative stress. *Antioxid Redox Signal.* (2014);21(11):1571-81.
98. Anedda A, Rial E, Gonzalez-Barroso MM. Metformin induces oxidative stress in white adipocytes and raises uncoupling protein 2 levels. *J Endocrinol.* (2008);199(1):33-40.
99. Wang Q, Zhang M, Liang B, Shirwany N, Zhu Y, Zou MH. Activation of AMP-activated protein kinase is required for berberine-induced reduction of atherosclerosis in mice: the role of uncoupling protein 2. *PLoS One.* (2011);6(9):e25436.
100. Berger J, Moller DE. The mechanisms of action of PPARs. *Annu Rev Med.* (2002);53:409-35.
101. Olefsky JM, Saltiel AR. PPAR gamma and the treatment of insulin resistance. *Trends Endocrinol Metab.* (2000);11(9):362-8.
102. Gurnell M. Peroxisome proliferator-activated receptor gamma and the regulation of adipocyte function: lessons from human genetic studies. *Best Pract Res Clin Endocrinol Metab.* (2005);19(4):501-23.
103. Staels B, Fruchart JC. Therapeutic roles of peroxisome proliferator-activated receptor agonists. *Diabetes.* (2005);54(8):2460-70.
104. Camirand A, Marie V, Rabelo R, Silva JE. Thiazolidinediones stimulate uncoupling protein-2 expression in cell lines representing white and brown adipose tissues and skeletal muscle. *Endocrinology.* (1998);139(1):428-31.
105. Strobel A, Siquier K, Zilberfarb V, Strosberg AD, Issad T. Effect of thiazolidinediones on expression of UCP2 and adipocyte markers in human PAZ6 adipocytes. *Diabetologia.* (1999);42(5):527-33.
106. Viguierie-Bascands N, Saulnier-Blache JS, Dandine M, Dauzats M, Daviaud D, Langin D. Increase in uncoupling protein-2 mRNA expression by BRL49653 and bromopalmitate in human adipocytes. *Biochem Biophys Res Commun.* (1999);256(1):138-41.
107. Chan SH, Wu CA, Wu KL, Ho YH, Chang AY, Chan JY. Transcriptional upregulation of mitochondrial uncoupling protein 2 protects against oxidative stress-associated neurogenic hypertension. *Circ Res.* (2009);105(9):886-96.
108. Brabazon F, Bermudez S, Shaughness M, Khayrullina G, Byrnes KR. The effects of insulin on the inflammatory activity of BV2 microglia. *PLoS One.* (2018);13(8):e0201878.
109. Ozturk F, Gul M, Esrefoglu M, Ates B. The contradictory effects of nitric oxide in caerulein-induced acute pancreatitis in rats. *Free Radic Res.* (2008);42(4):289-96.
110. Walthers EA, Bradford CS, Moore FL. Cloning, pharmacological characterization and tissue distribution of an ORL1 opioid receptor from an amphibian, the rough-skinned newt *Taricha granulosa*. *J Mol Endocrinol.* (2005);34(1):247-56.
111. Selimovic D, Hassan M, Haikel Y, Hengge UR. Taxol-induced mitochondrial stress in melanoma cells is mediated by activation of c-Jun N-terminal kinase (JNK) and p38 pathways via uncoupling protein 2. *Cell Signal.* (2008);20(2):311-22.
112. Liu D, Ma Z, Di S, Yang Y, Yang J, Xu L, et al. AMPK/PGC1alpha activation by melatonin attenuates acute doxorubicin cardiotoxicity via alleviating mitochondrial oxidative damage and apoptosis. *Free Radic Biol Med.* (2018);129:59-72.

- 1
2
3 113. Pagliaro B, Santolamazza C, Simonelli F, Rubattu S. Phytochemical Compounds and Protection from
4 Cardiovascular Diseases: A State of the Art. *Biomed Res Int.* (2015);2015:918069.
5
6 114. Clarke JD, Riedl K, Bella D, Schwartz SJ, Stevens JF, Ho E. Comparison of isothiocyanate metabolite levels
7 and histone deacetylase activity in human subjects consuming broccoli sprouts or broccoli supplement. *J Agric Food*
8 *Chem.* (2011);59(20):10955-63.
9
10 115. Kim SY, Yoon S, Kwon SM, Park KS, Lee-Kim YC. Kale juice improves coronary artery disease risk factors
11 in hypercholesterolemic men. *Biomed Environ Sci.* (2008);21(2):91-7.
12
13 116. Bhagani H, Nasser SA, Dakroub A, El-Yazbi AF, Eid AA, Kobeissy F, et al. The Mitochondria: A Target of
14 Polyphenols in the Treatment of Diabetic Cardiomyopathy. *Int J Mol Sci.* (2020);21(14).
15
16 117. Diao J, Wei J, Yan R, Fan G, Lin L, Chen M. Effects of resveratrol on regulation on UCP2 and cardiac
17 function in diabetic rats. *J Physiol Biochem.* (2019);75(1):39-51.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60**Table 1.** Pathological implications of UCP2 in animal models

Animal model	UCP2 modulation	Cellular effects	Phenotypes	References
Stroke				
MCAO mouse	downregulation	↑membrane hyperpolarization ↑ $\Delta\Psi$ ↑ROS	Ischemic stroke	[26]
JD-fed SHRSP	downregulation	↑ROS ↑inflammation	Ischemic stroke	[38, 48]
Heart diseases				
I/R rat model	downregulation	↑ROS ↑cell death ↓autophagy	Cardiac ischemia	[34, 61]
Wistar rat running on a treadmill	downregulation	↑extracellular matrix remodeling genes ↑cytoskeletal genes ↑protein synthesis genes ↑inflammatory genes	Compensated cardiac hypertrophy	[62]
Wistar rat with ligation of left anterior coronary artery	upregulation	↑cytoskeleton/extracellular matrix organization genes ↑cell death/apoptosis genes ↑cell growth/proliferation genes ↓ β -oxidation of fatty acids genes	Decompensated cardiac hypertrophy	[62, 63]
ET(A) receptor cardiac-specific knockout mice exposed to cold stress	upregulation	↑oxidative stress ↑apoptosis ↑autophagy	Cold stress-induced cardiac hypertrophy	[64, 65]
Obesity, diabetes, metabolic syndrome				
UCP2 knockout mice	downregulation	↑insulin secretion	Obesity	[23]
UCP2 obese mice	upregulation	β -cell dysfunction ↓insulin secretion	Diabetes	[23]
MS rat model treated with natural compounds	upregulation	↑PPAR- α ↓oxidative stress	↓dyslipidemia ↓blood pressure level ↓insulin resistance ↓obesity risk	[89, 91]

Abbreviations legends: MCAO=middle cerebral artery occlusion; SHRSP=stroke-prone spontaneously hypertensive rat; JD=Japanese diet; ($\Delta\Psi$)=membrane electrical potential; I/R=ischemia/reperfusion; PPAR- α =type α -peroxisome proliferator-activated receptor

Table2. Main *UCP2* SNPs and their pathological implications in human diseases.

SNP	Molecular effects	Clinical phenotypes	References
-866G>A SNP			
A allele	↑UCP2 ↓ROS		[80]
G allele	↓UCP2 ↑ROS		[57]
GG genotype	↑plasmatic hs-CRP	↓T2DM in obesity	[57, 80]
GA genotype	↑ myeloperoxidase level	CAD (reduced post-MI survival in T2DM) ↑Obesity and metabolic abnormalities in NAFLD patients	[69, 70] [92]
AA genotype		Associated with better functional outcome in IS patients after recanalization of proximal MCAO	[60]
	↑ myeloperoxidase level	CAD (reduced post-MI survival in T2DM)	[69, 70]
45 bp Ins/Del			
I/I genotype		Heart rate variability ↑blood pressure	[71]
D/D genotype	↑adiponectin level ↓leptin ↓TNFα ↓IL-6 levels after exercise training	Obesity, MS	[93]
Ala55Val SNP C>T			
C allele carriers		↑Obesity, ↑T2DM	[83]

Abbreviations legend: SNP=single nucleotide polymorphism; hs-CRP= high-sensitivity-CRP, CAD=coronary artery disease; T2DM=type 2 diabetes mellitus; NAFLD= non-alcoholic fatty liver disease; MS= metabolic syndrome

Legends to Figures

Figure 1

Schematic representation of electron transport chain (ETC), mitochondrial enzymatic complexes and UCP2 functions.

During the respiratory process, the energy generated from the oxidation of various substrates including glucose, is conserved in reduced molecules such as NADH (nicotinamide adenine dinucleotide) and FADH₂ (flavin adenine dinucleotide). Electron transporters and protonic pumps create a negative membrane potential and a proton-motive force across the inner mitochondrial membrane which activate the last enzyme of the chain, the ATP synthase. The latter, by the reaction between ADP (adenosine diphosphate) and inorganic phosphate (Pi) generates ATP (adenosine triphosphate). When the energy requirement of the cell is reduced, protons can return into the mitochondrial matrix across the UCP2 protein with reduction of ATP synthesis and heat release. ROS generated during the respiratory process are first reduced at H₂O₂ and then at H₂O from several antioxidant enzymes such as SOD1 (superoxide dismutase 1), SOD2 (superoxide dismutase 2) and GPx (glutathione peroxidase).

Figure 2

UCP2 gene/protein expression modulation in relation to cardiovascular and metabolic diseases.

Schematic representation of the modulation of UCP2 gene and protein expression by known agents. Both nutraceuticals and drugs can upregulate UCP2 level mainly through the AMPK/SIRT-1/PPARs pathway. Of interest, the epigenetic regulation can also be involved, as shown upon BO extract administration (see text). Increased UCP2 levels induce membrane mitochondrial depolarization, decreased ROS production, reduced cell death and inflammatory responses. Consequently, the occurrence of cardiovascular and metabolic diseases is significantly affected.




Figure 1

254x190mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2

254x190mm (300 x 300 DPI)