

# Stress Echocardiography in Italian Echocardiographic Laboratories: A Survey of the Italian Society of Echocardiography and Cardiovascular Imaging

Quirino Ciampi, Mauro Pepi<sup>1</sup>, Francesco Antonini-Canterin<sup>2</sup>, Andrea Barbieri<sup>3</sup>, Agata Barchitta<sup>4</sup>, Giorgio Faganello<sup>5</sup>, Sofia Miceli<sup>6</sup>, Vito Maurizio Parato<sup>7</sup>, Antonio Tota<sup>8</sup>, Giuseppe Trocino<sup>9</sup>, Massimiliana Abbate<sup>10</sup>, Maria Accadia<sup>11</sup>, Rossella Alemanni<sup>12</sup>, Andrea Angelini<sup>13</sup>, Francesco Anglano<sup>14</sup>, Maurizio Anselmi<sup>15</sup>, Iolanda Aquila<sup>16</sup>, Simona Aramu<sup>17</sup>, Enrico Avogadri<sup>18</sup>, Giuseppe Azzaro<sup>19</sup>, Luigi Badano<sup>19</sup>, Anna Balducci<sup>20</sup>, Flavia Ballocca<sup>21</sup>, Alessandro Barbarossa<sup>22</sup>, Giovanni Barbati<sup>23</sup>, Valentina Barletta<sup>24</sup>, Daniele Barone<sup>25</sup>, Francesco Becherini<sup>26</sup>, Giovanni Benfari<sup>27</sup>, Monica Beraldi<sup>28</sup>, Gianluigi Bergandi<sup>29</sup>, Giuseppe Bilardo<sup>30</sup>, Simone Maurizio Binno<sup>31</sup>, Massimo Bolognesi<sup>32</sup>, Stefano Bongiovi<sup>33</sup>, Renato Maria Bragato<sup>34</sup>, Gabriele Braggion<sup>35</sup>, Rossella Brancaleoni<sup>36</sup>, Francesca Bursi<sup>37</sup>, Christian Cadeddu Dessalvi<sup>38</sup>, Matteo Cameli<sup>39</sup>, Antonella Canu<sup>40</sup>, Mariano Capitelli<sup>41</sup>, Anna Clara Maria Capra<sup>42</sup>, Rosa Carbonara<sup>43</sup>, Maria Carbone<sup>44</sup>, Marco Carbonella<sup>45</sup>, Nazario Carrabba<sup>46</sup>, Grazia Casavecchia<sup>47</sup>, Margherita Casula<sup>48</sup>, Elena Chesi<sup>49</sup>, Sebastiano Cicco<sup>50</sup>, Rodolfo Citro<sup>51</sup>, Rosangela Cocchia<sup>52</sup>, Barbara Maria Colombo<sup>53</sup>, Paolo Colonna<sup>54</sup>, Maddalena Conte<sup>54</sup>, Giovanni Corrado<sup>55</sup>, Pietro Cortesi<sup>56</sup>, Lauro Cortigiani<sup>57</sup>, Marco Fabio Costantino<sup>58</sup>, Fabiana Cozza<sup>59</sup>, Umberto Cocchini<sup>60</sup>, Myriam D'Angelo<sup>61</sup>, Santina Da Ros<sup>62</sup>, Fabrizio D'Andrea<sup>63</sup>, Antonello D'Andrea<sup>64</sup>, Francesca D'Auria<sup>65</sup>, Giovanni De Caridi<sup>66</sup>, Stefania De Feo<sup>67</sup>, Giovanni Maria De Matteis<sup>68</sup>, Simona De Vecchi<sup>69</sup>, Carmen Del Giudice<sup>70</sup>, Luca Dell'Angela<sup>71</sup>, Lucrezia Delli Paoli<sup>72</sup>, Ilaria Dentamaro<sup>73</sup>, Paola Destefanis<sup>74</sup>, Gianluca Di Bella<sup>75</sup>, Maria Di Fulvio<sup>76</sup>, Renato Di Gaetano<sup>77</sup>, Giovanna Di Giannuario<sup>78</sup>, Angelo Di Gioia<sup>79</sup>, Luigi Flavio Massimiliano Di Martino<sup>80</sup>, Carmine Di Muro<sup>81</sup>, Concetta Di Nora<sup>82</sup>, Giovanni Di Salvo<sup>83</sup>, Claudio Dodi<sup>84</sup>, Sarah Dogliani<sup>85</sup>, Federica Donati<sup>86</sup>, Melissa Dottori<sup>87</sup>, Giuseppe Epitani<sup>88</sup>, Iacopo Fabiani<sup>89</sup>, Francesca Ferrara<sup>89</sup>, Luigi Ferrara<sup>90</sup>, Stefania Ferrua<sup>91</sup>, Gemma Filice<sup>92</sup>, Maria Fiorino<sup>93</sup>, Davide Forno<sup>94</sup>, Alberto Garini<sup>94</sup>, Gioachino Agostino Giarratana<sup>95</sup>, Giuseppe Gigantino<sup>96</sup>, Mauro Giorgi<sup>97</sup>, Elisa Giubertoni<sup>98</sup>, Cosimo Angelo Greco<sup>99</sup>, Michele Grigolato<sup>100</sup>, Walter Grosso Marra<sup>29</sup>, Anna Holzi<sup>101</sup>, Alessandra Iaiza<sup>102</sup>, Andrea Iannaccone<sup>103</sup>, Federica Ilardi<sup>104</sup>, Egidio Imbalzano<sup>105</sup>, Riccardo M. Inciardi<sup>106</sup>, Corinna Antonia Inerra<sup>107</sup>, Emilio Iori<sup>108</sup>, Annibale Izzo<sup>109</sup>, Giuseppe La Rosa<sup>110</sup>, Graziana Labanti<sup>111</sup>, Alberto Maria Lanzone<sup>112</sup>, Laura Lanzoni<sup>113</sup>, Ornella Lapetina<sup>114</sup>, Elisa Leiballi<sup>115</sup>, Mariateresa Librera<sup>116</sup>, Carmenita Lo Conte<sup>117</sup>, Maria Lo Monaco<sup>118</sup>, Antonella Lombardo<sup>119</sup>, Michelangelo Luciani<sup>120</sup>, Paola Lusardi<sup>121</sup>, Antonio Magnante<sup>122</sup>, Alessandro Malagoli<sup>123</sup>, Gelsomina Malatesta<sup>124</sup>, Costantino Mancusi<sup>125</sup>, Maria Teresa Manes<sup>126</sup>, Fiore Manganelli<sup>127</sup>, Francesca Mantovani<sup>128</sup>, Vincenzo Manuppelli<sup>129</sup>, Valeria Marchese<sup>129</sup>, Lina Marinacci<sup>130</sup>, Roberto Mattioli<sup>131</sup>, Civelli Maurizio<sup>132</sup>, Giuseppe Antonio Mazza<sup>133</sup>, Stefano Mazza<sup>134</sup>, Marco Melis<sup>135</sup>, Giulia Meloni<sup>136</sup>, Elisa Merli<sup>137</sup>, Alberto Milan<sup>138</sup>, Giovanni Minardi<sup>139</sup>, Antonella Monaco<sup>140</sup>, Ines Monte<sup>141</sup>, Graziano Montresor<sup>142</sup>, Antonella Moreo<sup>143</sup>, Fabio Mori<sup>144</sup>, Sofia Morini<sup>145</sup>, Claudio Moro<sup>146</sup>, Doralisa Morrone<sup>147</sup>, Francesco Negri<sup>82</sup>, Carmelo Nipote<sup>148</sup>, Fulvio Nisi<sup>149</sup>, Silvio Nocco<sup>150</sup>, Luigi Novello<sup>151</sup>, Luigi Nunziata<sup>152</sup>, Alessandro Paoletti Perini<sup>153</sup>, Antonello Parodi<sup>154</sup>, Emilio Maria Pasanisi<sup>155</sup>, Guido Pastorini<sup>156</sup>, Rita Pevrasini<sup>157</sup>, Daisy Pavoni<sup>162</sup>, Chiara Pedone<sup>158</sup>, Francesco Pelliccia<sup>159</sup>, Giovanni Pellicciari<sup>160</sup>, Elisa Pelloni<sup>161</sup>, Valeria Pergola<sup>162</sup>, Giovanni Perillo<sup>163</sup>, Enrica Petruccielli<sup>164</sup>, Chiara Pezzullo<sup>165</sup>, Gerardo Piacentini<sup>166</sup>, Elisa Picardi<sup>167</sup>, Giovanni Pinna<sup>168</sup>, Massimiliano Pizzarelli<sup>169</sup>, Alfredo Pizzutti<sup>170</sup>, Matteo Maria Poggi<sup>171</sup>, Alfredo Posteraro<sup>172</sup>, Carmen Privitera<sup>173</sup>, Debora Rampazzo<sup>174</sup>, Carlo Ratti<sup>175</sup>, Sara Rettegno<sup>176</sup>, Fabrizio Ricci<sup>177</sup>, Caterina Ricci<sup>178</sup>, Cristina Rolando<sup>179</sup>, Stefania Rossi<sup>180</sup>, Chiara Rovera<sup>167</sup>, Roberta Ruggieri<sup>181</sup>, Maria Giovanna Russo<sup>182</sup>, Nicola Sacchi<sup>183</sup>, Antonino Saladino<sup>184</sup>, Francesca Sani<sup>185</sup>, Chiara Sartori<sup>186</sup>, Virginia Scarabeo<sup>187</sup>, Angela Sciacqua<sup>8</sup>, Antonio Scillone<sup>188</sup>, Pasquale Antonio Scopelliti<sup>189</sup>, Alfredo Scorza<sup>190</sup>, Angela Scozzafava<sup>191</sup>, Francesco Serafini<sup>192</sup>, Walter Serra<sup>193</sup>, Sergio Severino<sup>194</sup>, Beatrice Simeone<sup>195</sup>, Domenico Sirico<sup>83</sup>, Marco Solari<sup>196</sup>, Gian Luca Spadaro<sup>23</sup>, Laura Stefani<sup>197</sup>, Antonio Strangio<sup>198</sup>, Francesca Chiara Surace<sup>199</sup>, Gloria Tamborini<sup>1</sup>, Nicola Tarquinio<sup>200</sup>, Eliezer Joseph Tassone<sup>201</sup>, Isabella Tavarozzi<sup>202</sup>, Bertrand Tchana<sup>203</sup>, Giuseppe Tedesco<sup>204</sup>, Monica Tinto<sup>205</sup>, Daniela Torzillo<sup>206</sup>, Antonio Totaro<sup>207,208</sup>, Oreste Fabio Triolo<sup>209</sup>, Federica Troisi<sup>73</sup>, Maurizio Tusa<sup>210</sup>, Federico Vancheri<sup>211</sup>, Vincenzo Varasano<sup>212</sup>, Amedeo Venezia<sup>213</sup>, Anna Chiara Vermi<sup>214</sup>, Bruno Villari, Giordano Zampi<sup>120</sup>, Jessica Zannoni<sup>210</sup>, Concetta Zito<sup>75</sup>, Antonello Zugaro<sup>215</sup>, Eugenio Picano<sup>216</sup>, Scipione Careri<sup>75</sup>

Cardiology Division, Fatebenefratelli Hospital, Benevento, <sup>1</sup>Cardiology Division, Centro Cardiologico Monzino, IRCCS, <sup>37</sup>Department of Health Sciences, Cardiology Division, University of Milan, San Paolo Hospital, ASST Santi Paolo e Carlo, <sup>132</sup>Cardiology Division, European Institute of Oncology, <sup>143</sup>De Gasperis Cardio Center, ASST Grande Ospedale Metropolitano Niguarda, Milano, <sup>2</sup>Department of Rehabilitative Cardiology, Rehabilitative Hospital High Speciality, Motta di Livenza, TV, <sup>3</sup>Department of Biomedical, Metabolic and Neural Sciences, Cardiology Division, University of Modena and Reggio Emilia, Policlinico di Modena, <sup>8</sup>Pascia Center, Polyclinic, <sup>89</sup>Internal Medicine Division, University Hospital Modena Polyclinic, <sup>123</sup>Division of Cardiology, Nephro-Cardiovascular Department, Baggiovara Hospital, University of Modena and Reggio Emilia, Modena, <sup>4</sup>Semi-Intensive Care Department, Padova University Hospital, <sup>83</sup>Pediatric Cardiology and Congenital Heart Disease Division, Padova University Hospital, <sup>162</sup>Cardiology Division, Padova University Hospital, Padova, <sup>5</sup>Cardiovascular Center, Maggiore Hospital, Trieste, <sup>6</sup>Geriatric Division, University Hospital Renato Dulbecco, <sup>16</sup>Cardiology Division, University Hospital Mater Domini, <sup>191</sup>Cardiology Division, Pugliese Hospital, Catanzaro, <sup>7</sup>Cardiology Division, Madonna del Soccorso Hospital, San Benedetto del Tronto, AP, <sup>8</sup>Cardiology Division, Polyclinic Hospital, <sup>43</sup>Cardiology Division, Maugeri Institute IRCCS, <sup>50</sup>Department of Precision and Regenerative Medicine and Ionian Area, Unit of Internal Medicine "G. Baccelli" and Unit of Hypertension "A.M. Pirrelli", University of Bari Aldo Moro Medical School, AUOC Policlinico di

**Address for correspondence:** Prof. Quirino Ciampi, Division of Cardiology, Fatebenefratelli Hospital, Viale Principe di Napoli, 12, I-82100, Benevento, Italy. E-mail: qciampi@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

Submitted: 05-Sep-2023 Accepted: 07-Sep-2023 Published: 20-Nov-2023

**How to cite this article:** Ciampi Q, Pepi M, Antonini-Canterin F, Barbieri A, Barchitta A, Faganello G, *et al*. Stress echocardiography in Italian echocardiographic laboratories: A survey of the Italian society of echocardiography and cardiovascular imaging. *J Cardiovasc Echography* 2023;33:125-32.

## Access this article online

### Quick Response Code:



Website:  
<https://journals.lww.com/JCEG>

DOI:  
10.4103/jcecho.jcecho\_48\_23

Bari, <sup>181</sup>Cardiology Division, Di Venere Hospital, Bari, <sup>9</sup>Non Invasive Cardiac Imaging Department, Fondazione IRCCS San Gerardo dei Tintori, Monza, <sup>10</sup>Cardiology Vanvitelli Division, AORN dei Colli, Monaldi Hospital, <sup>52</sup>Rehabilitative Cardiology, Cardarelli Hospital, <sup>70</sup>Cardiology Division, AORN dei Colli, Monaldi Hospital, <sup>104</sup>Cardiology Division, Federico II University Hospital, <sup>116</sup>Cardiology Division, Mediterranea Clinic, <sup>125</sup>Hypertension Center, Federico II University Hospital, <sup>182</sup>Pediatric Cardiology Division, AORN dei Colli, Monaldi Hospital, <sup>194</sup>Cardiology Division, Cotugno Hospital, Napoli, <sup>11</sup>Cardiology Division, Del Mare Hospital, Ponticelli, NA, <sup>12</sup>Cardiac Surgery Division, Casa Sollievo Della Sofferenza Hospital, San Giovanni Rotondo, <sup>13</sup>Cardiology Division, Cardinal Massaia Hospital, Asti, <sup>14</sup>Cardiology Division, Ravenna Medical Center, Ravenna, <sup>15</sup>Cardiology Division, Fracastoro Hospital, San Bonifacio, VR, <sup>17</sup>Cardiology Division, San Martino Hospital, Oristano, <sup>19</sup>Department of Medicine and Surgery, University Milano-Bicocca, Integrated Cardiovascular Diagnosi Unit, Istituto Auxologico Italiano, IRCCS, <sup>131</sup>Cardiology Division, IRCCS Multimedia Hospital, Sesto San Giovanni, <sup>34</sup>Echocardiography and Emergency Cardiovascular Care Division, Humanitas Clinical and Research Centre, Rozzano, <sup>107</sup>Cardiology Division, ASST-OVEST Milanese, Ospedale di Legnano, Legnano, <sup>149</sup>Anesthesia and Intensive Care Division, IRCCS Humanitas Research Hospital, Rozzano, MI, <sup>206</sup>Internal Medicine Division, L. Sacco Hospital, University of Milan, <sup>210</sup>Cardiology Division, St. Donato Polyclinic, San Donato Milanese, Milan, <sup>18</sup>Department of Rehabilitative Cardiology, SS Trinità Hospital, Fossano, CN, <sup>20</sup>Pediatric Cardiology Division, Policlinico S. Orsola-Malpighi IRCCS Hospital, <sup>111</sup>Cardiology Division, Bellaria Hospital, <sup>158</sup>Cardiology Division, Maggiore Hospital, Bologna, <sup>21</sup>Cardiology Division, Maria Vittoria Hospital, <sup>97</sup>Cardiology Division, Molinette Hospital - Città della Salute e della Scienza, <sup>103</sup>Internal Medicine Division, Ordine Mauriziano Hospital, <sup>121</sup>Cardiology and Cardiac Surgery Division, Maria Pia Hospital, <sup>133</sup>Pediatric Cardiology Division, Regina Margherita Hospital - Città della Salute e della Scienza, <sup>138</sup>Internal Medicine 4 Division, Molinette Hospital - Città della Salute e della Scienza, <sup>170</sup>Cardiology Outpatient Clinic, Koelliker Hospital, Torino, <sup>22</sup>Clinic of Cardiology and Arrhythmology, Marche University Hospital, <sup>87</sup>Cardiology Division, Marche University Hospital, <sup>124</sup>Cardiology Division, IRCCS INRCA Hospital, <sup>199</sup>Pediatric Cardiac Surgery and Cardiology Division, Marche University Hospital, Ancona, <sup>23</sup>Cardiology Division, St. Bortolo Hospital, Vicenza, <sup>25</sup>Cardiology Division, S. Andrea Hospital, La Spezia, <sup>24</sup>Cardiology 2 Division, Cardiac Vascular Thoracic Department, Pisa University Hospital, <sup>26</sup>Cardiology and Cardiovascular Medicine Division, Fondazione Toscana Gabriele Monasterio, <sup>147</sup>Cardiology Division, Cisanello University Hospital, <sup>216</sup>CNR, Institute of Clinical Physiology, Biomedicine Department, Pisa, <sup>27</sup>Cardiology Division, University of Verona, <sup>113</sup>Cardiology Division, Sacro Cuore Don Calabria IRCCS Hospital, Verona, <sup>28</sup>Cardiology Division, ASST Mantova, Mantova, <sup>29</sup>Cardiology Division, Civil Hospital, Ivrea, TO, <sup>30</sup>Cardiology Division, Civil Hospital Fetre, Feltre, BL, <sup>31</sup>Cardiology Division, Guglielmo da Saliceto Hospital, <sup>84</sup>Cardiology Division, San Antonino Clinic, Piacenza, <sup>32</sup>Center for Internal Medicine and Sports Cardiology, Local Health Unit of Romagna, Cesena, FC, <sup>33</sup>Cardiology Division, Immacolata Concezione Civil Hospital, Piove di Sacco, PD, <sup>35</sup>Cardiology Division, Santa Maria Regina degli Angeli Hospital, Adria, RO, <sup>36</sup>Cardiology Division, A. Costa Civil Hospital, Porretta Terme, BO, <sup>38</sup>Cardiology Division, University Hospital of Cagliari, <sup>139</sup>Cardiology Division, Brotzu Hospital, Cagliari, <sup>40</sup>Cardiology Division, Santissima Annunziata Hospital, <sup>39</sup>Cardiology Division, Polyclinic Le Scotte Hospital, Siena, <sup>136</sup>Center for Prevention, Diagnosis and Therapy of Arterial Hypertension and Cardiovascular Complications, St. Camillo Hospital, Sassari, <sup>41</sup>Internal Medicine Division, Pavullo Hospital, Pavullo nel Frignano, MO, <sup>42</sup>Cardiological Diagnostics Division, Synlab San Nicolò Diagnostic Center, Lecco, <sup>44</sup>Emergency Medicine Division, St. Anna and St. Sebastiano Hospital, <sup>108</sup>Cardiology Division, St. Anna and St. Sebastiano Hospital, Caserta, <sup>46</sup>Cardiology Division, SS Maria Addolorata Hospital, Eboli, SA, <sup>46</sup>Cardiology Division, Careggi University Hospital, <sup>144</sup>Non-invasive Cardiovascular Diagnostic Division, Careggi University Hospital, <sup>153</sup>Cardiology Division, St. Maria Nuova Hospital, <sup>171</sup>Interdisciplinary Internal Medicine Division, Careggi University Hospital, <sup>185</sup>Cardiology Division, St. Giovanni di Dio Hospital, <sup>197</sup>Sports Medicine Division, Careggi University Hospital, Firenze, <sup>47</sup>Cardiology Division, University Hospital Ospedali Riuniti, Foggia, <sup>48</sup>Cardiology Division, Nostra Signora di Bonaria Hospital, San Gavino Monreale, SU, <sup>49</sup>Neonatology Division, S. Maria Nuova Hospital, <sup>128</sup>Cardiology Division, Azienda USL- IRCCS di Reggio Emilia, Reggio Emilia, <sup>51</sup>Echocardiography Division, University Hospital San Giovanni di Dio e Ruggi d'Aragona, <sup>65</sup>Vascular - Endovascular Surgery Division, University Hospital San Giovanni di Dio e Ruggi d'Aragona, <sup>96</sup>Cardiology Division, University Hospital San Giovanni di Dio e Ruggi d'Aragona, Salerno, <sup>53</sup>Clinic of Emergency Medicine, IRCCS San Martino Polyclinic Hospital, Genoa, <sup>54</sup>Department of Translational Medical Sciences, University of Naples Federico II, Naples, <sup>55</sup>Cardiology Division, Valduce Hospital, Como, <sup>56</sup>Cardioncology Division, IRCCS Istituto Romagnolo per lo Studio dei Tumori "Dino Amadori", Meldola, FC, <sup>57</sup>Cardiology Division, San Luca Hospital, Lucca, <sup>58</sup>Cardiology Division, San Carlo Hospital, Potenza, <sup>59</sup>Cardiology Division, Poliambulanza Foundation Hospital, <sup>100</sup>Polycardiography Division, Civil Hospital, <sup>106</sup>Cardiology Division, Civil Hospital, Brescia, <sup>60</sup>Cardiology Division, San Bassiano Hospital, Bassano Del Grappa, VI, <sup>61</sup>Cardiology Division, Bonino Pulejo IRCCS Hospital, <sup>66</sup>Vascular Surgery Division, University Hospital Polyclinic G. Martino, University of Messina, <sup>105</sup>Internal Medicine Division, University Hospital Polyclinic G. Martino, University of Messina, <sup>75</sup>Cardiology Division, University Hospital Polyclinic G. Martino, University of Messina, <sup>68</sup>Division of Cardiology, Riuniti Padova Sud Hospital, Monselice, PD, <sup>63</sup>Cardiology Division, St. Andrea Hospital, <sup>68</sup>Cardiology Division, Sandro Pertini Hospital, <sup>102</sup>Cardiac Surgery Division, San Camillo-Fornalini Hospital, <sup>119</sup>Cardiology Division, Fondazione Policlinico A. Gemelli-IRCCS, Università Cattolica, <sup>139</sup>Echolab, Salvator Mundi International Hospital, <sup>159</sup>Cardiology Division, Umberto I Hospital, <sup>163</sup>Cardiology Division, Celio Military Polyclinic, <sup>166</sup>Fetal and Neonatal Cardiology Unit - Fatebenefratelli Isola Tiberina Gemelli Isola Hospital, <sup>168</sup>Neonatology and Neonatal Intensive Care Division, San Camillo-Fornalini Hospital, Roma, <sup>64</sup>Cardiology Division, Umberto I Hospital, Nocera Inferiore, SA, <sup>67</sup>Cardiology Division, P. Pederzoli Hospital, Peschiera del Garda, VR, <sup>69</sup>Cardiology Division, Major University Hospital of Charity, Novara, <sup>71</sup>Cardiology Division, Gorizia-Monfalcone Hospital, Gorizia, <sup>72</sup>San Michele Clinic, Cardiological Intensive Care Unit, Maddaloni, CE, <sup>73</sup>Cardiology Division, Miulli Hospital, Acquaviva delle Fonti, BA, <sup>74</sup>Cardiology Division, San Luigi Gonzaga University Hospital, Orbassano, <sup>76</sup>Cardiology-ICCU Division, Ss. Annunziata Hospital, <sup>177</sup>Cardiology Division, Ss. Annunziata Hospital, Chieti, <sup>77</sup>Cardiology Division, Bolzano Hospital, Bolzano, <sup>78</sup>Cardiology Division, Infermi Hospital, Rimini, <sup>79</sup>Cardiology Division, St. Giuliano Hospital, Gugliano in Campania, NA, <sup>80</sup>Cardiology Division, Santa Maria degli Angeli Hospital, Putignano, BA, <sup>81</sup>Sports Medicine Division, Livorno Hospital, <sup>155</sup>Cardiology Division, Livorno Hospital, Livorno, <sup>82</sup>Cardiology Division, Azienda Sanitaria Universitaria Friuli Centrale, Udine, <sup>86</sup>Cardiology Division, Ss. Annunziata Civil Hospital, Savignano, <sup>156</sup>Cardiology Division, Regina Montis Regalis Hospital, Mondovì, CN, <sup>89</sup>Internal Medicine Division, Camberlingo Hospital, Francavilla Fontana, BR, <sup>90</sup>Cardiology Division, Villa Dei Fiori Clinic, Acerra, <sup>152</sup>Cardiology Division, St. Maria della Pietà Hospital, Nola, NA, <sup>91</sup>Cardiology Division, Infermi Hospital, Rivoli, <sup>167</sup>Cardiology Division, Civic Hospital, Chivasso, <sup>176</sup>Cardiology Division, Hospital, Moncalieri, <sup>179</sup>Cardiology Division, Civil Hospital, Ciriè, TO, <sup>92</sup>Cardiology Division, Annunziata Hospital, <sup>188</sup>Intensive Cardiac Rehabilitation Unit, Villa del Sole Clinic, Cosenza, <sup>93</sup>Cardiology Division, ARNAS Civico Hospital, <sup>95</sup>Cardiac Surgery Division, Policlinico P. Giaccone Hospital, <sup>209</sup>Cardiology Division, Policlinico P. Giaccone Hospital, <sup>94</sup>Cardiology Division, Cremona Hospital, Cremona, <sup>98</sup>Cardiology Division, Civil Hospital, Guastalla, RE, <sup>99</sup>Cardiology Division, Veris Delli Ponti Hospital, Scorrano, LE, <sup>101</sup>Internal Medicine Division, Quisisana Clinic, <sup>157</sup>Cardiology Division, University Hospital of Ferrara, <sup>169</sup>Cardiology Outpatient Clinic, Quisisana Clinic, Ferrara, <sup>108</sup>Cardiology Division, New Civil Hospital, Sassuolo, <sup>175</sup>Cardiology Division, St. Maria Bianca Hospital, Mirandola, <sup>178</sup>Cardiology Outpatient Clinic, Casa della Salute "Regina Margherita", Castelfranco Emilia, MO, <sup>110</sup>Cardiology Division, St. Barbara Hospital, Gela, CL, <sup>112</sup>Cardiology Division, San Rocco Clinical Institute, Ome, <sup>142</sup>Cardiology Division, Civil Hospital, Gavardo, BS, <sup>114</sup>Cardiology Division, San Carlo Hospital, Melfi, PZ, <sup>115</sup>Cardiology and Rehabilitative Division, Azienda Sanitaria Friuli Occidentale (ASFO), Health Care, Sacile (Pd), <sup>117</sup>Cardiology Division, St. Ottone Frangipane Hospital, Ariano Irpino, AV, <sup>118</sup>Cardiology Division, Humanitas Gavazzeni Hospital, Bergamo, <sup>120</sup>Cardiology Division, Belcolle Hospital, Viterbo, <sup>122</sup>Cardiology Division, Madonna delle Grazie Hospital, Matera, <sup>126</sup>Cardiology Division, St. Francesco Hospital, Paola, CS, <sup>127</sup>Cardiology Division, St. Giuseppe Moscati Hospital, Avellino, <sup>129</sup>Cardiology Division, St. Maria della Speranza Hospital, Battipaglia, SA, <sup>130</sup>Cardiology Division, Civil Hospital, Città di Castello, <sup>183</sup>Medical Division, St. Agostino Hospital, Castiglione del Lago, PG, <sup>134</sup>Cardiology Division, Maggiore St. Andrea Hospital, Vercelli, <sup>137</sup>Cardiology Division, Degli Infermi Hospital, Faenza, RA, <sup>140</sup>Cardiology Outpatient Clinic, Cardiology Outpatient Clinic, Civitanova Marche, MC, <sup>141</sup>Cardiology Division, University Hospital Polyclinic "G. Rodolico-S. Marco", University of Catania, Catania, <sup>145</sup>Cardiology Division, Riuniti della Valdichiana Hospital, Montepulciano, SI, <sup>146</sup>Cardiology Division, Pio XI Hospital, Desio, MB, <sup>148</sup>Cardiology Division, Civil Hospital, Sant'Agata di Militello, ME, <sup>150</sup>Cardiology Division, Sirai Hospital, Carbonia, CI, <sup>151</sup>Geriatric Division, Valdagno Hospital, Arzignano, VI, <sup>154</sup>Cardiology Division, Padre Antero Micone Hospital, Genova, <sup>160</sup>Internal Medicine Division, Gruppioni Clinic, Pianoro, BO, <sup>161</sup>Cardiology Division, Parini Hospital, Aosta, <sup>164</sup>Cardiology Division, S. Giacomo Hospital, Monopoli, <sup>204</sup>Cardiology Division, Civil Hospital, Bitonto, <sup>213</sup>Geriatric Division, Miulli Hospital, Acquaviva delle Fonti, BA, <sup>165</sup>Cardiology Division, G.B. Grassi Hospital, Lido di Ostia, <sup>172</sup>Cardiology Division, St. Giovanni Evangelista Hospital, Tivoli, <sup>190</sup>Cardiology Division, Riuniti Anzio-Nettuno Hospital, Anzio, RM, <sup>173</sup>Pediatric Division, St. Chiara Hospital, Trento, <sup>174</sup>Cardiology Division, Madonna della Navicella Hospital, Chioggia, <sup>192</sup>Medical

Division, Dell'Angelo Hospital, Mestre, VE, <sup>180</sup>Cardiology Division, Civil Hospital, Lavagna, GE, <sup>184</sup>Cardiology Division, Giovanni Paolo II Hospital, Sciacca, AG, <sup>186</sup>Cardiology Division, Santi Antonio e Biagio e Cesare Arrigo Hospital, Alessandria, <sup>187</sup>Cardiology Division, Camposampiero Hospital, Camposampiero, PD, <sup>189</sup>Cardiology Division, Pesenti Fenaroli Hospital, Alzano Lombardo, BG, <sup>193</sup>Cardiology Division, University Hospital, <sup>203</sup>Pediatric Cardiology Division, University Hospital, Parma, <sup>195</sup>Cardiology Division, ICOT Marco Pasquali Clinic, Latina, <sup>196</sup>Cardiology Division, St. Giuseppe Hospital, Empoli, FI, <sup>198</sup>Cardiology Division, St. Giovanni di Dio Hospital, Crotone, <sup>200</sup>Internal Medicine Division, IRCCS INRCA Hospital, Osimo AN, <sup>201</sup>Echocardiography and Ergometry Laboratory, Medicare, Lamezia Terme, CZ, <sup>202</sup>Cardiology Division, Ferdinando Venezia Hospital, Isernia, <sup>205</sup>Cardiology Division, Mater Salutaris Hospital, Legnago, VR, <sup>207</sup>Department of Cardiovascular Sciences, Responsible Research Hospital, <sup>208</sup>Department of Medicine and Health Sciences "V. Tiberio", University of Molise, Campobasso, <sup>211</sup>Medical Division, St. Elia Hospital, Caltanissetta, <sup>212</sup>Internal and Emergency Medicine Division, Civil Hospital, Policoro MT, <sup>214</sup>Cardiology Division, Civil Hospital, Castel San Giovanni, PC, <sup>215</sup>Intensive Care Unit, St. Salvatore Hospital, L'Aquila, Italy

## Abstract

**Background:** The Italian Society of Echography and Cardiovascular Imaging (SIECVI) conducted a national survey to understand the volumes of activity, modalities and stressors used during stress echocardiography (SE) in Italy. **Methods:** We analyzed echocardiography laboratory activities over a month (November 2022). Data were retrieved through an electronic survey based on a structured questionnaire, uploaded on the SIECVI website. **Results:** Data were obtained from 228 echocardiographic laboratories, and SE examinations were performed in 179 centers (80.6%); 87 centers (47.5%) were in the northern regions of Italy, 33 centers (18.4%) were in the central regions, and 61 (34.1%) in the southern regions. We annotated a total of 4057 SE. We divided the SE centers into three groups, according to the numbers of SE performed: <10 SE (low-volume activity, 40 centers), between 10 and 39 SE (moderate volume activity, 102 centers) and  $\geq 40$  SE (high volume activity, 37 centers). Dipyridamole was used in 139 centers (77.6%); exercise in 120 centers (67.0%); dobutamine in 153 centers (85.4%); pacing in 37 centers (21.1%); and adenosine in 7 centers (4.0%). We found a significant difference between the stressors used and volume of activity of the centers, with a progressive increase in the prevalence of number of stressors from low to high volume activity ( $P = 0.033$ ). The traditional evaluation of regional wall motion of the left ventricle was performed in all centers, with combined assessment of coronary flow velocity reserve (CFVR) in 90 centers (50.3%); there was a significant difference in the centers with different volume of SE activity: the incidence of analysis of CFVR was significantly higher in high volume centers compared to low – moderate – volume (32.5%, 41.0% and 73.0%, respectively,  $P < 0.001$ ). The lung ultrasound (LUS) was assessed in 67 centers (37.4%). Furthermore for LUS, we found a significant difference in the centers with different volume of SE activity: significantly higher in high volume centers compared to low – moderate – volume (25.0%, 35.3% and 56.8%, respectively,  $P < 0.001$ ). **Conclusions:** This nationwide survey demonstrated that SE was significantly widespread and practiced throughout Italy. In addition to the traditional indication to coronary artery disease based on regional wall motion analysis, other indications are emerging with an increase in the use of LUS and CFVR, especially in high-volume centers.

**Keywords:** Coronary flow velocity reserve, lung ultrasound, stress echocardiography

## INTRODUCTION

Stress echocardiography (SE) is an efficient and cost-effective option for diagnosing and stratifying the risk of ischemic heart disease (coronary artery disease [CAD]).<sup>[1,2]</sup> Over time, SE has evolved beyond its traditional evaluation of regional wall motion analysis to encompass a broader scope of functional testing, including conditions such as valvular heart disease and cardiomyopathy.

The SE 2020 and SE 2030 multicenter studies,<sup>[3,4]</sup> endorsed by the Italian Society of Echography and Cardiovascular Imaging (SIECVI), have played a significant role in demonstrating the effectiveness of this new approach. These studies have validated emerging signs and integrated new information with established knowledge, leading to the standardization of procedures and the adoption of additional imaging evaluations beyond regional wall motion analysis. These new imaging evaluations include coronary flow velocity reserve (CFVR) on left anterior descending coronary artery (LAD)<sup>[5-7]</sup> and lung ultrasound (LUS),<sup>[8-12]</sup> making SE a more comprehensive and valuable tool in CAD and the diagnosis and assessment of various cardiac conditions including valvular heart disease and cardiomyopathy.<sup>[12-14]</sup>

With the advancements in SE and the expansion of its applications, the aim of this survey is to understand the current

volumes of activity, modalities, and stressors used during SE in Italy. Analyzing the new indications for SE both within and beyond CAD, as well as the implementation of the new imaging approach, can help in building next-generation SE labs and improving patient care.

## METHODS

We analyzed the activity of echocardiography laboratories in 1 month. November 2022 was chosen as an ideal reference month (30 days; away from holidays).

A list of accredited echocardiographic laboratories was obtained from SIECVI. Each member of SIECVI was contacted by mail. Data were retrieved through an electronic survey based on a structured questionnaire uploaded on the SIECVI website ([www.siec.it](http://www.siec.it)).

The methods of the survey were previously described in other survey of the SIECVI.<sup>[15-17]</sup>

For the allocation of the response, the questionnaire required general information, such as the name of the hospital, the investigator, and the interviewed person's name:

1. General information: date, hospital's name, department, name of the interviewed physician, city, and region of Italy



2. The number of SE performed
3. The stressors used
4. The number of SE with CFVR and LUS performed
5. The principal indications of SE.

### Statistical analysis

The categorical data are expressed in terms of the number of subjects and percentage, whereas continuous data are expressed as mean ± standard deviation or median (minimum-maximum) depending on the variables' distribution. For continuous variables, intergroup differences were tested with a one-way analysis of variance and inter-group comparison by Bonferroni or Kruskal-Wallis, followed by the Mann-Whitney test as appropriate. The Chi-square test or Fisher exact test was used to compare the distribution of categorical variables among groups.

All statistical calculations were performed using the SPSS for Windows, release 20.0 (Chicago, Illinois, USA).

## RESULTS

Data were obtained from 228 echocardiographic laboratories, and SE examinations were performed in 179 centers (80.6%): 87 centers (47.5%) were in the northern regions of Italy, 33 centers (18.4%) were in the central regions, and 61 (34.1%) in the southern regions.

During the month of observation, we annotated a total of 4057 SE.

We divided the SE centers in three groups, according to the numbers of SE performed: <10 SE (low-volume activity, 40 centers, 22%), between 10 and 39 SE (moderate volume activity, 102 centers, 57%) and ≥40 SE (high volume activity, 37 centers, 21%). The principal data of number and indication of SE are reported in the Table 1.

Dipyridamole was used in 139 centers (77.6%), dobutamine in 153 centers (85.4%); pacing in 37 centers (21.1%) and adenosine in 7 centers (4.0%) [Figure 1].

Exercise was performed in 120 centers (67.0%): treadmill in 20 centers (16.7%), semisupine bike in 82 centers (68.3%) and with both modality in 18 centers (15.0%) [Figure 1].




Figure 1: Percentage of stressors used in stress echocardiography

A single stressor was used in 29 centers (16.2%): only dipyridamole in 3 centers, only dobutamine in 9 centers and only exercise in 17 centers. Two stressor was used in 45 centers (25.1%) and 3 or more stressors in 105 centers (58.7%).

We found a significant difference between the stressors used and volume of activity of the centers, with a progressive increase in the prevalence of number of stressors from low-to-high volume activity [Figure 2].

The traditional evaluation of regional wall motion of the left ventricle was performed in all centers, with combined assessment of CFVR in 90 centers (50.3%). There was a significant difference in the centers with different volume of SE activity: the incidence of analysis of CFVR was significantly higher in high volume centers compared to low-to-moderate volume [Figure 3]. CFVR was evaluated routinely during traditional SE in 70 centers (77.8%), in patients with known CAD and previous coronary revascularization in 80 centers (88.9%) and in the evaluation of intermediate stenosis in 81 centers (90%).

The LUS was assessed in 67 centers (37.4%). We found also for LUS use a significant difference in the centers with different volume of SE activity: significantly higher in high volume centers compared to low-to-moderate volume [Figure 4]. The principal indications of LUS evaluation during SE were: heart failure with reduced ejection fraction in 61 centers (91.0%), heart failure with preserved ejection fraction in 59 centers (88.0%), routine evaluation in 51 centers (73.9%), and in post-COVID in 46 centers (68.7%) [Figure 5].

We found a significant difference between the three groups studied in the indications for aortic and mitral valve disease, with more frequent indication in high-volume center [Table 1].

## DISCUSSION

We described the use of SE in Italian echocardiographic laboratories. SE is a widely used diagnostic tool that combines echocardiography with physical or pharmacological SE, by




Figure 2: Numbers of stressors used during stress echocardiography in low, moderate, and high volume centers

**Table 1: Stress echocardiography and principal indications in the study groups**

	Overall (n=179)	Low-volume (n=40)	Moderate-volume (n=102)	High-volume (n=37)	P
SE (n)	22.7±24.3	4.6±2.2	15.8±6.7	61.0±28.0	<0.001
SE and CFVR evaluation (n)	8.9±17.2	1.3±2.1	5.3±7.0	26.8±29.9	<0.001
SE and CFVR evaluation, n (%)	90 (50.3)	13 (32.5)	57 (41.0)	27 (73.0)	0.001
SE and B-Lines evaluation (n)	6.0±13.8	1.5±3.2	3.6±7.2	17.6±24.7	<0.001
SE and CFVR evaluation, n (%)	67 (37.4)	10 (25.0)	36 (35.3)	21 (56.8)	0.013
Regional distribution in Italy, n (%)					
Northern	85 (47.5)	16 (40.4)	48 (47.1)	21 (56.8)	0.510
Center	33 (18.4)	10 (25.0)	19 (18.6)	4 (10.8)	
Southern	61 (34.1)	14 (35.0)	35 (34.3)	12 (32.4)	
Use of contrast, n (%)					
Never	83 (46.4)	21 (52.5)	46 (45)	16 (43.2)	0.611
Rare	55 (30.7)	13 (32.5)	31 (30.4)	11 (29.7)	
Frequent	26 (14.6)	5 (12.5)	15 (14.7)	6 (16.2)	
Routinely	15 (8.4)	1 (2.5)	10 (9.8)	4 (10.8)	
Indication CAD, n (%)					
Never	4 (2.2)	2 (5.0)	2 (2.0)	0	0.053
Rare	10 (5.6)	5 (12.5)	4 (3.9)	1 (2.7)	
Frequent	131 (73.2)	29 (82.5)	74 (72.5)	28 (75.7)	
Routinely	34 (19.0)	4 (10.0)	22 (21.6)	8 (21.6)	
Indication HCM, n (%)					
Never	55 (30.7)	16 (40.0)	33 (32.4)	6 (16.2)	0.134
Rare	81 (45.3)	17 (42.5)	47 (46.1)	4 (10.8)	
Frequent	41 (22.9)	6 (15.0)	21 (20.6)	14 (37.8)	
Routinely	2 (1.1)	1 (2.5)	1 (1.0)	0	
Indication HF, n (%)					
Never	47 (26.3)	14 (35.0)	26 (25.5)	7 (18.9)	0.058
Rare	82 (45.8)	22 (55.0)	47 (46.1)	13 (35.1)	
Frequent	45 (25.1)	3 (7.5)	26 (25.5)	16 (42.2)	
Routinely	5 (2.8)	1 (2.5)	3 (2.9)	1 (2.7)	
Mitral disease, n (%)					
Never	34 (19.0)	13 (32.5)	20 (19.6)	1 (2.7)	0.006
Rare	63 (35.2)	12 (30.0)	39 (38.2)	12 (32.4)	
Frequent	78 (43.5)	15 (37.5)	39 (38.2)	24 (64.8)	
Routinely	4 (2.2)	0	5 (3.9)	0	
Aortic disease, n (%)					
Never	22 (12.3)	7 (17.5)	13 (12.7)	2 (5.4)	0.050
Rare	73 (40.8)	17 (42.5)	45 (44.1)	11 (29.7)	
Frequent	82 (45.8)	10 (25.0)	42 (41.1)	24 (64.8)	
Routinely	2 (1.1)	0	2 (2.0)	0	

SE: Stress echocardiography, CFVR: Coronary flow velocity reserve, CAD: Coronary artery disease, HCM: Hypertrophic cardiomyopathy, HF: Heart failure

adhering to the SE recommendation, that to be familiar with all the forms of physical and pharmacological SE.<sup>[13]</sup>

One notable advancement in SE is the simultaneous assessment of CFVR on LAD and LUS. This approach significantly expands the diagnostic and prognostic potential of the traditional evaluation, which was primarily based on identifying regional wall motion abnormalities (RWMA).

### ABCDE stress echocardiography


The upgrade of SE to the ABCDE protocol represents a significant advancement in the field, aiming to provide a more comprehensive assessment of patients with ischemic heart disease and other cardiac conditions. The ABCDE protocol involves five steps that offer a more integrated evaluation

of patient vulnerability beyond the detection of anatomical CAD.<sup>[18,19]</sup>

Cardiac functional testing with ABCDE SE allows to gain a comprehensive insight on patient vulnerability still with an extraordinarily simple and feasible test with low cost, minimal risk, zero radiation, and near-zero environmental impact.

This approach is likely to enhance the diagnostic accuracy and prognostic value of SE in clinical practice, leading to better patient management and outcomes.<sup>[19]</sup>

This new approach was rapidly implemented by the Italian echocardiography laboratories and, as emerged from the survey, is now an integral part of the evaluation of the patient




**Figure 3:** Percentage of coronary flow velocity reserve evaluation during stress echocardiography in low, moderate, and high volume centers

in ischemic heart disease and beyond ischemic heart disease, especially in high-volume centers. It has also allowed an extension of the indications to SE, beyond ischemic heart disease, as in cardiomyopathies and valvulopathies.

There are three possible reasons of this rapid reshape of SE practice. First, the culture of SE has deep clinical and cultural roots in Italy, since several Italian laboratories were early adopters of the technique in the eighties and established the practice of SE well before it was recognized and endorsed by the international guidelines. Second, the last wave of SE innovation with the addition of B-lines and CFVR was again started in Italy in 2002 and 2004, and progressively accepted worldwide. Third and possibly more importantly, the new ABCDE protocol received the official endorsement of the SIECVI, which allowed a more efficient dissemination of the project, harmonization of protocols across different laboratories, and rapid uptake of the technique by leading edge laboratories. Stress echo 2030 is a flagship project of SIECVI, and over 20 Italian laboratories are currently active parts and recruiters of the study. When innovation starts in the clinically oriented laboratories, daily practice is aligned with the state of the art protocols, and the dissemination of innovation becomes easier and faster.

B-lines LUS identify pulmonary congestion at rest<sup>[8,9]</sup> and, more frequently, during stress<sup>[10,11]</sup> in a variety of cardiovascular conditions, characterized by the possible occurrence of increased pulmonary artery wedge pressure and accumulation of extravascular lung water. Stress B-lines were more frequent than rest B-lines, indicating that SE can be useful to unmask a condition of latent pulmonary congestion, undetectable at rest, and shared by different cardiovascular conditions: CAD, heart failure with preserved and reduced ejection fraction, valvulopathy.<sup>[9,11]</sup>

CFVR on LAD offers an integrated assessment of epicardial coronary artery stenosis and coronary microcirculation.<sup>[20]</sup> Large evidences supporting the usefulness of CFVR, especially



**Figure 4:** Percentage of lung ultrasound evaluation during stress echocardiography in low, moderate, and high volume centers

for risk stratification in CAD and HF and its endorsement in SE recommendations by the European Society of Echocardiography.<sup>[13]</sup> Starting 2016, CFVR was adopted in the ABCDE protocol of the stress echo 2020 first and after, stress echo 2030 study as the new clinical standard of the technique.<sup>[3,4]</sup>

CFVR during SE is feasible with high success with vasodilator than with dobutamine or exercise test.<sup>[5,21]</sup> Reduced CFVR is more prevalent in the patients with inducible RWMA or extensive CAD, but can be found also in patients with normal coronary arteries.<sup>[5]</sup> Reduced CFVR is a marker of altered coronary microvascular function and/or epicardial artery stenosis, which integrates and complements stress-induced RWMA which are more specific for a reduction of CFVR due to epicardial artery stenosis.<sup>[6,7]</sup> The risk is lowest for patients with preserved CFVR and no RWMA, intermediate in patients with only reduced CFVR and highest for patients with RWMA and reduced CFVR.<sup>[5]</sup>


### Comparison with previous studies

Compared to previous Italian SE survey of 2015,<sup>[22]</sup> we had more centers involved (179 vs. 125 centers, respectively): SE activity was present in 81% of Italian centers, higher data compared to previous Italian survey with 67% of the centers, 61% in UK<sup>[23]</sup> and 49% in Austria.<sup>[24]</sup>

Exercise was performed in 67% of labs, with an increase compared to the previous value of 2015 in Italy (56%), similar from UK practice, exercise was used in 67% of laboratories, while in Austria exercise was lower employed (only 26% of laboratories).

Dobutamine was widely used in 91% and vasodilators by 82% of the centers in the previous Italian survey, grossly comparable with 85% and 78%, respectively, of the centers involved in this survey.

Dobutamine was largely used in the UK (100%) and Austria (91%), whereas vasodilators were underused in



**Figure 5:** Percentage of stressors used in stress echocardiography (SE) on the center up. The symbol of stress echo 2030 on the right top. In the bottom on the left, an example of coronary flow evaluated at rest and at peak stress and percentage of coronary flow velocity reserve evaluation during SE in low, moderate and high volume centers. In the bottom on the center a picture of standard evaluation of regional wall motion with normal at rest and abnormal at peak stress. In the bottom on the right an example of lung ultrasound (LUS) evaluation at rest and at peak stress, and the percentage of LUS during SE in low-, moderate-, and high-volume centers

Austria (11%) and in UK (20%), a markedly lower percentage compared to the Italian data.

As an extraconsideration, the drug cost is not separately reimbursed in Italy, and therefore, the drug cost can represent an issue. In addition, in some European countries, intravenous dipyridamole is not commercially available.

The noninvasive pacemaker stress echo was used by 21% of laboratories, while only 6% in the previous Italian survey. The percentage was much higher in the UK (40%) and lower (2%) in Austria.

This option is particularly beneficial for patients with permanent pacemakers, as it allows for stress testing to be performed in a few minutes without the need for an intravenous line.<sup>[25]</sup> The described stress test is fast, safe, and requires minimal interaction with the electrophysiology outpatient lab. This means that it can be carried out efficiently and with fewer resources, making it a convenient and practical option for assessing cardiac function and response to stress in patients with pacemakers.

### Study limitations

We used the electronic mailing list of the Italian Society of Echocardiography, which covers most - but certainly not all - the SE activities in Italy.<sup>[15]</sup> In fact, certification is not mandatory, and it is also run in parallel and independently by the European Association of Cardiovascular Imaging, and many cardiologists are directly accredited by international societies and do not pass through the Italian society. Some of these extra-SIECVI centers are also of large volumes and high quality standards. Therefore, the survey might have underestimated the dissemination of SE activities in Italy but has likely mirrored the quality and pattern of practice in a

realistic fashion. As always in a survey, there are nonresponders for several reasons, including lack of time or unwillingness to participate to the study. No independent, external validation of the data provided by the cardiologist head of the participating lab was possible.<sup>[15-17]</sup>

### CONCLUSIONS

The survey described the state of the art of SE in Italy and the SE community, characterized as being open to innovation and efficient in integrating scientific evidence into everyday clinical practice with minimal time-lag. The framework established by this community is seen as culturally and logistically suitable for developing the new generation of SE, exploring various aspects of SE, both within and beyond CAD. The community is embracing new parameters, such as B-lines and CFVR, in addition to traditional RWMA.

Furthermore, SE is extending its applications beyond CAD to evaluate conditions such as dilated and hypertrophic cardiomyopathy and valvular heart disease. This expanded scope makes SE a more versatile and comprehensive tool for assessing various cardiac pathologies.

The stage is now set for prospective, large-scale, multicenter effectiveness studies, as SE2020 first and SE 2030 studies endorsed by SIECVI are crucial for determining the clinical utility and diagnostic accuracy of this multiparametric approach with ABCDE SE.

By embracing these new approaches, the SE community is striving to improve patient care, enhance diagnostic capabilities, and ultimately contribute to better patient outcomes.



As the field of SE continues to evolve and incorporate new findings, it reinforces the importance of evidence-based medicine and the collaborative efforts of researchers, clinicians, and the medical community to advance cardiovascular care.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

### REFERENCES

- Knuuti J, Wijns W, Saraste A, Capodanno D, Barbato E, Funck-Brentano C, *et al.* 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J* 2020;41:407-77.
- Pellikka PA, Arruda-Olson A, Chaudhry FA, Chen MH, Marshall JE, Porter TR, *et al.* Guidelines for performance, interpretation, and application of stress echocardiography in ischemic heart disease: From the American Society of Echocardiography. *J Am Soc Echocardiogr* 2020;33:1-41.e8.
- Picano E, Ciampi Q, Citro R, D'Andrea A, Scali MC, Cortigiani L, *et al.* Stress echo 2020: The international stress echo study in ischemic and non-ischemic heart disease. *Cardiovasc Ultrasound* 2017;15:3.
- Picano E, Ciampi Q, Cortigiani L, Arruda-Olson AM, Borguezan-Daros C, de Castro E Silva Pretto JL, *et al.* Stress echo 2030: The novel ABCDE-(FGLPR) protocol to define the future of imaging. *J Clin Med* 2021;10:3641.
- Ciampi Q, Zagatina A, Cortigiani L, Gaibazzi N, Borguezan Daros C, Zhuravskaya N, *et al.* Functional, anatomical, and prognostic correlates of coronary flow velocity reserve during stress echocardiography. *J Am Coll Cardiol* 2019;74:2278-91.
- Cortigiani L, Rigo F, Gherardi S, Sicari R, Galderisi M, Bovenzi F, *et al.* Additional prognostic value of coronary flow reserve in diabetic and nondiabetic patients with negative dipyridamole stress echocardiography by wall motion criteria. *J Am Coll Cardiol* 2007;50:1354-61.
- Cortigiani L, Rigo F, Bovenzi F, Sicari R, Picano E. The prognostic value of coronary flow velocity reserve in two coronary arteries during vasodilator stress echocardiography. *J Am Soc Echocardiogr* 2019;32:81-91.
- Picano E, Pellikka PA. Ultrasound of extravascular lung water: A new standard for pulmonary congestion. *Eur Heart J* 2016;37:2097-104.
- Picano E, Scali MC, Ciampi Q, Lichtenstein D. Lung ultrasound for the cardiologist. *JACC Cardiovasc Imaging* 2018;11:1692-705.
- Scali MC, Zagatina A, Ciampi Q, Cortigiani L, D'Andrea A, Daros CB, *et al.* Lung ultrasound and pulmonary congestion during stress echocardiography. *JACC Cardiovasc Imaging* 2020;13:2085-95.
- Merli E, Ciampi Q, Scali MC, Zagatina A, Merlo PM, Arbucci R, *et al.* Pulmonary congestion during exercise stress echocardiography in ischemic and heart failure patients. *Circ Cardiovasc Imaging* 2022;15:e013558.
- Lancellotti P, Pellikka PA, Budts W, Chaudhry FA, Donal E, Dulgheru R, *et al.* The clinical use of stress echocardiography in non-ischaemic heart disease: Recommendations from the European Association of Cardiovascular Imaging and the American Society of Echocardiography. *Eur Heart J Cardiovasc Imaging* 2016;17:1191-229.
- Sicari R, Nihoyannopoulos P, Evangelista A, Kasprzak J, Lancellotti P, Poldermans D, *et al.* Stress echocardiography expert consensus statement – Executive summary: European Association of Echocardiography (EAE) (a registered branch of the ESC). *Eur Heart J* 2009;30:278-89.
- Picano E, Pellikka PA. Stress echo applications beyond coronary artery disease. *Eur Heart J* 2014;35:1033-40.
- Ciampi Q, Pepi M, Antonini-Canterin F, Barbieri A, Barchitta A, Faganello G, *et al.* Organization and activity of Italian echocardiographic laboratories: A survey of the Italian Society of Echocardiography and cardiovascular imaging. *J Cardiovasc Echogr* 2023;33:1-9.
- Ciampi Q, Antonini-Canterin F, Barbieri A, Barchitta A, Benedetto F, Cresti A, *et al.* Remodeling of activities of Italian echocardiographic laboratories during the coronavirus disease 2019 lockdown: The SIECoVID study. *J Cardiovasc Med (Hagerstown)* 2021;22:600-2.
- Ciampi Q, Antonini-Canterin F, Barbieri A, Barchitta A, Benedetto F, Cresti A, *et al.* Reshaping of Italian echocardiographic laboratories activities during the second wave of COVID-19 pandemic and expectations for the post-pandemic Era. *J Clin Med* 2021;10:3466.
- Picano E, Zagatina A, Wierzbowska-Drabik K, Borguezan Daros C, D'Andrea A, Ciampi Q. Sustainability and versatility of the ABCDE protocol for stress echocardiography. *J Clin Med* 2020;9:3184.
- Ciampi Q, Zagatina A, Cortigiani L, Wierzbowska-Drabik K, Kasprzak JD, Haberka M, *et al.* Prognostic value of stress echocardiography assessed by the ABCDE protocol. *Eur Heart J* 2021;42:3869-78.
- Rigo F, Richieri M, Pasanisi E, Cutaia V, Zanella C, Della Valentina P, *et al.* Usefulness of coronary flow reserve over regional wall motion when added to dual-imaging dipyridamole echocardiography. *Am J Cardiol* 2003;91:269-73.
- Zagatina A, Zhuravskaya N. The additive prognostic value of coronary flow velocity reserve during exercise echocardiography. *Eur Heart J Cardiovasc Imaging* 2017;18:1179-84.
- Ciampi Q, Citro R, Severino S, Labanti G, Cortigiani L, Sicari R, *et al.* Stress echo in Italy: State-of-the-art 2015. *J Cardiovasc Med (Hagerstown)* 2017;18:637-9.
- Bhattacharyya S, Chehab O, Khattar R, Lloyd G, Senior R, British Society of Echocardiography. Stress echocardiography in clinical practice: A United Kingdom National Health Service Survey on behalf of the British Society of Echocardiography. *Eur Heart J Cardiovasc Imaging* 2014;15:158-63.
- Weidenauer D, Bartko P, Zach H, Zehetgruber M, Domanovits H, Graf S, *et al.* Stress-echocardiography is underused in clinical practice: A nationwide survey in Austria. *Wien Klin Wochenschr* 2015;127:514-20.
- Picano E, Alaimo A, Chubuchny V, Plonska E, Baldo V, Baldini U, *et al.* Noninvasive pacemaker stress echocardiography for diagnosis of coronary artery disease: A multicenter study. *J Am Coll Cardiol* 2002;40:1305-10.