



Article

# Experiencing forest therapy in the Italian landscape: bathing in the Selva of Castelfidardo

Giovanna Borriello<sup>1,4\*</sup>, Elisa Grazioli<sup>2</sup>, Paolo Zavarella<sup>3,5</sup>, Roberto Paolini<sup>3,5</sup>, Antonio Ianniello<sup>1</sup>, Dario Silvestri<sup>4</sup>, Claudia Cerulli<sup>2</sup>, Eliana Trinchita<sup>2</sup> and Attilio Parisi<sup>2</sup>

- <sup>1</sup> S. Andrea Hospital, University of Rome “Sapienza”, Via di Grottarossa1035, 00189 Roma
- <sup>2</sup> Department of Movement, Human and Health Science, University of Rome “Foro Italico”, Piazza Lauro De Bosis 6, 00135 Rome, Italy; elisa.grazioli@uniroma4.it
- <sup>3</sup> A.I.Me.F. Associazione Italiana di Medicina Forestale Via G.de Vecchi Pieralice 21, 00167 Roma; info@aimef.net
- <sup>4</sup> ASOMI College of Sciences, Block D G. Mitrovich Street, Pembroke, PBK 1014 Malta; dario.silvestri@acs-college.com
- <sup>5</sup> EDUCAM Institute - Education in Complementary and Alternative Medicine, Via G.de Vecchi Pieralice 21, 00167 Roma; info@educam.it
- \* Correspondence: PZ zavarella@airop.it Tel.: 00393358122955; GB giovanna.borriello@gmail.com Tel.: 003934779226262

**Abstract:** According to several evidence, forest environmental seems able to provide beneficial effects on functional and psychological parameters, related to cardiovascular, metabolic, respiratory functions as well depression and anxiety. The aim of this study is to investigate the effect of a one-day forest walking in the Selva of Castelfidardo (AN, Italy) on 37 participants aged 21-68, most of them living in either urban or suburban areas of large cities. We observed a statistically significant effect on sympatheticovagal balance by the means of heart rate, systolic and diastolic blood pressure, body temperature, skin temperature, skin conductance, HRV parameters (AVNN, SDNN, rMSSD, pNN50, LF, HF, LF/HF ratio), oxygen oximetry, PEF, FEV1. A significant difference was also detected on the Perceived Stress Scale responses (19.27 pre vs 13.81 post-immersion,  $p < 0,05$ ; -28,3% variation). Our data contribute to increase the body of literature about the effect of forest walking, adding data on an Italian area assigned to forest bathing.

**Citation:** Lastname, F.; Lastname, F.; Lastname, F. Title. *Int. J. Environ. Res. Public Health* **2022**, *19*, x. <https://doi.org/10.3390/xxxxx>

**Keywords:** forest bathing; forest medicine; human health

Academic Editor: Firstname Lastname

Received: date  
Accepted: date  
Published: date

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

In the last decades, several evidences suggest that being exposed to nature, such as walking in the forest or near waterfall, visiting or just looking at natural environments seems able to provide beneficial effects on human health [1]. According to the latest studies, functional and psychological parameters, such as cardiovascular, metabolic, respiratory functions and depression, stress, anxiety seems to improve after natural exposure, known as “forest bathing” [2–4]. This expression originates in Japan, and it was called “Shinrin-yoku” in Japanese. A forest bathing trip involves a short and leisurely visit to forest for relaxation and recreation while breathing in volatile substances, mostly wood essential oils called phytoncides, which are organic compounds obtained from trees, such as a-pinene and limonene [5]. It seems that the positive effects revealed after this trip could be attributed to the indirect effects of the organic compound, combined with an increased physical activity level, social interactions and exposure to sunlight [6]. More in deep, after the growing interest in ecology and health worldwide, a few studies have reported that natural and green spaces promote modern human health, increasing people quality of life [7-9]. The term “forest therapy,” or better “forest medicine”, describes the medically proven health effects resulting from exposure to forests. Latest

studies evidenced that the direct contact with nature, such as viewing natural landscapes or walking in the forest, seem to increase parasympathetic nervous activity and suppresses sympathetic nervous activity [10,11]. The Heart Rate Variability (HRV) is the non-invasive index of autonomic controls of the heart [12]. This parameter can provide objective evaluation about stress offering some advantages over alternative physiological measurements. In fact, HRV can be recorded continuously in a non-invasive manner [13]. The positive effect of nature exposure on this parameter is supported by studies that evidenced how negative emotions such as anxiety, depression, and tension are reduced, while positive emotions increase, as well as the psychological relaxation enhanced [14]. It seems that these relaxing effects of forest, and natural landscape, are produced by gaining information about the physical environment, such as air temperature, humidity, illuminance, sounds, etc., as well as about chemical environments such as phytoncides of forest through our five senses [15]. As far as we know, most of the studies that evidence the positive effect of forest-bathing on people quality of life have been performed in Asia, mostly Japan, and north Europe. Research on human health benefits associated with the immersion in nature needs to be implemented in south Europe. A specific field of research, conducted worldwide, is needed to produce evidence of the relationships between Shinrin-Yoku and clinical therapeutic effects. Nature therapy as a health-promotion method and potential universal health model is implicated for the reduction of then modern stressful lifestyle. The aim of this study is to investigate the effect on physiological and psychological parameters and ability to decrease undue stress of a one-day forest walking in the Selva of Castelfidardo (AN, Italy) on a non-selected, mixed group of 37 people habitually living in big cities, exposed to high level of urban traffic and fast pace of life and work. This pilot study will contribute to increase the body of literature about the effect of forest walking, adding data on an Italian area qualified for forest bathing.

## 2. Materials and Methods

This prospective, single-group, single-centre pilot study was approved by the University of Rome “Foro Italico” Committee (n. CAR 83/2021) and by Scientific Committee of ACS ASOMI College of Sciences - Higher Education Institution, Malta. Thirty-seven participants, both male and female, members of the association, were invited to participate to the bathing with their friends or relatives by mail and by poster on the official site of the Associazione Italiana di Medicina Forestale (A.I.Me.F.) on Facebook. Written informed consent was prepared and drafted encompassing the proposal of the data collection and signed by participants. The experience was free and open to all without any criteria of selection except for the adulthood and coming from an urban setting.

### 2.1. Demographic data

The demographics of the 37 participants are reported in Table 1. Overall, 15 men (40.5%) and 22 women (59.4%) participated, with 6 of them aged 21–30 years (16.2%), 7 aged 31–50 years (18.9%), 18 aged 51–65 years (48.6%), and 6 aged 66 years and older (16.2%). The mean age was 54,8 years, with a standard deviation of 12.7 years. Nearly 83% of participants had a college degree and are busy and currently employed for more than 8 hours/day. Most of the participants lived in either urban areas of large cities.

**Table 1.** Demographics of participants (n=37). Values are reported as mean  $\pm$  standard deviation (SD).

<b>Age</b>	<b>54,8 <math>\pm</math> 12,7</b>
<b>Weight (kg)</b>	<b>66,1 <math>\pm</math> 10,6</b>
<b>Height (cm)</b>	<b>166,1 <math>\pm</math> 6,9</b>
<b>Gender</b>	<b>22 F; 15 M</b>

Smokers

49%

92

## 2.2. Study Site

93

The Selva of Castelfidardo is a prehistoric forest, geographically known as Montoro Selva, covering the north side of the Monte Oro hill rising from an altitude of 15 meters above sea level to the summit, up to an altitude of 118 m. This mesophilic oak forest site of 36 hectares of sub-Mediterranean deciduous forests extends over the hills of the Conero Riviera, in the province of Ancona. The site is under the landscape restrictions of the Legislative Decree n.1497/1939, recognized since 1981 from the Marche Region as "Floristic protected Area" since 1974 and proposed as Site of Community Interest (S.I.C.) sine year 2000 [16]. Approximately 400 plants and herbal can be found, 37 mostly representative because endemic, rare, in danger of extinction, of great phytogeographical significance and peculiar expression of the habitat. The forest is not uniform, either pteridophyta than gymnospermae and angiospermae are presented [17] : in the upper we can found thermophilic and heliophilous species (*Quercus pubescens*, *Fraxinus ornus*, *Acer campestre*), a shrubby layer (*Carpinus orientalis*, *Ligustrum vulgare*, *Sorbus domestica*, *Sorbus torminalis*, *Prunus spinosa*), an herbaceous layer (*Buglossoides purpureoerulea*, *Rubia perigrina*, *Asparagus acutifolius*, *Smilax aspera*). In the lower part of the forest are predominant different kinds (*Quercus petraea*, *Quercus robur*, *carpinus betulus*), in the shrubby layer we can found *Ilex aquifolium*, *Corylus avellana* and *Iris foetidissima*, *hypericum androsaemum*, *Vinca minor*, *Campanula trachelium* as predominant herbal. Homogeneous distribution of others species is visible over the entire area, in particular *Quercus cerris*, *Laurus nobilis*, *Ostrya carpinifolia*, *Malus sylvestris*, *Ulmus minor*, *Cornus mas*, *Cornus sanguinea*, *Crataegus oxyachanta*, *Coronilla emerus*, *Erica arborea*, *Ruscus acukeatus*, *Daphne laureola*, *Lonicera etrusca*, *Robinia pseudoacacia*. Among the climbing vines, we appoint *Hedera helix*, *Tamus communis* and *Clematis vitalba*. Several herbal anf flowers as *Cyclamen repandum*, *cyclamen neapolitanum*, *Viola alba*, *Primula vulgaris*, *Allium pendulinum*, *Melittis melissophyllum*, *Solidago virgaurea*, *Ranunculus lanuginosus*. In the clearings are present bushes of *Rubus caesiu* , *Prunella vulgaris*, *Inula conyza*. Over 35 variety of mosses are present, the most widespread are *Brachytecium rutabulum*, *Rhynchostegium confertum*, *Eurhynchium stokesii*, *Mnium rostratum*, *Mnium undulatum*, *Eurhynchium striatum* [18-19]. The microclimate has allowed the development of a biodiversity which supported the choice from A.I.Me.F. to qualify it as a "Forest Bathing Center". The qualification procedure allows to demonstrate how spending time in a wooded area with extreme biodiversity produces beneficial effects that can be highlighted through biological and biopsychosocial investigations and tests. There are two steps (Phases) to qualify a site as a "Forest Bathing Center".

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

### Phase 1. Site quality analysis

127

It is the evaluation of the places (built and natural) of the location, using 10 indicators, each of which is assigned a variable index from 0 to 5 based on the presence or absence of quality characteristics for each indicator: air quality, water quality, quality of the earth, presence of trees / woods, presence of paths (easy to walk), gps and telephone network coverage, overnight accessibility, possibility of eating meals, area dedicated to forestry medicine, presence of paths. The site was found to be suitable for the assessments carried out in Phase 1, so it was subjected to the more specific assessments of Phase 2.

128

129

130

131

132

133

134

135

### Phase 2. Forest Immersion

136

The Forest Bathing was carried out on Sunday 12 July, from 10:00 to 18:00, all specific steps were reported in Table 2.

137

138

**Table 2.** Forest Immersion Schedule in Selva of Castelfidardo.

139

Phase	Activity	Duration
Participants Arrival	Registration and signature of the informed consent form; ID code assignment; PSS completion	30 minutes
Data Collection	At the beginning: Vital signs, HRV, PEF, FEV1, autonomous nervous system testing and oximetry	1 hour
Greeting to the Dojo of Nature	People are guided to the entrance to the Selva, where the “Greeting to the Dojo” is read and commented on	15 minutes
First Bath	First slow walk inside the Selva (500-600m). People are invited to observe nature	30 minutes
First Stop	First stop with a simple guided meditation paying attention to breathing. People find a place lying or sitting next to the trees	10 minutes
Second Bath	Second slow walk inside the Selva (500-600m). People are invited to observe nature and are provided more detailed information on plant recognition, plant neurobiology and phytosociology of the forest	30 minutes
Second Stop	Read of the “Eulogy of Silence”.	10 minutes
Third Bath	Participants are invited by the Tutors to take a walk slowly and silently in the Selva, without distraction (500-600m)	30 minutes
Third Stop	Description of the “Meditation of the 5 Senses”	10 minutes
Fourth Bath	Participants are invited by the Tutors to take a walk slowly and silently in the Selva, without distraction (500-600m)	10 minutes
Fourth Stop	Short briefing and description of the emotions / sensations experienced. Then the participants are invited to look for a Tree and get in direct contact: reading the “Meditation on the Trees”	15 minutes
Lunch break	Lunch break inside the Selva. Smoking (including e-cigarettes), coffee and alcohol are not allowed. Listening to the “Music of the Plants” with practice of the techniques of the “Plantfulness®” method and briefing	1 hour
Fifth Bath	Participants walking, heading towards the starting point but following another path (500-660m)	30 minutes
Fifth Stop	Historical and botanical description of the forest and its seasonal biodiversity	15 minutes
Sixth Bath	The participants walk always guided by the Tutors to notice the rich biodiversity	30 minutes
Sixth Stop	Guided practice of Yoga exercises, Barefooting, Grounding, Mindfulness etc.	30 minutes
Seventh Bath	participants walk to the exit of the Selva (500-600m)	
Seventh Stop	Outside the Selva participants read and comment on “Thanksgiving to Pachamama”	10 minutes
Post-bathing collection	At the end: Vital signs, HRV, PEF, FE1, autonomous nervous system testing and oximetry; PSS completion	1 hour

At the entrance and leaving from the forest we propose a brief thanksgiving to the Forest to acknowledge your presence and respectfully greet other practitioners. The 5 senses meditation helps people to connect to the present moment using sense of sight, sound, touch, taste and smell. We opted for use this meditation to ground participants in the here and now to avoid and to distance unwanted thoughts, feeling physical discomfort or uncomfortable emotions. Walking in silence, people are invited to focusing attention to the micro-cosmos all the way around, enhancing the contact with nature. Sharing emotional experiences with others is encouraged from the tutors to reactivate the emotion at a more symbolic level, all taking place as part of ensuing interpersonal interactions

140  
141  
142  
143  
144  
145  
146  
147  
148  
149

in the group, implementing bonding, to become closer to others and reduce feelings of loneliness, empathy and receiving attention from others, possibly also to entertain engaging others and facilitate social interactions inside the group. At the same way, we propose activities, games or exercises to carry out in group. Insights on biodiversity, herbal curiosities and other advices on plant, trees and biological compounds are provided from the experts in forest medicine from the staff. Barefooting or walking in minimalist shoes without risk of injury is recommended during immersion due to several benefits of walking barefoot, including a better control of foot position when it strikes the ground, an improvement in balance, proprioception, and body awareness, which can help with pain relief and to stronger leg muscles. On the walking back to the measurements, the participants can experience other emotional activities: a special electronic music machine call "music of the plants" it's played for the audience: this machine, that can detect and translate in sounds the difference in the electrical flow of plants with its two electrodes placed on leafs and roots, it's a powerful instrument to communicate and show to everyone how alive and aware are plants towards their own environment. Participants can interact with the plant connected to this musical instrument and hear a difference in the musical output. After this experience all participants are invited to hug one of the trees they choose: this practice, as well as helping release negative feelings, allows them to breathe close next to the bark of the tree where some of the heaviest BVOC are likely to be found and collected. If it meets the feeling of the group, tutor can finally act a greeting to Pachamama, a personification of the Mother Earth in Inca mythology, to underline prosperity, fertility and wondering Nature. To get an instant and warm feedback from the group, tutor can invite all the participants to take each other by the hands to create a circle where the tutor invite to imaging to through inside some of the feelings and the words that have remained etched in their memory regarding the experience of the forest therapy they just lived together. This practice called "wishing well" helps to consolidate the positive memories about nature and shows different perspectives about the experiences of all participants and lead the groups to a collective and sincere warm group hug.

#### *Measurements during Phase 2*

Before and after the one-day walking in Selva of Castelfidardo, all participants performed the following physiological measurements to evaluate sympathetic and parasympathetic nervous system activity, and sympathicovagal balance: heart rate (HR), systolic and diastolic blood pressure, body temperature, skin temperature, skin conductance, HRV parameters (AVNN, SDNN, rMSSD, pNN50, LF, HF, LF/HF ratio), oxygen oximetry, PEF, FEV1, catching data in a few minutes by portable and easy to use instruments. We used Mir Smart one (portable spirometer, individually set with single-use tube for expiratory act); Hylogy – digital upper arm blood pressure monitor; Lovia digital portable oximeter; Mindfield e-Sense Pulse to measure HRV parameters; Mindfield eSense Skin Response to measure skin conductance and Mindfield eSense Temperature - Biofeedback for cutaneous temperature variations. In the pre-test phase, all participants filled out a questionnaire, which included demographic information and questions regarding emotional status by the means of Perceived Stress Scale (PSS) from Sheldon Cohen, the most widely used psychological instrument for measuring the perception of stress. It is a measure of the degree to which situations in one's life are appraised as stressful. Items were designed to tap how unpredictable, uncontrollable, and overloaded respondents find their lives. The scale also includes a number of direct queries about current levels of experienced stress. The PSS was designed for use in community samples with at least a junior high school education. The items are easy to understand, and the response alternatives are simple to grasp. Moreover, the questions are of a general nature and hence are relatively free of content specific to any subpopulation group. The questions in the PSS ask about feelings and thoughts during the last month. PSS scores are obtained by reversing responses (e.g., 0 = 4, 1 = 3, 2 = 2, 3 = 1 & 4 = 0) to the four positively

stated items (items 4, 5, 7, & 8) and then summing across all scale items. A short 4 item scale can be made from questions 2, 4, 5 and 10 of the PSS 10 items.

### 2.3. Statistical Analysis

The SPSS® version 23.0 for Windows (IBM Corp., released 2015) was used to analyse the data. The Kolmogorov-Smirnov test was applied, before analysis, to test the normal distribution of data. Due to the normal distribution, parametric paired t-tests were performed to all variables, whilst mean and standard deviation scores were chosen as appropriate descriptors of measures of central tendency.

## 3. Results

The pre and post-immersion measures of physiological responses are reported in **Table 3** and displayed in figures.

**Table 3.** Pre- and post-immersion measures of physiological responses (Values are reported as mean,  $\pm$  SD and % of variation between pre and post data).

Parameters	Before (T0)	After (T1)	Var %	p-value
HR	71,03 $\pm$ 11,39	75,51 $\pm$ 13,05	6,3%	0,001
SBP	112,16 $\pm$ 18,53	112,03 $\pm$ 15,97	-0,1%	0,954
DBP	65,73 $\pm$ 11,67	65,95 $\pm$ 10,35	0,3%	0,883
Temperature min	30,04 $\pm$ 1,18	29,44 $\pm$ 0,91	-2,0%	0,002
LF/HF	1,08 $\pm$ 0,89	0,73 $\pm$ 0,33	-32,2%	<0,05
rMSSD/AVNN	0,04 $\pm$ 0,018	0,04 $\pm$ 0,017	-0,6%	<0,05
PEF	385,56 $\pm$ 129,7	429,67 $\pm$ 121,18	11,4%	0,000
FEV1	3,11 $\pm$ 0,80	3,39 $\pm$ 0,61	8,9%	0,000
Oximetry	95,76 $\pm$ 1,44	97,14 $\pm$ 1,03	1,4%	0,000
PSS	19,27 $\pm$ 5,87	13,81 $\pm$ 5,42	-28,3%	<0,05

In particular, regarding the HR of participants (**Figure 1**), they exhibited significantly higher post-test HRs than pre-test HRs median value (71.03 vs 75.51,  $p = 0.001$ ).

Despite the changes in blood pressure were not significant (112,16 vs 112.03 for SBP,  $p=0.954$ ; 65.73 vs 65.93 for DBP,  $p=0.883$ ), we observed a trend to normalize values (highest values versus normal range in particular) (**Figures 2 and 3**).

Regarding body temperature (**Figure 4**), we found significance only for Tmin (30.04 vs 29.44,  $p=0.002$ ).

The group showed significant pre-post-test difference in autonomic balance (LF/HF 1.08 vs 0.73, -32,2%,  $p<0,05$ ) (**Figure 5**).

Considering the HR analysis, rMSSD/AVNN ratio showed a variation (-0,6%) with  $p<0,05$  (**Figure 6**).

Ventilatory parameters showed an improvement in PEF (385.56 vs 429.67, +11.4%) and FEV1 (3.11 vs 3.39, +8.9%), both statistically significant (**Figures 7 and 8**).

The same observation arises for oximetry (95.76 vs 97.14, +1.4%) (**Figure 9**).

Finally, a significant difference on PSS responses median value during the post-immersion phase was detected (19.27 vs 13.81, -28,3% variation) (**Figure 10**).




Figure 1. Results for Heart Rate before (T0) and after (T1) the test.

232  
233




Figure 2. Results for Systolic Blood Pressure before (T0) and after (T1) the test.

234  
235




Figure 3. Results for Diastolic Blood Pressure before (T0) and after (T1) the test.

236  
237




Figure 4. Results for minimum Temperature before (T0) and after (T1) the test.

238

239




Figure 5. Results for Low Frequency to High Frequency ratio before (T0) and after (T1) the test.

240

241

242




Figure 6. Results for rMSSD to AVN ratio before (T0) and after (T1) the test.

243

244






Figure 7. Results for Peak Expiratory Flow before (T0) and after (T1) the test.

245

246




Figure 8. Results for Forced Expiratory Volume in 1 second (T0) and after (T1) the test.

247

248





Figure 9. Results for Oximetry before (T0) and after (T1) the test.

249

250



**Figure 10.** Results for the Perceived Stress Scale before (T0) and after (T1) the test.

#### 4. Discussion

Data from this single-day, accessible to everyone bathing in the Selva of Castelfidardo has confirmed the beneficial effect immediately perceived of Shinrin-Yoku on human health. Results from previous studies showed convincing data explaining the relationship between the natural environment and the relaxation effects on humans (e.g., decrease in blood pressure and pulse rates, inhibition of sympathetic nervous activity, enhancement of parasympathetic nervous activity, and decrease in cortisol concentration levels) [3,4,5]. Our observations confirmed that walking in a nature reserve fosters biological changes, indicating stress reduction involving fundamentally two physiological axes: the sympathetic adrenal-medullary and the hypothalamic-pituitary-adrenal one, as previously argued [5,20], with positive impact on potentially disabling chronic medical conditions [3]. In fact, results of previous studies performed on the physiological effects of the traditional Shinrin-Yoku showed how forest environments could lower concentrations of cortisol, lower pulse rate, lower blood pressure, increase parasympathetic nerve activity, and lower sympathetic nerve activity, in particular whereas citizens are involved in occasional or continuing forest bathing approaches with the aim of reducing stress [4,21,22]. In our study, the PSS Scores was significantly lower after walking in the forest, demonstrating the psychological benefits of forests, consistently with previous findings of the effects of viewing forest scenery or walking in forests [23,24,25]. A previous study, in agreement with our observation, found that participants with initially high blood pressure showed a decrease in blood pressure after walking in a forest, whereas those with initially low blood pressure showed an increase, suggesting that a forest environment can be used to help achieve an appropriate blood pressure, showing a truly person-oriented intervention [26]. Further studies, performed on Italian sites, are needed to better understand the positive impact of forest bathing on these parameters.

#### 5. Conclusions

Despite the heterogeneity of the sample and the small number of participants, our study contributes to underline the importance and beneficial effect of the forest medicine, which may be used to develop new strategies in particular regarding primary and secondary level strategies in the field of preventive medicine. Promoting a usual and consistent approach to forest therapy may implement institutional and local resources to avoid or reduce an incorrect lifestyle encouraging health promotion and chronic disease prevention, as interestingly demonstrated in a program involving a young population affected by cancer complicated by a syndrome called by the authors "connection deficit disorder" due to the actual COVID-19 pandemic [27]. Another point that we consider valuable to underline is the significance that studies, such as the one presented in this

article, may have to contribute to the dissemination of awareness among citizens of the health, preventive and therapeutic value of the woods and national parks in Italy.

**Author Contributions:** Conceptualization, GB, EG and PZ; methodology, EG and RP.; software, RP, AI; validation, PZ, DS and AP; formal analysis, RP; investigation, GB, PZ, CC.; resources, PZ, AI, CC.; data curation, RP, DS.; writing—original draft preparation, GB, EG.; writing—review and editing, GB, AI, EG, PZ; visualization, CC.; supervision, CC, AP, DS. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of The University of Rome “Foro Italico” (protocol code CAR 83/2021).

**Informed Consent Statement:** this study was approved by the University of Rome “Foro Italico” Committee (n. CAR 83/2021). Written informed consent has been obtained from the patients to publish this paper.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author.

**Acknowledgments:** The Authors acknowledge Dr. Gigliola Sigismundi, Dr. Carlo Giordano, Dr. Lorina Metaj, Dr. Anna Grazia Lentini, Dr. Chiara Maracci, Cristina Pietrolucci and Manuela Anzini for the fundamental support given in all the phases of the Qualification. A.I.Me.F. acknowledge the Fondazione Duca Roberto Ferretti di Castelferretto, Castelfidardo (AN) and the President Eugenio Paoloni for the technical and organizational support given during the Qualification and beyond.

**Conflicts of Interest:** All the authors declare no conflict of interest.

## Reference

- Catharine Ward Thompson, Peter Aspinall, Jenny Roe, Access to Green Space in Disadvantaged Urban Communities: Evidence of Salutogenic Effects Based on Biomarker and Self-report Measures of Wellbeing, *Procedia - Social and Behavioral Sciences*, Volume 153, 2014, Pages 10-22, <https://doi.org/10.1016/j.sbspro.2014.10.036>.
- Hansen MM, Jones R, Tocchini K. Shinrin-Yoku (Forest Bathing) and Nature Therapy: A State-of-the-Art Review. *Int J Environ Res Public Health*. 2017;14(8):851. Published 2017 Jul 28. doi:10.3390/ijerph14080851
- Chun MH, Chang MC, Lee SJ. The effects of forest therapy on depression and anxiety in patients with chronic stroke. *Int J Neurosci*. 2017 Mar;127(3):199-203. doi: 10.3109/00207454.2016.1170015. Epub 2016 Apr 10. PMID: 27033879.
- Han JW, Choi H, Jeon YH, Yoon CH, Woo JM, Kim W. The Effects of Forest Therapy on Coping with Chronic Widespread Pain: Physiological and Psychological Differences between Participants in a Forest Therapy Program and a Control Group. *Int J Environ Res Public Health*. 2016 Feb 24;13(3):255. doi: 10.3390/ijerph13030255. PMID: 26927141; PMCID: PMC4808918.
- Li, Q. Effect of forest bathing trips on human immune function. *Environ Health Prev Med* 15, 9–17 (2010). <https://doi.org/10.1007/s12199-008-0068-3>
- Antonelli M, Donelli D, Carlone L, Maggini V, Firenzuoli F, Bedeschi E. Effects of forest bathing (shinrin-yoku) on individual well-being: an umbrella review. *Int J Environ Health Res*. 2021 Apr 28:1-26. doi: 10.1080/09603123.2021.1919293. Epub ahead of print. PMID: 33910423.
- Braubach, M., A. Egorov, P. Mudu, T. Wolf, C. Ward Thompson, AND M. Martuzzi. Effects of urban green space on environmental health, equity, and resilience. Edition 1, Chapter 11, N. Kabish; A. Bonn, H. Korn, J. Stadler (ed.), *Nature-based solutions to climate change in urban areas - linkages of science, society and policy*. Springer, Heidelberg, Germany, 1:187-205, (2017). [https://doi.org/10.1007/978-3-319-56091-5\\_11](https://doi.org/10.1007/978-3-319-56091-5_11)
- Gascon M, Zijlema W, Vert C, White MP, Nieuwenhuijsen MJ. Outdoor blue spaces, human health and well-being: A systematic review of quantitative studies. *Int J Hyg Environ Health*. 2017 Nov;220(8):1207-1221. doi: 10.1016/j.ijheh.2017.08.004. Epub 2017 Aug 18. PMID: 28843736.
- Karjalainen E, Sarjala T, Raitio H. Promoting human health through forests: overview and major challenges. *Environ Health Prev Med*. 2010;15(1):1-8. doi:10.1007/s12199-008-0069-2
- Chorong Song, Harumi Ikei, Maiko Kobayashi, Takashi Miura, Qing Li, Takahide Kagawa, Shigeyoshi Kumeda, Michiko Imai, Yoshifumi Miyazaki, Effects of viewing forest landscape on middle-aged hypertensive men, *Urban Forestry & Urban Greening*, Volume 21, 2017, Pages 247-252, <https://doi.org/10.1016/j.ufug.2016.12.010>.
- Song C, Ikei H, Miyazaki Y. Physiological Effects of Nature Therapy: A Review of the Research in Japan. *Int J Environ Res Public Health*. 2016;13(8):781. Published 2016 Aug 3. doi:10.3390/ijerph13080781.

12. Kobayashi H, Ishibashi K, Noguchi H. Heart rate variability; an index for monitoring and analyzing human autonomic activities. *Appl Human Sci.* 1999 Mar;18(2):53-9. doi: 10.2114/jpa.18.53. PMID: 10388159. 342  
343
13. Kobayashi, Hiromitsu & Song, Chorong & Ikei, Harumi & Kagawa, Takahide & Miyazaki, Yoshifumi. (2015). Analysis of Individual Variations in Autonomic Responses to Urban and Forest Environments. Evidence-based complementary and alternative medicine: eCAM. 2015. 671094. 10.1155/2015/671094. 344  
345  
346
14. Jo H, Song C, Miyazaki Y. Physiological Benefits of Viewing Nature: A Systematic Review of Indoor Experiments. *Int J Environ Res Public Health.* 2019;16(23):4739. Published 2019 Nov 27. doi:10.3390/ijerph16234739. 347  
348
15. Yasuo Ohe, Harumi Ikei, Chorong Song, Yoshifumi Miyazaki, Evaluating the relaxation effects of emerging forest-therapy tourism: A multidisciplinary approach, *Tourism Management*, Volume 62, 2017, Pages 322-334, Blood pressure-lowering effect of Shinrin-yoku (Forest bathing): a systematic review and meta-analysis. 349  
350  
351
16. Introduction to the Selva of Castelfidardo. Fondazione Ferretti. Online site [www.fondazioneferretti.org](http://www.fondazioneferretti.org) 352
17. Ballelli S, Lucarini D, Orsomando E. Flora vascolare della Selva di Castelfidardo e del territorio circostante. Università di Camerino. Stampato presso Tipografia S. Giuseppe Pollenza Febbraio 2001 353  
354
18. Orsomando E, Balletti S, Catorci A, Formica E. Atlante iconografico di piante rare o significative della Selva di Castelfidardo. Area Floristica delle Marche. Tecnostampa Recanati 1992 355  
356
19. Aleffi M, Allegrini MC, Fuffa X, Muccichini S, Tacchi R. Briofite e licheni della Selva di Castelfidardo. Marche. Stampato presso Tipografia S. Giuseppe Pollenza Febbraio 2005 357  
358
20. Ideno Y, Hayashi K, Abe Y, Ueda K, Iso H, Noda M, Lee JS, Suzuki S. *BMC Complement Altern Med.* 2017 Aug 16;17(1):409. doi: 10.1186/s12906-017-1912-z. PMID: 28814305 359  
360
21. Laumann K, Garling T, Stormark KM. Selective attention and heart rate responses to natural and urban environments. *J Environ Psychol.* 2003;23(2):125–34. Psychological effects of forest environments on healthy adults: Shinrin-yoku (forest-air bathing, walking) as a possible method of stress reduction. 361  
362  
363
22. Morita E, Fukuda S, Nagano J, Hamajima N, Yamamoto H, Iwai Y, Nakashima T, Ohira H, Shirakawa T. *Public Health.* 2007 Jan;121(1):54-63. doi: 10.1016/j.puhe.2006.05.024. Epub 2006 Oct 20. PMID: 17055544  
<https://doi.org/10.1016/j.tourman.2017.04.010>. Psychosocial and Biological Outcomes of Immersive, Mindfulness-Based Treks in Nature for Groups of Young Adults and Caregivers Affected by Cancer: Results from a Single Arm Program Evaluation from 2016-2021. 364  
365  
366  
367  
368
23. Park B.J., Tsunetsugu Y., Kasetani T., Kagawa T., Miyazaki Y. The physiological effects of Shinrin-yoku (taking in the forest atmosphere or forest bathing): Evidence from field experiments in 24 forests across Japan. *Environ. Health Prev. Med.* 2010;15:18–26. doi: 10.1007/s12199-009-0086-9. 369  
370  
371
24. Tsunetsugu Y., Lee J., Park B.J., Tyrväinen L., Kagawa T., Miyazaki Y. Physiological and psychological effects of viewing urban forest landscapes assessed by multiple measurements. *Landsc. Urban Plan.* 2013;113:90–93. doi: 10.1016/j.landurbplan.2013.01.014. 372  
373  
374
25. Morita E., Fukuda S., Nagano J., Hamajima N., Yamamoto H., Iwai Y., Nakashima T., Ohira H., Shirakawa T. Psychological effects of forest environments on healthy adults: Shinrin-Yoku (forest-air bathing, walking) as a possible method of stress reduction. *Public Health.* 2007;121:54–63. doi: 10.1016/j.puhe.2006.05.024. 375  
376  
377
26. Song C., Ikei H., Miyazaki Y. Elucidation of a physiological adjustment effect in a forest environment: A pilot study. *Int. J. Environ. Res. Public Health.* 2015;12:4247–4255. doi: 10.3390/ijerph120404247. 378  
379
27. Victorson D, Doninger G, Victorson S, Victorson G, Hall L, Maletich C, Corr BR, Scortino K, Burns Z, Allen L, Rosa I, Quirk K, Adegbemi A, Strokoff J, Zuidema K, Sajdak K, Mckibben T, Roberts A, McDade TW, Boes A, McAlinden K, Arredondo K, Sauer C, Smith K, Salsman JM. Psychosocial and Biological Outcomes of Immersive, Mindfulness-Based Treks in Nature for Groups of Young Adults and Caregivers Affected by Cancer: Results from a Single Arm Program Evaluation from 2016–2021 *Int J Environ Res Public Health.* 2021 Dec; 18(23): 12622. doi: 10.3390/ijerph182312622. PMID: 34886348. 380  
381  
382  
383  
384  
385