


SUPPLEMENTARY DIGITAL MATERIAL 1

**Literature search strategies and PRISMA Flow Diagrams**

Supplementary Table I.—PICO 1 search strategies.


PubMed, searched on 01/06/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus celiacus" OR "truncus coeliacus"	196,368
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,556
#3	diagnosis	11,042,488
#4	CT angiography OR computed tomography OR computed tomography angiography	633,713
#5	#1 AND #2 AND #3 AND #4	6,577
#6	#1 AND #2 AND #3 AND #4 Filters: Clinical Trial, Comparative Study, Controlled Clinical Trial, Meta-Analysis, Observational Study, Randomized Controlled Trial, Review, Systematic Review, English, Italian	982
Cochrane Library, searched on 01/06/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus celiacus" OR "truncus coeliacus" (Word variations have been searched)	12,849
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms (Word variations have been searched)	5,592
#3	diagnosis (Word variations have been searched)	219,199
#4	CT angiography OR computed tomography OR computed tomography angiography (Word variations have been searched)	25,749
#5	#1 AND #2 AND #3 AND #4 in Cochrane Reviews and Trials	37



Supplementary Figure 1.—PICO 1 PRISMA flow diagram.

Supplementary Table II.—PICO 1 search strategies.


PubMed, searched on 16/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	196,307
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,278
#3	"Computed Tomography Angiography"[Mesh] OR computed tomography angiography OR "Magnetic Resonance Angiography"[Mesh] OR magnetic resonance angiography OR "Ultrasonography, Doppler, Duplex"[Mesh] OR ultrasonography doppler duplex OR "Angiography, Digital Subtraction"[Mesh] OR angiography digital subtraction	138,065
#4	screening OR prevention	7,980,053
#5	#1 AND #2 AND #3 AND #4	3,838
#6	#5 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	1,560
Cochrane Library, searched on 16/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	12,803
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,567
#3	computed tomography angiography OR magnetic resonance angiography OR ultrasonography doppler duplex OR angiography digital subtraction	4,608
#4	screening OR prevention	332,959
#5	#1 AND #2 AND #3 AND #4	33



Supplementary Figure 2.—PICO 2 PRISMA flow diagram.

Supplementary Table III.—PICO 3 search strategies.


PubMed, searched on 30/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	196,343
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,509
#3	associated disease OR concomitant disease OR comorbidity	4,544,676
#4	screening OR prevention	7,995,876
#5	#1 AND #2 AND #3 AND #4	8,398
#6	#1 AND #2 AND #3 AND #4 Filters: Clinical Trial, Comparative Study, Controlled Clinical Trial, Meta-Analysis, Observational Study, Randomized Controlled Trial, Review, Systematic Review, English, Italian	1,311
Cochrane Library, searched on 30/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus" (Word variations have been searched)	12,804
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms (Word variations have been searched)	5,567
#3	associated disease OR concomitant disease OR comorbidity (Word variations have been searched)	164,355
#4	screening OR prevention (Word variations have been searched)	332,968
#5	#1 AND #2 AND #3 AND #4 in Cochrane Reviews and Trials	93



Supplementary Figure 3.—PICO 3 PRISMA flow diagram.

Supplementary Table IV.—PICO 1 and 2 search strategies on renal artery aneurysms.

PubMed, searched on 17/05/2022		
Search	Query	Results
#1	"visceral artery"[Title] OR "visceral arteries"[Title] OR "renal artery"[Title] OR "renal arteries"[Title] OR "arteria renalis"[Title] OR (("arterias"[All Fields] OR "arteries"[MeSH Terms] OR "arteries"[All Fields] OR "arteria"[All Fields] OR "arteriae"[All Fields]) AND "renis"[Title])	9,585
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,289
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR reimplantations OR resection OR resections OR embolization OR embolizations OR aneurysmectomy OR aneurysmectomies OR bypass OR bypasses OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,197,583
#4	#1 AND #2 AND #3	1,785
#5	#4 NOT ("case report" OR "case reports" OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	660
Cochrane Library, searched on 17/05/2022		
Search	Query	Results
#1	visceral artery OR visceral arteries OR renal artery OR renal arteries OR arteria renalis OR arteria renis	5,166
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,011
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR reimplantations OR resection OR resections OR embolization OR embolizations OR aneurysmectomy OR aneurysmectomies OR bypass OR bypasses OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,453,624
#4	#1 AND #2 AND #3	293




Supplementary Figure 4.—PICO 1 and 2 PRISMA flow diagram on renal artery aneurysms.




Supplementary Table V.—PICO 1 and 2 search strategies on splenic artery aneurysms.

PubMed, searched on 24/05/2022		
Search	Query	Results
#1	splenic artery OR lienal artery OR (spleen AND artery)	11,791
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,414
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,190,840
#4	#1 AND #2 AND #3	2,012
#5	#4 NOT ("case report" OR "case reports" OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	622
Cochrane Library, searched on 24/05/2022		
Search	Query	Results
#1	splenic artery OR lienal artery OR (spleen AND artery)	311
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,567
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,469,946
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	16




Supplementary Figure 5.—PICO 1 PRISMA flow diagram on splenic artery aneurysms.



Supplementary Figure 6.—PICO 2 PRISMA flow diagram on splenic artery aneurysms.

Supplementary Table VI.—PICO 1 and 2 search strategies on celiac artery aneurysms.


PubMed, searched on 02/05/2022		
Search	Query	Results
#1	celiac artery OR coeliac artery OR celiac trunk OR truncus coeliacus OR celiac tripod OR coeliac tripod OR visceral	98,885
#2	endovascular OR open OR surgery OR surgical treatment OR emergent treatment OR urgent treatment OR elective treatment OR percutaneous OR stent OR stenting OR embolization OR coil OR coiling	6,547,722
#3	aneurysm OR pseudoaneurysm	180,208
#4	#1 AND #2 AND #3	4,114
Cochrane Library, searched on 19/05/2022		
Search	Query	Results
#1	celiac artery OR coeliac artery OR celiac trunk OR truncus coeliacus OR celiac tripod OR coeliac tripod OR visceral	228 reviews 47 protocols 6,778 trials 1 editorial 14 clinical answers
#2	endovascular OR open OR surgery OR surgical treatment OR emergent treatment OR urgent treatment OR elective treatment OR percutaneous OR stent OR stenting OR embolization OR coil OR coiling	8,419 reviews; 1,511 protocols; 454,171 trials; 84 editorials; 22 special collections; 2,884 clinical answers
#3	aneurysm OR pseudoaneurysm	294 reviews; 46 protocols; 5,052 trials; 2 editorials; 33 clinical answers
#4	#1 AND #2 AND #3	14 reviews 2 protocols 25 trials




Supplementary Figure 7.—PICO 1 and 2 PRISMA flow diagram on celiac artery aneurysms.

Supplementary Table VII.—Search strategies on gastropancreaticoduodenal artery aneurysms (PICO 1 and 2) and pseudoaneurysms (PICO 3 and 4).

PubMed, searched on 23/05/2022		
Search	Query	Results
#1	gastroduodenal OR gastro-duodenal OR pancreaticoduodenal OR pancreatico-duodenal	15,547
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,390
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,187,741
#4	#1 AND #2 AND #3	828
#5	#4 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	243
Cochrane Library, searched on 23/05/2022		
Search	Query	Results
#1	gastroduodenal OR gastro-duodenal OR pancreaticoduodenal OR pancreatico-duodenal	1,159
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,567
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,469,943
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	2




Supplementary Figure 8.—PICO 1 PRISMA flow diagram on gastropancreaticoduodenal artery aneurysms.




Supplementary Figure 9.—PICO 2 PRISMA flow diagram on gastropancreaticoduodenal artery aneurysms.






Supplementary Figure 10.—PICO 3 PRISMA flow diagram on gastropancreaticoduodenal artery pseudoaneurysms.




Supplementary Figure 11.—PICO 4 PRISMA flow diagram on gastropancreaticoduodenal artery pseudoaneurysms.

Supplementary Table VIII.—PICO 1, 2, 3, and 4 search strategies on hepatic artery aneurysms.


PubMed, searched on 10/05/2022		
Search	Query	Results
#1	hepatic OR liver OR "arteria hepatica"	1,309,705
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,150
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,145,515
#4	#1 AND #2 AND #3	3,932
#5	#4 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	1,554
Cochrane Library, searched on 10/05/2022		
Search	Query	Results
#1	hepatic OR liver OR "arteria hepatica"	74,269
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,011
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,453,344
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	160




Supplementary Figure 12.—PICO 1 PRISMA flow diagram on hepatic artery aneurysms.



Supplementary Figure 13.—PICO 2 PRISMA flow diagram on hepatic artery aneurysms.



Supplementary Figure 14.—PICO 3 PRISMA flow diagram on hepatic artery aneurysms.




Supplementary Figure 15.—PICO 4 PRISMA flow diagram on hepatic artery aneurysms.

Supplementary Table IX.—PICO 1 and 2 search strategies on mesenteric artery aneurysms.

PubMed, searched on 12/05/2022		
Search	Query	Results
#1	mesenteric artery OR mesenteric arteries	29,328
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,177
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,152,182
#4	#1 AND #2 AND #3	3,053
#5	#4 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	1,330
Cochrane Library, searched on 12/05/2022		
Search	Query	Results
#1	mesenteric artery OR mesenteric arteries	655
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,567
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,469,936
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	60







Supplementary Figure 16.—PICO 1 and 2 PRISMA flow diagram on mesenteric artery aneurysms.

Supplementary Table X.—PICO 1 and 2 search strategies on jejunal, ileal, and colic artery aneurysms.

PubMed, searched on 23/05/2022		
Search	Query	Results
#1	(jejunal arter*) OR (ileal arter*) OR (ileum arter*) OR (ileocolic arter*) OR (colic arter*)	7,023
#2	"endovascular procedures"[MeSH Terms] OR procedure* OR surg* OR technique* OR operation* OR intervention* OR therap* OR management* OR treatment OR treat* OR revascularization* OR reimplantation* OR resection* OR aneurysmectom* OR bypass* OR graft* OR prosthes* OR endovascular OR hybrid	17,823,701
#3	"aneurysm"[MeSH Terms] OR aneurysm* OR aneurism* OR "aneurysm, false"[MeSH Terms] OR pseudoaneurysm* OR pseudo-aneurysm*	181,881
#4	#1 AND #2 AND #3	452
#5	#1 AND #2 AND #3 Filters: English, Italian	375
Cochrane Library, searched on 23/05/2022		
Search	Query	Results
#1	((jejunal arter*) OR ((ileal arter*) OR (ileum arter*))) OR (ileocolic arter*) OR (colic arter*):ti,ab,kw (Word variations have been searched)	143
#2	(procedure* OR surg* OR technique* OR operation* OR intervention* OR therap* OR management* OR treatment OR treat* OR revascularization* OR reimplantation* OR resection* OR aneurysmectom* OR bypass* OR graft* OR prosthes* OR endovascular OR hybrid):ti,ab,kw (Word variations have been searched)	1,472,493
#3	((aneurysm*) OR (aneurism*) OR (pseudoaneurysm*) OR (pseudo-aneurysm*)):ti,ab,kw (Word variations have been searched)	5,147
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	2




Supplementary Figure 17.—PICO 1 PRISMA flow diagram on jejunal, ileal, and colic artery aneurysms.




Supplementary Figure 18.—PICO 2 PRISMA flow diagram on jejunal, ileal, and colic artery aneurysms.

Supplementary Table XI.—PICO 1 and 2 search strategies on isolated hypogastric artery aneurysms.

PubMed, searched on 12/05/2022		
Search	Query	Results
#1	hypogastric OR internal iliac	10,977
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,180
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	21,152,306
#4	#1 AND #2 AND #3	1,704
#5	#4 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	819
Cochrane Library, searched on 12/05/2022		
Search	Query	Results
#1	hypogastric OR internal iliac	994
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,567
#3	procedure OR procedures OR surgery OR surgeries OR technique OR techniques OR operation OR operations OR intervention OR interventions OR therapy OR therapies OR management OR treatment OR treatments OR correction OR corrections OR repair OR repairs OR revascularization OR revascularizations OR reimplantation OR resection OR resections OR aneurysmectomy OR bypass OR graft OR grafts OR prosthesis OR prostheses OR endovascular OR hybrid OR hybrids	1,469,935
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	80



Supplementary Figure 19.—PICO 1 PRISMA flow diagram on isolated hypogastric artery aneurysms.




Supplementary Figure 20.—PICO 2 PRISMA flow diagram on isolated hypogastric artery artery aneurysms.

Supplementary Table XII.—PICO 1 and 2 search strategies on medical therapy and follow-up.

PubMed, searched on 31/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	196,344
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	181,512
#3	"Computed Tomography Angiography"[Mesh] OR computed tomography angiography OR "Magnetic Resonance Angiography"[Mesh] OR magnetic resonance angiography OR "Ultrasonography, Doppler, Duplex"[Mesh] OR ultrasonography doppler duplex	130,315
#4	follow-up OR "follow up" OR surveillance OR monitoring	5,057,756
#5	#1 AND #2 AND #3 AND #4	1,604
#6	#5 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	976
Cochrane Library, searched on 31/05/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatico-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	12,849
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	5,592
#3	computed tomography angiography OR magnetic resonance angiography OR ultrasonography doppler duplex	4,265
#4	follow-up OR "follow up" OR surveillance OR monitoring	383,418
#5	#1 AND #2 AND #3 AND #4	42






Supplementary Figure 21.—PICO 1 and 2 PRISMA flow diagram on medical therapy and follow-up.

Supplementary Table XIII.—PICO 3 and 4 search strategies on medical therapy and follow-up.

PubMed, searched on 27/07/2022		
Search	Query	Results
#1	((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatoco-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"	197,271
#2	aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms	182,851
#3	antithrombotic OR anticoagulant OR antiplatelet OR heparin OR DOAC OR aspirin OR clopidogrel OR ticagrelor OR rivaroxaban OR apixaban OR dabigatran OR edoxaban OR statin OR betablockers OR "medical therapy" OR "drug therapy" OR "pharmacological therapy" OR "pharmacologic therapy" OR "medical treatment" OR "drug treatment" OR "pharmacological treatment" OR "pharmacologic treatment" OR "smoking cessation" OR antihypertensive OR angiotensin-converting enzyme inhibitors OR ACE-I OR angiotensin II receptor 1 antagonists OR ATIIR1 OR antidiabetic	3,482,874
#4	#1 AND #2 AND #3	1,531
#5	#4 NOT ("case report"[Title] OR "case reports"[Title] OR "case reports"[Publication Type] OR "letter"[Publication Type] OR "editorial"[Publication Type] OR "comment"[Publication Type]) Filters: English, Italian	663
Cochrane Library, searched on 27/07/2022		
Search	Query	Results
#1	((((visceral OR renal OR splenic OR lienal OR hepatic OR celiac OR coeliac OR gastric OR gastroepiploic OR mesenteric OR jejunal OR ileal OR iliac OR ilial OR colic OR pancreaticoduodenal OR gastroduodenal OR pancreatoco-duodenal OR gastro-duodenal OR hypogastric OR "internal iliac") AND (artery OR arteries)) OR "arteria coeliaca" OR "arteria gastrica" OR "arteria gastroduodenalis" OR "arteria gastroepiploica" OR "arteria hepatica" OR "arteria lienalis" OR "arteria mesenterica" OR "arteria renalis" OR "celiac axis" OR "celiac trunk" OR "coeliac axis" OR "coeliac trunk" OR "truncus coeliacus" OR "truncus coeliacus"):ti,ab,kw (Word variations have been searched)	11,450
#2	(aneurysm OR aneurysms OR microaneurysm OR microaneurysms OR pseudoaneurysm OR pseudoaneurysms):ti,ab,kw (Word variations have been searched)	5,311
#3	(antithrombotic OR anticoagulant OR antiplatelet OR heparin OR DOAC OR aspirin OR clopidogrel OR ticagrelor OR rivaroxaban OR apixaban OR dabigatran OR edoxaban OR statin OR betablockers OR "medical therapy" OR "drug therapy" OR "pharmacological therapy" OR "pharmacologic therapy" OR "medical treatment" OR "drug treatment" OR "pharmacological treatment" OR "pharmacologic treatment" OR "smoking cessation" OR antihypertensive OR angiotensin-converting enzyme inhibitors OR ACE-I OR angiotensin II receptor 1 antagonists OR ATIIR1 OR antidiabetic):ti,ab,kw (Word variations have been searched)	496,535
#4	#1 AND #2 AND #3 in Cochrane Reviews and Trials	98



Supplementary Figure 22.—PICO 3 and 4 PRISMA flow diagram on medical therapy and follow-up.

# Interpretation of selected evidence

## 1. DIAGNOSIS AND SCREENING

**PICO 1.1 Clinical question:** In patients with suspected aneurysm/pseudoaneurysm of visceral or renal artery, is Computed Tomography (CT) angiography more accurate than other modalities for the diagnosis and indication for treatment?

The literature search identified one non-analytical article <sup>8</sup> and five review articles <sup>9-13</sup> relevant to the proposed PICO. Although the included studies partially analyzed the same population of interest (patients with suspected aneurysm of the visceral or renal artery), they addressed considerably different aspects and did not draw a common conclusion regarding the accuracy of CT angiography. The five reviews, despite the lack of a dedicated design, suggested that CT angiography should be the first-choice imaging in both urgent and elective cases, due to its higher accuracy, lower invasiveness, and better ease of execution, compared with other imaging modalities.

**PICO 1.2 Clinical question:** Should patients diagnosed with visceral or renal artery aneurysms be screened with additional imaging to look for potential concomitant aneurysms in different locations to prevent complications, instead of forgoing further diagnostic examinations?

To date, no studies have found an answer to the proposed clinical question. There are no guidelines or studies supporting the need for additional investigations to detect concomitant aneurysms in other locations. CT angiography alone, widely employed nationwide, allows to diagnose and plan the treatment of the detected aneurysm. With the same exam, the presence of concomitant aneurysms in other locations within the same imaging scans can be evaluated. Given the widespread availability of this imaging modality, and the limited risks for the patients, the panel of experts decided to formulate a good clinical practice point recommendation in this regard.

**PICO 1.3 Clinical question:** Should patients with visceral or renal artery aneurysms be screened for associated underlying pathologies?

To date, literature has provided limited evidence in support of the screening for associated underlying pathologies. Despite analyzing the same population of interest (patients with visceral or renal artery aneurysms), the included studies addressed considerably different aspects, and did not draw a common conclusion regarding the possible screening for associated underlying conditions. One study, despite the low number of patients and the lack of a dedicated design, suggested that in individuals with visceral or renal artery aneurysms an ultrasound screening for popliteal aneurysms might be indicated <sup>14</sup>. The second selected article endorsed the assessment of fibromuscular dysplasia in patients with visceral or renal artery aneurysms <sup>15</sup>.

## 2. INDICATIONS AND TREATMENT OPTIONS

### 2.1 RENAL ARTERY ANEURYSMS

**PICO 2.1.1 Clinical question:** In a patient with an aneurysm/pseudoaneurysm of the renal arteries, when is it justified to propose a surgical/endovascular treatment compared to medical therapy/follow-up alone to improve the outcome?

There are no recent randomized controlled trials in the literature regarding the management of visceral aneurysms and renal arteries. The most significant meta-analysis and systematic review on the topic was published by Barrionuevo et al. in 2019 and analyzes all the works published between 1980 and 27 March 2017 concerning visceral and renal arteries aneurysms; this meta-analysis takes into consideration 1279 renal artery aneurysms, of which 358 treated with an endovascular approach and 921 treated with an open surgical approach<sup>16</sup>. The growth rate of renal artery aneurysms is very low (about 0.06 to 0.6 mm per year) and the natural history of aneurysms larger than 2 cm is associated with a low risk of rupture and a slow growth rate<sup>17-19</sup>. The most recent literature, indeed, has highlighted a threshold greater than or equal to 3 cm in diameter to give the indication for the elective treatment; however, in consideration of the characteristics of individual cases, treatment threshold may variate below 3 cm in diameter in the following cases: distal location, saccular morphology and rapid growth. From the analysis of selected studies, it emerged that the rupture of renal aneurysms is associated with a mortality equal to 10% in the general population, therefore emergency repair surgery is indicated in this case<sup>20, 21</sup>. Renal artery pseudoaneurysms, as well as all other visceral arteries pseudoaneurysms, have a higher risk of rupture than true aneurysms<sup>16</sup>, and for this reason literature generally recommends the treatment of a renal artery pseudoaneurysm regardless of its diameter. Pregnancy is associated with an increased risk of aneurysm rupture, and the maternal and fetal mortality rates associated with aneurysmal rupture reported in literature range from 56% to 84% and 82% to 100%, respectively. Evidence supports intervention in patients of childbearing age with renal artery aneurysm and acceptable operative risk even in case of diameters less than 3 cm<sup>22, 23</sup>.

**PICO 2.1.2 Clinical question:** Which intervention/procedure is preferable in terms of outcome in patients with renal artery aneurysm/pseudoaneurysm?

We identified a single systematic review and meta-analysis and two comparative cohort studies comparing endovascular treatment with open surgery<sup>16, 24, 25</sup>. Barrionuevo et al. and Gwon et al. showed substantial clinical similarity between endovascular and open surgery approaches, with only a reduced but not statistically significant rate of reinterventions in the open technique compared to the endovascular technique (0.16 [95% CI, 0.00-0.42] vs 0.03 [95% CI, 0.00-0.08])<sup>16, 24</sup>. Reintervention rate was substantially overlapping in the results reported by Li et al., as well as the rate of complications, and antihypertensive efficacy<sup>25</sup>. In terms of blood loss, duration of the procedure, intensive care unit stay, and hospitalization Li et al. reported the superiority of endovascular technique, although a lower reintervention rate seemed to characterize the open technique, which justifies a preferential surgical approach, especially considering that these patients are often young with a prolonged life expectancy<sup>25</sup>. Furthermore, literature showed satisfactory results regarding open surgery in the treatment of complex aneurysms<sup>26</sup>. On the other hand, the endovascular approach, when technically feasible, is preferred in patients with multiple comorbidities and a poor life expectancy<sup>17, 27, 28</sup>.

As demonstrated in literature, ex-vivo surgical technique is safe and effective in treating the majority of complex aneurysms<sup>29</sup>. Although outcomes were not significantly different when compared with open and endovascular surgery, autotransplant is more effective in preserving renal function<sup>24</sup>. Therefore, this technique should be preferred in complex aneurysms based on location or morphology, where prolonged ischemia time (>35 minutes) would expose to a higher risk of renal damage, and in patients with reduced renal function or those with a single functioning kidney<sup>24</sup>.

Literature review has shown encouraging results on the use of flow-modulating stents, micromesh stents, and in vivo overlapping stent technique. These therapeutic strategies have demonstrated vessel patency rates slightly below 94%, almost 90% complete sac thrombosis, and a 93.4% reduction/stabilization of the sac<sup>30</sup>. According to this systematic review and meta-analysis, the outcomes were generally reported for aneurysms of the abdominal visceral arteries (including 29 renal artery aneurysms out of a total of 225 visceral arteries aneurysms). Therefore, in selected situations based on anatomical characteristics, such as aneurysms or pseudoaneurysms involving the main branch or collateral branches that need to be preserved, the use of covered or uncovered stents should be taken into consideration<sup>30</sup>.

## 2.2 SPLENIC ARTERY ANEURYSMS

**PICO 2.2.1 Clinical Question:** When is it justified to propose surgical/endovascular treatment, compared with medical therapy/follow-up alone, to improve the outcomes of patients with a splenic artery aneurysm/pseudoaneurysm?

In almost all selected papers in case of ruptured aneurysms/pseudoaneurysms, authors have agreed that emergency surgical/endovascular treatment is recommended. In case of symptomatic aneurysms/pseudoaneurysms and in case of pseudoaneurysms, authors agree that surgical/endovascular treatment is recommended as soon as possible<sup>31-34</sup>. A systematic review and meta-analysis comparing management and outcome of open surgery versus endovascular treatment versus conservative treatment, conducted by Hogendoorn et al.<sup>35</sup>, based on data collected from 1321 patients in 47 articles, together with articles by other authors<sup>36,37</sup>, suggest active surveillance with both clinical and imaging-based follow-up in case of: a) splenic artery aneurysms less than 3 cm in diameter, b) dimensional stability at follow-up imaging, c) significant comorbidities, d) limited life expectancy. In a systematic review of 74 patients with splenic artery aneurysms, Batagini et al.<sup>38</sup> have highlighted that most splenic artery aneurysms remain stable and they have identified portal hypertension as the main risk factor for volume progression. In almost all selected papers, authors have indicated elective treatment in case of splenic artery aneurysms more than 3 cm in diameter<sup>16,35,36,39</sup>. Chaer et al.<sup>39</sup>, authors of the most recent American vascular surgery guidelines on the management of visceral aneurysms, propose observation based on a more controversial indication for the treatment of splenic artery aneurysms in asymptomatic patients ranging from 2 to 3 cm in diameter. On the other hand, Pulli et al.<sup>40</sup>, Venturini et al.<sup>41-43</sup>, Abbas et al.<sup>32</sup>, Wang et al.<sup>44</sup>, and Dorigo et al.<sup>45</sup> have suggested elective treatment. In the case of splenic artery aneurysms less than 2 cm in diameter, in the absence of other risk factors, elective treatment is suggested only in cases of demonstrated rapid volumetric growth<sup>16,39,46,47</sup>. The possible protective role of vessel wall calcifications is also being debated. According to Lakin et al.<sup>47</sup> and Sano et al.<sup>48</sup>, wall calcifications do have a protective role and may slow down aneurysm growth. However, such protective factor is questionable for other authors<sup>49</sup>. Because of the high risk of rupture, portal hypertension and liver transplant are significant objective elements in favour of treating splenic artery aneurysms, regardless of their size, as argued by Phan et al.<sup>50</sup> and Kobori et al.<sup>51</sup>. Another major risk factor for ruptured splenic artery aneurysms is pregnancy, as highlighted by Ha et al.<sup>52</sup> and Aung et al.<sup>53</sup>. Specifically, in a review on 32 ruptured aneurysms in pregnant women, Ha et al. report how the average diameter of half of the splenic artery aneurysms was less than 2 cm; however, according to other authors, such as Nanez et al.<sup>54</sup>, the risk of ruptured splenic artery aneurysms in pregnancy is relatively rare. The timing of possible treatment in pregnancy of splenic artery aneurysms is still a matter of debate. Surgical treatment is generally suggested during the second trimester, when embryogenesis is completed, though the size of the uterus has not yet precluded exposure of the aneurysm<sup>52</sup>. In the case of detection of splenic artery aneurysms in a young woman of childbearing age, close imaging monitoring (Computed Tomography, Magnetic Resonance Imaging, Doppler ultrasound) and treatment in case of rapid volumetric growth are suggested<sup>16,39,55-57</sup>.

**PICO 2.2.2 Clinical Question:** In a patient with a splenic artery aneurysm/pseudoaneurysm, which intervention/procedure should be carried out in order to obtain better outcomes?

In case of emergency/urgency (ruptured, symptomatic splenic artery aneurysms, pseudoaneurysms), the choice between surgical and endovascular treatment depends on many factors, including the feasibility of the two treatment options, which is also influenced by a) logistics, b) the type of splenic artery aneurysm (true or pseudoaneurysm), c) its location, d) its morphological features, e) patient's age, and f) patient's comorbidities. In general, the endovascular option is considered to be safer in terms of lower morbidity, mortality, and hospitalization time rates. Mortality rates and treatment costs are lower in both ruptured and intact splenic artery aneurysm treatments. In a systematic review and meta-analysis carried out by Hogendoorn et al.<sup>35</sup>, the mortality rate 30 days after surgical treatment versus endovascular treatment was found to be 5% versus 0.6% respectively, in case of intact splenic artery aneurysms (9% vs 2%, in case of ruptured cases). Also, the average hospitalization time after surgery was found to be significantly higher than after endovascular treatment (9.8 vs 2.0 days). Considering splenic artery aneurysms elective treatment, Hogendoorn et al.<sup>58</sup> pointed out, in another systematic review, that compared with open surgery, overall costs are lower in case of endovascular treatment (-\$3,384), which, however, has higher reintervention rates. Therefore, whenever possible, also based on clinical, anatomical, multidisciplinary, and logistic evaluations, endovascular treatment is recommended over surgical treatment because it is less invasive, it has lower morbidity and mortality rates, lower costs and shorter hospitalization times, both in elective treatment and in emergency settings<sup>31,42,43,59-66</sup>, even though surgical treatment has lower reintervention rates. In case of splenic artery aneurysms during pregnancy, the choice is more debated, partly because of issues related to the impact of radiation on the foetus<sup>52,53,67</sup>. In case of hilar splenic artery aneurysms, the choice of treatment is also debated. Hilar aneurysms may be treated endovascularly only by means of embolization with sacrifice of the target vessel if the aneurysms are fusiform, with a consequent higher risk of splenic infarction<sup>41,68,69</sup>. Again, based on clinical, anatomical, multidisciplinary, and logistical evaluations, although surgical treatment has been advocated for the treatment of giant

splenic artery aneurysms more than 5 cm in diameter, especially in case of compressive effect<sup>70</sup>, the endovascular option may also be considered<sup>71</sup>.

### 2.3 CELIAC ARTERY ANEURYSMS

**PICO 2.3.1 Clinical question:** In patients with celiac artery aneurysm/pseudoaneurysm, when is surgical and/or endovascular intervention indicated against medical therapy alone to reduce the risk of rupture?

In patients with celiac artery aneurysms, data collected indicate that intervention is recommended in asymptomatic patients when aneurysm size is 2 cm or greater<sup>16, 38, 72-94</sup>. Symptomatic celiac tripod aneurysms should be treated regardless of their size. Pseudoaneurysms of the celiac artery should always be carefully evaluated and, also in this case, treated regardless of their size<sup>16, 38, 72-92</sup>. In a single paper in which medical therapy was indicated (wait-and-see), aneurysms growth was almost zero at one-year follow-up<sup>38</sup>. In asymptomatic patients with known celiac artery aneurysms smaller than 2 cm in size, follow-up may be indicated and treatment should be performed when aneurysm size is greater than or equal to 2 cm. Literature suggests the treatment of celiac artery aneurysms in special cases as well (women of childbearing age, patients in whom liver transplantation is planned, etiology), even if without reporting a dedicated case history<sup>73, 75, 80</sup>. Regardless of the proposed intervention, data suggest a clinical benefit to patients in terms of quality of life, yet risks associated with surgery must be weighed against the possible benefit of conservative treatment considering the high risk of rupture and patient comorbidities<sup>88, 89</sup>. The overall number of patients evaluated allows a rather reliable assessment regarding this PICO and the data obtained from the authors' analysis was considered quite relevant to the target population. The risk of bias is considered unclear.

**PICO 2.3.2 Clinical question:** In patients with celiac artery aneurysm/pseudoaneurysm, with an indication for intervention, is endovascular treatment more suitable than open surgery to improve clinical success?

Many of the narrative reviews evaluated both procedures<sup>73, 79-84, 86, 93, 94</sup>, with only a couple of observational studies reporting a direct comparison between endovascular and open treatment<sup>85, 95</sup>; most observational studies reported results regarding a single treatment option<sup>72, 74, 75-77, 79, 96-103</sup>, whilst others were simple narrative reviews<sup>78, 87, 104, 105</sup>. A single systematic review compared the mortality and complications of the two techniques, commenting on the absence of direct comparisons<sup>16</sup>. Despite the heterogeneity of all these studies, the data obtained from the authors' evaluation were considered quite relevant to the target population. The risk of bias was considered unclear/low. The analysis performed on the population with celiac artery aneurysms showed that endovascular intervention may represent the first choice, especially in cases of favorable anatomy or severe comorbidities<sup>16, 72, 75, 82, 87, 94, 96, 102</sup>. The open surgical option might be considered the best option in patients with hostile anatomy, in case of failure of endovascular treatment or in emergency situations and in cases with severe hemodynamic instability<sup>16, 73, 77, 79, 81, 83-85, 93, 95, 99</sup>. Although there was a substantial overlap in efficacy in the medium term between the two interventions, the endovascular treatment offers the benefit of shorter hospital stay and a reduced number of postoperative complications over open surgery. The most severe complications, although statistically not significant, include possible organ ischemia secondary to stent or bypass occlusion or potential bleeding following artery rupture, regardless of the type of approach used<sup>16, 75, 76, 79, 82, 88-90, 97</sup>. Considering the greater invasiveness of the procedure itself, open surgical intervention has a potentially more severe impact on patients' lives, particularly on districts such as the intestinal tract (e.g., transit delay) and cardiac district (e.g., myocardial ischemia, heart failure)<sup>77, 85, 100</sup>. Regardless of the type of intervention, celiac artery revascularization using stents or bypass should be preferred versus ligation or embolization to preserve function<sup>81, 84, 93</sup>. Although complication rates differed considerably between the two interventions, treatment of celiac artery aneurysms is acceptable to both the patient and family members.

### 2.4 GASTROPANCREATICODUODENAL ARTERIES ANEURYSMS

**PICO 2.4.1 Clinical question:** In patients with aneurysms of the gastropancreaticoduodenal arteries, when is it justified to propose a surgical/endovascular treatment rather than medical therapy/follow up alone to improve the outcomes?

There are no randomised controlled trials responding to the PICO question. Among the most recent articles, two reviews<sup>106, 107</sup> and two case series<sup>108, 109</sup> are available. The analysed studies present overlapping results and conclusions. Specifically, it was found that the risk of rupture of aneurysms of the gastropancreaticoduodenal arteries was independent from their size and that the technical success of both endovascular and open elective treatment was satisfactory (90% according to Vandy et al.<sup>106</sup>, 100 % according to Michalinos et al.<sup>107</sup>). Therefore, a unanimous opinion emerged regarding the elective treatment of aneurysms of the gastropancreaticoduodenal arteries regardless of their size.

**PICO 2.4.2 Clinical question:** In patients with aneurysms of the gastropancreaticoduodenal arteries, which intervention/procedure is preferable in terms of outcomes?

Three reviews and three case series relative to the PICO question have recently been published. A first review from 2019 (Michalinos et al.<sup>107</sup>) underlines that the endovascular approach is characterised by a lower mortality and morbidity rate compared to the open treatment, even if technical success was similar. Furthermore, several procedures reveal that the endovascular treatment was proposed, when feasible, as the first-choice option, since it is characterized by a lower rate of complications, even if often subjected to a higher need for reinterventions<sup>16, 109</sup>. Vandy et al. report a very low mortality rate for patients with aneurysms of the gastropancreaticoduodenal arteries, relating it to the increasingly frequent use of endovascular treatments for incidentally found aneurysms<sup>106</sup>. Among the analysed studies, no one suggests the systematic revascularisation, either open or endovascular, of the coeliac artery when its stenosis is found as a possible aetiological cause of the aneurysm itself<sup>108, 109</sup>. On the other hand, the panel agrees in suggesting, as a good practice point, the revascularisation of the gastropancreaticoduodenal aneurysm when it is also associated with obstructive lesions affecting all three intestinal arteries (coeliac, superior and inferior mesenteric arteries). Finally, the authors agree in not contraindicating the use of the open surgical approach, although recognising the advantages of the endovascular technique, as already mentioned. In this sense, Bonardelli et al.<sup>110</sup> suggest a therapeutic approach to be customised according to the patient's characteristics, diagnosis, timing, and vascular anatomy.

**PICO 2.4.3 Clinical question:** In patients with pseudoaneurysms of the gastropancreaticoduodenal arteries, when is it justified to propose a surgical/endovascular treatment rather than medical therapy/follow up alone to improve the outcomes?

There are no randomised controlled trials responding to the PICO question. We identified two articles to make the recommendations, as they were adequate in terms of study design and responded to the PICO question. Specifically, a retrospective observational study<sup>110</sup> and a narrative review<sup>111</sup> were identified. They showed a weak scientific relevance (level of evidence 4) in relation to the study design and the few cases reported; therefore, the risk of bias might be considered. However, the studies are relevant for the target population and report similar results and conclusions. Although no relation has been reported between pseudoaneurysm size and risk of rupture, based on experts' opinions and on the aforementioned articles, both open and endovascular solutions are acceptable, in order to control bleeding in emergencies and to avoiding pseudoaneurysm rupture in the elective setting. The technical success of the interventions is satisfactory (90% for the endovascular treatment and 83% for the open treatment according to Bonardelli et al.<sup>110</sup>). Therefore, although with limited data, a unanimous opinion emerges regarding the need for treatment of pseudoaneurysms of the gastropancreaticoduodenal arteries regardless of size, both in emergencies and electively. Clinical trials comparing treatment and surveillance of non-ruptured pseudoaneurysms are desirable in order to identify whether there is a size threshold or a morphologic aspect for which the treatment-related risks do not outweigh the risks of pseudoaneurysm ruptures.



**PICO 2.4.4 Clinical question:** In patients with pseudoaneurysms of the gastropancreaticoduodenal arteries, which intervention/procedure is preferable in terms of outcomes?

The literature research led to the selection of eight articles for the formulation of the recommendations, as they were adequate in terms of study design and responded to the PICO question. Specifically, five retrospective observational studies<sup>110, 112-115</sup>, a narrative review<sup>111</sup>, a case series<sup>116</sup> and a cohort study<sup>117</sup>. Overall, all of them had a low methodological relevance in terms of study design and number of cases, therefore biases might exist. Furthermore, some of the reported outcomes do not specifically refer to the target population but include more generic outcomes that consider pseudoaneurysms of visceral arteries. Despite that, the evaluated studies are relevant for the target population and report similar results and conclusions. As reported in a retrospective observational study from Bonardelli et al.<sup>110</sup>, the endovascular treatment is to be preferred as a first approach if the location, size and anatomy of the vessel make the treatment feasible. Surgical treatment is to be preferred when the patient is haemodynamically unstable, if visceral resection is required, or in case the endovascular treatment is contraindicated or might have a high chance of failure. Referring to the narrative review by Kallamadi et al.<sup>111</sup>, the endovascular treatment has some advantages, such as less invasiveness and complications than surgical treatment. It allows a better pseudoaneurysm evaluation and selective embolization, sparing other vessels. It can be associated with the treatment of celiac stenosis when present. However, as also stated by Murata et al.<sup>115</sup>, the surgical treatment remains a viable option even after failure of endovascular treatment.

Gupta et al.<sup>116</sup> described some cases of bleeding gastropancreaticoduodenal arteries pseudoaneurysms in patients with chronic pancreatitis, suggesting that the endovascular treatment is indicated in most patients and may involve the use of thrombin if the aneurysm neck is narrow enough (less than 2 cm). Surgery is to be preferred depending on the location of the pseudoaneurysm and the underlying disease process. Clinical trials comparing surgical and endovascular treatment of ruptured pseudoaneurysms as well as electively treated pseudoaneurysms are still necessary to identify potential advantages of one over the other treatment options in terms of technical and clinical success.

## 2.5 HEPATIC ARTERY ANEURYSMS

**PICO 2.5.1 Clinical question:** In patients with aneurysms/pseudoaneurysms of the hepatic artery, when is it justified to propose a surgical/endovascular treatment rather than medical therapy/follow up alone to improve the outcomes?

According to literature, hepatic artery pseudoaneurysms account for 25% to 80% of reported cases and often occur after traumatic or iatrogenic traumas<sup>118, 119</sup>. Patient's clinical history and specific imaging findings allow to distinguish false aneurysms from true aneurysms. Imaging findings specific for pseudoaneurysms are focal arterial disruptions in a normal arterial setting and hematoma formation with inflammatory changes around the pseudoaneurysmatic sac. Most pseudoaneurysm cases are symptomatic at presentation with gastrointestinal bleeding and/or haemobilia, therefore differing from true aneurysm presentation<sup>76</sup>. Regarding pseudoaneurysm treatment, there are no randomized controlled trials, but only cohort studies are available. Due to a high propensity for rupture and significant disease-related mortality, treatment of all hepatic artery pseudoaneurysms is recommended, regardless of cause, as soon as diagnosed. Due to the rarity of these aneurysms, the natural history of hepatic artery aneurysms is unknown, making any recommendation for asymptomatic aneurysms controversial. The literature review process found mostly cohort studies, with low level of evidence given the retrospective setting. It was found that in the population with ruptured aneurysms the probability of rupture was higher when the diameter was  $>2$  cm<sup>38, 76, 118-123</sup>. Abbas et al. observed that, in a retrospective cohort of 36 patients with hepatic artery aneurysm managed non-operatively for 68 months, aneurysms enlarged in only 27% of cases, without evidence of any complications<sup>122</sup>. Given the high morbidity and mortality rate after hepatic artery aneurysm rupture (30% mortality rate described in one series<sup>122</sup>) and the low morbidity and mortality rate after elective hepatic artery aneurysm repair (0% mortality rate described in the same study<sup>122</sup>), the elective treatment is recommended in patients with aneurysms  $>2$  cm or with a high increasing size rate ( $>0.5$  cm/years), having been taken into account patient's comorbidities and life expectancy. Stark et al. demonstrated a 10-times higher incidence of rupture in women when compared to the male population, with a low rate of clinically asymptomatic patients (19% women versus 81% men). Moreover, female patients that presented with hepatic artery aneurysm rupture were older (more than 60 years of age)

and with a mean aneurysm diameter of  $2.5 \pm 1.2$  cm<sup>120</sup>. The presence of higher rupture rates in the female population requires indeed further investigations. Since hepatic artery aneurysms of non-atherosclerotic origin show higher rupture rate (incidence of 60% in patients with vasculitis and 50% in both Ehlers-Danlos syndrome type IV and alpha-1-antitrypsin deficiency patients), the elective treatment is recommended in patients with hepatic artery aneurysms even in case of diameters inferior to 2 cm<sup>124, 125</sup>.

**PICO 2.5.2 Clinical question:** In patients with an aneurysm/pseudoaneurysm of the hepatic artery, which intervention/procedure is preferable in terms of outcomes?

Hepatic artery aneurysms can be treated both open and endovascularly. The literature review found two systematic reviews and several observational cohort studies. The reviewed studies showed comparable long-term outcomes when treating hepatic artery aneurysms through open and endovascular repair; however, postoperative morbidity was significantly worse in case of open rather than endovascular approach. Since endovascular techniques improved and the morbidity rates associated with this type of treatment were relatively lower, in case of favorable anatomy the endovascular option should be preferentially offered to patients<sup>16, 126</sup>.

**PICO 2.5.3 Clinical question:** In patients with extra-hepatic aneurysms/pseudoaneurysms suitable for repair, is it preferable to maintain hepatic arterial circulation over endovascular vessel ligation/closure to avoid hepatic necrosis?

The literature review highlighted several systematic reviews and cohort studies, with a prevalence of observational studies. The ideal procedure should consist in aneurysm exclusion preserving liver circulation; this result could be obtained through stent-graft placement/coils embolization with endovascular exclusion of the aneurysm or by aneurysm resection and graft interposition. The use of stent-graft is mainly limited by anatomical factors, including a favorable anatomical setting that allows endovascular access to the aneurysm, an adequate "sealing zone" both proximal and distal and the absence of collateral pathways originating along or from the aneurysm itself<sup>16, 126, 127</sup>. In case of low-risk patients in which stent-graft placement might not be feasible, despite possible adequate collateral flow might be obtained through an endovascular exclusion as well, consider open surgical repair through autologous vein conduit to lower the chance of central necrosis<sup>128</sup>. In case of demonstrated patency of the pancreaticogastroduodenal arterial system, hepatic artery aneurysms located proximally to the origin of gastro-duodenal artery and to the right gastric artery could be considered for coil embolization with complete vascular exclusion of the main vessel<sup>129</sup>.

**PICO 2.5.4 Clinical question:** In patients with intra-hepatic aneurysms/pseudoaneurysms suitable for repair, is it preferable the endovascular treatment over surgical lobe resection to preserve hepatic function?

The literature review found several systematic reviews and observational cohort studies. Intra-hepatic aneurysms require the resection of the entire lobe where the aneurysm is located; however, given the numerous comorbidities associated with liver resection surgery, an endovascular approach, if anatomically feasible, would be a valuable treatment strategy for this type of vascular lesions. Complications related to intra-hepatic artery aneurysm embolization include liver ischemia, abscess/biloma formation, cholecystitis and possible recanalization<sup>16</sup>. The endovascular treatment is not recommended in patient with giant aneurysms involving a whole liver segment and/or lobe, given the high chance of massive liver necrosis related to the endovascular treatment; in these patients, the hepatic resection should be considered<sup>130</sup>.

## 2.6 MESENTERIC ARTERIES ANEURYSMS

**PICO 2.6.1 Clinical question:** In patients with a mesenteric artery aneurysm/pseudoaneurysm, when is it justified to propose surgical/endovascular treatment compared to medical therapy/follow-up alone to improve the outcomes?

Knowledge regarding the natural history of superior and inferior mesenteric artery aneurysms primarily derives from case series or retrospective cohort studies. There are no randomized studies comparing a conservative approach to corrective intervention. Another limitation of the existing literature is the frequent grouping of mesenteric artery aneurysms with cohorts of patients with aneurysms of other visceral arteries. Generally, the indication for surgical or endovascular treatment of a mesenteric artery aneurysm is justified when the risks associated with the natural history of the pathology (specifically, the risk of rupture) outweigh the risks associated with the intervention. True aneurysms of the mesenteric arteries can be asymptomatic or symptomatic with abdominal pain. Less common symptoms include nausea and gastrointestinal bleeding. The natural history of asymptomatic aneurysms is characterized by a tendency to increase in size potentially leading to rupture<sup>131</sup>. The evidence is consistent in concluding that the presence of asymptomatic true aneurysms of the superior or inferior mesenteric artery measuring >20 mm is an indication for treatment<sup>131, 132</sup>. One study<sup>133</sup> suggested using a cut-off diameter of 25 mm for intervention. Only one study conducted on a series of 21 superior mesenteric artery aneurysms suggested treatment regardless of size<sup>134</sup>. It is worth noting that recommendations regarding the cut-off diameter for treatment are primarily based on observations of the diameter of aneurysms that rupture at the time of presentation, as well as the high rate of major complications or mortality (38-58%) for emergency interventions compared to elective interventions (0-10%)<sup>133, 134</sup>. When comparing invasive treatment to monitoring, the analysed studies have a significant selection bias as larger aneurysms are predominantly treated. Furthermore, existing studies are unable to differentiate the natural history of saccular morphology aneurysms from fusiform ones. The presence of symptoms is usually associated with larger aneurysms, rapid growth, or rupture. Therefore, the presence of symptoms or signs of rupture represents a medical emergency that places the patient at immediate life-threatening risk and requires prompt correction. Mycotic aneurysms affecting the superior mesenteric artery are relatively common and are described in 15-20% of mesenteric artery aneurysm cases<sup>133-135</sup>. The most frequent cause is bacterial endocarditis. Dissecting aneurysms are characterized by isolated dissection of the superior mesenteric artery (SISMAD)<sup>136</sup>, which can progress to artery dilation. Mycotic<sup>137</sup> and dissecting aneurysms are considered to be at higher risk of rupture than asymptomatic true aneurysms; therefore, treatment is warranted in all cases regardless of diameter. Pseudoaneurysms of the mesenteric arteries can result from inflammatory/infectious processes (e.g., acute pancreatitis), iatrogenic or accidental trauma. Pseudoaneurysms carry a higher risk of complications compared to true aneurysms, therefore their treatment is indicated in all cases regardless of size<sup>135</sup>.

**PICO 2.6.2 Clinical question:** In patients with a mesenteric artery aneurysm/pseudoaneurysm, which intervention/procedure is preferable in terms of outcomes?

There are no randomised controlled trials in the literature comparing treatment strategies (open versus endovascular surgery) for mesenteric artery aneurysms. From the literature search, the meta-analysis and systematic review published by Barrionuevo et al.<sup>16</sup> was selected, which analyses 80 observational studies, with a total of 2845 visceral aneurysms and pseudoaneurysms of which 95 were of the superior mesenteric artery, 38 treated with endovascular approach, 57 with open surgery. Despite the small number of comparative studies, the data on short- and long-term mortality and perioperative complications are in favour of endovascular surgery as the first-line treatment choice. On the other hand, the rate of reinterventions was lower after open surgery. Five retrospective cohort studies<sup>130, 133, 138-140</sup> were identified, in overall agreement on results and conclusions: endovascular treatment is safe and effective, could be the treatment of choice in emergencies<sup>139</sup> and, when technically feasible, should be preferred as it is less invasive and associated with shorter hospital stays<sup>140</sup>. The literature search also identified nine observational studies/case series<sup>92, 134, 137, 141-146</sup> with low level of evidence, non-comparative and with small sample sizes; in some cases, the results reported aggregate data relating to all visceral aneurysms, not only specific for the mesenteric arteries. Despite these limitations, the studies tend to agree in their conclusions. The technique of choice in endovascular treatment should be left to the discretion of the operator: in the selected studies, the technical success was higher when using bare stent (dissecting aneurysm)<sup>142</sup>, covered

stent, coiling, double lumen balloon-assisted<sup>141</sup> and overlapping bare metal stent<sup>146</sup>. Endovascular treatment and open surgery are both characterised by high technical success and all patients with aneurysm of the superior mesenteric artery fit for repair can be considered for surgery<sup>134</sup>. Open surgery (ligation, aneurysmectomy and bypass) has acceptable morbidity and mortality<sup>137, 138</sup> and can be considered in cases of unfavourable anatomy, failure of endovascular treatment and as primary treatment in mycotic aneurysms.

## 2.7 JEJUNAL, ILEAL AND COLIC ARTERIES ANEURYSMS

**PICO 2.7.1 Clinical question:** In a patient with an aneurysm/pseudoaneurysm of the jejunal, ileal or colic artery, when is surgical/endovascular treatment indicated versus medical therapy or watchful waiting to improve outcome?

Five of the six publications selected are narrative reviews<sup>147-151</sup>, reporting clinical cases published in a very wide period ranging from 1937 to 2003. The observational study considered<sup>152</sup> includes patients treated from 1980 to 1998. No more recent publications including >5 patients responding to the PICO are available. Aneurysms of the colic arteries are the most commonly described (4 articles<sup>147, 148, 150, 151</sup> specifically for colic arteries); one article is specific for the jejunal arteries<sup>149</sup>; ileal aneurysms are described only in one publication<sup>152</sup>. Differentiation between aneurysms and pseudoaneurysms is difficult on the basis of the preoperative diagnostic investigations and some series do not differentiate them. In addition, the diameter of the aneurysm is not always reported. One article<sup>148</sup> includes all patients with segmental arterial mediolysis, while two other articles<sup>147, 152</sup> report more varied conditions: cystic dysplasia of the media, panarteritis nodosa, rheumatoid arthritis, Marfan syndrome, systemic lupus erythematosus, alpha-1 antitrypsin deficiency, tuberculosis. Colic aneurysms are the most reported (108 of 123), and several of them presented with rupture or shock regardless of their diameter.

**PICO 2.7.2 Clinical question:** In a patient with an aneurysm/pseudoaneurysm of the jejunal, ileal or colic artery, which intervention/procedure is preferable in terms of outcome?

The two publications considered are narrative reviews<sup>147, 149</sup>, reporting clinical cases published over a very wide time period (from 1937 to 2002). There are no more recent publications including >5 patients responding to the PICO. Considering the lack of recent articles included in the study, endovascular treatment of aneurysms may be underestimated compared to its current use in clinical practice. The aneurysms considered are colic or jejunal<sup>147, 149</sup>; none of the included articles consider ileal aneurysms. Differentiation between aneurysms and pseudoaneurysms is difficult on the basis of preoperative diagnostic investigations and some series do not differentiate the two pathologies.

The endovascular procedure has the advantage of avoiding laparotomy, thus for its lower invasiveness, it potentially reduces complications. For jejunal and colic aneurysms, the endovascular intervention could avoid procedures of bowel resection and its complications if the aneurysm develops in the proximity of the bowel wall. In the considered literature<sup>147, 149</sup>, no deaths were reported in the cases treated with endovascular techniques (0 of 7), while 3 of 39 patients died after open surgery. Although these data are scarce, they stress the less invasiveness of the endovascular treatment.

## 2.8 ISOLATED HYPOGASTRIC ARTERY ANEURYSMS

**PICO 2.8.1 Clinical question:** In patients with an isolated hypogastric artery aneurysm, when the endovascular/surgical treatment strategy is justifiable, instead of the conservative management/follow-up, to improve the outcomes?

The systematic review of the literature did not provide any prospective cohort study which investigated the natural history of isolated hypogastric artery aneurysms focusing in particular on their risk of rupture related to diameter or growth rate.

Traditionally, the threshold for treatment was 3 cm of diameter, following some observational studies that reported no ruptures for diameters below that value. However, those studies did not discriminate between aneurysms of the common iliac, external iliac or hypogastric arteries, or between isolated aneurysms or aneurysms associated with other iliac or abdominal aortic segments. This selection bias is common to most of the studies identified <sup>153-173</sup>, which provided surrogate data, being the information on diameters and rupture risk of isolated hypogastric artery aneurysms scarce and not easily extractable. In a study by Richardson et al. <sup>153</sup>, the authors indeed took generically into account aneurysms of the iliac arteries, and described as “aneurysm” dilatations with a diameter of 2.5 cm or above, not explaining the rationale behind this choice. Regarding the size cut-off between the elective treatment and the follow-up, the diameter set as reference in the discussion is still indicated at 3 cm. A study from Kliewer et al. <sup>154</sup> instead, examined patients with aneurysms of the hypogastric artery alone, isolated or in association with other aneurysms, and among the inclusion criteria there was a diameter of 1.6 cm or above, still not providing reasoning for this choice. Anyway, the authors stated that 1.6 cm was not the suggested cut-off for treatment indication, as they concluded that the critical dimensions were between 3 and 4 cm (the mean diameter of aneurysms treated in their study was 38.2 mm indeed). Smaller aneurysms were treated during the same endovascular procedure only if in association with aorto-iliac aneurysms. Considering the definition of a true arterial aneurysm as a permanent dilatation of a vessel above 50% of its normal diameter, the studies cited above <sup>153, 154</sup> demonstrate how a bias could still exist in the definition of aneurysm when it is isolated to the hypogastric artery. The reference measurements are in fact borrowed from other iliac districts, particularly from the common iliac artery that has an average diameter of about 1 cm, whereas for the hypogastric artery it is commonly 0.5 cm. In other two studies (Gao et al. <sup>155</sup> and Muradi et al. <sup>156</sup>) investigating isolated hypogastric artery aneurysms, the key-points considered indications for an elective treatment were: diameter of 3 cm or above and rapid growth, and a critical growth was considered as being an increase of 0.5 cm in size in 6 months <sup>155, 156</sup>, or 1 cm in 12 months <sup>156</sup>. These studies did not specify their rationale, as they just cited prior publications which almost exclusively included common iliac arteries aneurysms; hence, their conclusions are not applicable for the hypogastric arteries as well. A study from Laine et al. <sup>157</sup> included patients with isolated hypogastric artery aneurysms and other dilatations of the aorto-iliac axis, aiming to explain which the diameter of rupture of these aneurysms was and if it was proper considering 3 cm as being the cut-off. It was a retrospective analysis on 63 patients from 28 hospitals in 7 different countries. In this cohort, only 18 patients had an isolated hypogastric artery aneurysm, and all the patients were treated for a ruptured aneurysm of the hypogastric artery. The mean diameter of the ruptured aneurysms was 6.8 cm, with no significant difference between men and women. Considering only hypogastric artery aneurysms rupture, the mean diameter was 6.1 cm, while the diameter of non-isolated aneurysms of the hypogastric artery was 7.2 cm. The authors reported only one case of rupture with a diameter below 3 cm (1.6% of all ruptured aneurysms) and four cases with a diameter of less than 4 cm (6% of all ruptured aneurysms). Moreover, the reported 30-day mortality rate was 12.7%. Results from this study indicated that the risk of rupture of isolated hypogastric artery aneurysms of less than 4 cm of diameter could be similar to those of abdominal aortic aneurysms with a diameter less than 5.5 cm. However, an important difference emerged: data about abdominal aortic aneurysms come from multiple randomized control studies, with a level of evidence 1; whereas in this case we have a level of evidence 3. The authors emphasized how heterogeneous the measurements of aneurysm diameters were at the time of rupture, thus maybe underestimating the real dimensions compared to measurements taken pre-rupture. This condition suggested a less aggressive approach in the management of isolated hypogastric artery aneurysms, in which the treatment could be postponed until the diameter reaches 4 cm, which might be considered safe especially in elderly patients with a higher surgical risk. In a recent retrospective cohort study (Chen Rj et al. <sup>158</sup>) conducted on 23 patients with 31 hypogastric artery aneurysms, of whom 6 isolated hypogastric artery aneurysms, the authors reported that 4 out of 5 ruptured aneurysms showed a diameter greater than 4 cm, not specifying if this applied to isolated or combined aneurysms. Moreover, 6 aneurysms which were followed up during time showed an average annual growth rate of 2.4 mm. However, these data are not enough to motivate a less aggressive approach towards aneurysms within 4 cm of diameter, and to draw conclusions about their annual growth rate, because the annual growth rate sets around 1-4 mm/year, matching the growth rate shown by abdominal aortic aneurysms. The recent meta-analysis by Perini et al. <sup>174</sup>, analyzing studies dealing specifically with isolated hypogastric artery aneurysms (overall 192 patients with 202 aneurysms) applied the definition of aneurysm as a dilatation with a diameter of at least 0.8 cm. From this meta-analysis emerged the recommendation for a conservative management in case of patients with aneurysms of less than 3.5 cm diameter and less than 4 cm in the elderly. In those cases with a diameter of less than 3 cm, treatment might be considered only in young subjects and those patients with concomitant indication for other aorto-iliac aneurysms.

**PICO 2.8.2 Clinical question:** Which is the type of intervention/procedure to be preferred for patients with isolated hypogastric artery aneurysms?

Even if current literature is limited in terms of numbers, procedural standardization and outcomes (results of isolated iliac aneurysms, common iliac and hypogastric arteries are usually reported indistinctly), the endovascular solution represents

the first treatment option since it offers a higher technical success as well as better early and mid-term clinical success than open surgery. In a recent systematic literature review and meta-analysis, Perini et al.<sup>174</sup> analyze the results of 13 studies on surgical or endovascular repair of isolated hypogastric artery aneurysms. Out of 193 patients and 202 aneurysms, the results after endovascular and open repair were comparable in terms of technical success (open: 95% versus endovascular: 90%) with an overall incidence of buttock claudication of 14%. However, this complication was not reported in case of endovascular aneurysm repair consisting of hypogastric artery revascularization. Endovascular repair was associated to lower perioperative mortality (open: 8% versus endovascular: 3%) and shorter hospital stay (mean days: Open 13 versus endovascular 4) than open repair. The endovascular repair of isolated hypogastric artery aneurysms can be considered safe and effective even in emergency setting. In a single-center experience, Kobe et al.<sup>159</sup> reported that the endovascular approach can be considered the first technical choice even in cases of hypogastric aneurysm rupture. Obviously, according to their clinical presentation, results may differ in case of urgent clinical settings, in terms of technical success (elective: 100% vs urgent: 84%) and need for conversion to open surgery (4% only in urgent cases). Similar considerations can be summarized from the experience of Rana et al.<sup>160</sup>, which reported the results of surgical and endovascular repair of hypogastric artery aneurysms analyzing both urgent and elective scenarios. According to this experience, the endovascular repair had lower perioperative mortality (open: 2% versus endovascular: 0%), morbidity (open: 43% versus endovascular: 8%) and shorter hospitalizations (mean days: open 9 versus endovascular 1). The elective repair had a lower morbidity in terms of cardiological complications and bowel ischemia. The low invasiveness of endovascular repair was also reported by Patel et al.<sup>171</sup> and Chaer et al.<sup>172</sup>. Both these single-center experiences demonstrated that the endovascular repair of an isolated hypogastric artery aneurysm is associated with lower hospitalization and need of blood infusion than open surgical repair. Finally, several endovascular technical options were reported for both hypogastric embolization and revascularization<sup>174</sup>. Distal hypogastric artery revascularization was associated with a lower incidence of buttock claudication than proximal one<sup>174</sup>. In case of hypogastric artery embolization, the embolization of all distal aneurysm branches was associated with higher freedom from post-treatment aneurysm growth and reinterventions<sup>156</sup>.

### 3. MEDICAL THERAPY AND FOLLOW-UP

**PICO 3.1 Clinical question:** In patients with visceral or renal artery aneurysms who underwent corrective open/endovascular treatment, is CT angiography/Magnetic Resonance angiography superior to Doppler Ultrasound (DUS) for follow-up surveillance?

Literature research did not highlight randomized controlled trials, meta-analyses or observational studies correspondent to the PICO of interest, therefore recommendations relied on experts' opinion and current clinical practice. In case of patients treated endovascularly, it was established that CT angiography examinations should be performed within 3 months from index procedure and subsequently at 12 months to identify possible endoleaks or sac volume increase that might lead to aneurysm rupture. In case of open surgical treatment, CT angiography examination is suggested within 3 months from surgery and subsequently at 12 months; if no complications develop at 12 months, no further follow-up would be suggested.

**PICO 3.2 Clinical question:** In patients with visceral or renal artery aneurysms who did not undergo corrective treatment, is CT angiography/Magnetic Resonance angiography superior to Doppler Ultrasound (DUS) for aneurysm dimensions surveillance?

Literature research did not highlight randomized controlled trials, meta-analyses or observational studies correspondent to the PICO of interest, therefore recommendations relied on experts' opinion and current clinical practice. Considering the slow growth rate of these aneurysms, an annual DUS examination is suggested. In case of unfavorable anatomies which do not allow adequate ultrasound evaluation, it would be suggested to consider CT angiography or Magnetic Resonance angiography. It would be suggested to extended surveillance period up to 24-36 months in case of demonstrated volumetric stability.

**PICO 3.3 Clinical question:** In patients with visceral or renal artery aneurysms who did not undergo corrective treatment, is home medical therapy optimization indicated, compared to no therapy, to improve outcomes?

Literature research did not highlight randomized controlled trials, meta-analyses or observational studies correspondent to the PICO of interest, therefore recommendations relied on experts' opinion and current clinical practice. It was suggested to optimize medical therapy according to the etiopathogenetic cause of the aneurysm. Therefore, in patients with atherosclerotic visceral and/or renal artery aneurysms, modifiable risk factors treatment would be advisable, as well as medical therapy optimization according to current guidelines on atherosclerosis<sup>175-177</sup>. In patients with non-atherosclerotic non-inflammatory aneurysms (degenerative, connective tissue disorders or congenital diseases) antihypertensive therapy optimization and smoking cessations are desirable. In case of connective tissue disorders, antiplatelet therapy is suggested. In patients with visceral and/or renal artery inflammatory aneurysms, the use of steroids and/or immunosuppressants would be suggested to control inflammatory processes.

**PICO 3.4 Clinical question:** In patients with visceral or renal artery aneurysms who underwent corrective open/endovascular treatment, is home medical therapy optimization indicated, compared to no therapy, to improve outcomes?

Literature research did not highlight randomized controlled trials, meta-analyses or observational studies correspondent to the PICO of interest, therefore recommendations relied on experts' opinion and current clinical practice. It was suggested to optimize medical therapy according to the etiopathogenetic cause of the aneurysm. Therefore, in patients with atherosclerotic visceral and/or renal artery aneurysms, modifiable risk factors treatment would be advisable, as well as medical therapy optimization according to current guidelines on atherosclerosis<sup>175-177</sup>. In patients with non-atherosclerotic non-inflammatory aneurysms, antihypertensive therapy optimization and smoking cessations are desirable. In case of open surgical treatment, there is no evidence regarding the use of antiplatelet or anticoagulant therapy while, in case of endovascular treatment, it is suggested to evaluate short or long-term antiplatelet therapy according to the type of device used. Additionally, antiplatelet therapy might be considered for patients with connective tissue disorders. In patients with visceral and/or renal artery inflammatory aneurysms, the use of steroids and/or immunosuppressants would be suggested both in the pre- and post-operative surgical or endovascular period to control inflammatory processes.

## REFERENCES:

1. Scottish Intercollegiate Guidelines Network (SIGN). A guideline developer's handbook. Edinburgh: SIGN; 2019. (SIGN publication no. 50). [November 2019]. Available from URL: <http://www.sign.ac.uk> (ultimo accesso: 02/03/2023).
2. Centro Nazionale per l'Eccellenza Clinica, la Qualità e la Sicurezza delle Cure. Procedure di invio e valutazione di Linee Guida per la pubblicazione nell'SNLG – Manuale Operativo, versione 3.02 – febbraio 2020. Disponibile al sito web: [https://snlg.iss.it/wp-content/uploads/2020/02/MO\\_SNLG\\_v3.02\\_feb2020.pdf](https://snlg.iss.it/wp-content/uploads/2020/02/MO_SNLG_v3.02_feb2020.pdf) (ultimo accesso: 02/03/2023).
3. Centro Nazionale per l'Eccellenza Clinica, la Qualità e la Sicurezza delle Cure. Manuale metodologico per la produzione di linee guida di pratica clinica, versione 1.3.2 aprile 2019. Disponibile al sito web:

- [https://snlg.iss.it/wp-content/uploads/2019/04/MM\\_v1.3.2\\_apr\\_2019.pdf](https://snlg.iss.it/wp-content/uploads/2019/04/MM_v1.3.2_apr_2019.pdf) (ultimo accesso: 02/03/2023).
4. Cartabellotta A, Laganà AS. AGREE Reporting Checklist: uno strumento per migliorare il reporting delle linee guida. *Evidence* 2016;8(7): e1000146.
  5. AGREE Next Step Consortium. AGREE II. Checklist per la valutazione della qualità delle linee guida. Fondazione GIMBE: Bologna, aprile 2011. Disponibile a: [www.gimbe.org/agree](http://www.gimbe.org/agree) (ultimo accesso: 02/03/2023).
  6. Pratesi C, Esposito D, Apostolou D, Attisani L, Bellosta R, Benedetto F, et al. Guidelines on the management of abdominal aortic aneurysms: updates from the Italian Society of Vascular and Endovascular Surgery (SICVE). *J Cardiovasc Surg (Torino)*. 2022 Jun;63(3):328-352. doi: 10.23736/S0021-9509.22.12330-X. PMID: 35658387.
  7. Lanza G, Orso M, Alba G, Bevilacqua S, Capoccia L, Cappelli A, et al. Guideline on carotid surgery for stroke prevention: updates from the Italian Society of Vascular and Endovascular Surgery. A trend towards personalized medicine. *J Cardiovasc Surg (Torino)*. 2022 Aug;63(4):471-491. doi: 10.23736/S0021-9509.22.12368-2. PMID: 35848869.
  8. Pérez C, Llauger J, Pallardó Y, Sanchís E, Sabaté JM. Radiologic diagnosis of pseudoaneurysms complicating pancreatitis. *Eur J Radiol*. 1993 Feb;16(2):102-6. doi: 10.1016/0720-048x(93)90005-8. PMID: 8462572.
  9. Jesinger RA, Thoreson AA, Lamba R. Abdominal and pelvic aneurysms and pseudoaneurysms: imaging review with clinical, radiologic, and treatment correlation. *Radiographics*. 2013 May;33(3):E71-96. doi: 10.1148/rg.333115036. PMID: 23674782.
  10. Saba L, Anzidei M, Lucatelli P, Mallarini G. The multidetector computed tomography angiography (MDCTA) in the diagnosis of splenic artery aneurysm and pseudoaneurysm. *Acta Radiol*. 2011 Jun 1;52(5):488-98. doi: 10.1258/ar.2011.100283. Epub 2011 Mar 17. PMID: 21498313.
  11. Corvino F, Giurazza F, Ierardi AM, Lucatelli P, Basile A, Corvino A, et al. Splenic Artery Pseudoaneurysms: The Role of ce-CT for Diagnosis and Treatment Planning. *Diagnostics (Basel)*. 2022 Apr 17;12(4):1012. doi: 10.3390/diagnostics12041012. PMID: 35454060; PMCID: PMC9024490.
  12. Hagspiel KD, Flors L, Hanley M, Norton PT. Computed tomography angiography and magnetic resonance angiography imaging of the mesenteric vasculature. *Tech Vasc Interv Radiol*. 2015 Mar;18(1):2-13. doi: 10.1053/j.tvir.2014.12.002. Epub 2014 Dec 29. PMID: 25814198.
  13. Sidhu R, Lockhart ME. Imaging of renovascular disease. *Semin Ultrasound CT MR*. 2009 Aug;30(4):271-88. doi: 10.1053/j.sult.2009.04.002. PMID: 19711640.
  14. Lozano Sánchez FS, García-Alonso J, Torres JA, Velasco L, Salvador R, Peña R, et al. Decision-making and therapeutic options in intact splenic artery aneurysms: single-center experience and literature review. *Int Angiol*. 2020 Jun;39(3):241-251. doi: 10.23736/S0392-9590.20.04304-7. Epub 2020 Feb 13. PMID: 32057214.
  15. Olin JW, Sealove BA. Diagnosis, management, and future developments of fibromuscular dysplasia. *J Vasc Surg*. 2011 Mar;53(3):826-36.e1. doi: 10.1016/j.jvs.2010.10.066. Epub 2011 Jan 13. PMID: 21236620.
  16. Barrionuevo P, Malas MB, Nejim B, Haddad A, Morrow A, Ponce O, et al. A systematic review and meta-analysis of the management of visceral artery aneurysms. *J Vasc Surg*. 2019 Nov;70(5):1694-1699. doi: 10.1016/j.jvs.2019.02.024. Epub 2019 May 21. PMID: 31126761.
  17. Klausner JQ, Lawrence PF, Harlander-Locke MP, Coleman DM, Stanley JC, Fujimura N; Vascular Low-Frequency Disease Consortium. The contemporary management of renal artery aneurysms. *J Vasc Surg*. 2015 Apr;61(4):978-84. doi: 10.1016/j.jvs.2014.10.107. Epub 2014 Dec 20. PMID: 25537277.
  18. Wayne EJ, Edwards MS, Stafford JM, Hansen KJ, Corriere MA. Anatomic characteristics and natural history of renal artery aneurysms during longitudinal imaging surveillance. *J Vasc Surg*. 2014 Aug;60(2):448-52. doi: 10.1016/j.jvs.2014.03.006. Epub 2014 Apr 16. PMID: 24745940.
  19. Klausner JQ, Harlander-Locke MP, Plotnik AN, Lehrman E, DeRubertis BG, Lawrence PF. Current treatment of renal artery aneurysms may be too aggressive. *J Vasc Surg*. 2014 May;59(5):1356-61. doi: 10.1016/j.jvs.2013.11.062. Epub 2014 Jan 22. PMID: 24462256.
  20. henke
  21. Cohen JR, Shamash FS. Ruptured renal artery aneurysms during pregnancy. *J Vasc Surg*. 1987 Jul;6(1):51-9. doi: 10.1067/mva.1987.avs0060051. PMID: 3599281.
  22. Martin RS 3rd, Meacham PW, Ditesheim JA, Mulherin JL Jr, Edwards WH. Renal artery aneurysm: selective treatment for hypertension and prevention of rupture. *J Vasc Surg*. 1989 Jan;9(1):26-34. PMID: 2911140.
  23. Soliman KB, Shawky Y, Abbas MM, Ammary M, Shaaban A. Ruptured renal artery aneurysm during pregnancy, a clinical dilemma. *BMC Urol*. 2006 Aug 31;6:22. doi: 10.1186/1471-2490-6-22. PMID: 16945133; PMCID: PMC1569431.
  24. Gwon JG, Han DJ, Cho YP, Kim YH, Kwon TW. Role of heterotopic kidney auto-transplantation for renal artery aneurysms. *Medicine (Baltimore)*. 2018 Jun;97(23):e10856. doi: 10.1097/MD.00000000000010856. PMID: 29879018; PMCID: PMC5999450.
  25. Li Z, Zhao Z, Qin F, Wei X, Sun Y, Liu J, et al. Outcomes of Endovascular Treatment and Open Repair



- for Renal Artery Aneurysms: A Single-Center Retrospective Comparative Analysis. *J Vasc Interv Radiol*. 2018 Jan;29(1):62-70. doi: 10.1016/j.jvir.2017.08.020. PMID: 29102465.
26. Bilman V, Mascia D, Carta N, Santoro A, Saracino C, Chiesa R, et al. Contemporary Outcomes of in Situ Open Surgical Repair of Mid-Portion and Distal Renal Artery Aneurysms. *Ann Vasc Surg*. 2022 Jan;78:9-18. doi: 10.1016/j.avsg.2021.06.023. Epub 2021 Aug 28. PMID: 34464724.
  27. Laurin T, Borghese O, Branchereau J, Karam G, Brisard L, Corvec TL, et al. Single Centre Experience in Open and Endovascular Treatment of Renal Artery Aneurysms. *Ann Vasc Surg*. 2022 Feb;79:17-24. doi: 10.1016/j.avsg.2021.07.024. Epub 2021 Oct 10. PMID: 34644627.
  28. Brownstein AJ, Erben Y, Rajae S, Li Y, Rizzo JA, Mojibian H, et al. Natural history and management of renal artery aneurysms in a single tertiary referral center. *J Vasc Surg*. 2018 Jul;68(1):137-144. doi: 10.1016/j.jvs.2017.10.086. Epub 2018 Feb 2. PMID: 29398313.
  29. Machado M, Machado R, Almeida R. Renal Autotransplantation for The Treatment of Renal Artery Aneurysm. *Ann Vasc Surg*. 2022 Feb;79:226-232. doi: 10.1016/j.avsg.2021.07.048. Epub 2021 Oct 14. PMID: 34656716.
  30. Zhang Y, Xiang D, Lu Q, Wu M, Cui J. A systematic review and meta-analysis of the performance of flow-diverting stents in the treatment of peripheral and visceral artery aneurysms. *Catheter Cardiovasc Interv*. 2021 Feb 15;97(3):461-469. doi: 10.1002/ccd.29373. Epub 2020 Nov 11. PMID: 33175422.
  31. Loffroy R, Guiu B, Cercueil JP, et al. Transcatheter arterial embolization of splenic artery aneurysms and pseudoaneurysms: short- and long-term results. *Ann Vasc Surg* 2008; 22:618-626.
  32. Abbas MA, Stone WM, Fowl RJ, et al. Splenic artery aneurysms: two decades experience at Mayo clinic. *Ann Vasc Surg* 2002; 16:442-449.
  33. Berceli SA. Hepatic and splenic artery aneurysms. *Semin Vasc Surg* 2005; 18:196-201.
  34. Al-Habbal Y, Christophi C, Muralidharan V. Aneurysms of the splenic artery - a review. *Surgeon* 2010; 8:223-231.
  35. Hogendoorn W, Lavida A, Hunink MG, et al. Open repair, endovascular repair, and conservative management of true splenic artery aneurysms. *J Vasc Surg* 2014; 60:1667-1676.
  36. Lim HJ. A review of management options for splenic artery aneurysms and pseudoaneurysms. *Ann Med Surg (Lond)* 2020; 59:48-52.
  37. Keschenau PR, Kaisaris N, Jalaie H, Grommes J, Kotelis D, Kalder J, Jacobs MJ. Management strategies for true and dissecting visceral artery aneurysms. *J Cardiovasc Surg* 2020; 61:340-346.
  38. Batagini NC, Constantin BD, Kirksey L, et al. Natural History of Splanchnic Artery Aneurysms. *Ann Vasc Surg* 2021; 73:290-295.
  39. Chaer RA, Abularrage CJ, Coleman DM, Eslami MH, Kashyap VS, Rockman C, et al. The society for vascular surgery clinical practice guidelines on the management of visceral aneurysms. *J Vasc Surg* 2020; 72:3S-39S.
  40. Pulli R, Dorigo W, Troisi N, Pratesi G, Innocenti AA, Pratesi C. Surgical treatment of visceral artery aneurysms: a 25-year experience. *J Vasc Surg* 2008; 48:334-342.
  41. Venturini M, Marra P, Colombo M, et al. Endovascular treatment of visceral artery aneurysms and pseudoaneurysms in 100 patients: covered stenting vs transcatheter embolization. *J Endovasc Ther* 2017; 24:709-717.
  42. Venturini M, Marra P, Augello L, et al. Elective embolization of splenic artery aneurysms with an ethylene vinyl alcohol copolymer agent (Squid) and detachable coils. *J Vasc Interv Radiol* 2020; 31:1110-1117.
  43. Venturini M, Marra P, Colombo M, et al. Endovascular repair of 40 visceral artery aneurysms and pseudoaneurysms with the Viabahn stent-graft: technical aspects, clinical outcome and mid-term patency. *Cardiovasc Intervent Radiol* 2018; 41:385-397.
  44. Wang W, Chang H, Liu B, et al. Long-term outcomes of elective transcatheter dense coil embolization for splenic artery aneurysms: a two-center experience. *J Int Med Res* 2020; 48:300060519873256.
  45. Dorigo W, Pulli R, Azas L, et al. Early and intermediate results of elective endovascular treatment of true visceral artery aneurysms. *Ann Vasc Surg* 2016; 30:211-218.
  46. Venturini M, Piacentino F, Coppola A, et al. Visceral artery aneurysms embolization and other interventional options: state of the art and future perspectives. *J Clin Med* 2021; 10:2520
  47. Lakin RO, Bena JF, Sarac TP, et al. The contemporary management of splenic artery aneurysms. *J Vasc Surg* 2011; 53:958-964.
  48. Sano M, Hoshina K, Kawahara T, et al. Egg-shell like calcification as a protective factor for splenic artery aneurysm dilatation. *Ann Vasc Surg* 2020; 63:193-197.
  49. Clement M, Lareyre F, Loste A, et al. Vascular remodeling and immune cell infiltration in splenic artery aneurysms. *Angiology* 2021; 72:539-549.
  50. Phan D, Furtado R, Laurence JM, Pleass H. Splenic artery aneurysm management in the cirrhotic patient listed for liver transplantation: a systematic review. *Transplant Proc* 2022; 54:706-714.
  51. Kobori L, van der Kolk MJ, de Jong KP et al. Splenic artery aneurysms in liver transplant patients. *Liver Transplant Group. J Hepatol* 1997; 27:890-893.
  52. Ha JF, Phillips M, Faulkner K. Splenic artery aneurysm rupture in pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2009; 146:133-137.

53. Aung YYM, Berry C, Jayaram PR, Von Woon E. Splenic artery aneurysm in pregnancy: a systematic review. *Int J Gynaecol Obstet* 2023; 160:1-11.
54. Nanez L, Knowles M, Modrall JG, Valentine RJ. Ruptured splenic artery aneurysms are exceedingly rare in pregnant women. *J Vasc Surg* 2014; 60:1520-1523.
55. Noshier JL, Chung J, Brevetti LS, Graham AM, Siegel RL. Visceral and renal artery aneurysms: a pictorial essay on endovascular therapy. *Radiographics* 2006; 26: 1687-1704.
56. Chiesa R, Astore D, Guzzo G, et al. Visceral artery aneurysms. *Ann Vasc Surg* 2005; 19: 42-48.
57. Pitton MB, Dappa E, Jungmann F, et al. Visceral artery aneurysms: incidence, management, and outcome analysis in a tertiary care center over one decade, *Eur Radiol* 2015; 25: 2004-2014.
58. Hogendoorn W, Lavidia A, Hunink MG, et al. Cost-effectiveness of endovascular repair, open repair, and conservative management of splenic artery aneurysms. *J Vasc Surg* 2015; 61:1432-1440.
59. Venturini M, Marra P, Colarieti A, et al. Covered stenting and transcatheter embolization of splenic artery aneurysms in diabetic patients: A review of endovascular treatment of visceral artery aneurysms in the current era. *Pharmacol Res* 2018; 135:127-135.
60. Numoto I, Tsurusaki M, Oda T, Yagyu Y, Ishii K, Murakami T. Transcatheter arterial embolization treatment for bleeding visceral artery pseudoaneurysms in patients with pancreatitis or following pancreatic surgery. *Cancers* 2020; 12(10):2733.
61. Yamamoto S, Hirota S, Maeda H, Achiwa S, Arai K, Kobayashi K, Nakao N. Transcatheter coil embolization of splenic artery aneurysm. *Cardiovasc Intervent Radiol* 2008; 31:527-534.
62. Rossi M, Rebonato A, Greco L, Citone M, David V. Endovascular exclusion of visceral artery aneurysms with stent-grafts: technique and long-term follow-up. *Cardiovasc Intervent Radiol* 2008; 31:36-42.
63. Reed NR, Oderich GS, Manunga J, et al. Feasibility of endovascular repair of splenic artery aneurysms using stent grafts. *J Vasc Surg* 2015; 62:1504-1510.
64. Stella N, Palombo G, Taddeo C, Rizzo L, Taurino M. Stent-assisted coil embolization of a complex wide-neck splenic artery aneurysm. *Ann Vasc Surg* 2013; 27:1187.e5-8.
65. Colombi D, Bodini FC, Bossalini M, Rossi B, Michieletti E. Extracranial visceral artery aneurysms/pseudoaneurysms repaired with flow diverter device developed for cerebral aneurysms: preliminary results. *Ann Vasc Surg* 2018; 53:272.e1-272.e9.
66. Laganà D, Carrafiello G, Mangini M et al. Endovascular treatment of splenic artery aneurysms. *Rad Med* 2005; 110:77-87.
67. Lee SH, Yang S4, Park I, Im YC, Kim GY. Ruptured splenic artery aneurysms in pregnancy and usefulness of endovascular treatment in selective patients: a case report and review of literature. *World J Clin Cases* 2022; 10:9057-9063.
68. Marone EM, Peri A, Argenti F, Pugliese L, Rinaldi LF, Pietrabissa A. Robotic treatment of complex splenic artery aneurysms with deep hilar location: technical insights and midterm results. *Ann Vasc Surg* 2022; 68:50-56.
69. Illuminati G, Pizzardi G, Pasqua R. Open surgery for aneurysms of the splenic artery at the hilum of the spleen: Report of three cases. *Int J Surg Case Rep* 2018; 48:47-49.
70. Hamid HKS, Suliman AEA, Piffaretti G, et al. A systematic review on clinical features and management of true giant splenic artery aneurysms. *J Vasc Surg* 2020; 71:1036-1045.
71. Tipaldi MA, Krokidis M, Orgera G, et al. Endovascular management of giant visceral artery aneurysms. *Sci Rep* 2021; 11:700.
72. Mascia D, Salvati S, Carta N, Kahlberg A, Santoro A, Melissano G, et al. Endovascular Oriented Classification and Treatment of Celiac Trunk Aneurysms: 10 Years Experience. *Ann Vasc Surg.* 2022 Feb;79:219-225. doi: 10.1016/j.avsg.2021.07.012
73. Johal M, Kalaravy M, Ali F, Barve R, Ahmed A, Francis CT, et al. Evolving Diagnostic and Therapeutic Options for Visceral Artery Aneurysms. *Ann Vasc Surg.* 2021 Oct;76:488-499. doi: 10.1016/j.avsg.2021.03.012
74. Yuan FK, Xi HL, Qin RH, Tian ZL, Li C, Lu F. Endovascular treatment with stenting of celiac artery aneurysms. *Medicine (Baltimore).* 2020 Nov 25;99(48):e23448. doi: 10.1097/MD.00000000000023448
75. Guo B, Guo D, Xu X, Chen B, Shi Z, Luo J, et al. Early and intermediate results of endovascular treatment of symptomatic and asymptomatic visceral artery aneurysms. *J Vasc Surg.* 2016 Jul;64(1):140-8. doi: 10.1016/j.jvs.2016.02.037
76. Tulsyan N, Kashyap VS, Greenberg RK, Sarac TP, Clair DG, Pierce G, et al. The endovascular management of visceral artery aneurysms and pseudoaneurysms. *J Vasc Surg.* 2007 Feb;45(2):276-83; discussion 283. doi: 10.1016/j.jvs.2006.10.049
77. Popov P, Boskovic S, Sagic D, Radevic B, Ilijevski N, Nenezic D, et al. Treatment of visceral artery aneurysms: retrospective study of 35 cases. *Vasa.* 2007 Aug;36(3):191-8. doi: 10.1024/0301-1526.36.3.19.
78. Zhang W, Fu YF, Wei PL, E B, Li DC, Xu J. Endovascular Repair of Celiac Artery Aneurysm with the Use of Stent Grafts. *J Vasc Interv Radiol.* 2016 Apr;27(4):514-8. doi: 10.1016/j.jvir.2015.12.024.
79. Obara H, Kentaro M, Inoue M, Kitagawa Y. Correction to: Current management strategies for visceral artery aneurysms: an overview. *Surg Today.* 2020 Mar;50(3):320. doi: 10.1007/s00595-019-01947-x. Erratum for: *Surg Today.* 2020 Jan;50(1):38-49.

80. Sousa J, Costa D, Mansilha A. Visceral artery aneurysms: review on indications and current treatment strategies. *Int Angiol.* 2019 Oct;38(5):381-394. doi: 10.23736/S0392-9590.19.04194-4. Epub 2019 Jul 5. PMID: 31284707.
81. Hosn MA, Xu J, Sharafuddin M, Corson JD. Visceral Artery Aneurysms: Decision Making and Treatment Options in the New Era of Minimally Invasive and Endovascular Surgery. *Int J Angiol.* 2019 Mar;28(1):11-16. doi: 10.1055/s-0038-1676958. Epub 2019 Jan 8. PMID: 30880885; PMCID: PMC6417896.
82. Ibrahim F, Dunn J, Rundback J, Pellerito J, Galmer A. Visceral Artery Aneurysms: Diagnosis, Surveillance, and Treatment. *Curr Treat Options Cardiovasc Med.* 2018 Oct 26;20(12):97. doi: 10.1007/s11936-018-0696-x. PMID: 30367314.
83. Juntermanns B, Bernheim J, Karaindros K, Walensi M, Hoffmann JN. Visceral artery aneurysms. *Gefasschirurgie.* 2018;23(Suppl 1):19-22. doi: 10.1007/s00772-018-0384-x. Epub 2018 Apr 20. PMID: 29950792; PMCID: PMC5997106.
84. van Rijn MJ, Ten Raa S, Hendriks JM, Verhagen HJ. Visceral aneurysms: Old paradigms, new insights? *Best Pract Res Clin Gastroenterol.* 2017 Feb;31(1):97-104. doi: 10.1016/j.bpg.2016.10.017. Epub 2017 Jan 4. PMID: 28395793.
85. Batagini NC, El-Arousy H, Clair DG, Kirksey L. Open versus Endovascular Treatment of Visceral Artery Aneurysms and Pseudoaneurysms. *Ann Vasc Surg.* 2016 Aug;35:1-8. doi: 10.1016/j.avsg.2016.01.035. Epub 2016 May 27. PMID: 27238989.
86. Uberoi R, Chung D. Endovascular solutions for the management of visceral aneurysms. *J Cardiovasc Surg (Torino).* 2011 Jun;52(3):323-31.
87. Sachdev-Ost U. Visceral artery aneurysms: review of current management options. *Mt Sinai J Med.* 2010 May-Jun;77(3):296-303.
88. Saltzberg SS, Maldonado TS, Lamparello PJ, Cayne NS, Nalbandian MM, Rosen RJ, et al. Is endovascular therapy the preferred treatment for all visceral artery aneurysms? *Ann Vasc Surg.* 2005 Jul;19(4):507-15.
89. Stone WM, Abbas MA, Gloviczki P, Fowl RJ, Cherry KJ. Celiac arterial aneurysms: a critical reappraisal of a rare entity. *Arch Surg.* 2002 Jun;137(6):670-4
90. Graham LM, Stanley JC, Whitehouse WM Jr, Zelenock GB, Wakefield TW, Cronenwett JL, et al. Celiac artery aneurysms: historic (1745-1949) versus contemporary (1950-1984) differences in etiology and clinical importance. *J Vasc Surg.* 1985 Sep;2(5):757-64.
91. Messina LM, Shanley CJ. Visceral artery aneurysms. *Surg Clin North Am.* 1997 Apr;77(2):425-42.
92. Sessa C, Tinelli G, Porcu P, Aubert A, Thony F, Magne JL. Treatment of visceral artery aneurysms: description of a retrospective series of 42 aneurysms in 34 patients. *Ann Vasc Surg.* 2004 Nov;18(6):695-703.
93. Huang Y, Banga P, De Souza LR, Oderich GS. Endovascular Treatment of Visceral Artery Aneurysms. *J Cardiovasc Surg.* 2015 56:567-77.
94. Hemp Jh and Sabri SS. Endovascular Management of Visceral Artery Aneurysms. *Tech Vasc Interventional Radiology.* 2015 18: 14-23.
95. Shukla AJ, Eid R, Fish L, Avgerinos E, Marone L, Makaroun M, et al. Contemporary outcomes of intact and ruptured visceral artery aneurysms. *J Vasc Surg.* 2015 Jun;61(6):1442-7. doi: 10.1016/j.jvs.2015.01.005. Epub 2015 Mar 7. PMID: 25752692.
96. Qiu C, Liu Z, Huang L, Guo L, Lu W, Zhang H, et al. Covered Stents for Treatment of Visceral Artery Aneurysms: A Multicenter Study. *J Vasc Interv Radiol.* 2022 Jun;33(6):640-647. doi: 10.1016/j.jvir.2022.03.009. Epub 2022 Mar 18. PMID: 35314368.
97. Li X, Zhang W, Zhou M, Ding Y, Wang Y, Xie T, et al. A new classification and strategies for endovascular treatment of celiac artery aneurysms. *Vascular.* 2022 Oct;30(5):834-841. doi: 10.1177/17085381211032768. Epub 2021 Jul 15. PMID: 34263673.
98. Fankhauser GT, Stone WM, Naidu SG, Oderich GS, Ricotta JJ, Bjarnason H, et al; Mayo Vascular Research Center Consortium. The minimally invasive management of visceral artery aneurysms and pseudoaneurysms. *J Vasc Surg.* 2011 Apr;53(4):966-70. doi: 10.1016/j.jvs.2010.10.071. Epub 2011 Jan 8. PMID: 21216559.
99. Vohra R, Carr HM, Welch M, Tait WF, Durrans D, Walker MG. Management of coeliac artery aneurysms. *Br J Surg.* 1991 Nov;78(11):1373-5. doi: 10.1002/bjs.1800781133. PMID: 1760706.
100. Matsumoto K, Tanaka K, Ohsumi K, Nakamaru M, Obara H, Hayashi S, et al. Celiomesenteric anomaly with concurrent aneurysm. *J Vasc Surg.* 1999 Apr;29(4):711-4. doi: 10.1016/s0741-5214(99)70318-6. PMID: 10194500.
101. Waldenberger P, Bendix N, Petersen J, Tauscher T, Glodny B. Clinical outcome of endovascular therapeutic occlusion of the celiac artery. *J Vasc Surg.* 2007 Oct;46(4):655-61. doi: 10.1016/j.jvs.2007.05.033. Epub 2007 Aug 30. PMID: 17764875.
102. Xia FF, Fan ZQ, Huo XB, Fu YF, Xu YS. Endovascular stent repair of celiac arterial aneurysm. *Medicine (Baltimore).* 2019 Nov;98(48):e18203. doi: 10.1097/MD.00000000000018203. PMID: 31770276; PMCID: PMC6890336.
103. Cochennec F, Riga CV, Allaire E, Cheshire NJ, Hamady M, Jenkins MP, et al. Contemporary management of splanchnic and renal artery aneurysms: results of endovascular compared with open surgery from two

- European vascular centers. *Eur J Vasc Endovasc Surg.* 2011 Sep;42(3):340-6. doi: 10.1016/j.ejvs.2011.04.033. Epub 2011 May 31. PMID: 21628100.
104. Mohan IV, Stephen MS. Peripheral arterial aneurysms: open or endovascular surgery? *Prog Cardiovasc Dis.* 2013 Jul-Aug;56(1):36-56
  105. Jesinger RA, Thoreson AA, Lamba R. Abdominal and pelvic aneurysms and pseudoaneurysms: imaging review with clinical, radiologic, and treatment correlation. *Radiographics.* 2013 May;33(3):E71-96.
  106. Vandy FC, Sell KA, Eliason JL, Coleman DM, Rectenwald JE, Stanley JC. Pancreaticoduodenal and Gastroduodenal Artery Aneurysms Associated with Celiac Artery Occlusive Disease. *Ann Vasc Surg.* 2017; 41:32-40.
  107. Michalinos A, Schizas D, Ntourakis D, Filippou D, Troupis T. Arc of Bühler: the surgical significance of a rare anatomical variation. *Surg Radiol Anat.* 2019;41(5):575-581.
  108. Stoecker JB, Eddinger KC, Glaser JD, Wang GJ, Shlansky-Goldberg RD, Fairman RM, et al. A large series of true pancreaticoduodenal artery aneurysms. *J Vasc Surg.* 2022;75(5):1634-1642.
  109. Illuminati G, Hostalrich A, Pasqua R, Nardi P, Chaufour X, Ricco JB. Outcomes After Open and Endovascular Repair of Non-Ruptured True Pancreaticoduodenal and Gastroduodenal Artery Aneurysms Associated with Coeliac Artery Compression: A Multicentre Retrospective Study. *Eur J Vasc Endovasc Surg.* 2021;61(6):945-953.
  110. Bonardelli S, Spampinato B, Ravanelli M, Cuomo R, Zanotti C, Paro B, et al. The role of emergency presentation and revascularization in aneurysms of the peripancreatic arteries secondary to celiac trunk or superior mesenteric artery occlusion. *J Vasc Surg.* 2020;72(1S):46S-55S.
  111. Kallamadi R, DeMoya MA, Kalva SP. Inferior pancreaticoduodenal artery aneurysms in association with celiac stenosis/occlusion. *Semin Intervent Radiol.* 2009;26(3):215-223. doi:10.1055/S-0029-1225671
  112. Lee JH, Hwang DW, Lee SY, Hwang JW, Song DK, Gwon DI, et al. Clinical Features and Management of Pseudoaneurysmal Bleeding after Pancreatoduodenectomy. <https://doi.org/10.1177/000313481207800339>. 2012;78(3):309-317. doi:10.1177/000313481207800339
  113. Makowiec F, Riediger H, Euringer W, Uhl M, Hopt UT, Adam U. Management of delayed visceral arterial bleeding after pancreatic head resection. *J Gastrointest Surg.* 2005;9(9):1293-1299. doi:10.1016/J.GASSUR.2005.08.003
  114. Sharma S, Prasad R, Gupta A, Dwivedi P, Mohindra S, Yadav RR. Aneurysms of pancreaticoduodenal arcade: Clinical profile and endovascular strategies. *JGH Open.* 2020;4(5):923-928. doi:10.1002/JGH3.12365
  115. Murata S, Tajima H, Fukunaga T, Abe Y, Niggemann P, Onozawa S, et al. Management of pancreaticoduodenal artery aneurysms: results of superselective transcatheter embolization. *AJR Am J Roentgenol.* 2006;187(3). doi:10.2214/AJR.04.1726
  116. Gupta V, Irrinki S, Sakaray YR, Moond V, Yadav TD, Kochhar R, et al. Treatment strategies for bleeding from gastroduodenal artery pseudoaneurysms complicating the course of chronic pancreatitis-A case series of 10 patients. *Indian J Gastroenterol.* 2018;37(5):457-463. doi:10.1007/S12664-018-0897-Y
  117. Dohan A, Eveno C, Dautry R, Moond V, Yadav TD, Kochhar R, et al. Role and Effectiveness of Percutaneous Arterial Embolization in Hemodynamically Unstable Patients with Ruptured Splanchnic Artery Pseudoaneurysms. *Cardiovasc Intervent Radiol.* 2015;38(4):862-870. doi:10.1007/S00270-014-1002-2
  118. Berceci SA. Hepatic and splenic artery aneurysms. *Semin Vasc Surg* 2005;18:196-201. doi: 10.1053/j.semvascsurg.2005.09.005
  119. Fankhauser GT, Stone WM, Naidu SG, Oderich GS, Ricotta JJ, Bjarnason H, et al. The minimally invasive management of visceral artery aneurysms and pseudoaneurysms. *J Vasc Surg* 2011;53:966-70. doi: 10.1016/j.jvs.2010.10.071
  120. Stark JC, Eisenberg N, Mafeld S, McGilvray I, Roche-Nagle G, Howe KL. Assessment of open surgical and endovascular management of true hepatic artery aneurysms over 20 years highlights increased rupture risk in females. *J Vasc Surg.* 2022 Apr;75(4):1334-1342.e2. doi: 10.1016/j.jvs.2021.12.054
  121. Schick C, Ritter RG, Balzer JO, Thalhammer A, Vogl TJ. Hepatic artery aneurysm: treatment options. *Eur Radiol* 2004;14:157-9. doi: 10.1007/s00330-003-1881-0
  122. Abbas MA, Fowl RJ, Stone WM, Panneton JM, Oldenburg WA, Bower TC, et al. Hepatic artery aneurysm: factors that predict complications. *J Vasc Surg* 2003;38:41-5. doi: 10.1016/s0741-5214(03)00090-9
  123. Erskine JM. Hepatic artery aneurysm. *Vasc Surg* 1973;7: 106-25. doi: 10.1177/153857447300700205.
  124. Lal RB, Strohl JA, Piazza S, Aslam M, Ball D, Patel K. Hepatic artery aneurysm. *J Cardiovasc Surg (Torino)* 1989;30: 509-13.
  125. Koyama M, Tanaka M, Shimizu M, Nomura S, Kako N, Suzuki S, et al. Surgical treatment of mesenteric infarction, thoracoabdominal aortic aneurysm, and proper hepatic aneurysm in a middle-aged woman with Takayasu's arteritis. *J Cardiovasc Surg (Torino)* 1995;36:337-41.
  126. Kok HK, Asadi H, Sheehan M, Given MF, Lee MJ. Systematic Review and Single-Center Experience for Endovascular Management of Visceral and Renal Artery Aneurysms. *J Vasc Interv Radiol.* 2016 Nov;27(11):1630-1641. doi: 10.1016/j.jvir.2016.07.030;
  127. Sfyroeras GS, Dalainas I, Giannakopoulos TG, Antonopoulos K, Kakisis JD, Liapis CD. Flow-diverting

- stents for the treatment of arterial aneurysms. *J Vasc Surg.* 2012 Sep;56(3):839-46. doi: 10.1016/j.jvs.2012.04.020
128. Lumsden AB, Mattar SG, Allen RC, Bacha EA. Hepatic artery aneurysms: the management of 22 patients. *J Surg Res* 1996;60:345-50. doi: 10.1006/jsre.1996.0055.
129. Melissano G, Mascia D, Atique Gabriel S, Bertoglio L, Venturini M, DE Cobelli F, et al. Hepatic artery aneurysms: open and endovascular repair. *J Cardiovasc Surg (Torino)*. 2018 Feb;59(1):95-100. doi: 10.23736/S0021-9509.16.09220-X
130. Marone EM, Mascia D, Kahlberg A, Brioschi C, Tshomba Y, Chiesa R. Is open repair still the gold standard in visceral artery aneurysm management? *Ann Vasc Surg.* 2011 Oct;25(7):936-46. doi: 10.1016/j.avsg.2011.03.006
131. Pitcher GS, Cirillo-Penn NC, Mendes BC, Shuja F, DeMartino RR, Kalra M, et al. Aneurysms of the superior mesenteric artery and its branches. *J Vasc Surg.* 2022 Jul;76(1):149-157. doi: 10.1016/j.jvs.2022.02.047. Epub 2022 Mar 8. PMID: 35276263.
132. Erben Y, Brownstein AJ, Rajae S, Li Y, Rizzo JA, Mojibian H, et al. Natural history and management of splanchnic artery aneurysms in a single tertiary referral center. *J Vasc Surg.* 2018 Oct;68(4):1079-1087. doi: 10.1016/j.jvs.2017.12.057. Epub 2018 Mar 21. PMID: 29573962.
133. Corey MR, Ergul EA, Cambria RP, English SJ, Patel VI, Lancaster RT, et al. The natural history of splanchnic artery aneurysms and outcomes after operative intervention. *J Vasc Surg.* 2016 Apr;63(4):949-57. doi: 10.1016/j.jvs.2015.10.066. Epub 2016 Jan 11. PMID: 26792545.
134. Stone WM, Abbas M, Cherry KJ, Fowl RJ, Gloviczki P. Superior mesenteric artery aneurysms: is presence an indication for intervention? *J Vasc Surg.* 2002 Aug;36(2):234-7; discussion 237. doi: 10.1067/mva.2002.125027. PMID: 12170202.
135. Zilun L, Henghui Y, Yang Z, Mian W, Guangqi C, Shenming W. The Management of Superior Mesenteric Artery Aneurysm: Experience with 16 Cases in a Single Center. *Ann Vasc Surg.* 2017 Jul;42:120-127. doi: 10.1016/j.avsg.2016.11.014. Epub 2017 Mar 22. PMID: 28341504.
136. Shi Y, Ni G, Zhao B, Gu J, Huang H, Lu Z, et al. Management of Symptomatic Spontaneous Isolated Superior Mesenteric Artery Dissection: A Single Centre Experience with Mid Term Follow Up. *Eur J Vasc Endovasc Surg.* 2020 Dec;60(6):863-871. doi: 10.1016/j.ejvs.2020.08.010. Epub 2020 Oct 5. PMID: 33032925.
137. Jacobs CR, Fatima J, Scali ST, Hodges ZH, Back MR, Arnaoutakis DJ, et al. Surgical Treatment of True Superior Mesenteric Artery Aneurysms. *Ann Vasc Surg.* 2021 Feb;71:74-83. doi: 10.1016/j.avsg.2020.08.142. Epub 2020 Sep 14. PMID: 32941966.
138. DeCarlo C, Mohebalı J, Dua A, Conrad MF, Mohapatra A. Morbidity and mortality associated with open repair of visceral aneurysms. *J Vasc Surg.* 2022 Feb;75(2):632-640.e2. doi: 10.1016/j.jvs.2021.08.079. Epub 2021 Sep 22. PMID: 34560216.
139. Martinelli O, Giglio A, Irace L, Di Girolamo A, Gossetti B, Gattuso R. Single-Center Experience in the Treatment of Visceral Artery Aneurysms. *Ann Vasc Surg.* 2019 Oct;60:447-454. doi: 10.1016/j.avsg.2019.01.010. Epub 2019 Apr 19. PMID: 31009733.
140. Jiang J, Ding X, Su Q, Zhang G, Wang Q, Jian W, et al. Therapeutic management of superior mesenteric artery aneurysms. *J Vasc Surg.* 2011 Jun;53(6):1619-24. doi: 10.1016/j.jvs.2011.02.004. Epub 2011 Apr 30. PMID: 21531524.
141. Onal Y, Samanci C, Cicek ED. Double-Lumen Balloons, Are They Only Useful in Neurointerventions? Preliminary Outcomes of Double-Lumen Balloon-Assisted Embolization of Visceral Artery Aneurysms. *Vasc Endovascular Surg.* 2020 Apr;54(3):214-219. doi: 10.1177/1538574419896516. Epub 2019 Dec 27. PMID: 31878841.
142. Jia Z, Su H, Chen W, Ni G, Qi C, Gu J. Endovascular Treatment of Patients with Isolated Mesenteric Artery Dissection Aneurysm: Bare Stents Alone Versus Stent Assisted Coiling. *Eur J Vasc Endovasc Surg.* 2019 Mar;57(3):400-406. doi: 10.1016/j.ejvs.2018.08.057. Epub 2018 Oct 10. PMID: 30316568.
143. Kim SK, Lee J, Duncan JR, Picus DD, Darcy MD, Sauk S. Endovascular treatment of superior mesenteric artery pseudoaneurysms using covered stents in six patients. *AJR Am J Roentgenol.* 2014 Aug;203(2):432-8. doi: 10.2214/AJR.13.11644. PMID: 25055281.
144. Graham JM, McCollum CH, DeBakey ME. Aneurysms of the splanchnic arteries. *Am J Surg.* 1980 Dec;140(6):797. doi: 10.1016/0002-9610(80)90120-8. PMID: 7457704.
145. Huang YK, Hsieh HC, Tsai FC, Chang SH, Lu MS, Ko PJ. Visceral artery aneurysm: risk factor analysis and therapeutic opinion. *Eur J Vasc Endovasc Surg.* 2007 Mar;33(3):293-301. doi: 10.1016/j.ejvs.2006.09.016. Epub 2006 Nov 9. PMID: 17097898.
146. Zhang L, Yin CP, Li HY, Bao JM, Zhao ZQ, Qu LF, et al. Multiple overlapping bare stents for endovascular visceral aneurysm repair: a potential alternative endovascular strategy to multilayer stents. *Ann Vasc Surg.* 2013 Jul;27(5):606-12. doi: 10.1016/j.avsg.2012.07.013. Epub 2013 Mar 22. PMID: 23523448.
147. Huo CW. Middle colic artery aneurysm: a case report and review of the literature. *Ann Vasc Surg.* 2012 May;26(4):571.e1-6. doi: 10.1016/j.avsg.2011.07.023. Epub 2012 Feb 8. PMID: 22321489.
148. Inada K, Maeda M, Ikeda T. Segmental arterial mediolysis: unrecognized cases culled from cases of

- ruptured aneurysm of abdominal visceral arteries reported in the Japanese literature. *Pathol Res Pract*. 2007;203(11):771-8. doi: 10.1016/j.prp.2007.07.010. Erratum in: *Pathol Res Pract*. 2008;204(2):147. PMID: 17920781.
149. Asano M, Nushida H, Nagasaki Y, Tatsuno Y, Ueno Y. Rupture of a jejunal artery aneurysm. *Leg Med (Tokyo)*. 2008 Sep;10(5):268-73. doi: 10.1016/j.legalmed.2008.02.001. Epub 2008 Apr 18. PMID: 18378483.
  150. Shanley CJ, Shah NL, Messina LM. Uncommon splanchnic artery aneurysms: pancreaticoduodenal, gastroduodenal, superior mesenteric, inferior mesenteric, and colic. *Ann Vasc Surg*. 1996 Sep;10(5):506-15. doi: 10.1007/BF02000601. PMID: 8905073.
  151. Sarcina A, Bellosta R, Magnaldi S, Luzzani L. Aneurysm of the middle colic artery--case report and literature review. *Eur J Vasc Endovasc Surg*. 2000 Aug;20(2):198-200. doi: 10.1053/ejvs.1999.1076. PMID: 10942694.
  152. Tessier DJ, Abbas MA, Fowl RJ, Stone WM, Bower TC, McKusick MA, et al. Management of rare mesenteric arterial branch aneurysms. *Ann Vasc Surg*. 2002 Sep;16(5):586-90. doi: 10.1007/s10016-001-0271-9. Epub 2002 Aug 19. PMID: 12183776.
  153. Richardson JW, Greenfield LJ. Natural history and management of iliac aneurysms. *J Vasc Surg* 1988 Aug;8(2):165-71.
  154. Kliewer M, Plimon M, Taher F, Walter C, Hirsch K, Falkensammer J et al. Endovascular treatment of hypogastric artery aneurysms. *J Vasc Surg* 2019 Oct;70(4):1107-1114.
  155. Gao P, Dong D, Yang L, Yuan H, Wang M, Zhang J et al. Technical issues and clinical outcomes of endovascular repair of isolated iliac artery aneurysms: A single-center experience. *Vascular* 2018 Dec;26(6):591-599.
  156. Muradi A, Yamaguchi M, Okada T, Nomura Y, Idoguchi K, Ueshima E et al. Technical and outcome considerations of endovascular treatment for internal iliac artery aneurysms. *Cardiovasc Intervent Radiol* 2014 Apr;37(2):348-54.
  157. Laine MT, Björck M, Beiles CB, Szeberin Z, Thomson I, Altreuther M et al. Few internal iliac artery aneurysms rupture under 4 cm. *J Vasc Surg* 2017 Jan;65(1):76-81.
  158. Chen RJ, Vaes RHD, Qi SD, J Westcott M, Robinson DR. Modalities of endovascular management for internal iliac artery aneurysms. *ANZ J Surg* 2021 Nov;91(11):2397-2403.
  159. Kobe A, Andreotti C, Puipe G, Rancic Z, Kopp R, Lachat M et al. Primary Endovascular Elective Repair and Repair of Ruptured Isolated Iliac Artery Aneurysms Is Durable-Results of 72 Consecutive Patients. *J Vasc Interv Radiol* 2018 Dec;29(12):1725-1732.
  160. Rana MA, Kalra M, Oderich GS, de Grandis E, Głowiczki P, Duncan AA et al. Outcomes of open and endovascular repair for ruptured and nonruptured internal iliac artery aneurysms. *J Vasc Surg* 2014 Mar;59(3):634-44.
  161. Patel NV, Long GW, Cheema ZF, Rimar K, Brown OW, Shanley CJ. Open vs. endovascular repair of isolated iliac artery aneurysms: A 12-year experience. *J Vasc Surg* 2009 May;49(5):1147-53.
  162. Chaer RA, Barbato JE, Lin SC, Zenati M, Kent KC, McKinsey JF. Isolated iliac artery aneurysms: a contemporary comparison of endovascular and open repair. *J Vasc Surg* 2008 Apr;47(4):708-713.
  163. Yang M, Li L, Liu Y, Su Q, Dong Z, Li G et al. Therapeutic management of isolated internal iliac artery aneurysms. *J Vasc Surg* 2020 Dec;72(6):1968-1975.
  164. Pirvu A, Gallet N, Perou S, Thony F, Magne JL. Midterm results of internal iliac artery aneurysm embolization. *J Med Vasc* 2017 May;42(3):157-161.
  165. Bianchini Massoni C, Freyrie A, Gargiulo M, Tecchio T, Mascoli C, Gallitto E et al. Perioperative and Late Outcomes after Endovascular Treatment for Isolated Iliac Artery Aneurysms. *Ann Vasc Surg* 2017 Oct;44(4):83-93.
  166. Millon A, Paquet Y, Ben Ahmed S, Pinel G, Rosset E, Lermusiaux P. Midterm outcomes of embolisation of internal iliac artery aneurysms. *Eur J Vasc Endovasc Surg* 2013 Jan;45(1):22-7.
  167. Antoniou GA, Nassef AH, Antoniou SA, Loh CY, Turner DR, Beard JD. Endovascular treatment of isolated internal iliac artery aneurysms. *Vascular* 2011 Dec;19(6):291-300.
  168. Yamamoto H, Yamamoto F, Ishibashi K, Liu KX, Yamaura G, Chida Y et al. Long-term outcomes of open surgical repair for ruptured iliac artery aneurysms. *Ann Vasc Surg* 2011 Aug;25(6):740-7.
  169. Chemelli A, Hugl B, Klocker J, Thauerer M, Strasak A, Jaschke W et al. Endovascular repair of isolated iliac artery aneurysms. *J Endovasc Ther* 2010 Aug;17(4):492-503.
  170. Hu H, Takano T, Guntani A, Onohara T, Furuyama T, Inoguchi H et al. Treatment of solitary iliac aneurysms: clinical review of 28 cases. *Surg Today* 2008;38(3):232-6.
  171. Hiromatsu S, Hosokawa Y, Egawa N, Yokokura H, Akaiwa K, Aoyagi S. Strategy for isolated iliac artery aneurysms. *Asian Cardiovasc Thorac Ann* 2007 Aug;15(4):280-4.
  172. Boules TN, Selzer F, Stanziale SF, Chomic A, Marone LK, Dillavou ED et al. Endovascular management of isolated iliac artery aneurysms. *J Vasc Surg* 2006 Jul;44(1):29-37.
  173. McCready RA, Pairolo PC, Gilmore JC, Kazmier FJ, Cherry KJ Jr, Hollier LH. Isolated iliac artery aneurysms. *Surgery* 1983 May;93(5):688-93.
  174. Perini P, Mariani E, Fanelli M, Ucci A, Rossi, G, Bianchini Massoni C, et al. Surgical and endovascular

- management of isolated internal iliac artery aneurysms: a systematic review and meta-analysis. *Vascular Endovascular Surgery* 2021;(55):254-264.
175. Frank U, Nikol S, Belch J, Boc V, Brodmann M, Carpentier PH, et al. ESVM Guideline on peripheral arterial disease. *Vasa*. 2019;48(Suppl 102):1-79. doi:10.1024/0301-1526/a000834.
  176. Aboyans V, Ricco JB, Bartelink MEL, Björck M, Brodmann M, Cohnert T, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS). *Rev Esp Cardiol (Engl Ed)*. 2018;71(2):111. doi:10.1016/j.rec.2017.12.014.
  177. Gerhard-Herman MD, Gornik HL, Barrett C, Barshes NR, Corriere MA, Drachman DE, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines [published correction appears in *Circulation*. 2017 Mar 21;135(12 ):e791-e792]. *Circulation*. 2017;135(12):e726-e779. doi:10.1161/CIR.0000000000000471.
  178. Björck M, Koelemay M, Acosta S, Bastos Goncalves F, Kölbel T, Kolkman JJ, et al. Editor's Choice - Management of the Diseases of Mesenteric Arteries and Veins: Clinical Practice Guidelines of the European Society of Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg*. 2017 Apr;53(4):460-510. doi: 10.1016/j.ejvs.2017.01.010. PMID: 28359440.
  179. Wanhainen A, Verzini F, Van Herzele I, Allaire E, Bown M, Cohnert T, et al. Editor's Choice - European Society for Vascular Surgery (ESVS) 2019 Clinical Practice Guidelines on the Management of Abdominal Aorto-iliac Artery Aneurysms. *Eur J Vasc Endovasc Surg*. 2019 Jan;57(1):8-93. doi: 10.1016/j.ejvs.2018.09.020. Epub 2018 Dec 5. Erratum in: *Eur J Vasc Endovasc Surg*. 2020 Mar;59(3):494. PMID: 30528142.