Twisted Cohomotopy implies twisted String structure on M5-branes

Domenico Fiorenza, Hisham Sati, Urs Schreiber

March 18, 2021


Abstract

We show that charge-quantization of the M-theory C-field in J-twisted Cohomotopy implies emergence of a higher Sp(1)-gauge field on single heterotic M5-branes, which exhibits worldvolume $String^{c_2}$ -structure.

Towards an understanding of the elusive mathematical formulation of M-theory [Du96][Du99b][BBS06] the following assumption has been proposed [Sa13][FSS19b][FSS19c][SS19a][SS19b][SS21]:

Hypothesis H: The M-Theory C-field is charge-quantized in J-twisted Cohomotopy theory, i.e. the C-field flux densities G_4 , G_7 are in the image of the twisted/equivariant cohomotopical character map (from [FSS20b, §5.3][SS20c, Thm. 1.1]).

Assuming this, fields of the M5-brane sigma-model [FSS19d] are found to be encoded by classifying maps shown in green in the following homotopy-commutative diagram of topological spaces [FSS19c] (based on [FSS19b]):



Here we show how the purple maps come about, and what this means for the M5-brane. First some background:

The fundamental problem in string theory, which remains open¹, is to find its non-perturbative completion to a conjectural *M-theory* [Du99b][BBS06]. This subsumes the major open problem (e.g., [Mo12, p. 77][La12, p. 49][Hu13, p. 1][La19, 3.6]) of identifying the *higher gauge theory* ("non-abelian gerbe theory" [Wi02, p. 6, 15]), supposedly carried by *N* coincident M5-branes, either seen as the 6d SCFT right on the brane or seen in its vicinity ("AdS/CFT", see [Du99a]) in the non-perturbative *small N* limit (large 1/N limit). This is the problem we are concerned with here. Specifically, for $N \to 3$ this should, in turn, subsume ("AdS/QCD", e.g. [RZ16, §4]) the Millennium Problem (see [RS20][Ro21]) of formulating non-perturbative YM theory, hence confined QCD.

The beauty of the M-theory conjecture is that it reduces a zoo of physical fields to just a pair:² gravity and the C-field (review in [MiSc06]). This suggests that formulating M-theory is essentially tantamount to identifying the true mathematical nature of the C-field coupled to gravity. But this is tractable by an Ansatz like Hypothesis H, because at least the mathematical *type* of the C-field is known with some certainty:

The C-field is a higher gauge field with flux densities higher-degree differential forms [SSS08], and with gauge transformations accompanied by higher-order gauge-of-gauge transformations (*n*-group gauge symmetry [BaSc07] [FSS10][FSS13a][Sch13]). Such higher gauge fields are described, non-perturbatively, by *homotopy theory*, where they are embodied in *non-abelian generalized cohomology* theories (see [FSS20b, §2]). Coupling these to other fields, notably to gravity, means to consider *twisted* and *differential* (see (11)) non-abelian generalized cohomology:

The correspondence between higher gauge fields, generalized cohomology and geometric homotopy theory proceeds as follows [SSS12][FSS12a][Sch13][FSS20b][SS20b][SS20c][SS21] (see [FSS19a] for a gentle survey):

Non-perturbative Physics	Geometric (stacky) Homotopy theory	Twisted differential Cohomology	References	
Field configurations	$X \xrightarrow{\text{maps}} A \text{classifying space (∞-stack)}$	cocycles	[SSS12, Fig. 2]	
Gauge transformations	X homotopies A	coboundaries	[NSS12, §3] [FSS20b, §2.1] [SS20b, (22)] [SS21, §2.1]	
Gauge-of-gauge transformations	$X \Longrightarrow A$	cocorners	[3321, §2.1]	
Gauge-equivalence classes of field configuration	$A(X) := \left\{ X \xrightarrow{\text{maps}} A \right\}_{\text{homotopies}}$	cohomology		
Field configurations coupled to gravity	$X \xrightarrow{\text{maps}} A /\!\!/ \text{Spin}(n)$ $\downarrow^{\&} \text{homotopies} \text{quotient}$ classifying map of frame bundle $B\text{Spin}(n)$	tangentially twisted cocycle	[NSS12, §4] [FSS19b, §2] [SS20b, 2.94] [FSS20b, §2.2]	
Instanton/soliton numbers	$ \begin{array}{c} A \xrightarrow{\text{maps}} & B^n \mathrm{U}(1) \\ \text{classifying} & \text{Eilenberg-MacLane} \\ \text{space } (\infty\text{-stack}) & \text{space } (\infty\text{-stack}) \end{array} $	characteristic classes	[FSS10] [FSS12a, §3.1]	
Ordinary gauge instantons / gravitational instantons	gauge field $BU \xrightarrow{c_2} B^3U(1)$ $X \xrightarrow{\text{gravity}} BO \xrightarrow{\frac{1}{2}p_1} B^3U(1)$	2nd Chern class / 1st Pontrjagin class	[FSS12c] [FSS13a]	
Green-Schwarz mechanism	$BString^{c_2} \xrightarrow{\text{(pb)}} BU$ $\downarrow bomotopy \\ pullback$ $BO \xrightarrow{\frac{1}{2}p_1} B^3U(1)$	secondary invariants	[SSS12] [FSS12a]	

These are the building blocks of Diagram (1). We now explain how this pertains to the M5-brane gauge structure.

¹See, e.g.: [Du96, 6][HLW98, p. 2][Du98, p. 6][NH98, p. 2][Du99b, p. 330][Mo14, 12][CP18, p. 2][Wi19][Du19].

²Here we disregard the super-partner field – the gravitino – for focus of the exposition. In our context and under Hypothesis H the full super-exceptional multiplet of M-theory fields is discussed in [FSS19d][FSS20a].

The open problem of the gauge structure on the M5-brane ([Wi02, p. 6, 15][La19, 3.6]) has two aspects to it:

Q. 1 – Non-abelian gauge structure on M5-branes.

How does a non-abelian gauge group arise on coincident M5-branes?

(Given that the traditional argument for non-abelian gauge fields on D-branes, via perturbative open string scattering, does not apply.)

Q. 2 – Higher gauge structure on M5-branes.

How does this get promoted to a higher gauge group for a non-abelian gerbe gauge theory?

(Given that the gauge potential on the M5-brane is, locally, not a 1-form as in Yang-Mills theory, but a 2-form.)

But the putative higher non-abelian gauge field strength H_3 on the M5-brane is, in the abelian case, an incarnation on the brane of the bulk C_3 -field (see [BLNPST97, (4)]) – for whose description we now invoke *Hypothesis H*:

Hypothesis H in the diagrammatic language from p. 2 says that the C-field in M-Theory on 8-manifolds is classified by a map lifting the tangent bundle classifier to the homotopy quotient of the (homotopy type of) the 4-sphere by its canonical rotation action.

$$\mathbb{R}^{2,1} \times X^{\underbrace{8}} \xrightarrow{\text{M-theory C-field in } \atop \text{J-twisted Cohomotopy}} S^4 /\!\!/ \text{Spin}(5)$$

$$BSpin(8)$$

This hypothesis is motivated from hidden structures found in the "brane scan" [AETW87][Du88] or rather "brane bouquet" [FSS13b][HSS18, 2.1], which were revealed by a re-analysis of the κ -symmetric super p-brane sigma-models through the lens of super homotopy theory [FSS15][FSS16] (reviewed in [FSS19a, §7]). The hypothesis has since been checked to rigorously imply a fair number of phenomena expected to be characteristic of M-theory [FSS19b][FSS19c][SS19a][BSS19][SS19b][SS20][SS20c][SS21] (exposition in [Sch20]):

The single heterotic M5-brane... In particular, we proved in [FSS19d] that the free Perry-Schwarz action functional [PS97][Sc97][APPS97] for the single M5-brane in heterotic M-theory (Hořava-Witten theory [HW95][Wi96] [HW96][LLO97][LOW99][DOPW99][DOPW00][Ov02][Ov18]) *emerges* from *Hypothesis H* (following an analogous but much simpler derivation of the action functional for the single M2-brane in [HSS18, 6.2]).

...is already non-abelian. But the *heterotic* M5-brane is special in that even a single such brane is expected to carry a non-abelian (higher) gauge field, namely for gauge group the quaternionic unitary group $Sp(1) \simeq SU(2)$. This has been argued by identifying heterotic NS5-branes with "small instantons" in the $\mathfrak{so}(32)$ -heterotic gauge field [Wi95][AG96][AM97], and it has dually [Se99] been argued as an effect of gauge enhancement for coincident 5-branes on orientifolds [Mu97]: D5-branes in type I string theory [GP96], M5-branes in heterotic M-theory [GH96].

This second perspective says that a single heterotic M5-brane may be understood as consisting of two coincident M5-branes that are bound as mirror images of the orientifold group action, thus revealing the non-abelian Sp(1) gauge field on the brane as an instance of the general phenomenon expected on coincident M5-branes. Therefore it makes sense to investigate questions **Q. 1** and **Q. 2** already in the case of single heterotic M5-branes:

Coincident M5s → and their worldvolume fields:	N=1 M5	$N_{\text{HET}} = 1 \text{ M5}_{\text{HET}}$ $\parallel \qquad \qquad$	N > 2 M5	argued in [Wi96][AG96] [GP96][GH96]
Gauge field	none	Sp(1)-bundle	unclear	[GF90][GH90]
Higher gauge field	U(1)-gerbe = $BU(1)$ 2-bundle	nonabelian gerbe: String $^{c_2}(4)$ 2-bundle	unclear	derived in Theorem 1 from Hypothesis H

Results. We prove here that *Hypothesis H* implies, for the topological sector of single heterotic M5-branes:


- (1) the emergence of a non-abelian Sp(1)-gauge field structure on the brane worldvolume;
- (2) the emergence of a lift to a higher non-abelian gauge field (non-abelian gerbe field), specifically to a String^{c_2}-structure on the M5-brane worldvolume (as in [Sa11, 2.1], generalizing the notion of [CHZ11]);
- (3) the worldvolume 3-flux H_3 being the corresponding non-abelian Chern-Simons 2-gerbe [FSS13a][NSS12] (cf. [CJMSW05][Wa13]) reflecting a Green-Schwarz-type mechanism on the worldvolume [SSS12][FSS12a] spring

All these results follow from inspection of the single encapsulating homotopy-commutative diagram (1) in Theorem 1, using just basic homotopy theory (see pointers in [NSS12][FSS20b, §A]), the results of [FSS19b][FSS19c] and, for interpreting the factorization as String^{c2}-structure, the concepts developed in [FSS12a], as we now explain.

First we explain the existence of the outer part of diagram (1). All statements, proofs and references that we appeal to here are given in [FSS19b].


The classifying space of J-twisted 4-cohomotopy is the homotopy quotient $S^4 /\!\!/ O(5)$ of the 4-sphere by its canonical action of the orthogonal group. Up to homotopy equivalence this may be taken to be the Borel construction space $(S^4 \times EO(5))/O(5)$, where EO(5) is the total space of the universal O(5)-principal bundle. This is manifestly equipped with a map to ("fibered over") the classifying space BO(5) = (EO(5))/O(5). Pulling this back along $BSpin(5) \rightarrow BO(5)$ and then pushing forward along $BSpin(5) \rightarrow BSpin(8)$ yields the vertical map shown on the right.

A cocycle in J-twisted Cohomotopy on a spin-manifold X^8 is a continuous function c_3 to this classifying space, such that composition with this vertical map classifies (the frame bundle of) the tangent bundle TX^8 .




The quaternionic Hopf fibration $S^7 \xrightarrow{h_{\mathbb{H}}} S^4$ is equivariant with respect to this group action, via the exceptional isomorphism $\operatorname{Sp}(2) \simeq \operatorname{Spin}(5)$. Via Borel-equivariance this induces the map, shown on the right, between the classifying spaces for J-twisted Cohomotopy in degrees 7 and 4, all fibered over $B\operatorname{Spin}(8)$.

The full equivariance group of the quaternionic Hopf fibration is the central product group $Sp(2) \cdot Sp(1) \simeq Spin(5) \cdot Spin(3)$. This is hence singled out as the structure group of 8-manifolds which admit J-twisted Cohomotopy compatibly in degrees 4 and 7, related by the fully Borel-equivariant quaternionic Hopf fibration:



Here we restrict attention to the subgroup $\operatorname{Sp}(2) \simeq \operatorname{Sp}(2) \cdot \mathbb{Z}_2 \subset \operatorname{Sp}(2) \cdot \operatorname{Sp}(1)$ only for sake of exposition.


The homotopy equivalence of classifying spaces $S^4 /\!\!/ \mathrm{Spin}(5) \simeq B \mathrm{Spin}(4)$ is induced by the coset space realization of the 4-sphere $S^4 \simeq \mathrm{Spin}(5)/\mathrm{Spin}(4)$.




Under this equivalence, **the integral cohomology ring** $H^4(B\mathrm{Spin}(4);\mathbb{Z}) \simeq \mathbb{Z}[\frac{1}{2}p_1, \frac{1}{2}\mathcal{X}_4 + \frac{1}{4}p_1]$ is identified as containing the *shifted Euler class* $\widetilde{\Gamma}_4 := \frac{1}{2}\mathcal{X}_4 + \frac{1}{4}p_1$; the plain Euler class $\frac{1}{2}\mathcal{X}_4$ is the fiberwise volume element on S^4 . **Hypothesis H identifies** the C-field flux with the pullback of this class along the given cohomotopy cocycle c_3 .

This way the integral cohomology structure of BSpin(4) implies the shifted flux quantization of the C-field:


$$\tfrac{1}{2}\mathcal{X}_4 + \tfrac{1}{4}p_1 \in H^4(B\mathrm{Spin}(4);\mathbb{Z}) \qquad \Leftrightarrow \qquad G_4 + \tfrac{1}{4}p_1 \in H^4(X^8;\mathbb{Z}).$$





 $S^7 /\!\!/ Sp(2)$

The other homotopy equivalence of classifying spaces $S^7 /\!\!/ \operatorname{Sp}(2) \simeq B\operatorname{Sp}(1)$ is similarly induced by the coset space realization of the 7-sphere as $S^7 \simeq \operatorname{Sp}(2)/\operatorname{Sp}(1)$. This generalizes to the full equivariance group as the homotopy quotient $S^7 /\!\!/ (\operatorname{Sp}(2) \cdot \operatorname{Sp}(1)) \simeq B(\operatorname{Sp}(1) \cdot \operatorname{Sp}(1))$.



The homotopy-commutativity of the outer square

is seen as follows:

With the exceptional isomorphisms

$$Spin(4) \simeq Spin(3) \times Spin(3)$$


$$\simeq Sp(1) \times Sp(1)$$
(2)

and the induced cohomology identifica-

$$H^{\bullet}(B\mathrm{Sp}(1);\mathbb{Z}) \simeq \mathbb{Z}[c_2]$$

 $H^{\bullet}(B\mathrm{Spin}(4);\mathbb{Z}) \simeq \mathbb{Z}[c_2^L, c_2^R]$ (3)

the universal Pontrjagin class $\frac{1}{2}p_1$ on BSpin(4) is identified with the sum of the



second Chern classes of the two Sp(1)-factors (by [FSS19b, Lemma 3.9]), and the Borel-equivariant quaternionic Hopf fibration is identified with the inclusion of the left Sp(1)-factors (by [FSS19b, Prop. 2.22]), as shown here:

$$S^7/\!\!/\operatorname{Sp}(2) \underset{\operatorname{coset space realization}}{\overset{\text{coset space re$$


This makes it manifest that the pullback of the Pontrjagin class along the Borel-equivariant quaternionic Hopf fibration kills the right Chern class summand and retains the left Chern class summand, which is again the plain Chern class on the single Sp(1)-factor that is the domain of the Borel-equivariant quaternionic Hopf fibration:

$$(h_{\mathbb{H}} /\!\!/ \operatorname{Sp}(2))^* (\frac{1}{2}p_1) = c_2.$$
 (5)

Since $B^3U(1) \simeq K(\mathbb{Z},4)$, this equality of cohomology classes comes from a homotopy between their representative maps to classifying spaces, and this is the homotopy commutativity of the outer square above.

The pullback of the integral 4-flux class


 $\widetilde{\Gamma}_4$ along the Borel-equivariant quaternionic Hopf fibration is also seen, from (4), to equal the Chern class. The meaning of this for the C-field flux becomes transparent when equivalently considering the pullback of the difference $\widetilde{\Gamma}_4 - \frac{1}{2}p_1$: The last line of (4) shows that this difference *vanishes* after the pullback. Under the interpretation of vanishing of J-twisted 4-Cohomotopy as its factorization through $h_{\mathbb{H}} /\!\!/ \operatorname{Sp}(2)$, this is the statement that $\frac{1}{2}p_1$ is the *background charge* of the C-field [FSS19b, 3.5].



We now explain the factorization through twisted String structure in the inner part of the diagram (1). The terminology and conceptualization of twisted String structures and their relation to the Green-Schwarz mechanism [GS84] follows [Sa11][SSS12][FSS12a][FSS12b].

The homotopy fiber product of the classifying maps for the Pontrjagin- and Chern-classes (hence the homotopy-pullback of one along the other, denoted "(pb)" in the diagram) is, by definition, the classifying space for $c_2^{\mathrm{Sp}(1)}$ -twisted String(4)-structure [Sa11][SSS12, Def. 2.8]:

classifying space for
$$c_2$$
-twisted String(4)-structure Spin(4)-structure Spin(4)-structure Spin(2) $BString^{c_2}(4)$:= $BSpin(4) \times BSp(1) \times BSp$




This being a *homotopy*-fiber product means that its defining square diagram is filled by a universal homotopy H_3^{univ} , as shown above, that is a coboundary between cocycle representatives of these two characteristic classes after their pullback to the homotopy-fiber product space $B\text{String}^{c_2}(4)$. As such, H_3^{univ} is the homotopy-theoretic reflection of the *Green-Schwarz mechanism* [SSS12]: Upon enhancing all classifying spaces of topological structures in (6) to their corresponding *moduli stacks* of differential structures (according to [FSS10][SSS12][FSS12a][FSS12b], see (11) below) this homotopy is given, locally, by a 3-form flux H_3 whose de Rham differential is the heterotic Bianchi identity:

$$d \underbrace{H_3}_{\text{3-flux form}} = \underbrace{\text{Tr}(R \wedge R)}_{=\frac{1}{2}p_1(R)} - \underbrace{\text{Tr}(F \wedge F)}_{\text{ec}_2(F)}. \tag{7}$$


Here R and F denote, as usual, the local curvature forms of connections on the given Spin(4)- and Sp(1)-bundle, respectively, and we take the traces to be integrally normalized.

The dashed factorization in the inner part of (1) through the homotopy fiber product (6), shown on the right equivalently with domain $S^7 /\!\!/ Sp(2)$, is now the immediate consequence of the commutativity of the outer square, as established above (5), due to the defining universal property of homotopy-fiber products.


In conclusion, this shows that Sp(1) gauge- and $String^{c_2}(4)$ higher gauge structure emerges whenever a cocycle in J-twisted 4-Cohomotopy factors through the classifying space $S^7 /\!\!/ Sp(2)$ of J-twisted 7-Cohomotopy via the Borel-equivariant quaternionic Hopf fibration.



Next we explain, with [FSS15][FSS19c][FSS19d], how this space $S^7 /\!\!/ \mathrm{Sp}(2)$ is, under *Hypothesis H*, the classifying space for the higher gauge field in the M5-brane sigma-model with target space $\mathbb{R}^{2,1} \times X^8$.




The 3-sphere fibration over spacetime associated with the given cocycle c_3 in J-twisted Cohomotopy theory is the homotopy-pullback of the Borel-equivariant quaternionic Hopf fibration along c_3 . This is the direct analog in fibered topological spaces of the construction in rational superspaces (survey in [FSS19a]) which induces the super WZ term of the M5brane sigma model [FSS15], exhibiting 3-spherical T-duality of the M5 [FSS18] [SS18], and from that the full κ -symmetric Lagrangian of the M5sigma model [FSS19d].




In topological spaces

The 3-sphere fiber S^3 is identified with $Sp(1)_R$ under the coset space realizations (4). This follows using basic facts of homotopy theory (the pasting law for homotopy pullbacks, the commutativity of homotopy fibers with homotopy pullbacks, and the looping/delooping equivalence, see [NSS12] for background): First, since homotopy-pullback preserves homotopy fibers, it is identified with the homotopy-fiber of the Borel-equivariant quaternionic Hopf fibration, which by (4) is obtained as follows:



This implies that given any space $\widehat{\Sigma}$ equipped with a map ϕ to spacetime $\mathbb{R}^{2,1} \times X^8$, a lift b_2 , up to specified homotopy (which is notationally suppressed in the diagram on the right), of ϕ to the extended spacetime $\mathbb{R}^{2,1} \times \widehat{X}^8$ is locally on $\widehat{\Sigma}$ a map to $S^3 \simeq \mathrm{Sp}(1)_R$.

Such a map b_2 is a cocycle in twisted 3-Cohomotopy, with the twist being given by the pullback of the C-field cocycle c_3 in 4-Cohomotopy along the embedding field map ϕ . The pasting composition of the homotopy-commutative diagram shown on the right with the homotopy-commutative square shown above (4) identifies the pullback of the shifted integral C-field flux to $\hat{\Sigma}$ with that of the first fractional Pontrjagin class: Concretely, the image of this data in rational homotopy theory (see [FSS16, A] [BSS18, 2.1]) is:



such that
$$dH_3 = \phi^* \left(\widetilde{G}_4 - \frac{1}{2} p_1 \right). \tag{10}$$

This is the field content of the M5-brane sigma model with worldvolume $\widehat{\Sigma}$ and target space $\mathbb{R}^{2,1} \times X^8$. Thus $\mathbb{R}^{2,1} \times \widehat{X}^8$ is identified with the classifying space for M5-brane sigma-model fields, for given background C-field.


Now we are in position to state the main result of the present article:

If the worldvolume field ϕ is indeed an embedding of Spin manifolds, so that $\phi^* \frac{1}{2} p_1(TX^8) = \frac{1}{2} p_1(T\widehat{\Sigma})$, then **the integral lift of the trivialization** (10), **is a twisted String structure** on the M5-brane [Sa11, 2.1] namely a trivialization of $\frac{1}{2} p_1$ relative to a background 4-class $\phi^* \widetilde{G}_4$ in integral cohomology.

Here with *Hypothesis H* we see further substructure in this phenomenon, in that the pullback of the shifted integral C-field flux to the M5 worldvolume is identified with the second Chern class of an Sp(1)-gauge field (4) $\phi^*\widetilde{G}_4 \simeq c_2$, whence we have specifically c_2 -twisted String structure on the worldvolume. In direct analogy with Spin^c-structure, this is called String^{c2}-structure [Sa11, 2.1], with classifying space *B*String^{c2} (6).

In conclusion, the above discussion shows:

- (i) the homotopy-commutativity of the total outer part of the diagram (1);
- (ii) the existence of the dashed factorization map (6) induced by the universal property of the homotopy fiber product *B*String^{c2}, which identifies the field content (9) of the M5-brane sigma model with that of an embedding field together with a higher Sp(1)-gauge field with gauge 2-group String^{c2}.



Hence we have proven the following, for the mathematical formulation of the M-theory C-field subject to *Hypothesis H* according to [FSS19b], and the corresponding formulation of the B-field in the M5-brane sigma-model according to [FSS19c]:

Theorem 1. Assuming C-field charge-quantization in J-twisted 4-Cohomotopy for M-theory on 8-manifolds, then the induced B-field charge quantization in twisted 3-Cohomotopy on the M5-brane worldvolume is equivalently charge-quantization in $String^{c_2}(4)$ -cohomology, according to the homotopy-commutative diagram (1).

Differential String structure on M5-branes. We have focused here on discussion of the topological sector of all fields, classified by a non-abelian generalized but "topological" cohomology; while the full field content is in *differential non-abelian cohomology* [Sch13][FSS20b, §4.2]. For example, the classifying space BSpin(8) in the above discussion is to be promoted to the smooth moduli stack $BSpin(8)_{conn}$ of principal Spin connections [FSS10]. While we do not discuss such differential form data here, one impact of Theorem 1 is that it makes immediate how this discussion should proceed: namely directly by promoting (1) from a diagram in spaces to a diagram of the corresponding smooth ∞ -stacks according to [FSS10][FSS15] (reviewed in [FSS13a]); hence, in particular, promoting the topological twisted String structure classified by the space $BString^{c_2}(4)$ (6) to twisted *differential* string structure classified by a smooth 2-stack $BString^{c_2}_{conn}$, according to [SSS12][FSS12a][FSS12b]:

C	lassifying spaces	pass to shape (trad.: "geometric realization")	Moduli stacks	
for Spin bundles Spin structure	BSpin(n)		$\mathbf{B}Spin(n)_{conn}$	for Spin connections
for $B^{n-2}U(1)$ -bundles (ordinary) n -cohomology	$B^{n-1}U(1)$	promote →	$\mathbf{B}^{n-1}U(1)_{\mathrm{conn}}$	for abelian gerbe connections differential <i>n</i> -cohomology (ordinary)
for String ^c 2 2-bundles twisted String structure	B String $^{c_2}(n)$		B String $^{c_2}(n)_{\text{conn}}$	for String 2-connections twisted differential String structure
for $\Omega S^n \infty$ -bundles n -Cohomotopy	$B\Omega S^n \simeq S^n$		$\mathbf{B}\Omega S_{\mathrm{conn}}^n$	for $\Omega S^n \infty$ -connections differential <i>n</i> -Cohomotopy


Following the method of [SSS08][FSS10] for constructing such differential refinements, the idea of differential String structures on M5-branes has recently been explored in [SäS17] in an attempt to find a higher gauge theoretic interpretation of the action functionals for non-abelian D = 6, $\mathcal{N} = (1,0)$ gauge theories proposed in [SSW11]. But here the conceptual origin and precise flavor of the string gauge field on the M5 seems to have remained open.

Theorem 1 solves this issue by pinpointing specifically $String^{c_2}(4)$ -structure (6) and explaining how this connects to the broader structure of the M5-brane in M-theory, showing that this is the charge quantization on the

M5 that relates to the web of anomaly cancellation conditions in M-theory[FSS19b][FSS19c][SS20]. First consequences have been drawn in [Ro20].

With the stringy field content on the M5-brane in hand, we close by discussing the *Hopf WZ-term* in the M5-brane action functional [In00] *recast as a functional on a* String^{c2} *higher gauge field* (noticing that the Hopf WZ term induces the full Lagrangian density, by the method of [FSS19d]).

The cocycle c_6 in twisted 7-Cohomotopy, on the classifying space \widehat{X}^8 for M5-brane sigma-model fields, arises from the construction of the latter as a homotopy pullback of the Borel-equivariant quaternionic Hopf fibration. This c_6 is the *dual of the C-field*, with flux form G_7 , in the situation that the C-field itself trivializes [FSS19b], as it does after pullback along ϕ to the M5, by (10).



Euler 8-class vanishe

anishing Euler 8-cla


// Sp(2)

The *flux of the dual C-field* on X^8 must measure the *Page charge* [DS91, (41)] of solitonic M2-branes in the spacetime $\mathbb{R}^{2,1} \times X^8$, each stretched along the $\mathbb{R}^{2,1}$ factor. But this requires care:

By the *Poincaré-Hopf theorem* for 7-Cohomotopy [FSS19b, 2.6, 3.7], it follows that if the loci of these M2-branes are assumed to be removed from spacetime (as the locus of the magnetic monopole is in traditional Dirac charge-quantization [Fr11, 16.4e]) then the Euler 8-class χ_8 of χ_8 vanishes. This is the situation of M-theory on 8-manifolds [SVW96][FSS19b, 3.8 & Rmk. 3.1].

The vanishing of the Euler 8-class is homotopy-theoretically imposed [FSS19c, Def. 4.2] by homotopy pullback of the whole diagram (1) to the homotopy fiber $\widehat{BSp(2)}$ of the classifying map for χ_8 :


Thus in the case of vanishing Euler 8-class of X^8 , the cocycle c_6 in J-twisted 7-Cohomotopy on the classifying space for M5-brane sigma model fields lifts to the dotted map shown above.



The universal Hopf WZ-term of the M5-brane is, assuming *Hypothesis H*, a class Γ_7 in integral 7-cohomology on this space (12), as shown in green below. This is the result of [FSS19c, Theorem 4.8]:

Equivalently, this $\widetilde{\Gamma}_7$ is the universal *Page charge* density sourced by M2-branes, and as such is the "dual" of the integral shifted 4-flux $\widetilde{\Gamma}_4$. The integrality of $\widetilde{\Gamma}_7$ is crucial both for the Hopf WZ term to be well defined as a Wess-Zumino term for the M5-brane, as well as its interpretation as M2-brane charge being consistent.

Now, the dashed factorization in (1) directly implies, under pullback to $\widehat{BSp(2)}$ (12) that:



The Hopf WZ term descends to a class on String $_{\chi_o}^{c_2}$ higher gauge fields as shown in purple in the above.

Acknowledgement. This project started at the Jan. 2020 workshop *M-theory and Mathematics* supported by the NYUAD Research Institute. D. Fiorenza thanks New York University Abu Dhabi for hospitality during that time.

References

- [AETW87] A. Achúcarro, J. Evans, P. Townsend and D. Wiltshire, *Super p-Branes*, Phys. Lett. B 198 (1987) 441 [doi:10.1016/0370-2693(87)90896-3]
- [APPS97] M. Aganagic, J. Park, C. Popescu, and J. Schwarz, World-Volume Action of the M Theory Five-Brane, Nucl. Phys. **B 496** (1997), 191-214, [arXiv:hep-th/9701166].
- [AG96] P. Aspinwall and M. Gross, *The* SO(32) *Heterotic String on a K3 Surface*, Phys. Lett. **B387** (1996), 735-742, [arXiv:hep-th/9605131].
- [AM97] P. Aspinwall and D. Morrison, *Point-like Instantons on K3 Orbifolds*, Nucl. Phys. **B503** (1997), 533-564, [arXiv:hep-th/9705104].
- [BaSc07] J. Baez and U. Schreiber, *Higher gauge theory*, in A. Davydov et al. (eds.), *Categories in Algebra*, *Geometry and Mathematical Physics*, Contemp Math 431, AMS 2007, pp 7-30, [arXiv:math/0511710].
- [BLNPST97] I. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. Sorokin, and M. Tonin, *Covariant Action for the Super-Five-Brane of M-Theory*, Phys. Rev. Lett. **78** (1997), 4332-4334, [arXiv:hep-th/9701149].
- [BBS06] K. Becker, M. Becker and J. Schwarz, *String theory and M-theory: a modern introduction*, Cambridge University Press, 2006 [spire:744404].
- [BSS18] V. Braunack-Mayer, H. Sati, and U. Schreiber, *Gauge enhancement for Super M-branes via Parameterized stable homotopy theory*, Comm. Math. Phys. **371** (2019), 197-265, [arXiv:1806.01115].
- [BSS19] S. Burton, H. Sati, and U. Schreiber, *Lift of fractional D-brane charge to equivariant Cohomotopy theory*, J. Geom. Phys. **161**, 104034 (2021) [arXiv:1812.09679].
- [CJMSW05] A. L. Carey, S. Johnson, M. K. Murray, D. Stevenson, and B.-L. Wang, *Bundle gerbes for Chern-Simons and Wess-Zumino-Witten theories*, Commun. Math. Phys. **259** (2005), 577-613, [arXiv:math/0410013] [math.DG].
- [CHZ11] Q. Chen, F. Han, and W. Zhang, *Generalized Witten genus and vanishing theorems*, J. Diff. Geom. **88** (2011), 1-39, [arXiv:1003.2325] [math.DG].
- [CP18] S. M. Chester and E. Perlmutter, *M-Theory Reconstruction from* (2,0) *CFT and the Chiral Algebra Conjecture*, J. High Energy Phys. **2018** (2018) 116, [arXiv:1805.00892].
- [DOPW99] R. Donagi, B. Ovrut, T. Pantev, and D. Waldram, *Standard Models from Heterotic M-theory*, Adv. Theor. Math. Phys. **5** (2002), 93-137, [arXiv:hep-th/9912208].
- [DOPW00] R. Donagi, B. Ovrut, T. Pantev, D. Waldram, *Standard Model Vacua in Heterotic M-Theory*, Strings'99, Potsdam, Germany, 19 24 Jul 1999, [arXiv:hep-th/0001101].
- [Du88] M. Duff, Supermembranes: the first fifteen weeks, Class. Quant. Grav. 5 (1988) 189 [doi:10.1088/0264-9381/5/1/023]
- [Du96] M. Duff, M-Theory (the Theory Formerly Known as Strings), Int. J. Mod. Phys. A11 (1996), 5623-5642, [arXiv:hep-th/9608117].
- [Du98] M. Duff, A Layman's Guide to M-theory, Abdus Salam Memorial Meeting, Trieste, Italy, 19 22 Nov 1997, pp.184-213, [arXiv:hep-th/9805177].
- [Du99a] M. Duff, Anti-de Sitter space, branes, singletons, superconformal field theories and all that, Int. J. Mod. Phys. A14 (1999), 815-844, [arXiv:hep-th/9808100].
- [Du99b] M. Duff (ed.), *The World in Eleven Dimensions: Supergravity, Supermembranes and M-theory*, Institute of Physics Publishing, Bristol, 1999, [ISBN 9780750306720].
- [Du19] M. Duff, in: G. Farmelo, *The Universe Speaks in numbers*, interview 14, 2019, [grahamfarmelo.com/the-universe-speaks-in-numbers-interview-14] at 17:14.
- [DS91] M. Duff and K. Stelle, *Multi-membrane solutions of D* = 11 *supergravity*, Phys. Lett. **B 253** (1991), 113, [spire:299386].
- [FSS12a] D. Fiorenza, H. Sati, and U. Schreiber, *The E*₈ *moduli 3-stack of the C-field*, Comm. Math. Phys. **333** (2015), 117-151, [arXiv:1202.2455].

- [FSS12b] D. Fiorenza, H. Sati, and U. Schreiber, *Multiple M5-branes, String 2-connections, and 7d nonabelian Chern-Simons theory*, Adv. Theor. Math. Phys. **18** (2014), 229-321, [arXiv:1201.5277].
- [FSS12c] D. Fiorenza, H. Sati and U. Schreiber, *Extended higher cup-product Chern-Simons theories*, J. Geom. Phys. **74** (2013), 130-163, [doi:10.1016/j.geomphys.2013.07.011], [arXiv:1207.5449].
- [FSS13a] D. Fiorenza, H. Sati, U. Schreiber, *A higher stacky perspective on Chern-Simons theory*, in D Calaque et al. (eds.), Mathematical Aspects of Quantum Field Theories, Springer 2014, [arXiv:1301.2580].
- [FSS13b] D. Fiorenza, H. Sati, U. Schreiber, Super Lie n-algebra extensions, higher WZW models and super p-branes with tensor multiplet fields Int. J. Geom. Meth. Mod. Phys. 12 (2015) 1550018 [arXiv:1308.5264].
- [FSS15] D. Fiorenza, H. Sati and U. Schreiber, *The WZW term of the M5-brane and differential cohomotopy*, J. Math. Phys. **56** (2015), 102301, [arXiv:1506.07557].
- [FSS16] D. Fiorenza, H. Sati and U. Schreiber, *Rational sphere valued supercocycles in M-theory and type IIA string theory*, J. Geom. Phys. **114** (2017), 91-108, [arXiv:1606.03206].
- [FSS18] D. Fiorenza, H. Sati and U. Schreiber *Higher T-duality of super M-branes*, Adv. Theor. Math. Phys. **24** (2020), [arXiv:1803.05634].
- [FSS19a] D. Fiorenza, H. Sati, and U. Schreiber, *The rational higher structure of M-theory*, Proc. LMS-EPSRC Durham Symposium *Higher Structures in M-Theory*, Aug. 2018, Fortsch. Phys., 2019, [doi:10.1002/prop.201910017], [arXiv:1903.02834].
- [FSS19b] D. Fiorenza, H. Sati, and U. Schreiber, *Twisted Cohomotopy implies M-Theory anomaly cancellation on 8-manifolds*, Commun. Math. Phys. **377** (2020), 1961-2025, [doi:10.1007/s00220-020-03707-2], [arXiv:1904.10207].
- [FSS19c] D. Fiorenza, H. Sati, and U. Schreiber, *Twisted Cohomotopy implies level quantization of the full 6d Wess-Zumino-term of the M5-brane*, Comm. Math. Phys. 2021 (in print) [arXiv:1906.07417].
- [FSS19d] D. Fiorenza, H. Sati and U. Schreiber, *Super-exceptional geometry and super-exceptional embedding construction of M5*, J. High Energy Phys. **2020**, 107 (2020) [arXiv:1908.00042].
- [FSS20a] D. Fiorenza, H. Sati, and U. Schreiber, *Super-exceptional M5-brane model: Emergence of SU(2)-flavor sector*, [arXiv:2006.00012].
- [FSS20b] D. Fiorenza, H. Sati, and U. Schreiber, *The character map in (twisted differential) non-abelian coho-mology*, [arXiv:2009.11909].
- [FSS10] D. Fiorenza, U. Schreiber and J. Stasheff, Čech cocycles for differential characteristic classes, Adv. Theor. Math. Phys. 16 (2012), 149-250, [arXiv:1011.4735].
- [Fr11] T. Frankel, *The Geometry of Physics*, Camb. Univ. Press, 2011, [doi:10.1017/CB09781139061377].
- [GH96] O. J. Ganor and A. Hanany, *Small E*₈ *Instantons and Tensionless Non-critical Strings*, Nucl. Phys. **B474** (1996), 122-140, [arXiv:hep-th/9602120].
- [GP96] E. Gimon and J. Polchinski, *Consistency Conditions for Orientifolds and D-Manifolds*, Phys. Rev. **D54** (1996), 1667-1676, [arXiv:hep-th/9601038].
- [GS84] M. B. Green and J. H. Schwarz, *Anomaly cancellations in supersymmetric D* = 10 gauge theory and superstring theory, Phys. Lett. **B 149** (1984), 117-122, [doi.org/10.1016/0370-2693(84)91565-X].
- [HW95] P. Hořava and E. Witten, *Heterotic and Type I string dynamics from eleven dimensions*, Nucl. Phys. **B460** (1996), 506-5524, [arXiv:9510209] [hep-th].
- [HW96] P. Hořava, E. Witten, *Eleven dimensional supergravity on a manifold with boundary*, Nucl. Phys. **B475** (1996), 94-114, [hep-th/9603142].
- [HLW98] P. S. Howe, N. D. Lambert, and P. C. West, *The Self-Dual String Soliton*, Nucl. Phys. **B515** (1998), 203-216, [arXiv:hep-th/9709014].
- [Hu13] S. Hu, 6D (2,0) theory and M5 branes: A KK mode approach, doctoral dissertation, Texas A & M University, 2013, [hd1:1969.1/151094].
- [HSS18] J. Huerta, H. Sati, and U. Schreiber, *Real ADE-equivariant (co)homotopy of super M-branes*, Comm. Math. Phys. **371** (2019), 425-524, [arXiv:1805.05987].
- [In00] K. Intriligator, Anomaly Matching and a Hopf-Wess-Zumino Term in 6d, $\mathcal{N}=(2,0)$ Field Theories, Nucl. Phys. **B581** (2000), 257-273, [arXiv:hep-th/0001205].

- [LLO97] Z. Lalak, A. Lukas, and B. Ovrut, *Soliton Solutions of M-theory on an Orbifold*, Phys. Lett. **B425** (1998), 59-70, [arXiv:hep-th/9709214].
- [La12] N. Lambert, *M-Theory and Maximally Supersymmetric Gauge Theories*, Ann. Rev. Nucl. Part. Sc. **62** (2012), 285-313, [arXiv:1203.4244].
- [La19] N. Lambert, Lessons from M2's and Hopes for M5's, Proceedings of the LMS-EPSRC Durham Symposium: Higher Structures in M-Theory, August 2018, Fortschritte der Physik 67 (2019), 1910011, [arXiv:1903.02825].
- [LOW99] A. Lukas, B. Ovrut, and D. Waldram, *Five-Branes and Supersymmetry Breaking in M-Theory*, J. High Energy Phys. **9904** (1999), 009, [arXiv:hep-th/9901017].
- [MiSc06] A. Miemiec and I. Schnakenburg, *Basics of M-Theory*, Fortsch. Phys. **54** (2006), 5-72, [arXiv:hep-th/0509137].
- [Mo12] G. Moore, Applications of the six-dimensional (2,0) theories to Physical Mathematics, Felix Klein lectures, Bonn 2012, [http://www.physics.rutgers.edu/~gmoore/FelixKleinLectureNotes.pdf]
- [Mo14] G. Moore, *Physical Mathematics and the Future*, talk at Strings 2014, http://www.physics.rutgers.edu/~gmoore/PhysicalMathematicsAndFuture.pdf
- [Mu97] S. Mukhi, Orientifolds: The Unique Personality Of Each Spacetime Dimension, Workshop on Frontiers of Field Theory, Quantum Gravity and String Theory, Puri, Dec 1996 [arXiv:hep-th/9710004].
- [NH98] H. Nicolai and R. Helling, *Supermembranes and M(atrix) Theory*, In: M. Duff et. al. (eds.), Nonperturbative aspects of strings, branes and supersymmetry, World Scientific, Singapore, 1999, [arXiv:hep-th/9809103].
- [NSS12] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal* ∞-bundles general theory, J. Homotopy Related Structr. **10** (2015), 749–801, [arXiv:1207.0248].
- [Ov02] B. Ovrut, Lectures on Heterotic M-Theory, TASI 2001, [doi:10.1142/9789812702821_0007], [arXiv:hep-th/0201032].
- [Ov18] B. Ovrut, Vacuum Constraints for Realistic Heterotic M-Theories, Symmetry 10 (2018), 723, [www.mdpi.com/2073-8994/10/12/723], [arXiv:1811.08892].
- [PS97] M. Perry and J. Schwarz, *Interacting Chiral Gauge Fields in Six Dimensions and Born-Infeld Theory*, Nucl. Phys. **B 489** (1997), 47-64, [arXiv:hep-th/9611065].
- [RZ16] M. Rho and I. Zahed (eds.) The Multifaceted Skyrmion, World Scientific 2016 [doi:10.1142/9710]
- [Ro20] D. M. Roberts, *Topological sectors for heterotic M5-brane charges under Hypothesis H*, J. High Energy Phys. **2020** (2020), 52, [arXiv:2003.09832].
- [Ro21] C. Roberts, On Mass and Matter, AAPPS Bulletin 31 (2021), 6, [arXiv:2101.08340].
- [RS20] C. Roberts and S. M. Schmidt, *Reflections upon the Emergence of Hadronic Mass*, Eur. Phys. J. Special Topics **229** (2020), 3319-3340, [arXiv:2006.08782].
- [SäS17] C. Saemann and L. Schmidt, *Towards an M5-Brane Model I: A 6d Superconformal Field Theory*, J. Math. Phys. **59** (2018), 043502, [arXiv:1712.06623].
- [SSW11] H. Samtleben, E. Sezgin, and R. Wimmer, (1,0) superconformal models in six dimensions, J. High Energy Phys. 2011 (2011) 62, [arXiv:1108.4060].
- [Sa11] H. Sati, Geometric and topological structures related to M-branes part II: Twisted String- and String^c-structures, J. Australian Math. Soc. **90** (2011), 93-108, [arXiv:1007.5419].
- [Sa13] H. Sati, Framed M-branes, corners, and topological invariants, J. Math. Phys. **59** (2018), 062304, [arXiv:1310.1060] [hep-th].
- [SS18] H. Sati and U. Schreiber, *Higher T-duality in M-theory via local supersymmetry*, Phys. Lett. **B 781** (2018), 694-698, [arXiv:1805.00233].
- [SS19a] H. Sati and U. Schreiber, *Equivariant Cohomotopy implies orientifold tadpole cancellation*, J. Geom. and Phys. **156** (2020), 103775, [arXiv:1909.12277].
- [SS19b] H. Sati and U. Schreiber, Differential Cohomotopy implies intersecting brane observables via configuration spaces and chord diagrams, [arXiv:1912.10425].
- [SS20] H. Sati and U. Schreiber, Twisted Cohomotopy implies M5 anomaly cancellation, [arXiv:2002.07737]
- [SS20b] H. Sati and U. Schreiber, *Proper Orbifold Cohomology*, [arXiv:2008.01101].

- [SS20c] H. Sati and U. Schreiber, *The character map in equivariant twistorial Cohomotopy implies the Green-Schwarz mechanism with heterotic M5-branes* [arXiv:2011.06533].
- [SS21] H. Sati and U. Schreiber, *M/F-Theory as Mf-Theory*, [arXiv:2103.01877].
- [SSS08] H. Sati, U. Schreiber, and J. Stasheff, L_{∞} -algebra connections and applications to String- and Chern-Simons n-transport in: Quantum Field Theory, Birkhäuser (2009), 303-424 [arXiv:0801.3480].
- [SSS12] H. Sati, U. Schreiber, and J. Stasheff, *Twisted differential string and fivebrane structures*, Comm. Math. Phys. **315** (2012), 169–213, [arXiv:0910.4001].
- [Sch13] U. Schreiber, Differential cohomology in a cohesive infinity-topos, [arXiv:1310.7930] [math-ph].
- [Sch20] U. Schreiber, Microscopic brane physics from Cohomotopy theory, talk at: H. Sati (org.), M-Theory and Mathematics. NYUAD Research Institute, January 27-30, 2020, [ncatlab.org/schreiber/files/Schreiber-MTheoryMathematics2020-v200126.pdf]
- [Sc97] J. Schwarz, Coupling a Self-Dual Tensor to Gravity in Six Dimensions, Phys. Lett. **B 395** (1997), 191-195, [arXiv:hep-th/9701008].
- [Se99] A. Sen, *Duality symmetries in string theory*, Current Science 77 (1999), 1635-1645, [jstor:24104472].
- [SVW96] S. Sethi, C. Vafa, and E. Witten, *Constraints on Low-Dimensional String Compactifications*, Nucl. Phys. **B480** (1996), 213-224, [arXiv:hep-th/9606122].
- [Wa13] K. Waldorf, String Connections and Chern-Simons Theory, Trans. Amer. Math. Soc. **365** (2013), 4393-4432, [arXiv:0906.0117] [math.DG].
- [Wi95] E. Witten, *Small Instantons in String Theory*, Nucl. Phys. **B460** (1996), 541-559, [arXiv:hep-th/9511030].
- [Wi96] E. Witten, Strong Coupling Expansion Of Calabi-Yau Compactification, Nucl. Phys. **B 471** (1996), 135-158, [arXiv:hep-th/9602070].
- [Wi02] E. Witten, Conformal field theory in four and six dimensions, in: U. Tillmann (ed.), Topology, Geometry and Quantum Field Theory, Cambridge University Press, 2002, [doi:10.1017/CB09780511526398.017], [arXiv:0712.0157].
- [Wi19] E. Witten, in: G. Farmelo, *The Universe Speaks in numbers*, interview 5, 2019, [grahamfarmelo.com/the-universe-speaks-in-numbers-interview-5] at 21:15.

Domenico Fiorenza, Dipartimento di Matematica, La Sapienza Universita di Roma, Piazzale Aldo Moro 2, 00185 Rome, Italy.

fiorenza@mat.uniroma1.it

Hisham Sati, Mathematics, Division of Science, New York University Abu Dhabi, UAE. hsati@nyu.edu

Urs Schreiber, Mathematics, Division of Science, New York University Abu Dhabi, UAE; on leave from Czech Academy of Science, Prague. us13@nyu.edu