Determination of synthetic cannabinoids in oral fluid by dispersive liquid-liquid microextraction and liquid chromatography tandem mass spectrometry

Alessanda Gentili[a*], Pierpaolo Tomai[a], Roberta Curini[a], Rossella Gottardo[b], Franco Tagliaro[b], Salvatore Fanali[c]

[a] Department of Chemistry, Sapienza University, P.le Aldo Moro 5, 00185, Rome, Italy

[b] Department of Diagnostics and Public Health, University of Verona, Verona, Italy and Institute of Translational Medicine and Biotechnology, Sechenov First Moscow State Medical University, Moscow, Russia····

[c]Teaching Committee of Ph.D. School in Natural Science and Engineering University of Verona, Verona, Italy

[*]Corresponding author, e-mail: alessandra.gentili@uniroma1.it


JWH-018
Area (10^3)
3000
2500
2000
1500
1000
500
0
-500
0
10
20
30
40
50
60
ng/mL
y = 55,062x - 20,73
R² = 0,9992


JWH-019
Area (10^3)
1800
1600
1400
1200
1000
800
600
400
200
0
-200
0
10
20
30
40
50
60
ng/mL
y = 32,24x - 15,659
R² = 0,9988


JWH-073
Area (10^3)
1400
1200
1000
800
600
400
200
0
0
10
20
30
40
50
60
ng/mL
y = 24,024x + 2,9742
R² = 0,9985


JWH200
Area (10^3)
2500
2000
1500
1000
500
0
-500
0
10
20
30
40
50
60
ng/mL
y = 39,207x - 19,16
R² = 0,9914



Figure S1. Matrix-matched calibration curves for the six synthetic cannabinoids