



Promotion of healthy nutrition in primary and secondary cardiovascular disease prevention: a clinical consensus statement from the European Association of Preventive Cardiology

Vassilios S. Vassiliou ¹, Vasiliki Tsampasian¹, Ana Abreu ², Donata Kurpas ³, Elena Cavarretta ^{4,5}, Martin O’Flaherty ⁶, Zoé Colombet ⁶, Monika Siegrist ⁷, Delphine De Smedt ⁸, and Pedro Marques-Vidal ^{9*}

¹Department of Medicine, University of East Anglia, Norfolk and Norwich University Hospital, Colney Lane, Norwich NR4 7UG, UK; ²Centro Reabilitação Cardiovascular, Serviço de Cardiologia, Departamento Coração e Vasos, Centro Hospitalar Universitário Lisboa Norte, CAML, ISAMB, IMP, CCUL, Faculdade de Medicina, Universidade de Lisboa, Avenida Professor Egas Moniz MB, 1649-028 Lisbon, Portugal; ³Department of Family Medicine, Wrocław Medical University, 1 Syrokomli Street, 51-141 Wrocław, Poland; ⁴Department of Medical-Surgical Sciences and Biotechnologies, Sapienza University of Rome, Corso della Repubblica 79, 04100 Latina, Piazzale Aldo Moro 5, 00185 Roma, Italy; ⁵Mediterranea Cardiocentro, Via Orazio 2, 80122 Napoli, Italy; ⁶Department of Public Health, Policy and Systems, Institute of Population Health, Whelan Bld, Liverpool L69 3GB, UK; ⁷Department of Prevention and Sports Medicine, School of Medicine, University Hospital ‘rechts der Isar’, Technical University of Munich, Ismaninger Str. 22, 81675 Munich, Germany; ⁸Department of Public Health and Primary Care, Ghent University, Corneel Heymanslaan 10, 4K3, B-9000 Gent, Belgium; and ⁹Internal Medicine, Lausanne University Hospital and University of Lausanne, Rue du Bugnon 46, CH-1011 Lausanne, Switzerland

Received 23 June 2022; revised 15 February 2023; accepted 22 February 2023; online publish-ahead-of-print 27 February 2023

Background

Poor dietary habits are common and lead to significant morbidity and mortality. However, addressing and improving nutrition in various cardiovascular settings remain sub-optimal. This paper discusses practical approaches to how nutritional counselling and promotion could be undertaken in primary care, cardiac rehabilitation, sports medicine, paediatric cardiology, and public health.

Discussion

Nutrition assessment in primary care could improve dietary patterns and use of e-technology is likely to revolutionize this. However, despite technological improvements, the use of smartphone apps to assist with healthier nutrition remains to be thoroughly evaluated. Cardiac rehabilitation programmes should provide individual nutritional plans adapted to the clinical characteristics of the patients and include their families in the dietary management. Nutrition for athletes depends on the sport and the individual and preference should be given to healthy foods, rather than nutritional supplements. Nutritional counselling is also very important in the management of children with familial hypercholesterolaemia and congenital heart disease. Finally, policies taxing unhealthy foods and promoting healthy eating at the population or workplace level could be effective for prevention of cardiovascular diseases. Within each setting, gaps in knowledge are provided.


Conclusion

This clinical consensus statement contextualizes the clinician’s role in nutrition management in primary care, cardiac rehabilitation, sports medicine, and public health, providing practical examples of how this could be achieved.

* Corresponding author. Tel: +41 (0)21 314 09 34, Fax: +32 9 332 49 94, Email: Pedro-Manuel.Marques-Vidal@chuv.ch

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

Graphical Abstract



Keywords

Nutrition • E-counselling • Implementation • Prevention • Cardiovascular disease

Introduction

Poor dietary habits remain one of the leading causes of non-communicable diseases worldwide.¹ Switching to a healthier diet will help reduce cardiovascular risk factors² and cardiovascular events.³ Despite this, dietary recommendations are irregularly applied in clinical practice.⁴ This lack of implementation stems from the limited training in nutrition provided during medical school,⁵ lack of time or inadequate reimbursement of nutrition provision,⁶ insufficient resources and community-based support, and patient barriers.⁷

For healthy nutritional habits to be promoted in all levels of primary and secondary care, healthcare providers working together with the public need to be active participants in a cycle of training, education, and awareness. This will maximize the chances of implementing change and adopting important dietary measures that are crucial for the prevention of cardiovascular disease (Figure 1). In this clinical consensus statement, practical suggestions on providing nutritional counselling are given as proposals for public health professionals.

Healthy nutrition in primary care


Rationale

Sub-optimal nutrition leads to adverse effects in both acute and chronic diseases.⁸ An unhealthy diet is implicated in the pathogenesis of seven of

the ten leading causes of death worldwide, including heart disease, cancer, and diabetes.¹ On the contrary, better nutritional support and lifestyle approaches to patients in primary care can have substantial short- and long-term benefits both for the patients and the healthcare system under which they are looked after.⁹ In 2019, the World Health Organization issued a report on the importance of nutrition stating that appropriate investment in nutrition could save 3.7 million lives worldwide by the year 2025.¹⁰

Primary care physicians' motivation to provide nutritional care varies according to practice and experience. Several studies investigating barriers to nutritional counselling in primary care practice found that physicians feel insufficiently trained in this.¹¹ However, educational events (including continuing medical education courses), mentorship schemes, and policies created by professional and government organizations all help increase this provision by increasing competence and confidence.¹²

Whilst there are exceptions, the most commonly encountered practice in a primary care setting is to screen for patients in need of nutritional counselling.¹³ These are usually individuals with high risk or with already established chronic cardiovascular disease. Undoubtedly, offering such services to those in need is essential. However, primary care practices should aim to incorporate screening and counselling for all individuals. In this way, primary prevention will be at the centre of patients' healthcare, leading to significant benefits for them and their healthcare providers.^{14,15} Nutrition-focused education and public awareness are essential for the maintenance of healthy eating habits that are beneficial for one's physical health and disease prevention.^{16,17}



Such measures are important for everyone and even more so for individuals at increased risk of acute or chronic disease, such as elderly or frail people, pregnant women, and other groups vulnerable to illness.^{16,17}

Promotion of healthy dietary habits requires adopting a holistic approach towards a patient. Therefore, whilst dietitians and nutritionists routinely give nutritional advice, other team members such as nurses and allied healthcare professionals also play a vital role.¹⁸ A meaningful change towards healthy nutritional routines may significantly impact someone's life. A systematic review of 26 randomized controlled trials including 5500 adults found that primary care dietary consultation effectively improved dietary quality, glycaemic control, and weight control.¹⁹ These effects can have immense benefits, especially for patients with a high risk of cardiovascular disease.

Nutritional counselling in the primary care setting

Nutritional counselling is a two-way interaction; interpreting the dietary assessment results, identifying nutritional problems, and discussing goals with the patient and how to achieve them are all crucial facets of this. Its purpose is to help patients understand important information about the health effects of nutrition and focuses on practical measures to meet their nutritional needs. Moreover, it reinforces the importance of behaviour change towards healthy eating and optimal health.¹⁸ The complexity of counselling means that nutritional counsellors should be formally educated. The use of appropriate materials to increase understanding and retention, including illustrations, food models, home brochures, data collection forms, and referral forms, is encouraged.

The patient's current eating habits (e.g. during a 24-hour dietary recall, a short questionnaire or a tool for diet assessment such as the Healthy Eating Index²⁰) and knowledge should be evaluated when discussing nutrition. The primary care physician should provide education, ascertain patient willingness to change, agree on the pace of change, and support these changes. It should be appreciated that food plays many roles in people's lives and choosing what to eat can create emotional and social pleasures or stress. Furthermore, patients should be reminded that permanent changes occur slowly, following a continuous interaction between the primary care physician and the individual.²¹ The challenge is achieving meaningful clinical results, improving quality

of life, and encouraging a positive attitude to behaviour change by ensuring that patients understand the relevant information, accept the need for nutritional modification, and work towards a new goal. Adhering to change is essential to maintaining healthy nutrition. Ongoing feedback from both the patient and the physician and allowing the appropriate amount of time for serial consultations can significantly improve adherence to behavioural and lifestyle changes.¹⁸ In addition, personal, religious, cultural, economic, and psychological considerations should also be addressed and managed as these are essential factors that may impact on compliance and treatment continuation.

Face-to-face appointments may not always be an option, and e-counselling through virtual consultation has become increasingly used. The development of new technology in e-counselling provides an important opportunity for patients with limited access to consultations.¹⁸ Virtual consultations have the advantages of being easily accessible and often more convenient and can reduce barriers related to patient withdrawal, geographic distance, time constraints, and socio-economic status. In addition, smartphone applications have recently been used to improve nutrition knowledge and contribute to behavioural changes beyond weight loss.^{22,23} Such applications using artificial intelligence can provide accurate and near-real-time dietary assessments and positively influence chronic disease health outcomes. Nevertheless, these applications do not always provide personalized advice tailored to the individual and do not establish a rapport with the patient like the clinicians do. Therefore, they should be a complementary tool to the physician's assessment and not a replacement of it. Relying solely on apps might exclude specific tech-poor segments of the population, such as the elderly or deprived socio-economic groups which might not have the knowledge or the access to the technology. *Table 1* summarizes the key suggestions for primary healthcare professionals providing nutritional counselling.

Knowledge gaps

There is a lack of concise data collection from primary care practices across Europe regarding the implementation of nutritional counselling in cardiovascular disease prevention. Large national and international epidemiological studies are needed to better evaluate current practices and identify areas for improvement.

Most nutrition and diet apps focus on monitoring diet and estimating nutrient content. However, many apps have not been

Table 1 Key messages for effective nutritional counselling in primary care

Primary healthcare physicians should aim to:

- Seek formal and informal training in nutrition and nutritional counselling
- Attend educational events (including continuing medical education courses)
 - Participate in relevant mentorship schemes
 - Keep up to date with relevant literature, professional recommendations, and national and international guidance and policies
- Educate patients regarding the value of nutrition in cardiovascular health
- Ensure that there is appropriate time allowed for consultation when booking appointments, especially for patients at high risk of cardiovascular disease
 - When necessary, use appropriate materials (illustrations, food models, home brochures, and data collection forms) to increase understanding and retention
- Undertake a thorough and personalized evaluation of the patient's nutritional patterns
- Assess the patient's current eating habits and morphometric parameters such as weight, height, body mass index (BMI), and abdominal circumference
 - Discuss and agree with the patient on areas for improvement
 - Together with the patient, establish feasible goals that will lead to improvement of their daily nutritional intake
 - Encourage communication and ongoing feedback to identify and tackle issues that may occur
- Provide holistic approach
- Appreciate that changes in eating habits may cause emotional and psychological stress
 - Provide individualized support and encourage continuous interaction
 - Liaise effectively with other primary healthcare providers (including dietitians, nutritionists, and clinical psychologists) that may have valuable input in patients' care
 - Where appropriate, offer the option of e-counselling or refer the patient to smartphone applications (complementary to the consultation) for further reading, education, and monitoring

validated for primary care practice, and no specific apps are currently recommended. Research into the long-term impact of applications focusing on nutritional e-consulting (critical in primary care practice) is also limited²⁴ and further work in this area should be undertaken.

Nutritional care in cardiac rehabilitation programmes

Rationale

Cardiac rehabilitation programmes have seen an increasing number of complex patients with diabetes, hypertension, and dyslipidaemia, all exacerbated by poor nutrition. All patients starting a cardiac rehabilitation program would benefit from individualized assessment, evaluation, and precision intervention when it comes to nutritional advice in order to facilitate and maintain a successful outcome.^{25,26} If nutrition can be successfully addressed during rehabilitation, it is

expected to lead to a better quality of life, better cardiovascular risk management, and increased survival.²⁷

Despite the cardioprotective benefit of individualized nutritional plans in patients with established disease, adherence and compliance remain major issues in clinical practice.²⁸ Maintaining healthy dietary habits in the long run can be very difficult, with many patients relapsing to earlier eating habits as early as six months after participating in a cardiac rehabilitation programme.^{29,30} This is a recognized issue, which according to the European Association of Preventive Cardiology has five dimensions: the patient, the disease, the healthcare provider, the therapy, and the healthcare system. Each of these dimensions should be optimized simultaneously in order to achieve the best possible adherence to therapy and maximize the benefits of it.³¹

The success of achieving the best possible results not only for the short but also for the long term requires a multi-disciplinary team approach that will support the patients to make and maintain significant changes in their eating habits. All members of the multi-disciplinary team that comprises the cardiac rehabilitation programme, including physicians, nurse practitioners, clinical psychologists, and other allied health professionals have a vital role in this. Importantly, comprehensive nutritional assessment and counselling by a registered dietitian are invaluable in the evaluation, education, and management of patients with cardiovascular disease.^{32,33} Similarly to the assessment of the patient in primary care, adopting a holistic approach is essential, especially for the appropriate individualized management of patients with complex underlying cardiac pathology.

Nutrition as a core component of the cardiac rehabilitation programme

Before establishing a cardiac rehabilitation nutritional plan, information regarding the dietary intake and behaviours of the patient must be collected in addition to the diagnosis and comorbidities of the individual. This includes estimates of daily energy intake and food sources of saturated and *trans*-fat, cholesterol, sodium, and other micronutrients. Food intake habits, number of meals, snacks, frequency of meals outside the home, and alcohol consumption should be collected, for example using short questionnaires. Weight, height, and abdominal circumference should be measured to compute the body mass index and abdominal obesity, as well as other clinical conditions that might require specific dietary counselling, such as impaired renal function or weight modifications in sarcopenia. The collection of this information will allow assessing the targets for nutritional intervention.

Nutritional intervention should be individualized according to the identified target areas, like obesity, diabetes, hypertension, and dyslipidaemia and discussed with the patient, who always needs to be involved in the treatment decision process. Both the patient and their family/healthcare providers (where appropriate) should be educated regarding the nutritional plan. It will be fundamental to transmit how crucial adequate nutritional change is for cardiovascular risk factor control, one of the most challenging purposes of cardiac rehabilitation. Whenever necessary, it is recommended that the nutrition intervention includes models of behavioural change and adherence strategies. During the cardiac rehabilitation programme, cooking classes for patients and family members can be used (when possible) to provide tools in practice for healthy eating. Practitioners must be encouraged to adopt a flexible dietary approach, mindful of patient beliefs and preferences. Moreover, it should be acknowledged that other comorbidities may necessitate deviations from generic advice, with dietitians actively guiding the multi-professional team.²⁷

Regarding weight loss, the patient should be informed about the benefit of weight loss, targets to achieve, and proposed interventions. Adherence of the patient to the prescribed eating plan and/or food education is fundamental to obtain meaningful results. Interventions should combine eating, exercise, and behavioural programs to reduce

total energy intake, aiming for an energetic deficit (500–1000 kcal/day) to attain the desired weight, especially in patients with BMI > 25 kg/m² and/or abdominal circumference > 102 cm in men and 88 cm in women. Targets for weight loss could be >10% in some individuals, aiming for 0.5–1 kg/week for more than six months.^{25,34}

Table 2 provides recommendations for healthcare professionals involved in nutritional counselling of patients in cardiac rehabilitation programmes.

Knowledge gaps

Up to this day, there are no studies evaluating and comparing the effectiveness of the different methods used in cardiac rehabilitation (cooking classes, individual consultations, and group sessions) on adherence and maintenance of healthy dietary habits. As such, there is currently no guideline-recommended approach and future studies are needed to assess the most impactful intervention. The long-term effect of behavioural interventions also remains unclear as there is a lack of evidence for these.

Nutritional care in sports medicine

Rationale

Healthy nutrition and exercise have been recognized as strategic lifestyle components since Hippocrates. They act synergistically to enhance physical performance, reduce recovery time, and boost mental health.²³ Nutrition focuses on energy availability to provide substrate stores to meet the metabolic demand during exercise and recovery. In contrast, exercise training aims to increase metabolic efficiency and athletic skills. Although nutrition and exercise have been studied separately, their interaction is incompletely understood.

In sports medicine, nutrition has a central role in an athlete's life as it is crucial for repeated cycles of high-quality training, optimal performance, and adequate and fast recovery from exercise. Previous studies have shown that athletes who followed nutritional plans by sports nutritionists performed significantly better compared with those who followed a self-chosen nutritional strategy.^{35,36} Furthermore, systemic and local inflammation that follows excessive muscle damage may be ameliorated by a healthy nutritional status.³⁷

Importantly however, insufficient energy intake may have detrimental effects both for recreational and for competitive athletes. Relative energy deficiency in sport (RED-S) is a condition which reflects a significant impairment of physiological functioning and results in a multifaceted deleterious impact on the athlete's psychological, cardiovascular, endocrine, gastrointestinal, and haematological systems.^{38–40} In affected female athletes, amenorrhoea and hypoestrogenism are some of the features of RED-S that are linked with early atherosclerosis, endothelial dysfunction, and disruption of the renin–angiotensin–aldosterone axis.^{41,42} In the more severe form of the syndrome, anorexia nervosa may cause arrhythmias, pericardial effusion, and valvular abnormalities.⁴³

The International Olympic Committee, with the clearly stated goal to *protect the health of the athlete*, has generated a consensus statement in which clinical recommendations are provided for the management of the affected individuals.³⁸ It is of paramount importance that all healthcare providers are aware of the negative impact that chronic or severe energy deficiency can have on recreational or competitive athletes' health. More significantly, healthcare providers should be able to provide guidance for the prevention of this condition that can have a traumatic impact on an athlete's life.

Nutritional counselling for athletes

Education of athletes and, where appropriate, their next of kin is paramount in maintaining an adequate nutritional plan and sufficient energy intake in the long run. Nutrition education interventions have been shown to be effective methods of improving the athletes' sports nutrition knowledge both in individual and team sports.^{44,45} These can be provided by sports medicine experts or sports nutritionists and may be in the form of short classroom-based sessions, group activities, or interactive workshops.^{44,46} Participation in such activities has been shown to be an effective way of improving athletes' eating habits, which in turn has positive impacts on performance.^{47,48}

Both for recreational and for competitive athletes, tailored sport-specific requirements, athlete's characteristics, periodisation of training and competitions, and nutritional goals are essential for selecting the best nutritional strategy.⁴⁹ Therefore, in practice, nutritional plans can be individualized, considering the specific sport, performance goals, body composition, nutrient intake timing, competition planning, and food preferences, as 'one size does not fit all'.

The nutritional recommendations endorse a 'food first' approach with sufficient calories. This comprises macronutrients, mainly carbohydrates and proteins,^{50,51} scaled to the (lean) body mass, with vitamins, minerals, and other micronutrients primarily obtained from fresh vegetables and fruits.^{49,52} Restoring muscular glycogen is a critical factor in post-exercise recovery. Appropriate hydration and fresh fruits appear as effective as sugar sports beverages during recovery.⁵³ Recurrent injuries should prompt investigation into the appropriateness of or the adherence to the nutritional plan chosen.

Many superfoods, nutritional products, supplements, and nutraceuticals have been proposed on the market. Some have been tested in research studies to prove their efficacy; others are believed to be crucial, even without evidence-based data. However, the primary outcomes of studies, such as a lower degree of inflammation, oxidative stress reduction, or a faster metabolic recovery,^{52,54} may not be perceived as essential benefits nor as performance or recovery enhancers by the athletes

Table 2 Key messages for the promotion of effective nutritional counselling in cardiac rehabilitation

Healthcare practitioners, as part of the multi-disciplinary team, should aim to:

Undertake a detailed assessment of the patient's dietary habits

- Evaluate daily energy intake and preferred food sources of nutrients
- Assess number of meals, snacks, and, when applicable, cooking habits
- Explore alcohol consumption; where increased, explore psychosocial factors that may have an impact

Encourage and support behavioural changes that will enable healthy eating habits in the long run

- Educate the patient (and, where appropriate, their carers) on the importance of healthy nutrition for cardiovascular risk factor control, with an emphasis on their co-morbidities
- Provide individualized support tailored to the patient's beliefs and preferences
- When possible, cooking classes for patients and their next of kin can be provided

liaise effectively with other members of the team

- Physicians, nurse practitioners, dietitians, and clinical psychologists should collaborate to support the patient's needs and tackle issues that may occur
- Generate individualized nutritional plans that can be adopted in the patient's daily life

Table 3 Important considerations for the nutritional counselling for athletes

Healthcare practitioners should aim to:

Endorse a 'food first' approach with sufficient calories

- Assess athlete's body composition, specific sport, training plan, and performance goals
- Follow-up athletes regularly (weekly weight measurements) to ensure that there is an energy balance state and prevent significant changes in body composition/weight
- Ongoing communication with the athlete, coach, and next of kin is essential for maintenance of appropriate energy intake

Underpin the importance of appropriate nutritional strategy

- Educate the athlete and, when applicable the next of kin, on the significance of a healthy and tailored nutritional approach that covers their needs in energy demands
- Provide individualized support according to athlete's metabolic demands and dietary preferences

Identify and manage appropriately signs and symptoms of energy deficiency

- Recurrent injuries or illnesses should prompt investigation into the appropriateness of the chosen nutritional plan or adherence to it
- Seek appropriate expertise (sports medicine expert and sports nutritionist) for athletes with or at high risk of developing relative energy deficiency (RED-S)

and coaches. The debate is still ongoing if elite-level athletes need formulated supplementation and healthy nutrition to overcome the strenuous exercise workloads and reduce oxidative stress. This is because polyphenol-rich nutrition seems as effective as supplements.^{54,55} [Table 3](#) summarizes important elements of the nutritional counselling for athletes.

Knowledge gaps

The effect of nutraceuticals, supplements, and probiotics added to a healthy diet and correct nutrition on exercise performance or metabolic recovery has not been established.

One of the most significant challenges in sports nutrition studies is to consider differences in genetics, absorption, metabolism, and excretion, coupled with a high-quality methodology.⁵⁶ Nutrigenomics and nutrigenetics are rapidly growing experimental approaches that use the 'OMICS' technologies (metabolomics, lipidomics, and proteomics) together with genetic sequencing to analyse the individual athlete's response to nutrition and supplementation.⁵⁷ It is likely that implementation of both approaches will lead to a profound advance in sports nutrition; however, no firm evidence is available yet to confirm this.

Nutritional care in children with cardiovascular disease

Rationale

Healthy eating in childhood and adolescence with an appropriate supply of energy and nutrients ensures daily performance and enables growth processes and healthy development. Dietary recommendations for children often focus on obesity. However, healthy nutrition also plays an essential role in children with other chronic diseases. A high-quality

diet with healthy eating habits in childhood and adolescence is of paramount importance in the prevention of cardiovascular diseases and elimination of cardiovascular risk factors.⁵⁸

Children with established cardiovascular disease, such as congenital heart defects, are particularly prone to malnutrition, a serious issue that may result not only in failure to thrive but also in other complications such as heart failure symptoms, anaemia, and recurrent episodes of illnesses.⁵⁹ Ideally, the nutritional care of children and adolescents with cardiovascular disease should engage multiple members of the multi-disciplinary team including paediatricians, cardiologists, specialist nurses, paediatric dietitians, and pharmacists.⁶⁰ It is imperative that all physicians can identify these patients with cardiovascular disease and refer them early to a paediatric cardiology centre that will provide the appropriate support and nutritional counselling.

Nutritional counselling for children with familial hypercholesterolaemia

Familial hypercholesterolaemia is a common genetic disorder characterized by lifelong highly elevated low-density lipoprotein cholesterol levels. These changes are already present at birth leading to early atherosclerotic lesions and premature coronary heart disease. Therefore, screening and therapy should start early, including dietary advice from certified paediatric dietitians or nutritionists.⁶¹

Through individualized counselling, children and families should be supported in consuming less food and beverages with high cholesterol, saturated fat, and *trans*-fat content. Intake of fruit and vegetables, whole grains, fish, or lean meats according to the Mediterranean or heart-healthy diet and maintaining a healthy body weight should be encouraged.⁶² All nutrition advice should be age-adapted to ensure the healthy growth of children,⁶³ in addition to any necessary pharmacotherapy.

There is no information about special alternative diets which might be contraindicated in these children. All dietary strategies should be assessed regularly to ensure that all requirements for healthy growth are met.⁶³

Nutritional counselling for children with congenital heart disease

Congenital heart disease is a common birth defect, which often requires surgical interventions in early childhood. Children with congenital heart disease have a high risk for malnutrition or undernutrition, leading to growth restriction, delays in cardiac surgery, increased pre-operative morbidity, and post-operative complications.⁶⁴

Parents often reported feeding difficulties, including refusal, poor appetite, longer feeding times, and frequent feeding. In addition, structural anomalies of the gastrointestinal tract, gastroesophageal reflux, or neurological problems have additional adverse effects on feeding.^{64,65}

Human breast milk by breastfeeding, bottle, or feeding tubes is recommended for neonates^{64,65} and considered the 'gold standard' independently of the method delivered, in preference to formula milk. Perioperative nutrition, including enteral and parenteral nutrition, is vital for reaching sufficient energy intake and improving post-operative recovery.⁶⁴ After successful cardiac surgery, weight and growth improve immediately in many children. Children with multiple cardiac surgeries or early feeding disorders are at risk for long-term feeding disorders.⁶⁶ [Table 4](#) summarizes the most important messages for nutritional counselling for children and adolescents with cardiovascular disease.

Knowledge gaps

Optimal nutrition in children with congenital heart disease is essential to reduce mortality and to improve normal growth. However, there is a lack of large studies identifying the best nutrition strategies.⁶⁵

Table 4 Summary of key points essential for the nutritional counselling of children and adolescents with cardiovascular disease

Healthcare practitioners should aim to:

Acknowledge the importance of nutritional counselling in children with cardiovascular disease

- Provide nutritional advice that is tailored to the patient's age, body habitus, risk factors, and underlying disease
- Assess patients regularly to ensure that all requirements for healthy growth and development are met

Ensure that appropriate nutritional counselling is provided

- Refer patients and their caregivers early to a paediatric cardiology centre that will provide the appropriate support and nutritional counselling under the care of a specialist multi-disciplinary team
- When appropriate, liaise with certified paediatric dietitians or nutritionists for the selection of the most appropriate nutritional plan for the individual

Nutrition and public health in Europe

Nutritional counselling at the population level

In the current European policy landscape, most countries use information and education to improve dietary behaviours, such as curriculum and community-based nutrition education, developing cooking skills, training of cooking skills, training food service providers and teachers, and nutrition counselling in healthcare settings. Nutritional policies aim to result in a healthier dietary intake, with impact on a large spectrum of diseases including cardiovascular disease and obesity across all socio-economic strata.

The most commonly used nutrition policies are information-based policies providing nutrition information such as food-based dietary guidelines (in 23 European countries^{67,68}) and mass media and informational campaigns (e.g. the worldwide 5-A-Day campaign). However, whilst these can improve population health, they are likely to be failing the socially disadvantaged groups and, in doing so, widening inequalities in diet.^{69,70}

Tackling the food environment can result in more significant and equitable improvements in diet and health. Current popular actions—implemented in 32 European countries⁶⁸—are regulations on food health claims, mandatory nutrient lists on packaged food, and easy-to-understand labelling. A concrete example is France, Belgium, and Spain adopting the NutriScore label, classifying foods and drinks according to five categories of nutritional quality. However, these policies rely on consumer knowledge and behaviour and might result in widening dietary and social inequalities.⁷⁰ Food advertising and marketing restrictions exist in 11 European countries.⁶⁸ Portugal restricts the advertising of products high in energy, salt, sugar, saturated fat, and *trans*-fats before, during, and after children's programmes. The United Kingdom (UK) bans TV advertising of these products before 9 p.m.

Another approach developed in 17 European countries⁶⁸ is to improve the nutritional quality of the whole food supply through reformulation (reduction of salt, saturated fat, and sugar). The UK Food Standards Agency Salt reformulation programme was a success story, substantially decreasing the salt content and intake over a decade.

However, after England replaced it with the Public Health Responsibility Deal in 2011, a less demanding scheme for targeting and monitoring the salt content of food, annual declines in salt intake slowed significantly.⁷¹ Estimates suggest that this lack of robust target setting, monitoring, and enforcement might have resulted in 9900 additional cases of cardiovascular disease in England by 2018.⁷¹ Other policies aim to offer a healthier environment, especially in schools, through the distribution of fruit, vegetables, and milk products in the EU,⁷² bans on vending machines in France, offering healthy food options as default in food service outlets in Norway, and voluntary commitments to reduce portion sizes in Portugal, Spain, and the UK.

Finally, more countries are using fiscal tools to modify consumer food behaviours with health-related food taxes implemented in 12 European countries,⁶⁸ mainly targeting sugary drinks. These policies are not simply designed to reduce consumption but also to induce reformulation of food and drink products, as recently evidenced by the UK Soft Drinks Industry Levy.⁷³ European nutrition policies are greatly diverse and reflect governments' political will to tackle poor diets. However, most policies are information-based, relying on personal nutrition literacy, thus risking widening inequalities. More attention should be given to developing and implementing low-agency population policies, which are admittedly politically challenging.^{70,74} Low-agency population policies are interventions that require little or no involvement of the public, such as taxes on foods or changing the placement and availability of food products. Combining agency (i.e. interventions that require the individuals to engage such as modifying their nutrition) and low-agency policies as an integrated system approach will likely improve diets and narrow inequalities, as proposed by the European Food and Nutrition Action Plan 2015–2020.⁷⁵ A flowchart summarizing the steps for promoting healthy nutrition at the population level is provided in *Figure 2*. However, significant challenges remain to build an integrated food policy framework towards a healthy and sustainable food system.⁷⁶

Nutritional counselling in the workplace and schools

Workplace-based dietary interventions are large-scale strategies to reach as many individuals as possible for prolonged periods.^{77,78} As for nutrition policies, they aim at providing a healthy food environment in the workplace and not to specifically tackle cardiovascular diseases. A limited number of trials of workplace-based interventions seem to be effective in modifying some dietary habits, behaviour, and health outcomes.^{77,78} However, reviews have typically reported that workplace interventions targeting dietary behaviour yield modest improvements and are often poorly implemented, limiting their impact.⁷⁷ In Europe, most countries like the Netherlands, Norway, Germany, and Portugal have dietary guidelines for the workplace based on voluntary action.⁶⁸ For example, in the Netherlands, policies include putting the 'healthier' choices in the most visible places on the displayed range.⁶⁸ Furthermore, some European countries have mandatory standards influencing the food available in specific workplaces like Finland, Germany, Portugal, and UK.⁶⁸

Schools are a unique powerful tool through which nutrition education can be promoted in the society. According to a recent guideline published by the World Health Organization, provision of health education about nutrition is an essential intervention and component of school health services worldwide, as its influence for students can lead to meaningful and sustainable healthy behaviours.⁷⁹ Healthy dietary habits adopted by a young individual can not only reduce their risk of illness but also contribute to their emotional well-being and productivity.⁸⁰ A South Korean cross-sectional survey that included more than 65 000 school-going adolescents demonstrated that healthy dietary habits were associated with lower mental distress and higher psychological well-being, revealing in this way the important link between nutrition




Figure 2 Key steps for the promotion of healthy nutrition at the population level.

and welfare.⁸¹ This, in turn, can also lead to important long-term cost-effective advantages for the wider society. It is therefore imperative for nutritional counselling to be incorporated in primary and secondary education as its impact and benefits can be tremendous for all children and adolescents and, consequently, the society as a whole.

Knowledge gaps

Policymaking is a non-linear process and simply providing scientific evidence is not enough for implementation. Methods to engage with stakeholders and bring the voice of the public to understand the political, legal, and technical feasibility of new policies and interventions alongside public acceptance need to be further developed,⁸² and health practitioners should also advocate for population-level changes.⁸³

Nutritional counselling: is it cost-effective?

Knowledge of the cost-effectiveness of nutritional interventions could prioritise health policy decision-making.⁸⁴ Due to the scarce financial resources, policymakers must choose their investments wisely, focusing on the best value for money. It is well known that preventive interventions score rather well on their cost-effectiveness outcomes.^{85,86} Several studies focussing on the cost-effectiveness of nutritional interventions have been conducted. These often consider obesity or diabetes as intermediate outcomes in developing other chronic diseases such as cardiovascular disease.^{85,87,88}

Previous studies focussed mainly on community-based, school-based, or workplace-based programmes, often showing conflicting results. For example, according to a review, only five out of 23 strategies promoting fruit and vegetable intake in healthy adults were cost-effective.⁸⁹ Context and setting seem to be essential drivers in the cost-effectiveness outcome. A recent study focused on nutrition education and system-level dietary modification in a workplace setting. It resulted in a cost-effective outcome.⁹⁰ Importantly, most interventions are offered as comprehensive programmes focusing on healthy eating and

physical activity, making it difficult to assess the attributable impact of nutrition-related actions.

Recently, increased focus has been on the cost-effectiveness of food labelling (front-of-pack labelling), fiscal taxes, price reductions, and industry agreements with very favourable outcomes. Outstanding outcomes are seen in salt reduction strategies, with taxes and salt reduction by manufacturers and food labelling being cost-effective or even cost-saving.^{88,91} These results are confirmed for other nutrients, such as eliminating industrial *trans*-fats or using a fruit and vegetable subsidy, saturated fat, sugar and salt taxes, and junk food taxes.^{92–94} The magnitude of the tax seems critical here; taxes and subsidies should be used together, with the increased total food price being similar to the healthy food subsidy. Whilst the taxation level is expected to vary between countries, as well as the proposed plans of how this income is going to be used to aid healthier nutrition, it is anticipated that this should be in the region of 10–20%.^{93,95}

Knowledge gaps

The context and setting seem to be essential drivers in the cost-effectiveness outcome. Further research is needed into the prerequisites of cost-effective interventions and how to establish these boundary conditions in practice.

Areas of importance for future recommendations

The European Society of Cardiology Guidelines on cardiovascular disease prevention provide a comprehensive guide for all healthcare professionals who aim to reduce the burden of cardiovascular disease in individuals and at population level.⁹⁶ They reverberate the aforementioned statements regarding the effect of healthy dietary habits on cardiovascular disease prevention and risk factor modification. Policy suggestions for population-based approaches to healthy dietary habits are also in line with this document and highlight the need for appropriate legislation that can promote healthy nutrition in several community settings, such as workplace environment and schools.

Acknowledging that population-based interventions face challenges that may vary in different countries and settings, it would also be prudent to highlight the need for education of both the public and the healthcare professionals. For the former, incorporation of a taught module or optional cooking lessons for students in schools would be helpful. For the latter, implementation of nutritional counselling in the medical curriculum would be extremely beneficial.⁹⁷

It should also be highlighted that an individual's diet is linked with the environmental sustainability. For example, it has been shown that animal-based patterns of food purchases contribute significantly to the annual carbon footprint, whilst plant-based patterns made the smallest contribution to the carbon footprint of the purchases.⁹⁸ Individuals are highly motivated to adopt dietary habits that have an important beneficial impact on the environment.⁹⁹ Therefore, guidelines should focus on raising awareness of the environmental benefit of the nutritional advice and diets suggested.

Finally, nutrition represents not only an impactful driver of many cardiovascular risk factors such as obesity, dyslipidaemia, and diabetes but also a highly evolving and dynamic field. Future guidelines and recommendations should therefore reflect this rapidly changing domain so as to ensure the provision of optimal support in healthcare professionals.

Conclusion

Dietary prevention of cardiovascular disease should be targeted from the cradle to the grave, in the presence or absence of disease, and from an individual to a population level. Dietary recommendations and policies need to be well communicated and aimed to make the healthy choice the easy choice. There are no 'magic' foods, yet one should aspire to a diversified fresh diet, rich in seasonal fruit and vegetables, low in saturated fat meat products, and reduced in salty, sugary, fatty, or ultra-processed foods. Nevertheless, a healthy, environmentally sustainable, cost-effective diet is achievable provided patients, health professionals, food industries, and policymakers join efforts, enabling a reduction in cardiovascular adverse events. The provision of healthy foods and education, starting early in life, has the potential to create good eating habits which can be maintained and passed to the next generations.

Author contributions

Vassilios Vassiliou (MA, MBBS, PhD, FRCP Edin (FACC, FESC), Vasiliki Tsampasian (MSc (MD), Ana Abreu (MD, FESC), Donata Kurpas (MD), Elena Cavarretta (MD, PhD), Martin O'Flaherty (MD, PhD), Zoé Colombet (PhD), Monika Siegrist (PhD), Delphine de Smedt (MD, PhD), and Pedro Marques-Vidal (MD, PhD, FESC).

Supplementary material

Supplementary material is available at *European Journal of Preventive Cardiology* online.

Funding

This work did not receive specific funding.

Conflict of interest: None declared.

Data availability

This manuscript presents the review of previously published data, as such NO new data - hence no need for an availability statement.

References

1. Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama J, et al. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019;**393**:1958–1972.
2. Marques-Vidal P. Comparison of lifestyle changes and pharmacological treatment on cardiovascular risk factors. *Heart* 2020;**106**:852–862.
3. Rees K, Takeda A, Martin N, Ellis L, Wijesekara D, Vepa A, et al. Mediterranean-style diet for the primary and secondary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2019;**3**:3.
4. Marques-Vidal P, Jankowski P, de Bacquer D, Kotseva K. Dietary measures among patients with coronary heart disease in Europe. ESC EORP Euroaspire V. *Int J Cardiol* 2020;**302**:5–14.
5. Crowley J, Ball L, Hiddink GJ. Nutrition in medical education: a systematic review. *Lancet Planet Health* 2019;**3**:e379–e389.
6. Huy C, Diehm C, Schneider S. [Cardiovascular prevention at the general practitioner? First results of a study on attitudes, services, success and barriers in practice]. *Dtsch Med Wochenschr* 2012;**137**:17–22.
7. de Mestral C, Khalatbari-Soltani S, Stringhini S, Marques-Vidal P. Perceived barriers to healthy eating and adherence to dietary guidelines: nationwide study. *Clin Nutr* 2020;**39**:2580–2585.
8. Kris-Etherton PM, Akabas SR, Bales CW, Bistrian B, Braun L, Edwards MS, et al. The need to advance nutrition education in the training of health care professionals and recommended research to evaluate implementation and effectiveness. *Am J Clin Nutr* 2014;**99**:1153S–66S.
9. Polak R, Shani M, Dacey M, Tzuk-Onn A, Dagan I, Malatskey L. Family physicians prescribing lifestyle medicine: feasibility of a national training programme. *Postgrad Med J* 2016;**92**:312–317.
10. World Health Organisation. Essential Nutrition Actions Mainstreaming nutrition through the life-course, <https://www.who.int/publications/i/item/9789241515856> (accessed 6 January 2023).
11. Smith S, Seeholzer EL, Gullett H, Jackson B, Antognoli E, Krejci S, et al. Primary care residents' knowledge, attitudes, self-efficacy, and perceived professional norms regarding obesity, nutrition, and physical activity counseling. *J Grad Med Educ* 2015;**7**:388–394.
12. Crowley J, Ball L, Hiddink GJ. Nutrition care by primary-care physicians: advancing our understanding using the COM-B framework. *Public Health Nutr* 2020;**23**:41–52.
13. Aboueid S, Bourgeault I, Giroux I. Nutrition care practices of primary care providers for weight management in multidisciplinary primary care settings in Ontario, Canada—a qualitative study. *BMC Fam Pract* 2018;**19**:1–9.
14. Berkowitz SA, Terranova J, Randall L, Cranston K, Waters D, Hsu J. Association between receipt of a medically tailored meal program and health care use. *JAMA Intern Med* 2019;**179**:786–793.
15. Downer S, Berkowitz SA, Harlan TS, Olstad DL, Mozaffarian D. Food is medicine: actions to integrate food and nutrition into healthcare. *BMJ* 2020;**369**:m2482.
16. Bush CL, Blumberg JB, El-Sohemy A, Minich DM, Ordoñez JM, Reed DG, et al. Toward the definition of personalized nutrition: a proposal by the American Nutrition Association. *J Am Coll Nutr* 2019;**39**:5–15.
17. Sahyoun NR. Nutrition education for the healthy elderly population: isn't it time? *J Nutr Educ Behav* 2002;**34**:S42–S47.
18. Vasiloglou MF, Fletcher J, Poulia KA. Challenges and perspectives in nutritional counselling and nursing: a narrative review. *J Clin Med* 2019;**8**:1849.
19. Mitchell LJ, Ball LE, Ross LJ, Barnes KA, Williams LT. Effectiveness of dietetic consultations in primary health care: a systematic review of randomized controlled trials. *J Acad Nutr Diet* 2017;**117**:1941–1962.
20. Kennedy ET, Ohls J, Carlson S, Fleming K. The healthy eating index: design and applications. *J Am Diet Assoc* 1995;**95**:1103–1108.
21. Hever J. Plant-based diets: a physician's guide. *Perm J* 2016;**20**:93–101.
22. Pogosova NV, Yufereva YM, Yusubova AI, Ausheva AK, Starodubova AV, Allenov AM, et al. The effectiveness of preventive counseling with the use of remote technologies on medical awareness of cardiovascular risk factors in patients with high and very high cardiovascular risk. *Kardiologiia* 2019;**59**:31–40.
23. Schrauben SJ, Inamdar A, Yule C, Kwiczen S, Krekel C, Collins C, et al. Effects of dietary app-supported tele-counseling on sodium intake, diet quality, and blood pressure in patients with diabetes and kidney disease. *J Ren Nutr* 2022;**32**:39–50.
24. Rollo ME, Burrows T, Vincze LJ, Harvey J, Collins CE, Hutchesson MJ. Cost evaluation of providing evidence-based dietetic services for weight management in adults: in-person versus eHealth delivery. *Nutr Diet* 2018;**75**:35–43.
25. Ambrosetti M, Abreu A, Corrà U, Davos CH, Hansen D, Frederix I, et al. Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020;**28**:460–495.

26. Abreu A, Frederix I, Dendale P, Janssen A, Doherty P, Piepoli MF, et al. Standardization and quality improvement of secondary prevention through cardiovascular rehabilitation programmes in Europe: the avenue towards EAPC accreditation programme: a position statement of the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2020;**28**:496–509.
27. Butler T, Kerley CP, Altieri N, Alvarez J, Green J, Hinchliffe J, et al. Optimum nutritional strategies for cardiovascular disease prevention and rehabilitation (BACPR). *Heart* 2020;**106**:724–731.
28. Ma Y, Olendzki BC, Pagoto SL, Merriam PA, Ockene IS. What are patients actually eating: the dietary practices of cardiovascular disease patients. *Curr Opin Cardiol* 2010;**25**: 518–521.
29. Hämäläinen H, Paalosmaa-Puusa P, Seppänen R, Rastas M, Knuts LR, Voipio-Pulkki LM. Feasibility of, and success in adopting a low-fat diet in coronary patients. *Scand J Rehabil Med* 2000;**32**:180–186.
30. Twardella D, Merx H, Hahmann H, Wüsten B, Rothenbacher D, Brenner H. Long term adherence to dietary recommendations after inpatient rehabilitation: prospective follow up study of patients with coronary heart disease. *Heart* 2006;**92**:635–640.
31. Pedretti RFE, Hansen D, Ambrosetti M, Back M, Berger T, Ferreira MC, et al. How to optimize the adherence to a guideline-directed medical therapy in the secondary prevention of cardiovascular diseases: a clinical consensus statement from the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2022;**15**:1–18.
32. Andersen D, Baird S, Bates T, Chapel DL, Cline AD, Ganesh SN, et al. Academy of nutrition and dietetics: revised 2017 scope of practice for the registered dietitian nutritionist. *J Acad Nutr Diet* 2018;**118**:141–165.
33. Lara-Breitinger K, Lynch M, Kopecsky S. Nutrition intervention in cardiac rehabilitation: a review of the literature and strategies for the future. *J Cardiopulm Rehabil Prev* 2021;**41**: 383–388.
34. Yumuk V, Tsigos C, Fried M, Schindler K, Busetto L, Micic D, et al. European guidelines for obesity management in adults. *Obes Facts* 2015;**8**:402–424.
35. Hottenrott K, Hass E, Kraus M, Neumann G, Steiner M, Knechtle B. A scientific nutrition strategy improves time trial performance by ≈6% when compared with a self-chosen nutrition strategy in trained cyclists: a randomized cross-over study. *Appl Physiol Nutr Metab* 2012;**37**:637–645.
36. Hansen EA, Emanuelsen A, Gertsen RM, Sørensen SR. Improved marathon performance by in-race nutritional strategy intervention. *Int J Sport Nutr Exerc Metab* 2014;**24**: 645–655.
37. Close GL, Baar K, Sale C, Bermon S. Nutrition for the prevention and treatment of injuries in track and field athletes. *Int J Sport Nutr Exerc Metab* 2019;**29**:189–197.
38. Mountjoy M, Sundgot-Borgen J, Burke L, Ackerman KE, Blauwet C, Constantini N, et al. International Olympic Committee (IOC) consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Int J Sport Nutr Exerc Metab* 2018;**28**:316–331.
39. Desbrow B, Slater G, Cox GR. Sports nutrition for the recreational athlete. *Aust J Gen Pract* 2020;**49**:17–22.
40. Diplá K, Kraemer RR, Constantini NW, Hackney AC. Relative energy deficiency in sports (RED-S): elucidation of endocrine changes affecting the health of males and females. *Hormones* 2020;**20**:35–47.
41. O'Donnell E, Goodman JM, Mak S, Murai H, Morris BL, Floras JS, et al. Discordant orthostatic reflex renin-angiotensin and sympathoneural responses in premenopausal exercising-hypoestrogenic women. *Hypertension* 2015;**65**:1089–1095.
42. O'Donnell E, Goodman JM, Harvey PJ. Cardiovascular consequences of ovarian disruption: a focus on functional hypothalamic amenorrhea in physically active women. *J Clin Endocrinol Metab* 2011;**96**:3638–3648.
43. Spaulding-Barclay MA, Stern J, Mehler PS. Cardiac changes in anorexia nervosa. *Cardiol Young* 2016;**26**:623–628.
44. Foo W, Faghy MA, Sparks A, Newbury JW, Gough LA. The effects of a nutrition education intervention on sports nutrition knowledge during a competitive season in highly trained adolescent swimmers. *Nutrients* 2021;**13**:2713.
45. Sánchez-Díaz S, Yanci J, Castillo D, Scanlan AT, Raya-González J. Effects of nutrition education interventions in team sport players: A systematic review. *Nutrients* 2020;**12**:1–18.
46. Patton-Lopez MM, Manore MM, Branscum A, Meng Y, Wong SS. Changes in sport nutrition knowledge, attitudes/beliefs and behaviors following a two-year sport nutrition education and life-skills intervention among high school soccer players. *Nutrients* 2018;**10**:1636.
47. Kavouras SA, Arnaoutis G, Makrillos M, Garagouni C, Nikolaou E, Chira O, et al. Educational intervention on water intake improves hydration status and enhances exercise performance in athletic youth. *Scand J Med Sci Sports* 2012;**22**:684–689.
48. Rossi FE, Landreth A, Beam S, Jones T, Norton L, Cholewa JM. The effects of a sports nutrition education intervention on nutritional Status, sport nutrition knowledge, body composition, and performance during off season training in NCAA division I baseball players. *J Sports Sci Med* 2017;**16**:60.
49. Thomas DT, Erdman KA, Burke LM. American College of Sports Medicine joint position statement. Nutrition and athletic performance. *Med Sci Sports Exerc* 2016;**48**:543–568.
50. Chapman S, Chung HC, Rawcliffe AJ, Izard R, Smith L, Roberts JD. Does protein supplementation support adaptations to arduous concurrent exercise training? A systematic review and meta-analysis with military based applications. *Nutrients* 2021;**13**:1416.
51. Nielsen LLK, Lambert MNT, Jeppesen PB. The effect of ingesting carbohydrate and proteins on athletic performance: a systematic review and meta-analysis of randomized controlled trials. *Nutrients* 2020;**12**:1483.
52. Nieman DC, Gillitt ND, Chen GY, Zhang Q, Sha W, Kay CD, et al. Blueberry and/or banana consumption mitigate arachidonic, cytochrome P450 oxylipin generation during recovery from 75-km cycling: a randomized trial. *Front Nutr* 2020;**7**:121.
53. Nieman DC, Gillitt ND, Sha W, Esposito D, Ramamoorthy S. Metabolic recovery from heavy exertion following banana compared to sugar beverage or water only ingestion: a randomized, crossover trial. *PLoS One* 2018;**13**:e0194843.
54. Cavarretta E, Peruzzi M, del Vescovo R, Di Pilla F, Gobbi G, Serdoz A, et al. Dark chocolate intake positively modulates redox status and markers of muscular damage in elite football athletes: a randomized controlled study. *Oxid Med Cell Longev* 2018;**2018**:1–10.
55. Nieman DC, Gillitt ND, Knab AM, Shanely RA, Pappan KL, Jin F, et al. Influence of a polyphenol-enriched protein powder on exercise-induced inflammation and oxidative stress in athletes: a randomized trial using a metabolomics approach. *PLoS One* 2013;**8**:e72215.
56. Swinton PA, Hemingway BS, Saunders B, Gualano B, Dolan E. A statistical framework to interpret individual response to intervention: paving the way for personalized nutrition and exercise prescription. *Front Nutr* 2018;**5**:41.
57. Guest NS, Horne J, Vanderhout SM, El-Sohemy A. Sport nutrigenomics: personalized nutrition for athletic performance. *Front Nutr* 2019;**6**:8.
58. Funtikova AN, Navarro E, Bawaked RA, Fito M, Schröder H. Impact of diet on cardiometabolic health in children and adolescents. *Nutr J* 2015;**14**:1–11.
59. El-Koofy N, Mahmoud AM, Fattouh AM. Nutritional rehabilitation for children with congenital heart disease with left to right shunt. *Turk J Pediatr* 2017;**59**:442–451.
60. Marino LV, Johnson MJ, Davies NJ, Kidd CS, Fienberg J, Richens T, et al. Improving growth of infants with congenital heart disease using a consensus-based nutritional pathway. *Clin Nutr* 2020;**39**:2455–2462.
61. Nordestgaard BG, Chapman MJ, Humphries SE, Ginsberg HN, Masana L, Descamps OS, et al. Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease consensus statement of the European Atherosclerosis Society. *Eur Heart J* 2013;**34**:3478–3490.
62. Wiegman A, Gidding SS, Watts GF, Chapman MJ, Ginsberg HN, Cuchel M, et al. Familial hypercholesterolaemia in children and adolescents: gaining decades of life by optimizing detection and treatment. *Eur Heart J* 2015;**36**:2425–2437.
63. Ramaswami U, Humphries SE, Priestley-Barnham L, Green P, Wald DS, Capps N, et al. Current management of children and young people with heterozygous familial hypercholesterolaemia—HEART UK statement of care. *Atherosclerosis* 2019;**290**:1–8.
64. Herridge J, Tedesco-Bruce A, Gray S, Floh AA. Feeding the child with congenital heart disease: a narrative review. *Pediatr Med* 2021;**4**:1–15.
65. Martini S, Beghetti I, Annunziata M, Aceti A, Galletti S, Ragni L, et al. Enteral nutrition in term infants with congenital heart disease: knowledge gaps and future directions to improve clinical practice. *Nutrients* 2021;**13**:932.
66. Maurer I, Latal B, Geissmann H, Knirsch W, Bauersfeld U, Balmer C. Prevalence and predictors of later feeding disorders in children who underwent neonatal cardiac surgery for congenital heart disease. *Cardiol Young* 2011;**21**:303–309.
67. Hawkes C, Jewell J, Allen K. A food policy package for healthy diets and the prevention of obesity and diet-related non-communicable diseases: the NOURISHING framework. *Obes Rev* 2013;**14**:159–168.
68. World Cancer Research Fund International. NOURISHING policy database, https://policydatabase.wcrf.org/level_one?page=nourishing-level-one (accessed 13 January 2023).
69. Velardo S. The nuances of health literacy, nutrition literacy, and food literacy. *J Nutr Educ Behav* 2015;**47**:385–389.e1.
70. Adams J, Mytton O, White M, et al. Why are some population interventions for diet and obesity more equitable and effective than others? The role of individual agency. *PLoS Med* 2016;**13**:e1001990.
71. Laverty AA, Kyridemos C, Seferidis P, Vamos EP, Pearson-Stuttard J, Collins B, et al. Quantifying the impact of the public health responsibility deal on salt intake, cardiovascular disease and gastric cancer burdens: interrupted time series and microsimulation study. *J Epidemiol Community Health* 2019;**73**:881–887.
72. European Commission. School fruit, vegetables and milk scheme, <https://agriculture.ec.europa.eu/common-agricultural-policy/market-measures/school-fruit-vegetables-and-milk-scheme> (13 January 2023).
73. Scarborough P, Adhikari V, Harrington RA, Elhussein A, Briggs A, Rayner M, et al. Impact of the announcement and implementation of the UK Soft Drinks Industry Levy on sugar

- content, price, product size and number of available soft drinks in the UK, 2015–19: a controlled interrupted time series analysis. *PLoS Med* 2020;**17**:e1003025.
74. Capewell S, O'Flaherty M. Can dietary changes rapidly decrease cardiovascular mortality rates? *Eur Heart J* 2011;**32**:1187–1189.
 75. World Health Organization Regional Office for Europe. European Food and Nutrition Action Plan 2015–2020. <http://www.euro.who.int/pubrequest> (2015, accessed 13 January 2023).
 76. de Schutter O, Jacobs N, Clément C. A 'common food policy' for Europe: how governance reforms can spark a shift to healthy diets and sustainable food systems. *Food Policy* 2020;**96**:101849.
 77. Wolfenden L, Goldman S, Stacey FG, Grady A, Kingsland M, Williams CM, et al. Strategies to improve the implementation of workplace-based policies or practices targeting tobacco, alcohol, diet, physical activity and obesity. *Cochrane Database Syst Rev* 2018;**11**:CD012439.
 78. Glympi A, Chasioti A, Bälter K. Dietary interventions to promote healthy eating among office workers: a literature review. *Nutrients* 2020;**12**:3754.
 79. World Health Organization, Unesco. *WHO Guideline on school health services* Geneva: World Health Organization; 2021. p73.
 80. Baltag V, Sidaner E, Bundy D, Guthold R, Nwachukwu C, Engesveen K, et al. Realising the potential of schools to improve adolescent nutrition Valentina Baltag and colleagues argue that school health programmes have the potential to mitigate a growing epidemic of malnutrition in children and adolescents. *BMJ* 2022;**379**:e067678.
 81. Hong SA, Peltzer K. Dietary behaviour, psychological well-being and mental distress among adolescents in Korea. *Child Adolesc Psychiatry Ment Health* 2017;**11**:56.
 82. Lloyd-Williams F, Masters R, Hyseni L, Denny ES, O'Flaherty M, Capewell S. The QUEST for effective and equitable policies to prevent non-communicable diseases: co-production lessons from stakeholder workshops. *Int J Health Policy Manag* 2021;**10**:638.
 83. Chapman S. Advocacy for public health: a primer. *J Epidemiol Community Health* 2004;**58**:361–365.
 84. Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. Methods for the economic evaluation of health care programmes. In: Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, George W (eds.), *Torrance—Oxford University Press*. 4th ed. New York, USA: Oxford University Press; 2023.
 85. Ananthapavan J, Sacks G, Moodie M, Carter R. Economics of obesity—learning from the past to contribute to a better future. *Int J Environ Res Public Health* 2014;**11**:4007–4025.
 86. World Health Organization (WHO). The case for investing in public health. A public health summary report for EPHO 8. In: *Europe ROF*. Copenhagen, Denmark: Regional Office for Europe; 2020. p1–40.
 87. Brunner E, Cohen D, Toon L. Cost effectiveness of cardiovascular disease prevention strategies: a perspective on EU food based dietary guidelines. *Public Health Nutr* 2001;**4**:711–715.
 88. Schorling E, Niebuhr D, Kroke A. Cost-effectiveness of salt reduction to prevent hypertension and CVD: a systematic review. *Public Health Nutr* 2017;**20**:1993–2003.
 89. Cobiac LJ, Vos T, Lennert Veerman J. Cost-effectiveness of interventions to promote fruit and vegetable consumption. *PLoS One* 2010;**5**:e14148.
 90. Fitzgerald S, Murphy A, Kirby A, Geaney F, Perry IJ. Cost-effectiveness of a complex workplace dietary intervention: an economic evaluation of the Food Choice at Work study. *BMJ Open* 2018;**8**:e019182.
 91. Webb M, Fahimi S, Singh GM, Khatibzadeh S, Micha R, Powles J, et al. Cost effectiveness of a government supported policy strategy to decrease sodium intake: global analysis across 183 nations. *BMJ* 2017;**356**:i6699.
 92. Marklund M, Zheng M, Veerman JL, Wu JHY. Estimated health benefits, costs, and cost-effectiveness of eliminating industrial trans-fatty acids in Australia: a modelling study. *PLoS Med* 2020;**17**:e1003407.
 93. Niebylski ML, Redburn KA, Duhany T, Campbell NR. Healthy food subsidies and unhealthy food taxation: a systematic review of the evidence. *Nutrition* 2015;**31**:787–795.
 94. Cobiac LJ, Tam K, Veerman L, Blakely T. Taxes and subsidies for improving diet and population health in Australia: a cost-effectiveness modelling study. *PLoS Med* 2017;**14**:e1002232.
 95. Blakely T, Cleghorn C, Mizdrak A, Waterlander W, Nghiem N, Swinburn B, et al. The effect of food taxes and subsidies on population health and health costs: a modelling study. *Lancet Public Health* 2020;**5**:e404–e413.
 96. Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Bäck M, et al. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur J Prev Cardiol* 2022;**29**:5–115.
 97. Aspary KE, van Horn L, Carson JAS, Wylie-Rosett J, Kushner RF, Lichtenstein AH, et al. Medical nutrition education, training, and competencies to advance guideline-based diet counseling by physicians: a science advisory from the American Heart Association. *Circulation* 2018;**137**:e821–e841.
 98. Meinilä J, Hartikainen H, Tuomisto HL, Uusitalo L, Vepsäläinen H, Saarinen M, et al. Food purchase behaviour in a Finnish population: patterns, carbon footprints and expenditures. *Public Health Nutr* 2022;**25**:3265–3277.
 99. Culliford A, Bradbury J. A cross-sectional survey of the readiness of consumers to adopt an environmentally sustainable diet. *Nutr J* 2020;**19**:1–13.