







ORIGINAL RESEARCH

Differences in Postoperative Atrial Fibrillation Incidence and Outcomes After Cardiac Surgery According to Assessment Method and Definition: A Systematic Review and Meta-Analysis

Roberto Perezgrovas-Olaria , MD*; Talal Alzghari , MD*; Mohammed Rahouma , MD; Arnaldo Dimagli, MD; Lamia Harik, MD; Giovanni J. Soletti , MD; Kevin R. An , MD; Tulio Caldonazo, MD; Hristo Kirov , MD; Gianmarco Cancelli , MD; Katia Audisio, MD; Mohammad Yaghmour , MS; Hillary Polk, MS; Rajbir Toor, MS; Swetha Sathi , MS; Michelle Demetres , MLIS; Leonard N. Girardi, MD; Giuseppe Biondi-Zoccai, MD, MStat; Mario Gaudino , MD, MSCE, PhD

BACKGROUND: Postoperative atrial fibrillation (POAF) is the most frequent complication of cardiac surgery. Despite clinical and economic implications, ample variability in POAF assessment method and definition exist across studies. We performed a study-level meta-analysis to evaluate the influence of POAF assessment method and definition on its incidence and association with clinical outcomes.

METHODS AND RESULTS: A systematic literature search was conducted to identify studies comparing the outcomes of patients with and without POAF after cardiac surgery that also reported POAF assessment method. The primary outcome was POAF incidence. The secondary outcomes were in-hospital mortality, stroke, intensive care unit length of stay, and postoperative length of stay. Fifty-nine studies totaling 197 774 patients were included. POAF cumulative incidence was 26% (range: 7.3%–53.1%). There were no differences in POAF incidence among assessment methods (27%, [range: 7.3%–53.1%] for continuous telemetry, 27% [range: 7.9%–50%] for telemetry plus daily ECG, and 19% [range: 7.8%–42.4%] for daily ECG only; $P>0.05$ for all comparisons). No differences in in-hospital mortality, stroke, intensive care unit length of stay, and postoperative length of stay were found between assessment methods. No differences in POAF incidence or any other outcomes were found between POAF definitions. Continuous telemetry and telemetry plus daily ECG were associated with higher POAF incidence compared with daily ECG in studies including only patients undergoing isolated coronary artery bypass grafting.

CONCLUSIONS: POAF incidence after cardiac surgery remains high, and detection rates are variable among studies. POAF incidence and its association with adverse outcomes are not influenced by the assessment method and definition used, except in patients undergoing isolated coronary artery bypass grafting.

Key Words: assessment method ■ cardiac surgery ■ definition ■ incidence ■ postoperative atrial fibrillation

Correspondence to: Mario Gaudino, MD, MSCE, PhD, Department of Cardiothoracic Surgery, Weill Cornell Medicine, 525 E 68th St, New York, NY 10065. Email: mfg9004@med.cornell.edu

*R. Perezgrovas-Olaria and T. Alzghari contributed equally.

This article was sent to Luciano A. Sposato, MD, MBA, FRCPC, Associate Editor, for review by expert referees, editorial decision, and final disposition.

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.123.030907>

For Sources of Funding and Disclosures, see page 12.

© 2023 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- In this study-level meta-analysis, we found no difference in incidence of postoperative atrial fibrillation by assessment method, whether telemetry only, telemetry plus ECG, or ECG only were used, and there was also no difference in the incidence of postoperative atrial fibrillation irrespective of the definition used.

What Are the Clinical Implications?

- There was no difference among assessment methods in incidence of postoperative atrial fibrillation or the association of postoperative atrial fibrillation with adverse clinical outcomes.

Nonstandard Abbreviations and Acronyms

POAF postoperative atrial fibrillation

Postoperative atrial fibrillation (POAF) is the most frequent complication of cardiac surgery, with an incidence ranging from 15% to 40%.¹ Often regarded as a transient event triggered by inflammation after surgery, POAF has been associated with worse clinical outcomes (including mortality, stroke, and heart failure)^{2,3} and increased cost of care.^{4,5} Despite the proven success of medical⁶ and surgical⁷ strategies for POAF prevention, a high proportion of patients remains affected, resulting in efforts to elucidate the pathophysiologic mechanisms of, and risk factors for, POAF as potential targets for intervention.

The term POAF generally refers to new-onset atrial fibrillation during the postoperative hospitalization period.^{1,8,9} However, no consensus definition for POAF has been established by professional societies,^{10,11} leading to marked heterogeneity in POAF definitions across studies. Some groups have reported POAF episodes regardless of duration or need for treatment,^{12,13} others have used duration-based POAF definitions (with arbitrarily defined cutoffs to report the arrhythmia ranging from 30 seconds to 60 minutes),^{14,15} and others have reported only those POAF episodes that required treatment or intervention.^{16,17} In addition, there is variability in the POAF assessment methods employed across studies, ranging from monitoring with continuous telemetry during the whole postoperative hospitalization to the use of daily ECG only. The heterogeneity in POAF assessment method and definition opens questions on the interpretation and generalizability of published studies.

We performed a systematic review and meta-analysis to evaluate the incidence of POAF and its association with clinical outcomes according to the POAF assessment method and definition used in individual studies.

METHODS

The present review was registered in the National Institute for Health Research International Registry of Systematic Reviews (CRD42023399670). The article is compliant with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guideline.¹⁸ Institutional review board approval was waived. Informed consent was not required.

Search Strategy

A comprehensive literature search was performed by a medical librarian (M.D.) to identify studies that compared outcomes of patients who developed POAF versus patients who did not after cardiac surgery. Searches were run on July 6, 2022 and updated on May 31, 2023 in the following databases: Ovid MEDLINE (ALL; 1946 to present), Ovid EMBASE (1974 to present), and The Cochrane Library (Wiley; 1992 to present). The complete search strategy for Ovid MEDLINE is available in [Table S1](#).

Study Selection and Data Extraction

After deduplication, records were screened by 2 independent reviewers (L.H. and R.P.O.) using Microsoft Excel version 16.73. Any discrepancies were adjudicated by the senior author (M.G.). Titles and abstracts were reviewed against predefined inclusion and exclusion criteria. Studies were considered for inclusion if they compared outcomes of patients who developed POAF after noncongenital cardiac surgery versus patients who did not. Animal studies, abstracts, case reports, commentaries, editorials, expert opinions, conference presentations, and studies that did not report POAF assessment method were excluded. The full text of the selected articles was pulled for a second round of eligibility screening. The reference lists were also reviewed for relevant studies not captured by the original search. The methodological quality of the included studies was assessed by 2 reviewers (L.H. and R.P.O.) in 3 domains: (1) cohort selection and comparability, (2) reporting of POAF assessment method and POAF incidence, and (3) reporting of POAF definition. This assessment was based on the Newcastle-Ottawa Scale. Details of POAF definition in each study are provided in [Table S2](#).

Two investigators (T.C. and R.P.O.) independently performed data extraction, and the accuracy was verified by the senior author (M.G.). The variables

included were study characteristics (publication year, institution, country of origin, study period, type of surgery, sample size, POAF definition, POAF incidence, and POAF assessment method), patients' demographic characteristics (age, sex, left ventricular ejection fraction, hypertension, diabetes, chronic obstructive pulmonary disease, prior cerebrovascular accident, prior myocardial infarction, preoperative use of beta blockers, prior cardiac surgery, and chronic kidney disease), in-hospital mortality, stroke, intensive care unit (ICU) length of stay (LOS) and postoperative LOS.

Outcomes

The primary outcome was POAF incidence according to the POAF assessment method used. Three different assessment methods were identified: (1) continuous telemetry until hospital discharge, (2) continuous telemetry during ICU stay followed by daily ECG while patients were in the regular ward, and (3) daily ECG only.

The secondary outcomes were in-hospital mortality, stroke, ICU LOS, and postoperative LOS according to the POAF assessment method used.

Secondary Analyses

In the secondary analyses, the primary and secondary outcomes were analyzed based on the definition of POAF used in individual studies: (1) intervention-based definition, which included only POAF episodes requiring treatment, and (2) nonintervention-based definition, which included POAF regardless of episode duration or need for treatment. Details of the assigned POAF definition category in each study are provided in [Table 1](#).

Subgroup and Additional Analyses

A subgroup analysis for the primary outcome was performed in studies including only patients undergoing isolated coronary artery bypass grafting (CABG). An additional analysis looking at the trend in POAF incidence over the study period was also performed.

Statistical Analysis

Categorical variables were extracted as numbers and continuous variables were extracted as mean and SD.

For each assessment method, the incidence of POAF across studies was pooled as an overall proportion (overall number of events/total number of patients) using an inverse variance method, which takes into account the weight of each study relative to the study sample size. Both common and random effects estimates were reported. Ninety-five percent CIs were estimated using the Clopper–Pearson interval.

Subsequently, pooled proportions were compared between the different types of monitoring using a standard test for heterogeneity across the subgroup results, as previously described.¹⁹ Similarly, categorical outcomes were compared by assessment methods.

The proportion of POAF across monitoring methods was also compared using chi-square test among the subgroups.

Trend in the postoperative incidence of POAF during the study years was investigated using the locally estimated scatterplot smoothing.

For continuous outcomes (ICU and postoperative LOS), the LOS across studies was pooled as an overall mean using an inverse variance method and subsequently compared using a test for subgroup differences.

Statistical heterogeneity was assessed with I^2 , which describes the percentage of the variability in the effect estimates due to heterogeneity rather than sampling error. Low, moderate, and high heterogeneity were defined as $I^2 < 25%$, $25%$ to $50%$, and $> 50%$, respectively.²⁰ Tau-squared using DerSimonian–Laird model was used to estimate the between-study variance.

Funnel plot and Egger's test were used to assess for publication bias graphically and quantitatively.

Univariable, random-effects meta-regression was used to explore the association between POAF incidence and the rigor of the assessment method used. Compared with the main analysis, which is based on pairwise comparisons, the meta-regression uses the monitoring methods as an ordinal, 3-level variable, where continuous telemetry until hospital discharge is considered more rigorous than telemetry during ICU stay followed by daily ECG in the regular ward, which is considered more rigorous than daily ECG only. Moreover, logistic regression with robust SE was performed for the association of POAF and categorical variables after reproducing the individual level data.

Statistical analyses were performed in R version 4.0.3 (R Foundation for Statistical Computing) using the package *meta*.

Data Availability

Data collected for the study will be made available by the corresponding author upon reasonable request after publication.

RESULTS

Study and Patient Characteristics

Among the 8974 identified (6212 screened) articles, a total of 59 studies published between 1993 and 2022 were included in the present analysis.^{3–5,7,12–17,21–69} The Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram outlining the study

Table 1. Characteristics of the Included Studies

Study, year	Institution or trial name	Country	Study period	Type of surgery	Total number of patients	Incidence of POAF	Assessment method	POAF definition category (duration, if applicable)
Creswell, 1993 ²¹	Washington University Medical Center	United States	1986–1991	Multiple	3983	34.6%	Telemetry	Not reported
Aranki, 1996 ¹⁶	Brigham and Women's Hospital	United States	1993–1994	CABG	570	33.2%	Telemetry	Intervention-based
Stamou, 2000 ¹⁷	Washington Hospital Center	United States	1987–1999	CABG	969	21.3%	Telemetry	Intervention-based
Tamris, 2000 ²²	St. Luke's Hospital	United States	1992–1994	CABG	216	25.5%	Telemetry+ECG	Nonintervention-based (≥30min)
Hakala, 2002 ²³	Kuopio University Hospital	Finland	1992–1996	CABG	3676	31%	ECG	Not reported
Silva, 2004 ²⁴	Institute of Cardiology of Rio Grande/Fundacion Universitaria de Cardiologia	Brazil	2002	Multiple	158	28.5%	Telemetry+ECG	Nonintervention-based (≥15 min)
Villareal, 2004 ¹²	Texas Heart Institute	United States	1993–1999	CABG	6475	15.4%	ECG	Nonintervention-based (any duration)
Kalavrouziotis, 2007 ²⁵	Maritime Heart Center	Canada	1995–2003	CABG±AVR	7347	27.9%	Telemetry+ECG	Intervention-based
Mariscalco, 2007 ²⁶	Umea University Hospital	Sweden	1994–2004	Multiple	8434	25.6%	Telemetry+ECG	Nonintervention-based (≥15 min)
Nisanoglu, 2007 ¹³	Turgut Ozal Medical Center	Turkey	2001–2005	CABG	426	21.4%	Telemetry+ECG	Nonintervention-based (any duration)
Mariscalco, 2008 ²⁷	Varese University Hospital/Monizo Cardiology Center	Italy	2000–2005	CABG	1832	31.1%	Telemetry	Nonintervention-based (≥15 min)
Ahlsson, 2010 ²⁸	Örebro University Hospital	Sweden	1999–2000	CABG	571	28.9%	Telemetry+ECG	Nonintervention-based (≥1 min)
Bramer, 2010 ²⁹	Catharina Hospital	Netherlands	2003–2007	CABG	5098	22%	Telemetry+ECG	Nonintervention-based (≥30min)
Shirzad, 2010 ³⁰	Tehran Heart Center	Iran	2002–2008	Multiple	15580	7.2%	Telemetry	Nonintervention-based (≥5 min)
Attaran, 2011 ³¹	Liverpool Heart and Chest Hospital	United Kingdom	1998–2009	Multiple	17379	28.7%	ECG	Nonintervention-based (any duration)
Bramer, 2011 ³²	Catharina Hospital	Netherlands	2003–2010	Mitral valve repair/replacement ± CABG±tricuspid valve repair/replacement	856	42.2%	Telemetry+ECG	Nonintervention-based (≥30min)
Girerd, 2012 ³³	Laval Hospital	Canada	2000–2007	CABG	6728	27.8%	Telemetry+ECG	Intervention-based
Helgadottir, 2012 ³⁴	Landspítali Hospital	Iceland	2002–2006	CABG or AVR	744	43.8%	Telemetry+ECG	Nonintervention-based (≥5 min)

(Continued)

Table 1. Continued

Study, year	Institution or trial name	Country	Study period	Type of surgery	Total number of patients	Incidence of POAF	Assessment method	POAF definition category (duration, if applicable)
Saxena, 2012 ³⁵	Australasian Society of Cardiac and Thoracic Surgeons Cardiac Surgery Database	Australia	2001–2009	CABG	19947	27.8%	Telemetry+ECG	Intervention-based
Horwich, 2013 ³⁶	Queen Elizabeth II Health Sciences Centre	Canada	1995–2009	CABG	8058	27.5%	Telemetry	Intervention-based
O'neal, 2013 ³⁷	East Carolina Heart Institute	United States	1992–2011	CABG	13 165	22.1%	Telemetry	Intervention-based
Saxena, 2013 ³⁸	Australasian Society of Cardiac and Thoracic Surgeons Cardiac Surgery Database	Australia	2001–2009	AVR	2065	35.1%	Telemetry+ECG	Intervention-based
Ivanovic, 2014 ³⁹	Clinical Center of Serbia	Serbia	2006–2009	CABG	460	22.4%	Telemetry+ECG	Nonintervention-based (≥15 min)
Philip, 2014 ⁴⁰	Cleveland Clinic	United States	1993–2005	CABG	5135	29%	Telemetry+ECG	Nonintervention-based (≥2 min)
Pivatto, 2014 ⁴¹	Cardiology Institute/ University Foundation of Cardiology	Brazil	2000–2011	AVR	348	32.8%	Telemetry+ECG	Nonintervention-based (any duration)
Weidinger, 2014 ⁴²	University of Maryland	Austria and United States	2001–2010	CABG	384	15.4%	Telemetry	Intervention-based
Junior, 2015 ⁴³	Santa Isabel Hospital	Brazil	2011–2013	CABG	230	16.1%	ECG	Not reported
Melduni, 2015 ⁴⁵	Mayo Clinic	United States	2000–2005	Multiple	603	37.5%	Telemetry	Nonintervention-based (≥30s)
Tsai, 2015 ⁴⁴	Tri-Service General Hospital, National Defense Medical Center	Taiwan	2009–2012	CABG	266	47.4%	Telemetry+ECG	Intervention-based
Tulla, 2015 ⁴⁵	Kuopio University Hospital	Finland	2000–2010	CABG	276	50%	Telemetry+ECG	Nonintervention-based (≥5 min)
Omer, 2016 ⁴⁶	Michael E. DeBakey VA Medical Center	United States	2006–2013	CABG	1248	17.2%	Telemetry	Not reported
Sahin, 2016 ⁴⁷	Kolan International Hospital	Turkey	2008–2012	CABG	149	36.9%	Telemetry+ECG	Not reported
Ismail, 2017 ⁴⁸	King Faisal Specialist Hospital and Research Center	Saudi Arabia	2013–2015	CABG	252	33.3%	Telemetry	Intervention-based
Lee, 2017 ⁴⁹	Severance Cardiovascular Hospital	Korea	2005–2011	CABG	1664	24.8%	Telemetry+ECG	Nonintervention-based (any duration)
Park, 2017 ⁵⁰	Gachon University Gil Medical Center	Korea	1999–2010	Multiple	938	22.1	Telemetry+ECG	Nonintervention-based (≥1 min)

(Continued)

Table 1. Continued

Study, year	Institution or trial name	Country	Study period	Type of surgery	Total number of patients	Incidence of POAF	Assessment method	POAF definition category (duration, if applicable)
Swinkels, 2017 ¹⁴	St. Antonius Hospital, Academic Medical Center	Netherlands	1990–1994	AVR±CABG	569	42.4%	ECG	Nonintervention-based (>120 min)
Kosmidou, 2018 ⁵¹	Evaluation of XIENCE versus Coronary Artery Bypass Surgery for Effectiveness of Left Main Revascularization trial	International	2010–2014	CABG	893	18%	Telemetry+ECG	Nonintervention-based (≥30 s)
Schwann, 2018 ⁵²	University of Toledo	United States	1994–2012	CABG±carotid reconstruction	8807	22.6%	Telemetry	Intervention-based
Almassi, 2019 ⁵	Randomized On/Off Bypass Trial	United States	2002–2008	CABG	2103	26.2%	Telemetry	Nonintervention-based (≥30 min)
Carier-storch, 2019 ⁵³	Odense University Hospital	Denmark	2014–2016	AVR±CABG	96	53.1%	Telemetry	Nonintervention-based (any duration)
Filardo, 2019 ⁵⁴	Baylor University Medical Center, The Heart Hospital at Baylor Plano, Emory University, Washington University	United States	2002–2010	CABG	9203	31.5%	Telemetry	Nonintervention-based (any duration)
Hernández-leiva, 2019 ⁴	Institute of Cardiology	Colombia	Not reported	Multiple	44	50%	Telemetry	Nonintervention-based (≥30 s)
Kato, 2019 ⁵⁵	Sakakibara Heart Institute, Kishiwada Tokushukai Hospital, Lobe City Medical Center General Hospital, The Cardiovascular Institute, Fukuyama Cardiovascular Hospital, St. Luke's International Hospital, Kitano Hospital, Shizuoka Medical Center, Higashi Takarazuku Satoh Hospital, Sakakibara Heart Institute of Okayama	Japan	2015–2016	Multiple	302	21.9%	Telemetry	Nonintervention-based (≥5 min)
Benedetto, 2020 ³	Arterial Revascularization Trial	International	2004–2007	CABG	3023	24.3%	Telemetry+ECG	Nonintervention-based (≥30 s)
Cole, 2020 ⁵⁶	Liverpool Heart and Chest Hospital	United Kingdom	2013–2018	Multiple	5588	24.8%	Telemetry+ECG	Not reported

(Continued)

Table 1. Continued

Study, year	Institution or trial name	Country	Study period	Type of surgery	Total number of patients	Incidence of POAF	Assessment method	POAF definition category (duration, if applicable)
Fragao, 2020 ⁵⁷	University Of Porto	Portugal	2014–2015	AVR	379	42.2%	Telemetry+ECG	Nonintervention-based (≥30s)
Krishna, 2020 ⁵⁸	Kasturba Medical College	India	2015	CABG	99	20.2%	ECG	Intervention-based
Malhotra, 2020 ⁵⁹	Sanjay Gandhi Postgraduate Institute of Medical Sciences	India	2018–2019	CABG	263	9.1%	ECG	Not reported
Thoren, 2020 ⁶⁰	Uppsala University Hospital	Sweden	1996–2012	CABG	7145	30.6%	Telemetry+ECG	Intervention-based
Fan, 2021 ⁶¹	Peking University People's Hospital	China	2012–2015	CABG	165	15.2%	Telemetry	Nonintervention-based (≥30s)
Gaudino, 2021 ⁷	Weill Cornell Medicine-New York Presbyterian Hospital	United States	2017–2021	Multiple	420	24.5%	Telemetry	Nonintervention-based (≥30s)
Hsu, 2021 ⁶²	National Taiwan University Hospital	Taiwan	2007–2017	Multiple	6267	32.2%	Telemetry+ECG	Nonintervention-based (≥30s)
Lee, 2021 ⁶³	Sejong General Hospital	Korea	2015–2017	CABG	507	18.5%	Telemetry	Not reported
Omar, 2021 ⁶⁴	Cairo University Hospitals	Egypt	2019–2020	CABG	1000	7.8%	ECG	Nonintervention-based (≥30s)
Wang, 2021 ⁶⁵	National University Heart Centre Singapore, National Heart Centre Singapore	Singapore	2008–2012	Multiple	2740	20.9%	Telemetry+ECG	Nonintervention-based (≥60 min)
Zhao, 2021 ⁶⁶	Fuwai Hospital	China	2012–2019	Total arch repair	1271	32.3%	Telemetry	Nonintervention-based (≥5 min)
Musa, 2022 ⁶⁷	Institute Jantung Negera	Malaysia	2019–2021	CABG±valve	242	36.4%	Telemetry	Nonintervention-based (≥30s)
Orali, 2022 ⁶⁸	Tehran Heart Center	Iran	2012–2016	CABG	9310	12.9%	Telemetry+ECG	Nonintervention-based (≥30s)
Potdar, 2022 ⁶⁹	Rabindranath Tagore International Institute of Cardiac Sciences	India	2018	CABG	1108	7.9%	Telemetry+ECG	Intervention-based

AVR indicates aortic valve replacement; CABG, coronary artery bypass grafting; and POAF, postoperative atrial fibrillation.

selection process and the checklist are detailed in [Figure S1](#) and [Table S3](#), respectively. All studies were considered of high or good methodological quality for the purpose of this meta-analysis ([Table S4](#)). Eighteen studies (30.5%) were from Asia, 16 (27.1%) from North America, 15 (25.4%) from Europe, 4 (6.8%) from South America, 3 (5.1%) from multiple regions, 2 (3.4%) from Oceania, and 1 (1.7%) from Africa. Thirty-six studies (61%) included patients undergoing isolated CABG, 20 (33.9%) included more than 1 cardiac surgical procedure, 2 (3.4%) included patients undergoing isolated aortic valve replacement, and 1 (1.7%) included patients undergoing total arch repair ([Table 1](#)).

A total of 197 774 patients were included in the pooled analysis. The number of patients in each study ranged from 44 to 19947 with a median sample size of 969 (interquartile range: 364–5362). The cumulative incidence of POAF was 26% (range: 7.3%–53.1%; [Figure S2](#)). Twenty-one (35.6%) studies reported POAF incidence based on continuous telemetry until hospital discharge, 30 (50.9%) based on telemetry during ICU stay followed by daily ECG in the regular floor, and 8 (13.6%) based on daily ECG only. Fifteen (25.4%) studies had an intervention-based POAF definition, 36 (61%) a nonintervention-based definition, and 8 (13.6%) did not report POAF definition ([Table 1](#)).

The mean age range was 53.7 to 77.4 years in patients with POAF, and 48.0 to 76.5 years in patients without POAF. Female patients ranged from 0.9% to 49.1% in the POAF group and 1.1% to 43.6% in the non-POAF

group. The mean left ventricular ejection fraction range was 43.2% to 65.6% in patients with POAF and 44.4% to 64.0% in patients without POAF. The prevalence of hypertension ranged from 35.1% to 95.3% in patients with POAF and 19.7% to 97.0% in patients without POAF. The prevalence of diabetes ranged from 3.2% to 80.0% in patients with POAF, and 3.4% to 60.4% in patients without POAF. The prevalence of chronic obstructive pulmonary disease ranged from 0% to 41.8% in patients with POAF and 0.8% to 36.7% in patients without POAF. Preoperative use of beta blockers ranged from 26.2% to 91.7% in patients with POAF and 24% to 95% in patients without POAF. Demographic data of the patient population in each study are summarized in [Table S5](#).

Meta-Analysis Primary Outcome

The cumulative incidence of POAF was 26% (range: 7.3%–53.1%). POAF incidence in the group that used continuous telemetry until hospital discharge was 27% (range: 7.3%–53.1%), compared with 27% (range: 7.9%–50%) for the telemetry plus daily ECG group and 19% (range: 7.8%–42.4%) for the group that only used daily ECG. No difference in POAF incidence was found between any of the assessment methods (continuous telemetry versus telemetry plus daily ECG: $P=0.89$; continuous telemetry versus daily ECG only: $P=0.12$; telemetry plus daily ECG versus daily ECG only: $P=0.09$; [Table 2](#); [Figure 1A through D](#) and [2](#); [Table S6](#)).

Table 2. Summary of Primary and Secondary Outcomes Based on Assessment Method

Primary outcome				
Outcome	Comparison group	Pooled estimates (range)	P value	Tau-squared
Postoperative atrial fibrillation incidence	Telemetry vs telemetry+ECG	27% (7.3%–53.1%) vs 27% (7.9%–50%)	0.89	0.30 vs 0.25
	Telemetry vs ECG only	27% (7.3%–53.1%) vs 19% (7.8%–42.4%)	0.12	0.30 vs 0.48
	Telemetry+ECG vs ECG only	27% (7.9%–50%) vs 19% (7.8%–42.4%)	0.09	0.26 vs 0.48
Secondary outcomes				
Outcome	Comparison group	Pooled estimates (95% CI)	P value	
Mortality	Telemetry vs telemetry+ECG	4% (3%–5%) vs 3% (2–4%)	0.29	0.24 vs 0.68
	Telemetry vs ECG only	4% (3%–5%) vs 2% (1–4%)	0.16	0.24 vs 0.76
	Telemetry+ECG vs ECG only	3% (2%–4%) vs 2% (1–4%)	0.47	0.68 vs 0.76
Stroke	Telemetry vs telemetry+ECG	3% (2%–4%) vs 2% (2–3%)	0.30	0.13 vs 0.73
	Telemetry vs ECG only	3% (2%–4%) vs 2% (1–4%)	0.32	0.13 vs 0.17
	Telemetry+ECG vs ECG only	2% (2%–3%) vs 2% (1–4%)	0.88	0.17 vs 0.73
Intensive care unit LOS	Telemetry vs telemetry+ECG	3.7 d (2.1–5.2) vs 3.2 d (2.3–4.1)	0.63	3.56 vs 1.50
	Telemetry vs ECG only	3.7 d (2.1–5.2) vs 3.0 d (1.8–4.3)	0.55	3.56 vs 1.92
	Telemetry+ECG vs ECG only	3.2 d (2.3–4.1) vs 3.0 d (1.8–4.3)	0.82	1.92 vs 1.50
Postoperative LOS	Telemetry vs telemetry+ECG	13.6 d (9.1–18.1) vs 11.0 d (9.4–12.6)	0.29	56.50 vs 12.43
	Telemetry vs ECG only	13.6 d (9.1–18.1) vs 10.1 d (7.7–12.6)	0.18	56.50 vs 8.75
	Telemetry+ECG vs ECG only	11.0 d (9.4–12.6) vs 10.1 d (7.7–12.6)	0.56	12.43 vs 8.75

LOS indicates length of stay.




Figure 1. Comparison of postoperative atrial fibrillation incidence by assessment method. A, Telemetry vs telemetry plus ECG group, $P=0.89$. B, Telemetry vs ECG only, $P=0.12$. C, Telemetry plus ECG vs ECG only, $P=0.09$. D, Grouped funnel plot based on the monitoring approaches with untransformed proportion and standard error (as a measure of precision).

Secondary Outcomes

No differences in in-hospital mortality, stroke, ICU LOS, and postoperative LOS were found between the 3 POAF assessment methods. (Table 2; Figure 2, Tables S7 and S8; Figures S2 through S6).

Secondary Analyses

No difference in POAF incidence was found between intervention- and nonintervention-based POAF definitions (26% [range: 7.9%–47.4%] versus 27% [range: 7.3%–53.1%], respectively; $P=0.67$). No differences in in-hospital mortality, stroke, ICU LOS, and postoperative LOS were found between both definition categories (Table 3; Table S9; Figures S7 through S11).

There was no evidence of publication bias (Egger’s intercept test $P=0.86$; Figure S12).

Subgroup Analysis

In CABG studies significant differences in POAF incidence between continuous telemetry and daily ECG (25% [range: 15.2%–33.3%] versus 15% [range: 7.8%–31%], respectively; $P=0.02$), and between telemetry plus daily ECG compared with daily ECG only (26% [range: 7.9–50%] versus 15% [range: 7.8%–31%], respectively; $P=0.02$) were found (Table 4; Figure S13A through C).

Trend in POAF Incidence

There was no significant change in POAF incidence during the study period (P for trend=0.54; Figure 3).




Figure 2. Summary figure of the main results. ICU indicates intensive care unit; LOS, length of stay; and POAF, postoperative atrial fibrillation.

Meta-Regression

No association between POAF incidence and increasing rigor of the assessment method was found when all cardiac surgeries were considered (beta coefficient 0.17 [95% CI, -0.04 to 0.39, *P*=0.10]); however, an association was found in studies including only patients undergoing isolated CABG where an increase in the rigor of the POAF assessment method was associated with higher POAF detection rates (beta coefficient 0.27 [95% CI, 0.04–0.50], *P*=0.02; Tables S10 and S11).

DISCUSSION

In the present meta-analysis of 59 studies, we found no significant difference in the incidence of POAF after cardiac surgery or its association with adverse outcomes based on the POAF definition or assessment

method used; however, in studies including only patients undergoing isolated CABG, continuous telemetry in the ICU and telemetry in the ICU plus daily ECG in the regular ward were associated with higher POAF incidence compared with daily ECG only.

Prior evidence has suggested that continuous telemetry during the complete postoperative hospitalization is associated with higher POAF incidence compared with other assessment methods⁷⁰ and this finding was seen also in studies in patients who had noncardiac surgery.⁷¹ Our results did not support this finding when all cardiac surgeries were considered; however, this was the case when considering studies that included only patients undergoing isolated CABG. In this subgroup, the use of telemetry at any point postoperatively (either throughout the postoperative stay or only in the ICU) was associated with higher POAF incidence compared with daily ECG use only. The lack of

Table 3. Summary of Secondary Analyses Based on Postoperative Atrial Fibrillation Definition

Outcome	Intervention-based definition (95% CI, unless noted otherwise)	Nonintervention-based definition (95% CI, unless noted otherwise)	<i>P</i> value
Postoperative atrial fibrillation incidence	26% (range 7.9%–47.4%)	27% (range: 7.3%–53.1%)	0.67
Mortality	2% (1%–3%)	2% (1%–2%)	0.69
Stroke	1% (1%–2%)	2% (1%–2%)	0.43
Intensive care unit LOS	2.7 d (1.2–4.2)	2.3 d (1.7–2.8)	0.57
Postoperative LOS	9.7 d (7.1–12.4)	9.5 d (8–10.9)	0.86

LOS indicates length of stay.

Downloaded from http://ahajournals.org by on March 3, 2024

Table 4. Summary of the Subgroup Analyses in Studies That Included Only Patients Undergoing Isolated Coronary Artery Bypass Grafting According to Postoperative Atrial Fibrillation Assessment Method

Outcome	Comparison group	Pooled estimates (range)	P value
Postoperative atrial fibrillation incidence	Telemetry vs telemetry+ECG	25% (15.2%–33.3%) vs 26% (7.9%–50%)	0.81
	Telemetry vs ECG only	25% (15.2%–33.3%) vs 15% (7.8%–31%)	0.02
	Telemetry+ECG vs ECG only	26% (7.9%–50%) vs 15% (7.8–31%)	0.02

difference in POAF incidence between complete stay telemetry and ICU-only telemetry could be explained by the characteristics of POAF, in that over 70% of POAF episodes occur within 72 hours of surgery⁷² and the mean ICU LOS in both groups was approximately 72 hours, suggesting the preponderance of POAF is captured by in-ICU telemetry.

The lack of difference in POAF incidence between intervention- and nonintervention-based POAF definitions could be explained by heterogeneity in the individual study definitions of what is considered treatment of POAF. For example, one study counted POAF episodes only if the arrhythmia required either medical or electrical cardioversion,³³ whereas others limited the description of the definition to *any POAF episode*

requiring treatment^{48,52} without further elaboration. The range of possible interventions in the last setting includes rate control treatment with beta blockers (received by virtually all patients with POAF in the absence of contraindications), to anticoagulation (with variability in treatment recommendations from different professional societies^{11,73,74}) and cardioversion (reserved for patients with hemodynamic instability or resistance/contraindications to medical treatment).

Methodological and Clinical Implications

Our findings have methodological implications for the selection of POAF detection methods in future studies. The main finding is that in patients who had cardiac




Figure 3. Trends in postoperative atrial fibrillation incidence over the study period. P for trend=0.54. POAF indicates postoperative atrial fibrillation.

Downloaded from http://ahajournals.org by on March 3, 2024

surgery POAF incidence and its association with adverse outcomes are not influenced by the assessment method and definition used. However, in studies that included only patients undergoing isolated CABG, the increasing sensitivity of the POAF assessment method was associated with higher POAF detection rates. While the reason for this difference is unclear, it is possible that the less invasive nature of isolated CABG compared with other cardiac surgeries results in less inflammation and shorter POAF episodes (lasting <24 hours),¹ making POAF less likely to be captured by daily ECG. It is also possible that the shorter duration of ICU stay after CABG may have played a role in the reported difference.

From a clinical standpoint, although POAF rates are not affected by the assessment methods, it must be noted that continuous telemetry monitoring outside of the ICU may allow detection of other clinically relevant arrhythmias and is consistent with recommendations from professional societies.⁷⁵

Limitations

This study must be interpreted considering its limitations. Although our systematic review identified the best available evidence comparing outcomes of patients with and without POAF after cardiac surgery, POAF assessment methods were nonrandomized in all studies, creating the possibility for biases and confounding. Additionally, not all studies reported the secondary outcomes of interest, decreasing the power of some comparisons. Moreover, heterogeneity was high for all the outcomes, and it is also possible that studies that used daily ECG only for POAF assessment come from low-resource centers or are using data from an older era.

CONCLUSIONS

POAF incidence remains high after cardiac surgery, although detection rates exhibit variability among different studies. No differences in POAF incidence, in-hospital mortality, stroke, ICU LOS, and postoperative LOS were found between assessment methods and POAF definitions. POAF incidence was higher in studies that included only patients undergoing isolated CABG that used telemetry for POAF assessment (regardless of whether it was during the whole postoperative stay or only in the ICU) compared with daily ECG only.

ARTICLE INFORMATION

Received May 8, 2023; accepted August 22, 2023.

Affiliations

Department of Cardiothoracic Surgery, Weill Cornell Medicine, New York, NY (R.P.-O., T.A., M.R., A.D., L.H., G.J.S., K.R.A., G.C., K.A., M.Y., H.P., R.T., S.S.,

L.N.G., M.G.); Division of Cardiac Surgery, Department of Surgery, University of Toronto, ON, Canada (K.R.A.); Department of Cardiothoracic Surgery, Friedrich Schiller University Jena, Jena, Germany (T.C., H.K.); Samuel J. Wood Library and C.V. Starr Biomedical Information Centre, Weill Cornell Medicine, New York, NY (M.D.); Department of Medical-Surgical Sciences and Biotechnologies, Sapienza University of Rome, Latina, Italy (G.B.-Z.); and Mediterranean Cardiocentro, Naples, Italy (G.B.-Z.).

Sources of Funding

None.

Disclosures

Giuseppe Biondi-Zoccai has consulted for Amarin, Balmed, Cardionovum, Crannmedical, Endocore Lab, Eukon, Guidotti, Innovheart, Meditrial, Microport, Opsens Medical, Terumo, and Translumina, outside the present work. The remaining authors have no disclosures to report.

Supplemental Material

Tables S1–S11

Figures S1–S13

REFERENCES

- Dobrev D, Aguilar M, Heijman J, Guichard J-B, Nattel S. Postoperative atrial fibrillation: mechanisms, manifestations and management. *Nat Rev Cardiol*. 2019;16:417–436. doi: 10.1038/s41569-019-0166-5
- Caldonazo T, Kirov H, Rahouma M, Robinson NB, Demetres M, Gaudino M, Doenst T, Dobrev D, Borger MA, Kiehnopf M, et al. Atrial fibrillation after cardiac surgery: a systematic review and meta-analysis. *J Thorac Cardiovasc Surg*. 2021;1:94–103.e24.
- Benedetto U, Gaudino MF, Dimagli A, Gerry S, Gray A, Lees B, Flather M, Taggart DP, Westaby S, Cook J, et al. Postoperative atrial fibrillation and long-term risk of stroke after isolated coronary artery bypass graft surgery. *Circulation*. 2020;142:1320–1329. doi: 10.1161/CIRCULATIONAHA.120.046940
- Hernández-Leiva E, Alvarado P, Dennis RJ. Postoperative atrial fibrillation: evaluation of its economic impact on the costs of cardiac surgery. *Braz J Cardiovasc Surg*. 2019;34:34. doi: 10.21470/1678-9741-2018-0218
- Almassi GH, Hawkins RB, Bishawi M, Shroyer AL, Hattler B, Quin JA, Collins JF, Bakaeen FG, Ebrahimi R, Grover FL, et al. New-onset postoperative atrial fibrillation impact on 5-year clinical outcomes and costs. *J Thorac Cardiovasc Surg*. 2021;161:1803–1810.e3. doi: 10.1016/j.jtcvs.2019.10.150
- Arsenault KA, Yusuf AM, Crystal E, Healey JS, Morillo CA, Nair GM, Whitlock RP. Interventions for preventing post-operative atrial fibrillation in patients undergoing heart surgery. *Cochrane Database Syst Rev*. 2013;2013:CD003611. doi: 10.1002/14651858.CD003611.pub3
- Gaudino M, Sanna T, Ballman KV, Robinson NB, Hameed I, Audisio K, Rahouma M, Di Franco A, Soletti GJ, Lau C, et al. Posterior left pericardiectomy for the prevention of atrial fibrillation after cardiac surgery: an adaptive, single-centre, single-blind, randomised, controlled trial. *Lancet*. 2021;398:2075–2083. doi: 10.1016/S0140-6736(21)02490-9
- Gaudino M, Di Franco A, Rong LQ, Piccini J, Mack M. Postoperative atrial fibrillation: from mechanisms to treatment. *Eur Heart J*. 2023;44:1020–1039. doi: 10.1093/eurheartj/ehad019
- Gaudino M, Di Franco A, Rong LQ, Cao D, Pivato CA, Soletti GJ, Chadow D, Cancelli G, Perezgrovas Olaria R, Gillinov M, et al. Pericardial effusion provoking atrial fibrillation after cardiac surgery. *J Am Coll Cardiol*. 2022;79:2529–2539. doi: 10.1016/j.jacc.2022.04.029
- Frendl G, Sodickson AC, Chung MK, Waldo AL, Gersh BJ, Tisdale JE, Calkins H, Aranki S, Kaneko T, Cassivi S, et al. 2014 AATS guidelines for the prevention and management of perioperative atrial fibrillation and flutter for thoracic surgical procedures. *J Thorac Cardiovasc Surg*. 2014;148:e153–e193. doi: 10.1016/j.jtcvs.2014.06.036
- Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella M, Diener H-C, Heidbuchel H, Hendriks J, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J*. 2016;37:2893–2962. doi: 10.1093/eurheartj/ehw210
- Villareal RP, Hariharan R, Liu BC, Kar B, Lee V-V, Elyada M, Lopez JA, Rasekh A, Wilson JM, Massumi A. Postoperative atrial fibrillation and mortality after coronary artery bypass surgery. *J Am Coll Cardiol*. 2004;43:742–748. doi: 10.1016/j.jacc.2003.11.023

13. Nisanoglu V, Erdil N, Aldemir M, Ozgur B, Berat Cihan H, Yologlu S, Battaloglu B. Atrial fibrillation after coronary artery bypass grafting in elderly patients: incidence and risk factor analysis. *Thorac Cardiovasc Surg.* 2007;55:32–38. doi: 10.1055/s-2006-924711
14. Swinkels BM, de Mol BA, Kelder JC, Vermeulen FE, ten Berg JM. New-onset postoperative atrial fibrillation after aortic valve replacement: effect on long-term survival. *J Thorac Cardiovasc Surg.* 2017;154:492–498. doi: 10.1016/j.jtcvs.2017.02.052
15. Melduni RM, Schaff HV, Bailey KR, Cha SS, Ammash NM, Seward JB, Gersh BJ. Implications of new-onset atrial fibrillation after cardiac surgery on long-term prognosis: a community-based study. *Am Heart J.* 2015;170:659–668. doi: 10.1016/j.ahj.2015.06.015
16. Aranki SF, Shaw DP, Adams DH, Rizzo RJ, Couper GS, VanderVliet M, Collins JJ, Cohn LH, Burstin HR. Predictors of atrial fibrillation after coronary artery surgery: current trends and impact on hospital resources. *Circulation.* 1996;94:390–397. doi: 10.1161/01.CIR.94.3.390
17. Stamou SC, Dargas G, Hill PC, Pfister AJ, Dullum MKG, Boyce SW, Bafi AS, Garcia JM, Corso PJ. Atrial fibrillation after beating heart surgery. *Am J Cardiol.* 2000;86:64–67. doi: 10.1016/S0002-9149(00)00829-8
18. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;2021:n71. doi: 10.1136/bmj.n71
19. Borenstein M, Higgins JPT. Meta-analysis and subgroups. *Prev Sci.* 2013;14:134–143. doi: 10.1007/s11121-013-0377-7
20. Higgins JPT. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327:557–560. doi: 10.1136/bmj.327.7414.557
21. Creswell LL, Schuessler RB, Rosenbloom M, Cox JL. Hazards of postoperative atrial arrhythmias. *Ann Thorac Surg.* 1993;56:539–549. doi: 10.1016/0003-4975(93)90894-N
22. Tamis JE, Steinberg JS. Atrial fibrillation independently prolongs hospital stay after coronary artery bypass surgery. *Clin Cardiol.* 2000;23:155–159. doi: 10.1002/clc.4960230305
23. Hakala T, Pitkänen O, Hippeläinen M. Feasibility of predicting the risk of atrial fibrillation after coronary artery bypass surgery with logistic regression model. *Scand J Surg.* 2002;91:339–344. doi: 10.1177/145749690209100406
24. Silva RGD, Lima GGD, Laranjeira A, Costa ARD, Pereira E, Rodrigues R. Fatores de risco e morbimortalidade associados à fibrilação atrial no pós-operatório de cirurgia cardíaca. *Arq Bras Cardiol.* 2004;83:99–104. doi: 10.1590/S0066-782X2004001400002
25. Kalavrouziotis D, Buth KJ, Ali JS. The impact of new-onset atrial fibrillation on in-hospital mortality following cardiac surgery. *Chest.* 2007;131:833–839. doi: 10.1378/chest.06-0735
26. Mariscalco G, Engström KG. Atrial fibrillation after cardiac surgery: risk factors and their temporal relationship in prophylactic drug strategy decision. *Int J Cardiol.* 2008;129:354–362. doi: 10.1016/j.ijcard.2007.07.123
27. Mariscalco G, Klersy C, Zanobini M, Banach M, Ferrarese S, Borsani P, Cantore C, Biglioli P, Sala A. Atrial fibrillation after isolated coronary surgery affects late survival. *Circulation.* 2008;118:1612–1618. doi: 10.1161/CIRCULATIONAHA.108.777789
28. Ahlsson A, Fengsrud E, Bodin L, Englund A. Postoperative atrial fibrillation in patients undergoing aortocoronary bypass surgery carries an eightfold risk of future atrial fibrillation and a doubled cardiovascular mortality. *Eur J Cardio-Thorac Surg.* 2010;37:1353–1359. doi: 10.1016/j.ejcts.2009.12.033
29. Brammer S, Van Straten AHM, Soliman Hamad MA, Berreklouw E, Martens EJ, Maessen JG. The impact of new-onset postoperative atrial fibrillation on mortality after coronary artery bypass grafting. *Ann Thorac Surg.* 2010;90:443–449. doi: 10.1016/j.athoracsur.2010.03.083
30. Shirzad M, Karimi A, Tazik M, Aramin H, Ahmadi SH, Davoodi S, Marzban M. Determinants of postoperative atrial fibrillation and associated resource utilization in cardiac surgery. *Rev Esp Cardiol.* 2010;63:1054–1060. doi: 10.1016/S1885-5857(10)70209-7
31. Attaran S, Shaw M, Bond L, Pullan MD, Fabri BM. Atrial fibrillation postcardiac surgery: a common but a morbid complication. *Interact Cardiovasc Thorac Surg.* 2011;12:772–777. doi: 10.1510/icvts.2010.243782
32. Brammer S, Van Straten AHM, Soliman Hamad MA, Van Den Broek KC, Maessen JG, Berreklouw E. New-onset postoperative atrial fibrillation predicts late mortality after mitral valve surgery. *Ann Thorac Surg.* 2011;92:2091–2096. doi: 10.1016/j.athoracsur.2011.06.079
33. Girerd N, Pibarot P, Daleau P, Voisine P, O'Hara G, Després J-P, Mathieu P. Statins reduce short- and long-term mortality associated with postoperative atrial fibrillation after coronary artery bypass grafting: impact of postoperative atrial fibrillation and statin therapy on survival. *Clin Cardiol.* 2012;35:430–436. doi: 10.1002/clc.21008
34. Helgadottir S, Sigurdsson MI, Ingvarsdottir IL, Arnar DO, Gudbjartsson T. Atrial fibrillation following cardiac surgery: risk analysis and long-term survival. *J Cardiothorac Surg.* 2012;7:87. doi: 10.1186/1749-8090-7-87
35. Saxena A, Dinh DT, Smith JA, Shardey GC, Reid CM, Newcomb AE. Usefulness of postoperative atrial fibrillation as an independent predictor for worse early and late outcomes after isolated coronary artery bypass grafting (multicenter Australian study of 19,497 patients). *Am J Cardiol.* 2012;109:219–225. doi: 10.1016/j.amjcard.2011.08.033
36. Horwich P, Buth KJ, Légaré J-F. New onset postoperative atrial fibrillation is associated with a long-term risk for stroke and death following cardiac surgery. *J Cardiac Surg.* 2013;28:8–13. doi: 10.1111/jocs.12033
37. O'Neal WT, Efid JT, Davies SW, O'Neal JB, Anderson CA, Ferguson TB, Chitwood WR, Kypson AP. The impact of postoperative atrial fibrillation and race on long-term survival after coronary artery bypass grafting. *J Cardiac Surg.* 2013;28:484–491. doi: 10.1111/jocs.12178
38. Saxena A, Shi WY, Bappayya S, Dinh DT, Smith JA, Reid CM, Shardey GC, Newcomb AE. Postoperative atrial fibrillation after isolated aortic valve replacement: a cause for concern? *Ann Thorac Surg.* 2013;95:133–140. doi: 10.1016/j.athoracsur.2012.08.077
39. Ivanovic B, Tadic M, Bradic Z, Zivkovic N, Stanisavljevic D, Celic V. The influence of the metabolic syndrome on atrial fibrillation occurrence and outcome after coronary bypass surgery: a 3-year follow-up study. *Thorac Cardiovasc Surg.* 2014;62:561–568. doi: 10.1055/s-0034-1372349
40. Philip F, Becker M, Galla J, Blacksone E, Kapadia SR. Transient post-operative atrial fibrillation predicts short and long term adverse events following CABG. *Cardiovasc Diagn Ther.* 2014;4:365–372. doi: 10.3978/j.issn.2223-3652.2014.09.02
41. Pivatto Júnior F, Teixeira Filho GF, Sant'anna JR, Py PM, Prates PR, Nesralla IA, Kalil RA. Advanced age and incidence of atrial fibrillation in the postoperative period of aortic valve replacement. *Rev Bras Cirurgia Cardiovasc.* 2014;29:45–50. doi: 10.5935/1678-9741.20140010
42. Weidinger F, Schachner T, Bonaros N, Hofauer B, Lehr EJ, Vesely M, Zimrin D, Bonatti J. Predictors and consequences of postoperative atrial fibrillation following robotic totally endoscopic coronary bypass surgery. *Eur J Cardio-Thorac Surg.* 2014;45:318–322. doi: 10.1093/ejcts/ezt282
43. Bohatch Júnior MS, Matkovski PD, Giovanni FJD, Fenili R, Varella EL, Dietrich A. Incidence of postoperative atrial fibrillation in patients undergoing on-pump and off-pump coronary artery bypass grafting. *Rev Bras Cirurgia Cardiovasc.* 2015;30:316–324. doi: 10.5935/1678-9741.20150040
44. Tsai YT, Lai C, Loh SH, Lin CY, Lin YC, Lee CY, Ke HY, Tsai CS. Assessment of the risk factors and outcomes for postoperative atrial fibrillation patients undergoing isolated coronary artery bypass grafting. *Acta Cardiol Sinica.* 2015;31:436–443. doi: 10.6515/acs20150609a
45. Tulla H, Hippeläinen M, Turpeinen A, Pitkänen O, Hartikainen J. New-onset atrial fibrillation at discharge in patients after coronary artery bypass surgery: short- and long-term morbidity and mortality. *Eur J Cardio-Thorac Surg.* 2015;48:747–752. doi: 10.1093/ejcts/ezu526
46. Omer S, Cornwell LD, Bakshi A, Rachlin E, Preventza O, Rosengart TK, Coselli JS, LeMaire SA, Petersen NJ, Pattakos G, et al. Incidence, predictors, and impact of postoperative atrial fibrillation after coronary artery bypass grafting in military veterans. *Tex Heart Inst J.* 2016;43:397–403. doi: 10.14503/THIJ-15-5532
47. Sahin M. Incidence and predisposing factors of atrial fibrillation after coronary artery bypass surgery. *J Clin Anal Med.* 2016;7. doi: 10.4328/JCAM.3839
48. Ismail MF, El-mahrouk AF, Hamouda TH, Radwan H, Haneef A, Jamjoom AA. Factors influencing postoperative atrial fibrillation in patients undergoing on-pump coronary artery bypass grafting, single center experience. *J Cardiothorac Surg.* 2017;12:40. doi: 10.1186/s13019-017-0609-1
49. Lee S-H, Lee H, Park J-K, Uhm J-S, Kim J-Y, Pak H-N, Lee M-H, Yoon H-G, Joung B. Gender difference in the long-term clinical implications of new-onset atrial fibrillation after coronary artery bypass grafting. *Yonsei Med J.* 2017;58:1119–1127. doi: 10.3349/ymj.2017.58.6.1119
50. Park YM, Cha MS, Park C-H, Choi CH, Jeon YB, Kang WC, Choi IS, Park KY. Newly developed post-operative atrial fibrillation is associated with an increased risk of late recurrence of atrial fibrillation in patients who underwent open heart surgery: long-term follow up. *Cardiol J.* 2017;24:633–641. doi: 10.5603/CJ.a2017.0073

51. Kosmidou I, Chen S, Kappetein AP, Serruys PW, Gersh BJ, Puskas JD, Kandzari DE, Taggart DP, Morice M-C, Buszman PE, et al. New-onset atrial fibrillation after PCI or CABG for left main disease. *J Am Coll Cardiol*. 2018;71:739–748. doi: 10.1016/j.jacc.2017.12.012
52. Schwann TA, Al-Shaar L, Engoren MC, Bonnell MR, Goodwin M, Schwann AN, Habib RH. Effect of new-onset atrial fibrillation on cause-specific late mortality after coronary artery bypass grafting surgery. *Eur J Cardio-Thorac Surg*. 2018;54:294–301. doi: 10.1093/ejcts/ezy028
53. Carter-Storch R, Dahl JS, Christensen NL, Pecini R, Søndergård EV, Øvrehus KA, Møller JE. Postoperative atrial fibrillation after aortic valve replacement is a risk factor for long-term atrial fibrillation. *Interact Cardiovasc Thorac Surg*. 2019;29:378–385. doi: 10.1093/icvts/ivz094
54. Filardo G, Ailawadi G, Pollock BD, Da Graca B, Phan TK, Thourani V, Damiano RJ. Postoperative atrial fibrillation: sex-specific characteristics and effect on survival. *J Thorac Cardiovasc Surg*. 2020;159:1419–1425. e1. doi: 10.1016/j.jtcvs.2019.04.097
55. Kato M, Saitoh M, Kawamura T, Iwata K, Sakurada K, Okamura D, Tahara M, Yuguchi S, Kamisaka K, Oura K, et al. Postoperative atrial fibrillation is associated with delayed early rehabilitation after heart valve surgery: a multicenter study. *Phys Ther Res*. 2019;22:1–8. doi: 10.1298/ptr.E9957
56. Cole OM, Tosif S, Shaw M, Lip GYH. Acute kidney injury and post-operative atrial fibrillation in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth*. 2020;34:1783–1790. doi: 10.1053/j.jvca.2019.12.048
57. Fragão-Marques M, Mancio J, Oliveira J, Falcão-Pires I, Leite-Moreira A. Gender differences in predictors and long-term mortality of new-onset postoperative atrial fibrillation following isolated aortic valve replacement surgery. *Ann Thorac Cardiovasc Surg*. 2020;26:342–351. doi: 10.5761/atcs.0a.19-00314
58. Krishna V, Patil N, Nileswhar A. Prospective evaluation of the utility of CHA₂DS₂-VASc score in the prediction of postoperative atrial fibrillation after off-pump coronary artery bypass surgery—an observational study. *Ann Cardiac Anaesth*. 2020;23:122. doi: 10.4103/aca.ACA_161_18
59. Malhotra P, Pande S, Mahindru S, Thukral A, Kotwal A, Gupta R, Tewari P, Agarwal S. Postoperative atrial fibrillation in coronary artery bypass grafting herald poor outcome. *Ann Cardiac Anaesth*. 2021;24:464. doi: 10.4103/aca.ACA_30_20
60. Thorén E, Wernroth M-L, Christersson C, Grinnemo K-H, Jidéus L, Ståhle E. Compared with matched controls, patients with postoperative atrial fibrillation (POAF) have increased long-term AF after CABG, and POAF is further associated with increased ischemic stroke, heart failure and mortality even after adjustment for AF. *Clin Res Cardiol*. 2020;109:1232–1242. doi: 10.1007/s00392-020-01614-z
61. Fan G, Liu J, Dong S, Chen Y. Postoperative atrial fibrillation after minimally invasive direct coronary artery bypass: a single-center, 5-year follow-up study: POAF after MIDCAB. *Heart Surg Forum*. 2021;24:E456–E460. doi: 10.1532/hcf.3621
62. Hsu J-C, Huang C-Y, Chuang S-L, Yu H-Y, Chen Y-S, Wang C-H, Lin L-Y. Long term outcome of postoperative atrial fibrillation after cardiac surgery—a propensity score-matched cohort analysis. *Front Cardiovasc Med*. 2021;8:650147. doi: 10.3389/fcvm.2021.650147
63. Lee H, Kim HJ, Yoo JS, Kim DJ, Yeom SY, Cho KR. Early pharmacologic conversion of atrial fibrillation after off-pump coronary artery bypass grafting. *J Thorac Dis*. 2021;13:4072–4082. doi: 10.21037/jtd-21-466
64. Omar A, Elshihy EM, Singer M, Zarif D, Dawoud O. Perioperative risk factors predisposing to atrial fibrillation after CABG surgery. *Heart Surg Forum*. 2021;24:E402–E406. doi: 10.1532/hsf.3759
65. Wang KKP, Liu W, Chew STH, Ti LK, Shen L. New-onset atrial fibrillation after cardiac surgery is a significant risk factor for long-term stroke: an eight-year prospective cohort study. *J Cardiothorac Vasc Anesth*. 2021;35:3559–3564. doi: 10.1053/j.jvca.2021.07.003
66. Zhao M, Woodward M, Vaartjes I, Millett ERC, Klipstein-Grobusch K, Hyun K, Carcel C, Peters SAE. Sex differences in cardiovascular medication prescription in primary care: a systematic review and meta-analysis. *J Am Heart Assoc*. 2020;9:e014742. doi: 10.1161/JAHA.119.014742
67. Musa AF, Dillon J, Taib MEM, Yunus AM, Sanusi AR, Nordin MN, Smith JA. Incidence and outcomes of postoperative atrial fibrillation after coronary artery bypass grafting of a randomized controlled trial: a blinded end-of-cycle analysis. *Rev Cardiovasc Med*. 2022;23:122. doi: 10.31083/j.rcm2304122
68. Oraili A, Masoudkabar F, Pashang M, Jalali A, Sadeghian S, Mortazavi SH, Ghorbanpour Landy M, Pourhosseini H, Salarifar M, Mansourian S, et al. Effect of postoperative atrial fibrillation on early and mid-term outcomes of coronary artery bypass graft surgery. *Eur J Cardiothorac Surg*. 2022;62:ezac264. doi: 10.1093/ejcts/ezac264
69. Potdar SP, Shales S, Baviskar M, Sharma M, Kapoor L, Narayan P. Incidence, predictors, and outcome for post-operative atrial fibrillation in Indian patients undergoing off-pump coronary artery bypass grafting—a prospective observational study. *Indian J Thorac Cardiovasc Surg*. 2022;38:366–374. doi: 10.1007/s12055-022-01358-7
70. Maesen B, Nijs J, Maessen J, Alessie M, Schotten U. Post-operative atrial fibrillation: a maze of mechanisms. *Europace*. 2012;14:159–174. doi: 10.1093/europace/eur208
71. McIntyre WF, Belley-Côté EP, Vadakken ME, Rai AS, Lengyel AP, Rochweg B, Bhatnagar AK, Deif B, Um KJ, Spence J, et al. High-sensitivity estimate of the incidence of new-onset atrial fibrillation in critically ill patients. *Crit Care Explor*. 2021;3:e0311. doi: 10.1097/CCE.0000000000000311
72. Perezgrovas-Olaria R, Chadow D, Lau C, Rahouma M, Soletti GJ, Cancelli G, Harik L, Dimagli A, Rong LQ, Gillinov M, et al. Characteristics of post-operative atrial fibrillation and the effect of posterior pericardiectomy. *Ann Thorac Surg*. 2023;116:615–622. doi: 10.1016/j.athoracsur.2022.11.007
73. Andrade JG, Aguilar M, Atzema C, Bell A, Cairns JA, Cheung CC, Cox JL, Dorian P, Gladstone DJ, Healey JS, et al. The 2020 Canadian Cardiovascular Society/Canadian Heart Rhythm Society comprehensive guidelines for the management of atrial fibrillation. *Can J Cardiol*. 2020;36:1847–1948. doi: 10.1016/j.cjca.2020.09.001
74. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, et al. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:e199–e267. doi: 10.1161/CIR.0000000000000041
75. Sandau KE, Funk M, Auerbach A, Barsness GW, Blum K, Cvach M, Lampert R, May JL, McDaniel GM, Perez MV, et al. Update to practice standards for electrocardiographic monitoring in hospital settings: a scientific statement from the American Heart Association. *Circulation*. 2017;136:e273–e344.