



Interfaces with other disciplines

## Monotonicity of equilibria in nonatomic congestion games

Roberto Cominetti<sup>a</sup>, Valerio Dose<sup>b,\*</sup>, Marco Scarsini<sup>c</sup><sup>a</sup> Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Diagonal las Torres 2640, Peñalolén, 7910000, Región Metropolitana, Chile<sup>b</sup> Dipartimento di Ingegneria Informatica, Automatica e Gestionale, "Sapienza" Università di Roma, Via Ariosto 25, Roma, 00185, Italy<sup>c</sup> Dipartimento di Economia e Finanza, Luiss University, Viale Romania 32, Roma, 00197, Italy

### ARTICLE INFO

#### Keywords:

Game theory  
Comonotonicity  
Singleton congestion games  
Wardrop equilibrium

### ABSTRACT

This paper studies the monotonicity of equilibrium costs and equilibrium loads in nonatomic congestion games, in response to variations of the demands. The main goal is to identify conditions under which a paradoxical non-monotone behavior can be excluded. In contrast with routing games with a single commodity, where the network topology is the sole determinant factor for monotonicity, for general congestion games with multiple commodities the structure of the strategy sets plays a crucial role.

We frame our study in the general setting of congestion games, with a special focus on singleton congestion games, for which we establish the monotonicity of equilibrium loads with respect to every demand. We then provide conditions for comonotonicity of the equilibrium loads, i.e., we investigate when they jointly increase or decrease after variations of the demands. We finally extend our study from singleton congestion games to the larger class of constrained series–parallel congestion games, whose structure is reminiscent of the concept of a series–parallel network.

### 1. Introduction

Decision making in a multi-agent strategic context is prone to various paradoxes that are impossible in a single-agent framework. For instance, expanding the feasible choice set produces a better outcome in single-agent optimization, but, in a game, it may give rise to an equilibrium that is worse for all players. Analogously, more information is beneficial in single-agent decision making under risk, but may induce worse Bayes-Nash equilibria in a game.

Several paradoxes arise in routing games. These games represent situations where roads to go from one origin to the corresponding destination are chosen strategically by travelers in a way that minimizes their traveling time. The nonatomic version of these games is a good approximation of situations with a large number of travelers. In nonatomic games the standard equilibrium concept, due to Wardrop (1952), prescribes that, for each OD pair, only the paths with the smallest traveling time are used and they all have the same traveling time. A famous paradox in routing games, due to Braess (1968), Braess et al. (2005), shows that adding an edge to a network can make the traveling time worse for all players. Other paradoxes arise in this class of games. For instance, although one could expect that an increase in traffic demand would make the traveling time higher across the network, this is not always the case. In fact, while Hall (1978) proved that – *ceteris paribus* – an increase in the demand of one OD pair increases the traveling time of this OD, Fisk (1979) showed that an

increase of traffic demand of one OD pair can be beneficial for some other OD pair by decreasing its traveling time. Even in networks with a single OD pair, an increment in the traffic demand may decrease the equilibrium load on some edges in the network. These paradoxes will be examined in detail in Examples 3 and 4.

Networks in which the equilibrium loads of all the edges increase with the travel demand of every OD pair are more predictable and easier to handle for a social planner, because an edge is never used below a certain level of demand and is always used above that level. The goal of this paper is precisely to understand when the equilibrium travel times and edge loads are monotone in the demand, so that the paradoxical phenomena observed in the above examples cannot happen. Rather than focusing on routing games, we will state our results for the wider class of congestion games, of which routing games are a significant but particular example.

#### 1.1. Our results

Nonatomic congestion games are defined by a finite set of resources and a finite set of commodities. Each commodity has a demand that can be satisfied by different strategies in a strategy set, where each strategy is a subset of the resource set. In a Wardrop equilibrium each resource has a nonnegative load (a fraction of the total demand), which varies with the demand vector.

\* Corresponding author.

E-mail addresses: [roberto.cominetti@uai.cl](mailto:roberto.cominetti@uai.cl) (R. Cominetti), [valerio.dose@uniroma1.it](mailto:valerio.dose@uniroma1.it) (V. Dose), [marco.scarsini@luiss.it](mailto:marco.scarsini@luiss.it) (M. Scarsini).

The first part of our paper (Section 3) focuses on singleton congestion games, in which every strategy contains only one resource. We start by proving an equilibrium selection result for this class of games: [Theorem 6](#) shows that, even when there exist multiple equilibrium flows, one can always select one equilibrium whose corresponding resource loads are monotone increasing with respect to each demand.

We then use the notion of comonotonicity, which captures the idea that different resource loads jointly increase or decrease upon variations of the demands. [Theorem 12](#) provides some structural results about the demand regions where different subsets of resources are used in equilibrium and how these resources become active or inactive as the demands vary. This analysis allows us to identify regions of the space of demands where the equilibrium loads are comonotonic.

The following section is devoted to games that are more general than singleton congestion games. [Proposition 19](#) shows that every congestion game can be suitably represented as a routing game that is subject to some restrictions, i.e., not every path from an origin to a destination is feasible. Then [Theorem 23](#) extends the monotonicity properties of Section 3 to a class of games that is obtained from singleton congestion games by applying the series and parallel operations. Finally [Theorem 27](#) relates constrained series–parallel games to routing games. These results shed light on the features that produce the non-monotonicity paradoxes, and highlights the difference between the single- and multiple-OD networks: for routing games with a single OD pair, the network topology is the sole relevant factor that guarantees the monotonicity of equilibrium loads, whereas for multiple ODs the structure of the set of feasible routes plays a crucial role.

## 1.2. Related work

Several authors studied the sensitivity of Wardrop equilibria in routing games with respect to changes in the demand. [Hall \(1978\)](#) observed that, when the costs are strictly increasing, the equilibrium loads depend continuously on the demands. [Patriksson \(2004\)](#) and [Josefsson and Patriksson \(2007\)](#) studied the directional differentiability (or lack thereof) of equilibrium costs and loads, whereas [Cominetti et al. \(2023\)](#) studied differentiability along a curve in the space of demands. Specific cases of differentiability, were also considered in [Pradeau \(2014\)](#).

As mentioned previously, [Hall \(1978\)](#) proved that the equilibrium cost of an OD pair increases when the demand of that OD pair grows.

Some positive results concerning the monotonicity of equilibrium loads in series–parallel single-commodity networks can be found in [Klimm and Warode \(2022\)](#) for piece-wise linear costs and in [Cominetti et al. \(2021\)](#) for general nondecreasing costs.

Traffic equilibria in routing games exhibit a multitude of paradoxes. The most famous, due to [Braess \(1968\)](#), shows that removing an edge from a network could actually improve the equilibrium cost for all players (see [Fig. 2](#)). Also surprising is the fact observed by [Fisk \(1979\)](#) that an OD can reduce its cost and benefit from an increase in the demand of a different OD, even after doubling all the demands. [Fisk and Pallottino \(1981\)](#) showed that such paradoxical phenomena could be observed in real life in the City of Winnipeg, Manitoba, Canada. [Dafermos and Nagurney \(1984\)](#) studied how equilibrium costs are affected by changes in the travel demand or addition of new routes under a more general non-separable cost structure. A related paradoxical phenomenon was studied by [Mehr and Horowitz \(2020\)](#) in a model with both regular and autonomous vehicles: despite the fact that autonomous vehicles are more efficient by allowing shorter headways and distances, replacing regular with autonomous vehicles may increase the total network delay.

A particularly simple class of congestion games is the one of singleton congestion games where each strategy comprises a single resource. Different variants of these type of games have been considered in the literature, including atomic weighted and unweighted players, with splittable or unsplittable loads, as well as nonatomic games.

For *atomic splittable* singleton games, [Harks and Timmermans \(2017\)](#) developed a polynomial time algorithm to compute a Nash equilibrium with player-specific affine costs. In a different direction, [Bilò and Vinci \(2017\)](#) investigated how the structure of the players' strategy sets affects the efficiency in singleton load balancing games. Atomic splittable singleton games have also been used to model the charging strategies of a population of electric vehicles ([Deori et al., 2017](#); [Ma et al., 2013](#); [Nimalsiri et al., 2020](#)). In a related but different direction, [Castiglioni et al. \(2019\)](#) studied the computational complexity of finding Stackelberg equilibria in games where one player acts as leader and the others as followers.

For *atomic unsplittable* singleton games, [Gairing and Schoppmann \(2007\)](#) provided upper and lower bounds on the price of anarchy, distinguishing between restricted and unrestricted strategy sets, weighted and unweighted players, and linear vs. polynomial costs. [Fotakis et al. \(2009\)](#) studied the combinatorial structure and computational complexity of Nash equilibria, including the problems of deciding the existence of pure equilibria, computing pure/mixed equilibria, and computing the social cost of a given mixed equilibrium. [Gairing et al. \(2010\)](#) studied weighted atomic unsplittable routing games on a parallel-edge network where each user can only route over a restricted set of edges. They developed a polynomial time algorithm for the model where the edge costs are identical and linear, and both player weights, and edge capacities are integer. [Harks and Klimm \(2012\)](#) characterized the classes of cost functions that guarantee the existence of pure equilibria for weighted routing games and singleton congestion games.

Finally, in the *nonatomic* setting, which is the focus of our paper, [Gonczarowski and Tennenholtz \(2016\)](#) used a clever hydraulic system representation to study asymmetric singleton congestion games, presenting applications in the home internet and cellular markets, as well as in cloud computing. Another recent application of nonatomic singleton congestion games to hospital choice in healthcare systems is discussed in [van de Klundert et al. \(2023\)](#). In the special case of routing games, singleton games correspond to parallel networks. Despite its simple topology they are nevertheless of interest in the literature (see, e.g., [Acemoglu & Ozdaglar, 2007](#); [Harks et al., 2019](#); [Wan, 2016](#)). [Fujishige et al. \(2017\)](#) considered nonatomic congestion games and used matroid theory to characterize games for which two forms of Braess's paradox cannot occur. A similar problem was considered by [Verbree and Cherukuri \(2023\)](#), who – among other things – studied the effect of Braess's paradox at different levels of the demand in single-OD routing games with affine costs.

In Section 4 we use the concept of comonotonicity. Although its definition is purely analytic and concerns real functions defined on an arbitrary space, the idea originated in various applications in actuarial science ([Borch, 1962](#)), economic theory ([Arrow, 1970](#); [Wilson, 1968](#)), and decision theory ([Schmeidler, 1989](#); [Yaari, 1987](#)). A mathematical treatment of the concept – in connection with Choquet capacities – can be found in [Dellacherie \(1971\)](#), who uses the term “même tableau de variation” and [Schmeidler \(1986\)](#), who – to the best of our knowledge – was the first to use the term comonotonic in his preprint [Schmeidler \(1984\)](#) (there exists a previous version, [Schmeidler \(1982\)](#), which we could not access, so we do not know whether the term was used there or not). A recent application of comonotonicity to game theory can be found in [Kocyiğit et al. \(2022\)](#). The reader is referred to [Dhaene et al. \(2002\)](#), [Puccetti and Scarsini \(2010\)](#) for a more thorough discussion and further references.

## 1.3. Organization of the paper

The paper is organized as follows. Section 2 recalls the standard model of non-atomic congestion games and reviews the basic properties of equilibria. This section includes the definition of monotonic equilibrium selection and comonotonicity. Sections 3 and 4 both deal with singleton congestion games. Section 3 contains the central monotonicity result, whereas Section 4 discusses comonotonicity and the structure

of the domains associated to different sets of resources. Section 5 studies the monotonicity properties of more complex congestion games beyond the case of singleton strategies. Section 6 summarizes the results of our paper and proposes some open problems. Appendix A includes some supplementary proofs. Appendix B contains a list of the symbols used throughout the paper.

## 2. Congestion games and equilibria

In this section we recall the basic concepts and properties of nonatomic congestion games, and we fix the notations used throughout the paper. The basic structural elements are:

- a finite set  $\mathcal{R}$  of resources and, for each  $r \in \mathcal{R}$ , a continuous nondecreasing cost function  $c_r : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ , where  $c_r(x_r)$  represents the cost of resource  $r$  under a workload  $x_r$ ; and
- a finite set  $\mathcal{H}$  of commodities and, for each  $h \in \mathcal{H}$ , a family  $S^h \subset 2^{\mathcal{R}} \setminus \emptyset$  of feasible strategies, where every  $s \in S^h$  is a nonempty subset of resources  $s \subset \mathcal{R}$ .

These elements define a congestion game structure  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  with  $c := (c_r)_{r \in \mathcal{R}}$  the vector of cost functions and  $\mathcal{S} := \times_{h \in \mathcal{H}} S^h$  the set of strategy profiles.

Every vector  $\mu := (\mu^h)_{h \in \mathcal{H}}$  of demands  $\mu^h \geq 0$ , determines a nonatomic congestion game  $(\mathcal{G}, \mu)$  as follows. For each commodity  $h \in \mathcal{H}$ , a feasible flow is a vector  $f^h := (f_s^h)_{s \in S^h}$  satisfying

$$\mu^h = \sum_{s \in S^h} f_s^h, \quad f_s^h \geq 0, \text{ for all } s \in S^h. \tag{2.1}$$

A family  $f := (f^h)_{h \in \mathcal{H}}$ , where each  $f^h$  is a feasible flow satisfying (2.1), induces aggregate loads  $x = (x_r)_{r \in \mathcal{R}}$  over the resources, given by

$$x_r := \sum_{h \in \mathcal{H}} \sum_{s \in S^h} f_s^h \mathbb{1}_{\{r \in s\}}, \quad \forall r \in \mathcal{R}, \tag{2.2}$$

which in turn induce strategy costs, defined as

$$c_s(x) := \sum_{r \in s} c_r(x_r), \quad \forall s \subset \mathcal{R}. \tag{2.3}$$

We call  $\mathcal{F}_\mu$  the set of feasible pairs  $(f, x)$  satisfying (2.1) and (2.2). We also write  $\mathcal{X}_\mu$  for the projection of  $\mathcal{F}_\mu$  on the  $x$  variables, that is, the set of load profiles  $x$  induced by all feasible flow vectors  $f$ .

The concept of Wardrop equilibrium is based on the assumption that for each commodity only the strategies with the smallest possible cost are actually used. A feasible pair  $(f, x) \in \mathcal{F}_\mu$  is a Wardrop equilibrium if there exists a nonnegative vector  $\lambda := (\lambda^h)_{h \in \mathcal{H}}$ , such that

$$\forall h \in \mathcal{H}, \quad \begin{cases} c_s(x) = \lambda^h & \text{for all } s \in S^h \text{ with } f_s^h > 0, \\ c_s(x) \geq \lambda^h & \text{for all } s \in S^h \text{ with } f_s^h = 0. \end{cases} \tag{2.4}$$

The quantity  $\lambda^h$  is called the equilibrium cost of commodity  $h \in \mathcal{H}$ . A strategy  $s \in S^h$  is said to be active if  $c_s(x) = \lambda^h$ . Similarly, a resource  $r \in \mathcal{R}$  is active for commodity  $h \in \mathcal{H}$  if it belongs to some active strategy. Clearly, the equilibrium equation implies that every strategy carrying a strictly positive flow  $f_s^h > 0$  is necessarily active. Note, however, that a strategy with zero flow may still be active as long as its cost matches the minimum.

As shown by Beckmann et al. (1956), the set of load profiles induced by equilibrium flows coincides with the set of optimal solutions of the minimization problem

$$\min_{x \in \mathcal{X}_\mu} \sum_{r \in \mathcal{R}} C_r(x_r), \tag{2.5}$$

where  $C_r(x_r) := \int_0^{x_r} c_r(z) dz$ . Since the cost functions  $c_r$  are continuous and nondecreasing, the above objective function is convex and differentiable. Thus, since  $\mathcal{X}_\mu$  is a bounded polytope, for every  $\mu$  there exists at least one optimal solution.

For an equilibrium load profile  $\hat{x}$ , we define the equilibrium resource costs  $\tau_r := c_r(\hat{x}_r)$ . By using Fenchel's duality theory (see e.g.,




Fig. 1. Fisk's network.

Remark 30 in Appendix A, or Fukushima (1984) for the special case of nonatomic routing games), we can prove that the equilibrium resource costs are optimal solutions of the strictly convex dual program

$$\min_{\tau} \sum_{r \in \mathcal{R}} C_r^*(\tau_r) - \sum_{h \in \mathcal{H}} \left( \mu^h \min_{s \in S^h} \sum_{r \in s} \tau_r \right), \tag{2.6}$$

where  $C_r^*(\cdot)$  is the Fenchel conjugate of  $C_r(\cdot)$ , which is strictly convex.

Thus, for each  $\mu$  the equilibrium resource costs  $\tau_r$  are uniquely defined and are the same for all equilibrium loads. This implies that the strategy costs  $c_s = \sum_{r \in s} \tau_r$  and equilibrium costs  $\lambda^h = \min_{s \in S^h} \sum_{r \in s} \tau_r$  depend only on  $\mu$  and not on the particular equilibrium flow under consideration. Thus, also the active strategies and active resources only depend on  $\mu$ .

The active regime at demand  $\mu$  is defined as  $\hat{\mathcal{R}}(\mu) := (\hat{\mathcal{R}}^h(\mu))_{h \in \mathcal{H}}$  with  $\hat{\mathcal{R}}^h(\mu)$  the set of active resources for commodity  $h \in \mathcal{H}$ . We also let  $\mu \mapsto \lambda(\mu)$  denote the equilibrium cost map, whose basic properties are summarized in the next proposition.


**Proposition 1.** Let  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  be a congestion game structure. Then the equilibrium cost map  $\mu \mapsto \lambda(\mu)$  is continuous and monotone in the sense that  $\langle \lambda(\mu_1) - \lambda(\mu_2), \mu_1 - \mu_2 \rangle \geq 0$  for every  $\mu_1, \mu_2 \in \mathbb{R}_+^{\mathcal{H}}$ . In particular, each component  $\lambda^h(\mu)$  is nondecreasing with respect to its own demand  $\mu^h$ . Moreover, the equilibrium resource costs  $\tau_r(\mu)$  are uniquely defined and continuous.

Proposition 1 is a simple extension of Cominetti et al. (2021, Proposition 3.1) to the multi-commodity setting. See also Hall (1978) for the case of strictly increasing costs. For the sake of completeness, we include a proof of Proposition 1 in Appendix A.

**Remark 2.** When the cost functions are strictly increasing, thus invertible, Proposition 1 implies that the equilibrium load vector  $x(\mu)$  is unique for every  $\mu \in \mathbb{R}_+^{\mathcal{H}}$ , and the map  $\mu \mapsto x(\mu)$  is continuous. If the costs are just nondecreasing, the equilibrium loads may be non-unique. Here we point out that there exists some literature about the characterization of games having the so-called uniqueness property (see, e.g., Konishi, 2004; Meunier & Pradeau, 2014; Milchtaich, 2000, 2005). A natural question for the case of multiple equilibria is whether there exists a continuous selection  $\mu \mapsto x(\mu)$ .

Routing games are an important instance of congestion games. In this class of games there is a finite network in the background with a finite set of OD pairs (the commodities of the game); edges are the resources and paths from one origin to the corresponding destination are the strategies of the game.

Hall (1978) proved that in routing games an increase of traffic demand for one OD pair – when the remaining demands are kept fixed – weakly increases the traveling time of this OD pair. The following example, due to Fisk (1979), shows that an increase in the traffic demand of one OD pair may actually reduce the traveling time of another OD pair. Fisk (1979) showed that it is also possible for the social cost  $SC(\mu) = \sum_{h \in \mathcal{H}} \mu^h \lambda^h(\mu)$  to decrease along a direction where the total demand  $\sum_{h \in \mathcal{H}} \mu^h$  increases.



(a) Wheatstone network

	$O v_1 v_2 D$	$O v_1 D$	$O v_2 D$
$\mu \in [0, 1]$	$\mu$	0	0
$\mu \in [1, 2]$	$2 - \mu$	$\mu - 1$	$\mu - 1$
$\mu \in [2, +\infty)$	0	$\mu/2$	$\mu/2$

(b) Equilibrium flows for different values of the demand  $\mu$

Fig. 2. In the Wheatstone network with three paths and a single OD pair, the equilibrium load on the vertical edge  $(v_1, v_2)$  equals the equilibrium flow on the path  $O v_1 v_2 D$  and is decreasing for  $\mu \in [1, 2]$ .

**Example 3.** Consider the network depicted in Fig. 1 with three OD pairs  $(a, b)$ ,  $(b, c)$ ,  $(a, c)$ . Let the initial demands be  $\mu^{(a,b)} = 1$ ,  $\mu^{(a,c)} = 20$ ,  $\mu^{(b,c)} = 100$ , and let the cost functions be as in Fig. 1. The equilibrium loads are  $x_{(a,b)} = 4$ ,  $x_{(a,c)} = 17$ ,  $x_{(b,c)} = 103$ , and the corresponding equilibrium costs are

$$\lambda^{(a,b)} = 4, \quad \lambda^{(a,c)} = 107, \quad \lambda^{(b,c)} = 103.$$

If we now let the demand  $\mu^{(a,b)}$  rise from 1 to 4, the new equilibrium loads are  $x_{(a,b)} = 6$ ,  $x_{(a,c)} = 18$ ,  $x_{(b,c)} = 102$ , and the corresponding equilibrium costs are

$$\lambda^{(a,b)} = 6, \quad \lambda^{(a,c)} = 108, \quad \lambda^{(b,c)} = 102.$$

That is, the increase of  $\mu^{(a,b)}$  increases the cost of the edge  $ab$  and pushes the  $(a, c)$  pair to favor the use of the direct edge  $ac$ . This reduces the load on the edge  $bc$ , which ultimately benefits the pair  $(b, c)$  by reducing its cost.

Perhaps more surprising is the fact that this phenomenon may even occur when all the demands increase by the same factor: with demands  $\mu^{(a,b)} = 60$ ,  $\mu^{(a,c)} = 30$ ,  $\mu^{(b,c)} = 6$  the equilibrium cost for the third OD is  $\lambda^{(b,c)} = 24$ , and when all the demands are doubled it decreases to  $\lambda^{(b,c)} = 18$ .

**Example 4.** Even in networks with a single OD pair, where the equilibrium cost increases with the demand, it may happen that the load on some edges decrease after a surge in the demand. This can be observed in the classical Wheatstone network depicted in Fig. 2 (see Braess, 1968; Braess et al., 2005, for the famous paradox that uses this network).

In what follows, we want to determine if a congestion game has an equilibrium selection such that the resource loads are monotone with respect to an increase in any demand. This is made precise in the following definition.

**Definition 5.** A congestion game structure  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  is said to have a *monotonic equilibrium selection* (MES) if there exists an equilibrium load vector  $x(\mu)$  such that for every resource  $r \in \mathcal{R}$  the map  $\mu \mapsto x_r(\mu)$  is nondecreasing with respect to each component  $\mu^h$  of the demand vector  $\mu$ .

In mixed scenarios where some demands increase and other decrease, one may naturally expect that the same holds for the induced equilibrium loads. However, it is still of interest to identify groups of resources whose equilibrium loads vary in the same direction, regardless whether  $\mu$  and  $\mu'$  are comparable or not. In such a case, observing an increase/decrease in the load of a specific resource one can infer that all the remaining loads in the group move in the same direction. This property is captured by the notion of comonotonicity: a family of functions  $\{\psi_i : \Omega \rightarrow \mathbb{R}\}_{i \in A}$  is *comonotonic* if for all  $i, j \in A$  we have

$$\forall \omega_1, \omega_2 \in \Omega, \quad (\psi_i(\omega_1) - \psi_i(\omega_2))(\psi_j(\omega_1) - \psi_j(\omega_2)) \geq 0. \quad (2.7)$$

For singleton congestion games, we will identify subsets of resources whose equilibrium loads exhibit such comonotonic behavior in specific regions of the space of demands  $\mathbb{R}_+^H$ . Informally, we will show that a group of commodities that share the same equilibrium cost behave as a single commodity, and the loads on the resources used by this group are comonotonic.

### 3. Monotonicity in singleton congestion games

In a *singleton congestion game* each strategy corresponds to a single resource. Thus, for every commodity  $h \in \mathcal{H}$  the set of feasible strategies  $\mathcal{S}^h$  can be viewed as a subset  $\mathcal{R}^h \subset \mathcal{R}$  of the set of resources. The following result shows that the MES property holds in this case.

**Theorem 6.** Every singleton congestion game structure  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  has a MES.

**Proof.** We first prove the result for strictly increasing cost functions, and we then use a regularization argument to address the general case of nondecreasing costs.

Suppose first that the costs  $c_r(\cdot)$  are strictly increasing. We will prove the existence of a MES locally by showing that for every demand vector  $\mu_0 \in \mathbb{R}_+^H$  and every commodity  $h \in \mathcal{H}$ , there exists  $\varepsilon > 0$  such that  $x_r(\mu_0 + t e^h) \geq x_r(\mu_0)$  for all  $t \in [0, \varepsilon]$ , where  $e^h$  is the  $h$ -th vector of the canonical basis of  $\mathbb{R}^H$ . The global MES property throughout the space of demands then follows from the continuity of the map  $\mu \mapsto x(\mu)$  (see Remark 2).

Let  $\mathcal{R}_0$  be the set of resources such that  $c_r(x_r(\mu_0)) = \lambda^h(\mu_0)$ . This set contains the active resources for commodity  $h$  but may also include resources used by other commodities and that are not feasible for  $h$ . By continuity of the equilibrium costs (Proposition 1), there exists  $\varepsilon > 0$  such that an increase in the demand for commodity  $h$  by an amount  $t$  smaller than  $\varepsilon$  can only affect the equilibrium loads of resources in  $\mathcal{R}_0$ , and therefore for  $r \notin \mathcal{R}_0$  and  $t \in [0, \varepsilon]$  we have  $x_r(\mu_0 + t e^h) = x_r(\mu_0)$ . Let us then focus on the resources  $r \in \mathcal{R}_0$ . Fix an arbitrary  $t \in [0, \varepsilon]$  and partition  $\mathcal{R}_0$  into the three subsets

$$\mathcal{R}_0^+ := \{r \in \mathcal{R}_0 : x_r(\mu_0 + t e^h) > x_r(\mu_0)\}, \quad (3.1)$$

$$\mathcal{R}_0^- := \{r \in \mathcal{R}_0 : x_r(\mu_0 + t e^h) < x_r(\mu_0)\}, \quad (3.2)$$

$$\mathcal{R}_0^= := \{r \in \mathcal{R}_0 : x_r(\mu_0 + t e^h) = x_r(\mu_0)\}. \quad (3.3)$$

Suppose by contradiction that  $\mathcal{R}_0^-$  is not empty. Since the total demand at  $\mu_0 + t e^h$  is strictly larger than the total demand at  $\mu_0$ , whereas the total flow on the resources  $\mathcal{R}_0^- \cup \mathcal{R}_0^=$  decreases, some flow must have been transferred from  $\mathcal{R}_0^- \cup \mathcal{R}_0^=$  to  $\mathcal{R}_0^+$ . This implies the existence of a commodity  $h'$  which has feasible resources in both  $\mathcal{R}_0^- \cup \mathcal{R}_0^=$  and  $\mathcal{R}_0^+$ , and which sends a positive flow along a resource in  $\mathcal{R}_0^+$  at demand  $\mu_0 + t e^h$ . This contradicts the equilibrium condition for that commodity because

the cost of all resources in  $\mathcal{R}_0^+$  is strictly higher than the cost of the resources in  $\mathcal{R}_0^- \cup \mathcal{R}_0^0$ . This establishes the existence of a MES for the case of strictly increasing costs.

When costs  $c_r(x_r)$  are assumed to be just nondecreasing, we perturb them as  $c_r^\epsilon(x_r) := c_r(x_r) + 2\epsilon x_r$  with  $\epsilon > 0$ , to make them strictly increasing, and then consider the limit as  $\epsilon$  approaches zero. As recalled in Section 2, the equilibrium flow  $x(\mu, \epsilon)$  for the congestion game structure  $\mathcal{G}^\epsilon := (\mathcal{R}, c^\epsilon, \mathcal{S})$  is the unique solution of the Beckmann problem (2.5), which in this case has the form

$$\min_{x \in \mathcal{X}_\mu} \sum_{r \in \mathcal{R}} C_r(x_r) + \epsilon \|x\|^2, \tag{3.4}$$

with  $C_r(x_r) := \int_0^{x_r} c_r(z) dz$ . Tikhonov regularization (see, e.g., Attouch, 1996, section 1.1) tells us that  $x(\mu, \epsilon)$  converges, as  $\epsilon$  approaches zero, to the minimal norm equilibrium  $x_0(\mu)$  of the original unperturbed game  $\mathcal{G}$ . From the previous case of strictly increasing costs, for each  $\epsilon > 0$  the map  $\mu \mapsto x(\mu, \epsilon)$  is nondecreasing with respect to each demand  $\mu^h$ , and this property is inherited by  $\mu \mapsto x_0(\mu)$  in the limit as  $\epsilon \downarrow 0$ , providing a MES as claimed.  $\square$

**Remark 7.** The quadratic regularizer  $\epsilon \|x\|^2$  was introduced by Tikhonov in the study of ill-posed inverse problems (Tikhonov, 1943, 1963; Tikhonov & Arsenin, 1977). It is also the basis of ridge regression in statistics (Hoerl, 1959, 1962; Hoerl & Kennard, 1970). In our setting this is just one choice among others, and can be replaced by a separable regularizer  $\epsilon \sum_{i=1}^n g_i(x_i)$  with  $g_i(\cdot)$  strictly increasing. Every such regularizer selects a specific optimal solution in the limit when  $\epsilon \downarrow 0$  (see Attouch (1996, theorem 2.1) and Auslender et al. (1997, proposition 2.5). Moreover, one can verify that the previous proof is still valid and yields a monotone selection of the set of Wardrop equilibria. In particular,  $\epsilon \sum_{i=1}^n x_i \log(x_i)$  selects the Wardrop equilibrium of maximal entropy. A similar entropic regularization was used in Rossi et al. (1989) to select one among multiple flow decompositions of a Wardrop equilibrium (see Borchers et al., 2015, for a survey of related work). In our case we deal with multiple equilibria and the regularization is used to obtain a selection with monotonicity properties. As alternatives one may consider general penalty schemes of the form  $\epsilon \sum_{i=1}^n \theta(x_i/\epsilon)$ , including the classical log-barrier  $\theta(x) = -\log(x)$ , the inverse-barrier  $\theta(x) = 1/x$ , the exponential penalty  $\theta(x) = \exp(-x)$ , and more (see Cominetti, 1999). Let us also mention the multi-scale regularizer  $\sum_{i=1}^n \epsilon^i x_i^2$ , which yields a hierarchical selection principle: select the Wardrop equilibria that have the smallest first coordinate  $x_1^2$ , among these the ones with smallest  $x_2^2$ , and inductively with  $x_3^2, \dots, x_n^2$ .

**Remark 8.** Theorem 6 is related to results in Fujishige et al. (2017), which investigates Braess’s paradox in the context of nonatomic matroid congestion games, where the strategy set for each commodity  $h$  is the set  $\mathcal{B}^h$  of bases of some matroid  $M^h = (\mathcal{R}, \mathcal{I}^h)$ , defined over a common ground set  $\mathcal{R}$  of resources. Among other results, lemma 3.2 in that paper establishes the monotonicity of the resource costs at equilibrium, from which one can readily deduce the monotonicity of the loads when the cost functions are strictly increasing.

**4. Comonotonicity and active regimes in singleton congestion games**

Theorem 6 shows that the equilibrium loads in singleton congestion games respond monotonically when all the demands increase or stay the same. In mixed cases where some demands increase and others decrease, one can still identify groups of resources that behave comonotonically in specific regions of the space of demands. A trivial example is when all commodities can use every resource  $\mathcal{R}^h \equiv \mathcal{R}$ , so they can be treated as a single commodity and the equilibrium loads are just nondecreasing functions of the total demand  $\mu_H = \sum_{h \in \mathcal{H}} \mu^h$ . More generally, we will show that a subset  $C \subset \mathcal{H}$  of commodities that have the same equilibrium cost, behave as if they were a single-commodity

on a smaller congestion game restricted to a subset  $\mathcal{R}_C$  of resources, and the equilibrium loads of these resources are nondecreasing functions of the aggregate demand  $\mu_C$  of the group, so that they are comonotonic.

To state our result precisely, given a singleton congestion game structure  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$ , we partition the space of demands  $\mathbb{R}_+^{\mathcal{H}}$  into different regions  $\Gamma^\preceq$  characterized by the order in which the commodities are ranked by equilibrium cost. In order to understand the geometry of such regions, we further decompose them into sub-regions corresponding to different active regimes.

**Definition 9.** For any fixed weak order  $\preceq$  on  $\mathcal{H}$  we call  $\Gamma^\preceq$  the set of demands that rank the commodities exactly in this order, that is,

$$\Gamma^\preceq = \left\{ \mu \in \mathbb{R}_+^{\mathcal{H}} : \lambda^h(\mu) \leq \lambda^{h'}(\mu) \iff h \preceq h' \text{ for all } h, h' \in \mathcal{H} \right\}, \tag{4.1}$$

and we call  $\Gamma_\rho^\preceq$  the sub-region with active regime  $\rho := (\rho^h)_{h \in \mathcal{H}}$  with  $\rho^h \subset \mathcal{R}^h$ , that is,

$$\Gamma_\rho^\preceq := \left\{ \mu \in \Gamma^\preceq : \widehat{\mathcal{R}}(\mu) = \rho \right\}. \tag{4.2}$$

We recall that the equivalence relation and strict order associated with  $\preceq$  are defined by

$(h' \sim h)$  if and only if  $(h \preceq h')$  and  $(h' \preceq h)$ ,

$(h' > h)$  if and only if  $(h \preceq h')$  and  $\neg(h' \preceq h)$ .


The relation  $\sim$  partitions  $\mathcal{H}$  into equivalence classes, called cost classes: two commodities are in the same cost class if and only if  $h \sim h'$ , that is to say, if and only if  $\lambda^h(\mu) = \lambda^{h'}(\mu)$  for all  $\mu \in \Gamma^\preceq$ . To each cost class  $C$  we associate the subset  $\mathcal{R}_C$  of all the resources  $r \in \mathcal{R}$  that are feasible for some commodity  $h \in C$ , excluding those which are also feasible for higher ranked commodities  $h' > h$ , that is

$$\mathcal{R}_C = \left( \bigcup_{h \in C} \mathcal{R}^h \right) \setminus \left( \bigcup_{h' > C} \mathcal{R}^{h'} \right). \tag{4.3}$$


**Definition 10.** Let  $C$  be a cost class for a weak order  $\preceq$  on  $\mathcal{H}$ . We let  $\mathcal{G}_C := (\mathcal{R}_C, c, \mathcal{S}_C)$  denote the singleton congestion game structure with a single commodity whose strategy set  $\mathcal{S}_C$  comprises all the singletons in  $\mathcal{R}_C$ .

The regions  $\Gamma^\preceq$  can be empty for some orders  $\preceq$  (e.g., if  $h, h' \in \mathcal{H}$  are such that  $\mathcal{R}^h \subseteq \mathcal{R}^{h'}$  we cannot have  $\lambda^h(\mu) < \lambda^{h'}(\mu)$ ). We stress that each commodity  $h \in \mathcal{H}$  belongs to a unique cost class  $C$ , whereas each resource  $r$  belongs to the cost class of the highest ranked commodity among those for which  $r$  is feasible.

**Example 11.** Consider a singleton congestion game structure with three resources  $\mathcal{R} = \{r_1, r_2, r_3\}$  with affine costs  $c_1(x) = x + 1$ ,  $c_2(x) = x$ ,  $c_3(x) = x + 2$ , and two commodities  $\alpha$  and  $\beta$  with  $\mathcal{R}^\alpha = \{r_1, r_2\}$  and  $\mathcal{R}^\beta = \{r_2, r_3\}$ . For visualization, Fig. 3(a) represents this as a routing game on a parallel network, where both commodities have to move traffic between the two vertices, but each of them is allowed to use only certain edges. In Fig. 3(b) the horizontal axis represents the demand of commodity  $\alpha$  and the vertical axis the demand of  $\beta$ . The three colors represent the regions  $\Gamma^\preceq$  corresponding to the possible orders of  $\lambda^\alpha$  and  $\lambda^\beta$ . In the top-left region in green  $\lambda^\alpha < \lambda^\beta$ , in the bottom-right region in orange  $\lambda^\alpha > \lambda^\beta$ , whereas in the middle region in purple  $\lambda^\alpha = \lambda^\beta$ . Hence, in the top-left and bottom-right regions we have two cost classes  $C_1 = \{\alpha\}$  and  $C_2 = \{\beta\}$ , each containing one commodity. However, in the top-left region the corresponding resource sets are  $\mathcal{R}_{C_1} = \{r_1\}$  and  $\mathcal{R}_{C_2} = \{r_2, r_3\}$ , whereas in the bottom-right region  $\mathcal{R}_{C_1} = \{r_1, r_2\}$  and  $\mathcal{R}_{C_2} = \{r_3\}$ . The region in purple has a single cost class  $C = \{\alpha, \beta\}$  with  $\mathcal{R}_C = \{r_1, r_2, r_3\}$ . The sub-regions delimited by horizontal and diagonal lines within a colored region, correspond to different active regimes. In the purple region characterized by  $\lambda^\alpha = \lambda^\beta$ , with a single cost class  $C = \{\alpha, \beta\}$  and  $\mathcal{R}_C = \{r_1, r_2, r_3\}$ , there are three sub-regions depending on the value of the total demand  $\mu_C = \mu^\alpha + \mu^\beta$ . When  $\mu_C \in (0, 1)$  both  $\alpha$  and  $\beta$  use only the central edge with active



(a) A routing game with one commodity that uses the two top edges, and a second commodity that uses the bottom two.



(b) The colors represent the regions  $\Gamma^{\succsim}$  for the three possible orders  $\succsim$  of the equilibrium costs. These regions are further decomposed into polyhedral sub-regions  $\Gamma_{\rho}^{\succsim}$  that correspond to different active regimes.

Fig. 3. A singleton congestion game with affine costs.

regime  $\rho^{\alpha} = \rho^{\beta} = \{r_2\}$  and equilibrium costs  $\lambda^{\alpha} = \lambda^{\beta} = \mu_C$ . For  $\mu_C \in [1, 3)$  we have  $\lambda^{\alpha} = \lambda^{\beta} = (1 + \mu_C)/2$ , with  $\beta$  using the central edge  $\rho^{\beta} = \{r_2\}$ , whereas  $\alpha$  splits the flow between the top and central edge with  $\rho^{\alpha} = \{r_1, r_2\}$ . Finally for  $\mu_C \geq 3$  the active regime is  $\rho^{\alpha} = \{r_1, r_2\}$  and  $\rho^{\beta} = \{r_2, r_3\}$  with equilibrium costs  $\lambda^{\alpha} = \lambda^{\beta} = 1 + \mu_C/3$ . Similarly, the green region is characterized by  $\lambda^{\alpha} < \lambda^{\beta}$  with cost classes  $C_1 = \{\alpha\}$  and  $C_2 = \{\beta\}$ . Throughout this green region the active regime for  $\alpha$  is constant  $\rho^{\alpha} = \{r_1\}$ , whereas  $\rho^{\beta} = \{r_2\}$  if  $\mu^{\beta} < 2$  and  $\rho^{\beta} = \{r_2, r_3\}$  if  $\mu^{\beta} \geq 2$ .

Our next result describes the equilibrium within a cost class  $C$ : we show that the loads on the resources in  $\mathcal{R}_C$  coincide with those of the single-commodity game  $\mathcal{G}_C$ . In other words, in terms of equilibrium loads the commodities in  $C$  behave as if they were a single commodity. This allows in turn to analyze the regions on which the equilibrium loads on  $\mathcal{R}_C$  are comonotone. Moreover, part (c) further analyzes the geometry of the regions of comonotonicity, as observed in Example 11. The simple structure exhibited by the sub-regions in Example 11 holds more generally: even if the cost functions are nonlinear, the sub-regions are separated by hyperplanes defined by the aggregate demand of some cost class. We recall that a *break point* in a single commodity game is a demand  $\bar{\mu}$  at which the set of active resources changes, i.e., this set is not constant on any interval  $(\bar{\mu} - \epsilon, \bar{\mu} + \epsilon)$  with  $\epsilon > 0$  (see Cominetti et al., 2021, definition 3.4).

**Theorem 12.** Let  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  be a singleton congestion game structure, and  $\Gamma^{\succsim}$  the region associated with a weak order  $\succsim$  on  $\mathcal{H}$ . Then, for each cost class  $C$  for  $\succsim$  we have:

- (a) For all  $\mu \in \Gamma^{\succsim}$  and every equilibrium load  $x$  of  $(\mathcal{G}, \mu)$ , the vector  $\bar{x} = (x_r)_{r \in \mathcal{R}_C}$  is an equilibrium in the single-commodity game  $(\mathcal{G}_C, \mu_C)$  with aggregate demand  $\mu_C := \sum_{h \in C} \mu^h$ .
- (b) If  $\mathcal{G}_C$  has a unique equilibrium for each demand in  $\mathbb{R}_+$ , then for  $\mu \in \Gamma^{\succsim}$  the equilibrium loads  $x_r(\mu)$  with  $r \in \mathcal{R}_C$  can be expressed as nondecreasing functions of the aggregate demand  $\mu_C$ , which is equivalent to the fact that the equilibrium loads of the resources in  $\mathcal{R}_C$  are comonotonic in the region  $\Gamma^{\succsim}$ .
- (c) If the costs are strictly increasing, then the boundary between the sub-regions  $\Gamma_{\rho}^{\succsim}$  coincides with the points  $\mu \in \Gamma^{\succsim}$  satisfying at least

one of the linear equations

$$\sum_{h \in C} \mu^h = \bar{\mu},$$

where  $\bar{\mu}$  is a break point in the single-commodity game  $\mathcal{G}_C$ .

**Proof.** (a) Let  $x$  be an equilibrium load vector of demand  $\mu \in \Gamma^{\succsim}$ . We note that every commodity  $h \in C$  allocates traffic only through resources in  $\mathcal{R}_C$ . Indeed, if a commodity  $h \in C$  has a feasible resource also in  $\mathcal{R}_{C'}$  with  $C' \neq C$ , then, because of (4.3), we have  $h' > h$  for every  $h' \in C'$ , which is equivalent to  $\lambda^{h'}(\mu) > \lambda^h(\mu)$ , because  $\mu \in \Gamma^{\succsim}$ . For this reason, all the commodities  $h \in C$  have the same equilibrium cost  $\lambda^h(\mu) =: \lambda_C(\mu)$ , which implies that for every  $r, r' \in \mathcal{R}_C$  we have

$$x_r > 0 \implies c_r(x_r) = \lambda_C(\mu) \leq c_r'(x_r').$$

Since  $\sum_{r \in \mathcal{R}_C} x_r = \sum_{h \in C} \mu^h = \mu_C$ , the vector  $\bar{x} = (x_r)_{r \in \mathcal{R}_C}$  is a single-commodity equilibrium for  $\mathcal{G}_C$  with demand  $\mu_C$ . It follows that  $\lambda_C(\mu)$  is in fact a function of the aggregate demand  $\mu_C$  and so we can write it as  $\lambda_C(\mu_C)$ .

(b) By the result in (a), for each  $r \in \mathcal{R}_C$  and  $\mu \in \Gamma^{\succsim}$  the equilibrium load  $x_r(\mu)$  coincides with the unique equilibrium in the single-commodity game  $\mathcal{G}_C$  with demand  $\mu_C$ , and therefore it is a function of the aggregate demand  $\mu_C$ . Now, according to Cominetti et al. (2021, proposition 3.12) every single-commodity game on a SP network has a nondecreasing selection of equilibria, so that  $x_r(\mu)$  is a nondecreasing function of  $\mu_C$ . The equivalence with the comonotonicity of the maps  $\mu \mapsto x_r(\mu)$  for  $r \in \mathcal{R}_C$  throughout the region  $\mu \in \Gamma^{\succsim}$ , then follows from a known result (see e.g., Dellacherie (1971) and Landsberger and Meilijson (1994)). Since we could not find a proof of this latter result in the literature, we include one in Lemma 31 in Appendix A.

(c) Consider any demand  $\mu \in \Gamma^{\succsim}$ . By (a), the equilibrium loads can be partitioned by cost classes  $(x_r(\mu))_{r \in \mathcal{R}_C}$ , the latter being an equilibrium in the single-commodity game  $\mathcal{G}_C$ . The equilibrium cost for  $\mathcal{G}_C$  is a strictly increasing function  $\mu_C \mapsto \lambda_C(\mu_C)$  of the aggregate demand  $\mu_C = \sum_{h \in C} \mu^h$ . It then follows that each load  $x_r(\mu) = c_r^{-1}(\lambda_C(\mu_C))$  for  $r \in \mathcal{R}_C$  is also a strictly increasing function of  $\mu_C$ .

If  $\mu \in \Gamma^{\succsim}$  is on the boundary between two or more sub-regions  $\Gamma_{\rho}^{\succsim}$ , the set of active resources changes locally at  $\mu$  and then there must




Fig. 4. An example with quadratic costs. The first commodity uses the two top edges, and the second commodity uses the bottom two. The three colors represent the regions  $\Gamma^{\preceq}$  for the possible orders  $\preceq$  of the equilibrium costs. The straight lines within each region separate sub-regions corresponding to different active regimes. The regions  $\Gamma^{\preceq}$  are not convex, but the boundary between sub-regions is still affine.

exist a cost class  $C$  whose set of active resources also changes locally at  $\mu_C$ , which is therefore a break point in the single-commodity game  $\mathcal{G}_C$ .  $\square$

**Example 13.** Consider the variant of Example 11 with quadratic costs as in Fig. 4. The regions  $\Gamma^{\preceq}$  are no longer convex, but the sub-regions for different active regimes are still delimited by the hyperplanes described in Theorem 12(c). In the purple region where  $\lambda^\alpha = \lambda^\beta$  with a single cost class  $C = \{\alpha, \beta\}$  and  $\mathcal{R}_C = \{r_1, r_2, r_3\}$ , the equilibrium loads of all the resources are strictly increasing functions of the total demand  $\mu_C = \mu^\alpha + \mu^\beta$ , and the active regimes present break points at  $\mu_C = 1$  and  $\mu_C = 1 + \sqrt{2}$ , that is,

$$\begin{cases} \rho^\alpha = \{r_2\} \text{ and } \rho^\beta = \{r_2\} & \text{if } \mu_C \in [0, 1), \\ \rho^\alpha = \{r_1, r_2\} \text{ and } \rho^\beta = \{r_2\} & \text{if } \mu_C \in [1, 1 + \sqrt{2}), \\ \rho^\alpha = \{r_1, r_2\} \text{ and } \rho^\beta = \{r_2, r_3\} & \text{if } \mu_C \in [1 + \sqrt{2}, \infty). \end{cases} \quad (4.4)$$

Similarly, in the green region where  $\lambda^\alpha < \lambda^\beta$  with cost classes  $C_1 = \{\alpha\}$  and  $C_2 = \{\beta\}$ , we have  $\rho^\beta = \{r_2\}$  for  $\mu^\beta < \sqrt{2}$  and  $\rho^\beta = \{r_2, r_3\}$  for  $\mu^\beta \geq \sqrt{2}$ , whereas  $\rho^\alpha = \{r_1\}$  is constant.

**Remark 14.** By Remark 2, having strictly increasing costs ensures the uniqueness of equilibria for  $\mathcal{G}_C$ , as required in Theorem 12(b). Actually, it suffices that no two resources in  $\mathcal{R}_C$  have cost functions that are constant and equal on some (possibly different) non-degenerate intervals. Moreover, for strictly increasing costs the equilibrium loads  $x_r(\mu)$  for  $r \in \mathcal{R}_C$  and  $\mu \in \Gamma^{\preceq}$  are strictly increasing with  $\mu_C$ . Indeed, since  $\sum_{r \in \mathcal{R}_C} x_r(\mu) = \mu_C$ , a strict increase of  $\mu_C$  implies that some load  $x_r(\mu)$  and its corresponding cost  $c_r(x_r(\mu))$  must strictly increase. However, across  $\Gamma^{\preceq}$  the equilibrium costs of all the resources  $r \in \mathcal{R}_C$  remain equal, so that all their loads  $x_r(\mu)$  must strictly increase simultaneously.

**Remark 15.** Theorem 12(b) implies that comonotonicity fails across different cost classes  $C \neq C'$ : if  $\mu_C$  increases and  $\mu_{C'}$  decreases, the equilibrium loads of the resources  $\mathcal{R}_C$  and  $\mathcal{R}_{C'}$  will move in opposite directions. On the contrary, if both aggregate demands move in the same direction, the same holds for the corresponding equilibrium loads.

**Remark 16.** The comonotonicity in Theorem 12(b) may fail when  $\mathcal{G}_C$  has multiple equilibria. Consider for instance a variant of Example 11 with costs  $c_1(x) = c_3(x) = 1$  and  $c_2(x) = x$ . When the demand is  $\mu = (2, 0)$  the equilibrium sends 1 unit of flow through  $r_1$  and  $r_2$ , and

zero on  $r_3$ . Instead, at demand  $\nu = (0, 2)$  nothing is sent through  $r_1$ , with 1 unit of traffic on both  $r_2$  and  $r_3$ . Hence, despite the fact that at both  $\mu$  and  $\nu$  all three resources have the same equilibrium cost equal to 1, the load on resource  $r_1$  decreases when moving from  $\mu$  to  $\nu$ , whereas the load on resource  $r_3$  increases, so these loads are not comonotonic. Theorem 12(b) does not apply here because the single-commodity game  $\mathcal{G}_C$  on the three resources and aggregate demand 2 has multiple equilibria.

**Remark 17.** For single-commodity routing games on SP networks the number of active regimes is at most the number of paths. This bound does not hold for multiple commodities and there can be as many as  $\prod_{h \in \mathcal{H}} (2^{|\mathcal{R}^h|} - 1)$  potential combinations for  $\mathcal{R}(\mu)$ , (see Appendix A.3).

### 5. Beyond singleton congestion games

In this section we provide some monotonicity results that go beyond the class of singleton congestion games studied in Section 3 and that also extend some known theorems for routing games with single OD pair. We recall that in a standard routing the commodities coincide with OD pairs and, moreover, the feasible strategies for each OD pair are all the possible paths connecting the corresponding origin and destination.

Cominetti et al. (2021, proposition 3.12) proved that in a single-OD routing game over a series-parallel network the equilibrium load of each edge is nondecreasing in the traffic demand. Every network that is not series-parallel contains a Wheatstone subnetwork (see Milchtaich, 2006); therefore, as shown in Example 4, there exist costs for which the equilibrium loads of some edges are decreasing in some demand interval. This implies that the series-parallel nature of the network is the best topological assumption that guarantees monotonicity of the equilibrium loads in a single-OD setting.

Unfortunately, for multi-OD routing games the network topology alone does not provide a criterion for the monotonicity of equilibrium loads. To obtain some useful results, we consider the following class of constrained routing games.

**Definition 18.** A constrained routing game (CRG) is a tuple  $(G, \mathcal{H}, c, \mathcal{P}, \mu)$  where

- $G = (\mathcal{V}, \mathcal{E})$  is a directed multigraph with vertex set  $\mathcal{V}$  and edge set  $\mathcal{E}$ ,
- $\mathcal{H}$  is a finite family of commodities,
- $c = (c_e)_{e \in \mathcal{E}}$  is a vector of edge cost functions,

- $\mathcal{P} = (\mathcal{P}^h)_{h \in \mathcal{H}}$ , with  $\mathcal{P}^h$  a nonempty set of paths between an origin  $O^h \in \mathcal{V}$  and a destination  $D^h \in \mathcal{V}$ ,
- $\mu = (\mu^h)_{h \in \mathcal{H}}$  is a demand vector.

This defines a congestion game structure with resource set  $\mathcal{R} = \mathcal{E}$ , commodity set  $\mathcal{H}$ , costs  $c = (c_e)_{e \in \mathcal{E}}$ , and strategy sets  $S^h = \mathcal{P}^h$ . Notice that in a constrained routing game the commodities are distinguished by their different strategy sets  $\mathcal{P}^h$ , although they might share the same OD pair and may also have some paths in common. This is in contrast with standard routing games where each OD pair is identified as a single commodity and  $\mathcal{P}^h$  includes all the paths from  $O^h$  to  $D^h$ . All the examples in Section 4 are in fact constrained routing games.

Although restricting the paths to a subset might seem a minor detail, it is in fact a flexible feature that allows us to represent any congestion game as a constrained routing game. Furthermore, we can also turn this routing game into a *common-OD* where all commodities have the same origin and destination, by


- adding a super-source  $O$  connected to each  $O^h$  by a zero-cost edge  $(O, O^h)$ ,
- adding a super-sink  $D$  connected by zero-cost edges  $(D^h, D)$ , and
- appending the edges  $(O, O^h)$  and  $(D^h, D)$  to each path of commodity  $h$ .

The next proposition shows that every congestion game is equivalent to a common-OD routing game over an extremely simple network, and all the complexity of the game is in fact encoded into the feasible sets of paths.

Formally, two congestion game structures  $\mathcal{G}$  and  $\check{\mathcal{G}}$  are said to be *equivalent* if there exist one-to-one correspondences  $h \leftrightarrow \check{h}$  between their commodities and  $s \leftrightarrow \check{s}$  between strategies, such that for each demand  $\mu$  and each feasible flow  $f$  of the first game, the flow  $\check{f}$  defined as  $\check{f}_{\check{s}} = f_s$  is feasible in the second game and the strategy costs coincide  $\check{c}_{\check{s}}(\check{f}) = c_s(f)$ . In this case the equilibria of both games are also in one-to-one correspondence.

**Proposition 19.** *Every congestion game is equivalent to a common-OD constrained routing game over a SP network.*

**Proof.** Consider a congestion game structure with resources  $\mathcal{R} = \{r_1, \dots, r_m\}$ . Consider the SP network in the figure below,



where each resource is represented by two parallel edges: one of them has the original resource cost  $c_r(\cdot)$ , and the other edge provides a bypass with zero cost. Any strategy  $s \subseteq \mathcal{R}$  can be represented as a path joining  $O$  to  $D$  that takes the top edge for each resource in  $s$ , and the bypass otherwise. We can then represent the commodities of the congestion game in the routing game by prescribing that they all have the same origin  $O$  and same destination  $D$ , whereas the feasible paths correspond to their feasible strategies in the original congestion game.  $\square$

Regarding the previous result, one may naturally ask whether a given nonatomic congestion game is equivalent to an *unconstrained* nonatomic routing game. We are not aware of any result on this question, apart from the somewhat related result by Milchtaich (2013), who showed that every finite game can be represented as a *weighted* atomic routing game.

As mentioned above, for standard multi-commodity routing games a SP network topology does not suffice to guarantee the monotonicity of the equilibrium loads. Indeed, Examples 20 and 21 below show that there exist common-OD constrained routing games such that:

- $G$  is SP;

- every commodity uses paths  $\mathcal{P}^h$  that form a SP subnetwork;
- the equilibrium loads  $x(\mu)$  are unique; but
- the map  $\mu \mapsto x(\mu)$  is not a MES.

**Example 20.** Consider Fisk’s network in Fig. 5(a) with  $c_{e_1}(x) = c_{e_2}(x) = x$ ,  $c_{e_3}(x) = x + 90$ , as in Example 3, and add bypass edges  $e_4, e_5$ , as in Fig. 5(b), with  $c_{e_4}(x) = c_{e_5}(x) = 0$ , producing commodities  $h_1, h_2, h_3$  where  $O^h = a$ ,  $D^h = c$  for every commodity  $h$ , and  $\mathcal{P}^{h_1} = \{(e_1, e_5)\}$ ,  $\mathcal{P}^{h_2} = \{(e_4, e_2)\}$ , and  $\mathcal{P}^{h_3} = \{(e_1, e_2), e_3\}$ . This defines an equivalent common-OD constrained routing game. As noted in Example 3, an increment in the demand of  $h_1$  pushes commodity  $h_3$  to divert more flow towards the direct path  $e_3$ , thus reducing the load on  $e_2$  (see Fisk, 1979).

**Example 21.** Monotonicity can also fail in a common-OD constrained routing game, even on a SP graph. Indeed, the standard Braess’s routing game in Fig. 2 corresponds to the single-commodity congestion game structure  $(\mathcal{E}, c, \{O v_1 v_2 D, O v_1 D, O v_2 D\})$ . Using Proposition 19 this is equivalent to a common-OD constrained routing game on a SP network, for which the MES property fails.

These examples show that, in addition to a SP topology, we need to impose further conditions on how the commodities overlap. To this end we introduce the following operations of series and parallel connection of congestion game structures.

**Definition 22.** Let  $\mathcal{G}_1 = (\mathcal{R}_1, c_1, \mathcal{S}_1)$  and  $\mathcal{G}_2 = (\mathcal{R}_2, c_2, \mathcal{S}_2)$  be two congestion game structures with disjoint resource sets  $\mathcal{R}_1 \cap \mathcal{R}_2 = \emptyset$ . The series and parallel game structures are both defined on the resource set  $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$  with their original cost functions. Specifically:

- The *series* game structure  $\mathcal{G}_1 \times \mathcal{G}_2$  has commodities  $(h_1, h_2) \in \mathcal{H}_1 \times \mathcal{H}_2$ , with corresponding strategy set  $S^{(h_1, h_2)} = \{s_1 \cup s_2 : (s_1, s_2) \in S_1^{h_1} \times S_2^{h_2}\}$ .
- The *parallel* game structure  $\mathcal{G}_1 \cup \mathcal{G}_2$  has commodity set  $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2$  and the original strategy sets  $S_1^{h_1}$  for  $h_1 \in \mathcal{H}_1$  and  $S_2^{h_2}$  for  $h_2 \in \mathcal{H}_2$ .
- A *constrained series-parallel* (CSP) congestion game structure is constructed starting from singleton congestion game structures and applying a finite number of series or parallel connections to game structures already constructed.

Whereas the parallel connection is a simple superposition of disjoint commodities, its combination with the series connection and the possibility of imposing constraints in the set of resources, provides a flexible tool to distinguish different types of commodities and to represent complex strategy sets that result from sequential processes. Consider for instance a family of different job classes, each one representing a commodity  $h \in \mathcal{H}$ , which must be processed in a series of stages  $k \in K$ . At every stage there is a set of machines  $M_k$  that work in parallel to perform the given task, while the jobs of type  $h$  can only be processed in a subset  $M_k^h \subset M_k$ . A commodity  $h$  can then be identified with a particular sequence of feasible machines  $(M_k^h)_{k \in K}$ , and its strategy set corresponds to the strategy set for a connection in series of singleton congestion game structures, one for each stage. On the other hand, some job classes might not require some processing stages, which can be modeled as a bypass strategy using the parallel operation. As an illustration, the graph in Fig. 6 can represent simultaneously commodities that must go sequentially over all four processing stages, possibly with restrictions on the machines that are allowed for each of them, as well as commodities that only perform the first and fourth stages, and skip the two intermediate stages. The bypass strategy can also be replaced by a series of alternative processing stages. Using such series and parallel operations one can model complex processing paths for different job classes. This construction gives rise to an SP graph, however the main additional ingredient is which combinations of commodities are allowed along the construction.






Fig. 5. Fisk's multi-commodity network can be embedded in a SP network with a common-OD, by adding two edges with zero cost.




Fig. 6. The SP graph for a job-processing game.

**Theorem 23.** Every CSP congestion game structure has a MES.

**Proof.** By induction and Theorem 6, it suffices to show that the MES property is preserved under series and parallel operations on game structures. To this end, let  $\mathcal{G}_1$  and  $\mathcal{G}_2$  be two congestion game structures with MES's  $\mu_1 \mapsto x_1(\mu_1)$  and  $\mu_2 \mapsto x_2(\mu_2)$  respectively. Then we prove the two parts:

(a) *The series game structure  $\mathcal{G}_1 \times \mathcal{G}_2$  has a MES.* Let  $\mu = (\mu^{(h_1, h_2)})_{h_1 \in \mathcal{H}_1, h_2 \in \mathcal{H}_2}$  with  $\mu^{(h_1, h_2)}$  the demand for the commodity  $(h_1, h_2)$  in a game whose structure is  $\mathcal{G}_1 \times \mathcal{G}_2$ . Define

$$\forall h_1 \in \mathcal{H}_1, \mu_1^{h_1} = \sum_{h_2 \in \mathcal{H}_2} \mu^{(h_1, h_2)}; \quad \forall h_2 \in \mathcal{H}_2, \mu_2^{h_2} = \sum_{h_1 \in \mathcal{H}_1} \mu^{(h_1, h_2)}, \tag{5.1}$$

$\mu_1 = (\mu_1^{h_1})_{h_1 \in \mathcal{H}_1}$ , and  $\mu_2 = (\mu_2^{h_2})_{h_2 \in \mathcal{H}_2}$ . An equilibrium for  $\mu$  can be obtained by superposing  $x_1(\mu_1)$  on the resources  $\mathcal{R}_1$  and  $x_2(\mu_2)$  on the resources  $\mathcal{R}_2$ . Since an increase of any demand  $\mu^{(h_1, h_2)}$  induces an increase in the demands  $\mu_1^{h_1}$  and  $\mu_2^{h_2}$ , the loads in  $x_1(\mu_1)$  and  $x_2(\mu_2)$  increase, so that this superposed equilibrium provides a MES for the series game.

(b) *The union game structure  $\mathcal{G}_1 \cup \mathcal{G}_2$  has a MES.* For each demand  $\mu = (\mu_1, \mu_2)$  in a game with structure  $\mathcal{G}_1 \cup \mathcal{G}_2$  we can directly find an equilibrium by superposing  $x_1(\mu_1)$  on the resources  $\mathcal{R}_1$  and  $x_2(\mu_2)$  on the resources  $\mathcal{R}_2$ . Since the loads in these two equilibria are monotone with respect to each individual demand, the same holds for their superposition which provides a MES for the parallel game structure.  $\square$

**Remark 24.** The common-OD constrained routing game of Example 20, in which Fisk's network is embedded, does not have a CSP structure: the strategy set of  $h_3$  cannot be obtained as a strategy set of a previously present commodity when constructing a parallel game structure. For a different reason, the common-OD SP routing game in Example 21 does not have a CSP game structure either. Indeed, its network would be made of 5 two-edge parallel network in series, each one associated to a resource, i.e., an edge of the Wheatstone network. The classical Braess's routing game has a single commodity with three strategies, and a strategy set with cardinality 3 cannot be obtained as the Cartesian product of the strategy sets of the 5 two-edge parallel networks connected in series. A series of Pigou games has strong limitations, for example, a commodity with a number of paths divisible by a prime larger than 2 is not constructible as in Definition 22.

**Remark 25.** It is somewhat odd that the class of CSP's does not include single-OD routing games over an SP graph, in which there is

a single commodity and every path is allowed. This happens because the parallel connection does not allow to merge commodities and they are kept separated. However, if in the construction of CSP's instead of taking singleton congestion games as the initial atoms, one replaces each singleton strategy with a series-parallel routing game structure with a single commodity, then Theorem 23 remains true for this larger class which trivially includes series-parallel routing games.

**Remark 26.** Every CSP game structure as defined above is a matroid game. In fact, if  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are two matroid game structures, then the parallel game structure  $\mathcal{G}_1 \cup \mathcal{G}_2$  is trivially a matroid game structure, whereas the strategy sets  $\mathcal{S}^{(h_1, h_2)}$  in the series game structure  $\mathcal{G}_1 \times \mathcal{G}_2$  correspond to the direct sum operation on the original matroid bases  $\mathcal{S}_1^{h_1}$  and  $\mathcal{S}_2^{h_2}$ . As a consequence, for strictly increasing costs the previous result can also be derived from Fujishige et al. (2017, lemma 3.2).

Whereas CSP congestion game structures were defined for general congestion games, they can also be described as constrained routing games with a specific structure. The following representation is also more natural compared to the one in Proposition 19.

**Theorem 27.** Every CSP congestion game structure  $\mathcal{G} = (\mathcal{R}, c, \mathcal{S})$  is equivalent to a common-OD constrained routing game structure  $(G, c, \mathcal{P})$  such that

- (i) the graph  $G$  is SP,
- (ii) for each commodity  $h$  all the paths in  $\mathcal{P}^h$  visit the same vertices in the same order,
- (iii) for every two paths  $p_1, p_2 \in \mathcal{P}^h$  and edges  $e_1 \in p_1$  and  $e_2 \in p_2$  connecting two subsequent vertices, the paths obtained from  $p_1$  and  $p_2$  by exchanging  $e_1$  with  $e_2$  also belong to  $\mathcal{P}^h$ .

Furthermore, every common-OD constrained routing game satisfying (i), (ii), (iii), has a CSP congestion game structure.

**Proof.** Every CSP game structure is built starting with singleton congestion games and applying a finite number of series or parallel operations. We start by noting that every singleton congestion game is equivalent to a constrained routing game on a parallel network with two vertices connected by edges corresponding to the resources of the singleton congestion game, and any such game satisfies (i), (ii), (iii). Hence, it suffices to show that these properties are preserved under series and parallel operations.

Consider two congestion game structures  $\mathcal{G}_1 = (\mathcal{R}_1, c_1, \mathcal{S}_1)$  and  $\mathcal{G}_2 = (\mathcal{R}_2, c_2, \mathcal{S}_2)$  which are respectively equivalent to some common-OD constrained routing games  $(G_1, c_1, \mathcal{P}_1)$  and  $(G_2, c_2, \mathcal{P}_2)$  satisfying (i), (ii), (iii).

The series game structure  $G_1 \times G_2$  is then equivalent to the constrained routing game structure  $(\tilde{G}, \tilde{c}, \tilde{\mathcal{P}})$  where  $\tilde{G}$  is obtained by joining in series the graphs  $G_1$  and  $G_2$ , the costs  $\tilde{c}$  are the cost functions given by  $c_1$  and  $c_2$  on the corresponding edges, and the commodities are given by the sets of paths obtained choosing commodities  $h_1$  for  $(G_1, c_1, \mathcal{P}_1)$  and  $h_2$  for  $(G_2, c_2, \mathcal{P}_2)$ , and joining every path in  $\mathcal{P}^{h_1}$  with every path in  $\mathcal{P}^{h_2}$  to construct paths in  $\tilde{\mathcal{P}}$ . Moreover,  $(\tilde{G}, \tilde{c}, \tilde{\mathcal{P}})$  satisfies (i), (ii), (iii) because  $(G_1, c_1, \mathcal{P}_1)$  and  $(G_2, c_2, \mathcal{P}_2)$  do.

Similarly, the parallel game structure  $G_1 \cup G_2$  is equivalent to the constrained routing game structure  $(\bar{G}, \bar{c}, \bar{\mathcal{P}})$  where  $\bar{G}$  is obtained joining in parallel  $G_1$  and  $G_2$ , the costs  $\bar{c}$  are the cost functions given by  $c_1$  and  $c_2$  on the corresponding edges, and the commodities are given by the commodities of  $(G_1, c_1, \mathcal{P}_1)$  and  $(G_2, c_2, \mathcal{P}_2)$ . Also in this case, the routing game  $(\bar{G}, \bar{c}, \bar{\mathcal{P}})$  satisfies (i), (ii), (iii) because  $(G_1, c_1, \mathcal{P}_1)$  and  $(G_2, c_2, \mathcal{P}_2)$  do.

This completes the proof of the first claim of the theorem.

Conversely, notice that every SP graph  $G$  is constructed starting with parallel networks and joining them in series or in parallel for a finite number of times. Suppose that a common-OD constrained routing game  $(G, c, \mathcal{P})$  structure satisfies (i), (ii), (iii).

If the graph  $G$  is obtained by joining in series two graphs  $G_1$  and  $G_2$ , we can endow them with cost functions that associate costs to edges as in  $c$ . Furthermore, given a commodity  $h$  for  $(G, c, \mathcal{P})$  we can define commodities  $h_1$  on  $G_1$  and  $h_2$  on  $G_2$  by determining for  $i = 1, 2$  the set of paths

$$\mathcal{P}^{h_i} = \{p \text{ path in } G_i \text{ s.t. } p \text{ is part of a path in } \mathcal{P}^h\}.$$

Since  $(G, c, \mathcal{P})$  satisfies (iii), we have  $\mathcal{P}^h = \mathcal{P}^{h_1} \times \mathcal{P}^{h_2}$ , so that  $(G, c, \mathcal{P})$  is the series game structure of the two constrained routing games just defined on  $G_1$  and  $G_2$ .

If the graph  $G$  is obtained by joining in parallel two graphs  $G_1$  and  $G_2$ , then we can assume that the direct edges from the origin and the destination of  $G$  are all contained in one of the two. We can again endow  $G_1$  and  $G_2$  with cost functions that associate costs to edges as in  $c$ . Furthermore because of property (ii), for every commodity  $h$  of  $(G, c, \mathcal{P})$  the paths in  $\mathcal{P}^h$  all belong to one between  $G_1$  and  $G_2$ . This allows us to define for each commodity of  $(G, c, \mathcal{P})$ , a commodity either in  $G_1$  or  $G_2$ , so that  $(G, c, \mathcal{P})$  is the parallel game structure of the two constrained routing games just defined on  $G_1$  and  $G_2$ .  $\square$

**Remark 28.** Note that Braess’s classical example in Fig. 2 satisfies (ii) and (iii), but does not satisfy (i). Fisk’s network embedding of Example 20 satisfies (i) and (iii) but does not satisfy (ii). Finally, the constrained routing game of Example 21, obtained by embedding Braess’s game in a SP graph as in Proposition 19, satisfies (i) and (ii), but not (iii).

**Remark 29.** Conditions (i), (ii), (iii) in Theorem 27 can be equivalently stated by requiring that all feasible paths for a commodity  $h$  visit a specific ordered sequence of nodes; between successive nodes only a specific subset of parallel edges are allowed; and  $\mathcal{P}^h$  includes all possible paths in this subnetwork. Still another equivalent description is to require that for any two paths  $p_1, p_2 \in \mathcal{P}^h$  the mixed path where we follow  $p_1$  up to an intermediate node and then continue with  $p_2$  is also in  $\mathcal{P}^h$ .

## 6. Summary and open problems

This paper studied the monotonicity of equilibrium travel times and equilibrium loads in response to variations of the demands, identifying conditions under which the paradoxical phenomena of non-monotonicity cannot happen. We considered the general setting of congestion games, with a special focus on singleton congestion games with multiple commodities for which we established in Theorem 6 the existence of a selection of the equilibrium loads which monotonically increase with respect to the demand of every commodity.

We next explored the notion of comonotonicity, which captures the idea that different resource loads jointly increase or decrease after variations of the demands. Theorem 12 described how comonotonicity is connected to the structure of equilibria in terms of how the commodities are ranked by cost and how the resources become active or inactive as the demands vary. We complemented this finding by a structural result on the regions of the demand space for which the same sets of resources are used at equilibrium.

Theorem 23 extended the study of monotonicity from singleton congestion games to the larger class of congestion games having a CSP structure, reminiscent of the concept of a SP network. We also derived an embedding that maps congestion games into constrained routing games (see Proposition 19) and characterized the classes of congestion games with good monotonicity properties by embedding them into routing games (see Theorem 27). This last result sheds light on the features that produce the paradoxes and showcases the difference between single and multiple OD networks. When the network has a single OD pair, its topology is the sole relevant factor to guarantee the monotonicity of equilibrium loads. In the multiple OD case the structure of the paths that are in each OD pair also plays a crucial role.

A first open question not addressed in this paper, and which will be interesting to explore, is how the structural results on the regions  $\Gamma^\approx$  and sub-regions  $\Gamma_\theta^\approx$  for the different active regimes might be exploited to devise an algorithm for building a curve of equilibria along a demand curve, analog to the path-following method for piece-wise affine costs developed by Klimm and Warode (2022). A basic question here is to investigate the geometry of the regions  $\Gamma^\approx$  for specific classes of cost functions. For the special case of Bureau of Public Roads (BPR) costs, we conjecture that the boundaries between these regions are asymptotic to straight lines through the origin. This would imply that when the demands are scaled proportionally, the regimes will not repeat and the curve will eventually enter into a particular asymptotic region  $\Gamma^\approx$  and remain there forever. The latter could inspire a path following algorithm to build a curve of equilibria.

A second open problem is to find an algorithm to recognize CSP congestion game structures. In this regard, one could be tempted to use the equivalent game in Proposition 19 for which (i) and (ii) in Theorem 27 hold trivially, so that only (iii) would need to be checked. Unfortunately, the CSP property is not preserved under equivalence: for instance, a singleton congestion game with only one commodity is CSP by definition, but its equivalent representation in Proposition 19 is not because property (iii) fails. This suggests that recognizing CSP game structures is not straightforward. As a possible starting point to address this question, one might try to adapt the existing algorithms for recognizing SP networks (see Eppstein, 1992; He & Yesha, 1987; Valdes et al., 1982).

## Acknowledgments

We thank the reviewers for their careful reading and insightful comments. We also thank Tobias Harks for pointing out the connection of our results with the paper by Fujishige et al. (2017), and Tzachi Gilboa and Ludger Rüschemdorf for some historical insights about comonotonicity. Valerio Dose and Marco Scarsini are members of GNAMPA-INdAM. Their work was partially supported by the GNAMPA project CUP\_E53C22001930001 “Limiting behavior of stochastic dynamics in the Schelling segregation model” and by the MIUR PRIN project 2022EKNE5K “Learning in Markets and Society”. Roberto Cominetti’s research was supported by Proyecto Anillo ANID/PIA/ACT192094.

**Appendix A. Supplementary proofs**

*A.1. Missing proof*

**Proof of Proposition 1.** Let  $(\mathcal{R}, c, S)$  be a nonatomic congestion game structure. For every demand  $\mu \in \mathbb{R}_+^H$ , let  $V(\mu)$  be the minimum value of the Beckmann potential as in (2.5), that is,

$$V(\mu) = \min_{x \in X_\mu} \sum_{r \in \mathcal{R}} C_r(x_r). \tag{A.1}$$

We obtain the result as a consequence of convex duality. Consider the function  $\varphi_\mu : \mathbb{R}^S \times \mathbb{R}^H \rightarrow \mathbb{R} \cup \{+\infty\}$  given by

$$\begin{aligned} \varphi_\mu(f, z) &= \begin{cases} \sum_{r \in \mathcal{R}} C_r(\sum_{s' \ni r} f_{s'}) & \text{if } f \geq 0, \sum_{s \in S^h} f_s = \mu^h + z^h \text{ for every } h \in H, \\ +\infty & \text{otherwise,} \end{cases} \end{aligned} \tag{A.2}$$

which is a proper closed convex function. Letting  $v_\mu(z)$  denote the optimal value function of the primal problem

$$(P_\mu) \quad \inf_f \varphi_\mu(f, z), \tag{A.3}$$

we have  $V(\mu + z) = v_\mu(z)$  and, in particular,  $V(\mu) = v_\mu(0)$ .

Since  $\varphi_\mu$  is convex, we have that  $z \mapsto v_\mu(z) = V(\mu + z)$  is also convex, from which we deduce that  $\mu \rightarrow V(\mu)$  is convex. Moreover, the perturbed function  $\varphi_\mu$  yields a corresponding dual

$$(D_\mu) \quad \min_{\lambda \in \mathbb{R}^H} \varphi_\mu^*(0, \lambda), \tag{A.4}$$

where  $\varphi_\mu^*$  is the Fenchel conjugate function, that is,

$$\begin{aligned} \varphi_\mu^*(0, \lambda) &= \sup_{f, z} \langle 0, f \rangle + \langle \lambda, z \rangle - \varphi_\mu(f, z) \\ &= \sup_{f \geq 0} \sum_{h \in H} \left( \lambda^h \left( \sum_{s \in S^h} f_s - \mu^h \right) \right) - \sum_{r \in \mathcal{R}} C_r \left( \sum_{s' \ni r} f_{s'} \right). \end{aligned} \tag{A.5}$$

Since  $V(\mu')$  is finite for all  $\mu' \in \mathbb{R}_+^H$ , it follows that  $v_\mu(z) = V(\mu + z)$  is finite for  $z$  in some interval around  $0$ , and then the convex duality theorem implies that there is no duality gap and the subgradient  $\nabla v_\mu(0)$  at  $z = 0$  coincides with the optimal solution set  $S(D_\mu)$  of the dual problem, that is,  $\nabla V(\mu) = \nabla v_\mu(0) = S(D_\mu)$ .

We claim that the dual problem has a unique solution, which is exactly the vector of equilibrium costs  $\lambda(\mu)$ . Indeed, fix an optimal solution  $\hat{f}$  for  $v_\mu(0) = V(\mu)$  and recall that this is just a Wardrop equilibrium. The dual optimal solutions are precisely the  $\lambda$ 's in  $\mathbb{R}^H$  such that

$$\varphi_\mu(\hat{f}, 0) + \varphi_\mu^*(0, \lambda) = 0.$$

This equation can be written explicitly as

$$\sum_{r \in \mathcal{R}} C_r \left( \sum_{s \ni r} \hat{f}_s \right) + \sup_{f \geq 0} \sum_{h \in H} \left( \lambda^h \left( \sum_{s \in S^h} f_s - \mu^h \right) \right) - \sum_{r \in \mathcal{R}} C_r \left( \sum_{s' \ni r} \hat{f}_{s'} \right) = 0,$$

from which it follows that  $f = \hat{f}$  is an optimal solution in the latter supremum. The corresponding optimality conditions are

$$\begin{aligned} \lambda^h - \sum_{r \in s} c_r \left( \sum_{s' \ni r} \hat{f}_{s'} \right) &= 0, \quad \text{if } \hat{f}_s > 0, h \in H, s \in S^h, \\ \lambda^h - \sum_{r \in s} c_r \left( \sum_{s' \ni r} \hat{f}_{s'} \right) &\leq 0, \quad \text{if } \hat{f}_s = 0, h \in H, s \in S^h, \end{aligned}$$

which imply that  $\lambda^h$  is the equilibrium cost of the OD pair  $h$  for the Wardrop equilibrium, that is,  $\lambda^h = \lambda^h(\mu)$  for every  $h \in H$ . It follows that the subgradient  $\nabla V(\mu) = \{\lambda(\mu)\}$  so that  $\mu \mapsto V(\mu)$  is not only convex but also differentiable with gradient  $\nabla V(\mu) = \lambda(\mu)$ . The conclusion follows by noting that every convex differentiable function

is automatically of class  $C^1$  and its gradient is monotone, in the sense that  $\langle \nabla V(\mu_1) - \nabla V(\mu_2), \mu_1 - \mu_2 \rangle \geq 0$  for every  $\mu_1, \mu_2 \in \mathbb{R}_+^H$ , which in particular implies that  $\lambda^h$  is nondecreasing in the variable  $\mu^h$ .

The continuity of the equilibrium resource costs  $\tau_r = \tau_r(\mu)$  is a consequence of Berge's maximum theorem (see, e.g., Aliprantis & Border, 2006, Section 17.5). Indeed, as explained in Fukushima (1984), the equilibrium resource costs are optimal solutions for the strictly convex dual program (2.6). Hence, since the objective function is jointly continuous in  $(\tau, \mu)$ , Berge's theorem implies that the optimal solution correspondence is upper-semicontinuous. However, in this case the optimal solution is unique, so that the optimal correspondence is single-valued, and, as a consequence, the equilibrium resource costs  $\tau_r(\mu)$  are continuous.  $\square$

**Remark 30.** A similar analysis where we reformulate the primal problem by including resource load variables  $x_r$  and considering perturbations in the flow balance equations  $x_r = \sum_{s \ni r} f_s + z_r$ , yields the dual problem (2.6), which characterizes the equilibrium costs  $\tau_r$ .

Perhaps a more direct argument is as follows. Let us rewrite the flow balance equations  $\hat{x}_r = \sum_{s \ni r} \hat{f}_s$  in vector form as  $\hat{x} = \sum_{s \in S} \hat{f}_s \eta^s$  where  $\eta^s = (\eta_r^s)_{r \in \mathcal{R}}$  denotes the indicator vector with

$$\eta_r^s = \mathbb{1}_{\{r \in s\}}. \tag{A.6}$$

Since  $\mu_h = \sum_{s \in S^h} \hat{f}_s$ , by letting  $\hat{\alpha}_s^h = \hat{f}_s / \mu^h$  for all  $s \in S^h$  we have that  $\sum_{s \in S^h} \hat{\alpha}_s^h = 1$  and  $\hat{\alpha}_s^h \geq 0$ , the latter inequality being strict only for the optimal strategies for commodity  $h$ . With these notations, we can write

$$\hat{x} = \sum_{s \in S} \hat{f}_s \eta^s = \sum_{h \in H} \mu_h \sum_{s \in S^h} \hat{\alpha}_s^h \eta^s. \tag{A.7}$$

Now, for each  $h \in H$  the super-differential of the concave function  $\Theta_h(\tau) := \min_{s \in S^h} \sum_{r \in s} \tau_r$  is given by convex hull of the indicators of optimal strategies, that is,

$$\partial \Theta_h(\tau) = \text{co} \left\{ \eta^s : s \in S^h, \sum_{r \in s} \tau_r = \Theta_h(\tau) \right\}, \tag{A.8}$$

so that from (A.7) we derive

$$\hat{x} = \sum_{h \in H} \mu_h \sum_{s \in S^h} \hat{\alpha}_s^h \eta^s \in \sum_{h \in H} \mu_h \partial \Theta(\tau). \tag{A.9}$$

Finally, letting  $\Phi(x) := \sum_{r \in \mathcal{R}} C_r(x_r)$  and  $\tau_r = c_r(\hat{x}_r)$  we clearly have  $\tau = \nabla \Phi(\hat{x})$ , which is equivalent to  $\hat{x} \in \partial \Phi^*(\tau)$  where the Fenchel's conjugate is given by  $\Phi^*(\tau) = \sum_{r \in \mathcal{R}} C_r^*(\tau_r)$ . Since all the involved functions are finite and continuous, using standard subdifferential calculus rules, (A.9) is equivalent to  $0 \in \partial \Psi(\tau)$  for the convex function  $\Psi(\tau) = \Phi^*(\tau) - \sum_{h \in H} \mu_h \Theta_h(\tau)$  which is precisely the objective function in (2.6).

*A.2. Characterization of comonotonicity*

For the sake of completeness we include the following characterization of comonotonicity. This is a folk result (see e.g., Landsberger and Meilijson (1994)), but its proof is not easy to find in the literature.

**Lemma 31.** Consider a finite family of functions  $\psi_i : \Omega \rightarrow \mathbb{R}$  for  $i = 1, \dots, m$  and let  $s(\omega) := \sum_{i=1}^m \psi_i(\omega)$ . Then, the family  $\{\psi_i : i = 1, \dots, m\}$  is comonotonic if and only if there exist nondecreasing functions  $F_i : \mathbb{R} \rightarrow \mathbb{R}$  such that  $\psi_i(\omega) = F_i(s(\omega))$  for all  $\omega \in \Omega$  and  $i = 1, \dots, m$ .

**Proof.** Since the “if” implication holds trivially, it suffices to prove the “only if”. Suppose that the  $\psi_i$ 's are comonotonic. For  $z \in \mathbb{R}$  define  $F_i(z) = \sup_{\omega \in \Omega \{ \psi_i(\omega) : s(\omega) \leq z \}}$  if there is some  $\omega \in \Omega$  with  $s(\omega) \leq z$ , and  $F_i(z) = \inf_{\omega \in \Omega \{ \psi_i(\omega) \text{ otherwise} \}}$ . Clearly the functions  $F_i$  are nondecreasing and  $\psi_i(\omega) \leq F_i(s(\omega))$ , whereas comonotonicity implies that the latter holds with equality for all  $i = 1, \dots, m$  and  $\omega \in \Omega$ .

It remains to show that the  $F_i$ 's are everywhere finite. Indeed, if  $F_i(z) = \infty$  for some  $z \in \mathbb{R}$  we can find a sequence  $\omega_n \in \Omega$  with  $s(\omega_n) \leq z$

such that  $\psi_i(\omega_n)$  increases to  $\infty$ . However, by comonotonicity, the latter implies  $s(\omega_n) \rightarrow \infty$  which is a contradiction. Now, if  $F_i(z) = -\infty$  we must be in the case  $F_i(z) = \inf_{\omega \in \Omega} \psi_i(\omega) = -\infty$  and comonotonicity implies  $\inf_{\omega \in \Omega} s(\omega) = -\infty$ , so we may find  $\omega \in \Omega$  with  $s(\omega) \leq z$  which yields the contradiction  $F_i(z) \geq \psi_i(\omega) > -\infty$ .  $\square$

A.3. A remark on the number of active regimes

The monotonicity result in Cominetti et al. (2021, proposition 3.12) implies that the number of active regimes in a single-commodity routing game on a SP network is at most the number of paths. This bound does not hold for multiple commodities. In a singleton congestion game there are  $\prod_{h \in \mathcal{H}} (2^{|\mathcal{R}^h|} - 1)$  potential combinations for  $\hat{\mathcal{R}}(\mu)$ , and this bound may be attained (see Example 32 below). This is not the case for single-commodity routing games: if we consider a subnetwork composed by only two paths, it is always SP and only two of the three nonempty subsets of paths can actually correspond to an active regime  $\hat{\mathcal{R}}(\mu)$  for some  $\mu \in [0, +\infty)$ .

**Example 32.** Let us build a multi-commodity routing game that attains the maximal bound for the number of active regimes. Take  $m$  a positive integer and consider a routing game on a parallel network with  $m$  resources (edges)  $\mathcal{R} = \{1, \dots, m\}$  with cost functions

$$\forall i \in \{1, \dots, m\}, \quad c_i(x_i) = x_i + i,$$

and  $m + 1$  commodities where each commodity  $i = 1, \dots, m$  can only use one specific resource  $\mathcal{R}^i = \{i\}$ , whereas commodity  $(m + 1)$  can use all the resources  $\mathcal{R}^{m+1} = \mathcal{R}$ .

We claim that  $\hat{\mathcal{R}}(\mu)$  assumes the maximum number  $\prod_{i=1}^{m+1} (2^{|\mathcal{R}^i|} - 1) = 2^m - 1$  of possible active regimes as the demands  $\mu$  vary. Indeed, for each commodity  $i \leq m$  the active regime is always  $\{i\}$ , whereas every nonempty subset  $\rho^{m+1} \subset \mathcal{R}$  is the active regime of the  $(m + 1)$ -th commodity for some demand  $\mu$ . Namely, let  $i_{\max} = \max\{i \in \rho^{m+1}\}$  and consider the demand

$$\forall i \leq m, \quad \mu^i = \begin{cases} 0 & \text{if } i \in \rho^{m+1}, \\ i_{\max} & \text{if } i \notin \rho^{m+1}, \end{cases}$$

$$\mu^{m+1} = \sum_{i \in \rho^{m+1}} (i_{\max} - i).$$

Then, the unique equilibrium is such that commodity  $(m + 1)$  allocates  $i_{\max} - i$  to each resource  $i \in \rho^{m+1}$  with cost  $i_{\max}$ , whereas every resource  $i \notin \rho^{m+1}$  has a cost  $i_{\max} + i > i_{\max}$ , so that the active regime for commodity  $(m + 1)$  is exactly  $\rho^{m+1}$ .

Appendix B. List of symbols

$c_r$	cost function of resource $r$
$c_s$	cost function of strategy $s$ , defined in (2.3)
$c$	vector of cost functions; it can be indexed both by elements in $\mathcal{E}$ or $\mathcal{P}$
$c_r^\varepsilon(x_r)$	regularized cost $c_r(x_r) + 2\varepsilon x_r$
$C_r(x_r)$	primitive of costs $\int_0^{x_r} c_r(z) dz$
$C_r^*(\cdot)$	Fenchel conjugate of $C_r(\cdot)$
$\mathcal{C}$	subset of commodities having the same equilibrium cost
$D^h$	destination for OD pair $h$
$(D_\mu)$	dual problem, defined in (A.4)
$\mathcal{E}$	set of edges
$e^h$	$h$ -th vector of the canonical basis of $\mathbb{R}^{\mathcal{H}}$
$f_s^h$	flow on strategy $s$ in commodity $h$
$f^h$	$h$ -th commodity flow vector $(f_s^h)_{s \in \mathcal{S}^h}$
$f$	flow vector $(f^h)_{h \in \mathcal{H}}$
$F_\mu$	set of feasible pairs $(x, f)$ for the demand vector $\mu$

$G$	directed multigraph
$\mathcal{G}$	$(\mathcal{R}, c, \mathcal{S})$ , congestion game structure
$\mathcal{G}^\varepsilon$	$(\mathcal{R}, c^\varepsilon, \mathcal{S})$ , perturbed congestion game structure
$\mathcal{G}_C$	$(\mathcal{R}_C, c, \mathcal{S}_C)$ , single commodity game defined in Definition 10
$\mathcal{G}_1 \times \mathcal{G}_2$	series game
$\mathcal{G}_1 \cup \mathcal{G}_2$	parallel game
$h$	commodity
$\mathcal{H}$	set of commodities
$O^h$	origin for OD pair $h$
$(P_\mu)$	primal problem, defined in (A.3)
$\hat{\mathcal{P}}(\mu)$	set of paths that attain equilibrium cost at equilibrium with demand $\mu$
$\mathcal{P}^h$	the set of paths of commodity $h$
$\mathcal{P}$	$(\mathcal{P}^h)_{h \in \mathcal{H}}$
$r$	resource
$\mathcal{R}$	set of resources
$\hat{\mathcal{R}}^h(\mu)$	set of active resources for commodity $h \in \mathcal{H}$
$\hat{\mathcal{R}}(\mu)$	$(\hat{\mathcal{R}}^h(\mu))_{h \in \mathcal{H}}$ , active regime
$\mathcal{R}_0$	set of resources such that $c_r(x_r(\mu_0)) = \lambda^h(\mu_0)$
$\mathcal{R}_0^+$	$\{r \in \mathcal{R}_0 : x_r(\mu_0 + te^h) > x_r(\mu_0)\}$ , defined in (3.1)
$\mathcal{R}_0^-$	$\{r \in \mathcal{R}_0 : x_r(\mu_0 + te^h) < x_r(\mu_0)\}$ , defined in (3.2)
$\mathcal{R}_0^=$	$\{r \in \mathcal{R}_0 : x_r(\mu_0 + te^h) = x_r(\mu_0)\}$ , defined in (3.3)
$\mathcal{R}_C$	$(\cup_{h \in \mathcal{C}} \mathcal{R}^h) \setminus (\cup_{h' > \mathcal{C}} \mathcal{R}^{h'})$ , defined in (4.3)
$\mathcal{S}^h$	set of feasible strategies for commodity $h$
$\mathcal{S}$	$\times_{h \in \mathcal{H}} \mathcal{S}^h$ , set of strategy profiles
$S(D_\mu)$	optimal solution set of the dual problem
$SC(\mu)$	$\sum_{h \in \mathcal{H}} \mu^h \lambda^h(\mu)$ , social cost
$v_\mu(z)$	$\inf_f \varphi_\mu(f, z)$ , defined in (A.3)
$V(\mu)$	$\min_{x \in \mathcal{X}_\mu} \sum_{r \in \mathcal{R}} C_r(x_r)$ , defined in (A.1)
$\mathcal{V}$	set of vertices
$x_r$	load of resource $r$ , defined in (2.2)
$x$	$(x_r)_{r \in \mathcal{R}}$ , load vector
$\mathcal{X}_\mu$	projection of the set of feasible pairs $F_\mu$ onto the $x$ variables
$\Gamma^{\lesssim}$	demand regions induced by a given order $\lesssim$ , defined in (4.1)
$\Gamma_\rho^{\lesssim}$	demand subregions for a given order $\lesssim$ and active regime $\rho$ , defined in (4.2)
$\eta_r^s$	$\mathbb{1}_{\{r \in s\}}$ , defined in (A.6)
$\eta^s$	$(\eta_r^s)_{r \in \mathcal{R}}$
$\lambda^h$	equilibrium cost of commodity $h$ , defined in (2.4)
$\lambda$	$(\lambda^h)_{h \in \mathcal{H}}$ , equilibrium cost vector
$\mu^h$	demand for commodity $h$
$\mu$	$(\mu^h)_{h \in \mathcal{H}}$ demand vector
$\mu_C$	aggregate demand on $\mathcal{C}$
$\rho$	$(\rho^h)_{h \in \mathcal{H}}$ , regime
$\rho^h$	subset of $\mathcal{P}^h$
$\tau_r$	equilibrium cost of resource $r$
$\varphi_\mu$	defined in (A.2)
$\varphi_\mu^*$	Fenchel conjugate of $\varphi_\mu$ , defined in (A.5)

References

Acemoglu, D., & Ozdaglar, A. (2007). Competition and efficiency in congested markets. *Mathematics of Operations Research*, 32(1), 1–31.

Aliprantis, C. D., & Border, K. C. (2006). *Infinite dimensional analysis* (third ed.). (p. xxii+703). Berlin: Springer.

Arrow, K. J. (1970). *Essays in the theory of risk-bearing* (p. vii+278). Amsterdam-London: North-Holland Publishing Co..

Attouch, H. (1996). Viscosity solutions of minimization problems. *SIAM Journal on Optimization*, 6(3), 769–806.

Auslender, A., Cominetti, R., & Haddou, M. (1997). Asymptotic analysis for penalty and barrier methods in convex and linear programming. *Mathematics of Operations Research*, 22(1), 43–62.

- Beckmann, M. J., McGuire, C. B., & Winsten, C. B. (1956). *Studies in the economics of transportation*. New Haven, CT: Yale University Press.
- Bilò, V., & Vinci, C. (2017). On the impact of singleton strategies in congestion games. In *LIPICs. Leibniz int. proc. inform.: vol. 87, 25th European symposium on algorithms*. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, Art. No. 17, 14.
- Borch, K. (1962). Equilibrium in a reinsurance market. *Econometrica*, 30(3), 424–444.
- Borchers, M., Breuewsma, P., Kern, W., Sloatbeek, J., Still, G., & Tibben, W. (2015). Traffic user equilibrium and proportionality. *Transportation Research Part B*, 79, 149–160.
- Braess, D. (1968). Über ein Paradoxon aus der Verkehrsplanung. *Unternehmensforschung*, 12, 258–268.
- Braess, D., Nagurney, A., & Wakolbinger, T. (2005). On a paradox of traffic planning. *Transportation Science*, 39(4), 446–450.
- Castiglioni, M., Marchesi, A., Gatti, N., & Coniglio, S. (2019). Leadership in singleton congestion games: what is hard and what is easy. *Artificial Intelligence*, 277, 103177, 31.
- Cominetti, R. (1999). Nonlinear averages and convergence of penalty trajectories in convex programming. In M. Théra, & R. Tichatschke (Eds.), *Ill-posed variational problems and regularization techniques* (pp. 65–78). Berlin, Heidelberg: Springer.
- Cominetti, R., Dose, V., & Scarsini, M. (2021). The price of anarchy in routing games as a function of the demand. *Mathematical Programming*.
- Cominetti, R., Dose, V., & Scarsini, M. (2023). *Phase transitions of the price-of-anarchy function in multi-commodity routing games: Technical report*, arXiv:2305.03459.
- Dafermos, S., & Nagurney, A. (1984). On some traffic equilibrium theory paradoxes. *Transportation Research Part B*, 18(2), 101–110.
- Dellacherie, C. (1971). Quelques commentaires sur les prolongements de capacités. In *Lecture notes in math.: vol. 191, Séminaire de probabilités, V (Univ. Strasbourg, Année universitaire 1969–1970)* (pp. 77–81). Springer, Berlin.
- Deori, L., Margellos, K., & Prandini, M. (2017). On the connection between Nash equilibria and social optima in electric vehicle charging control games. *IFAC-PapersOnLine*, 50(1), 14320–14325.
- Dhaene, J., Denuit, M., Goovaerts, M. J., Kaas, R., & Vyncke, D. (2002). The concept of comonotonicity in actuarial science and finance: theory. *Insurance: Mathematics & Economics*, 31(1), 3–33, 5th IME Conference (University Park, PA, 2001).
- Eppstein, D. (1992). Parallel recognition of series-parallel graphs. *Information and Computer*, 98(1), 41–55.
- Fisk, C. (1979). More paradoxes in the equilibrium assignment problem. *Transportation Research Part B*, 13(4), 305–309.
- Fisk, C., & Pallottino, S. (1981). Empirical evidence for equilibrium paradoxes with implications for optimal planning strategies. *Transportation Research Part A*, 15(3), 245–248.
- Fotakis, D., Kontogiannis, S., Koutsoupias, E., Mavronicolas, M., & Spirakis, P. (2009). The structure and complexity of Nash equilibria for a selfish routing game. *Theoretical Computer Science*, 410(36), 3305–3326.
- Fujishige, S., Goemans, M. X., Harks, T., Peis, B., & Zenklusen, R. (2017). Matroids are immune to Braess' paradox. *Mathematics of Operations Research*, 42(3), 745–761.
- Fukushima, M. (1984). On the dual approach to the traffic assignment problem. *Transportation Research Part B*, 18(3), 235–245.
- Gairing, M., Lücking, T., Mavronicolas, M., & Monien, B. (2010). Computing Nash equilibria for scheduling on restricted parallel links. *Theory of Computing Systems*, 47(2), 405–432.
- Gairing, M., & Schoppmann, F. (2007). Total latency in singleton congestion games. In X. Deng, & F. C. Graham (Eds.), *Internet and network economics* (pp. 381–387). Berlin, Heidelberg: Springer.
- Gonczarowski, Y. A., & Tennenholtz, M. (2016). A hydraulic approach to equilibria of resource selection games. In *Proceedings of the 2016 ACM conference on economics and computation* (p. 477). New York, NY, USA: Association for Computing Machinery.
- Hall, M. A. (1978). Properties of the equilibrium state in transportation networks. *Transportation Science*, 12(3), 208–216.
- Harks, T., & Klimm, M. (2012). On the existence of pure Nash equilibria in weighted congestion games. *Mathematics of Operations Research*, 37(3), 419–436.
- Harks, T., Schröder, M., & Vermeulen, D. (2019). Toll caps in privatized road networks. *European Journal of Operational Research*, 276(3), 947–956.
- Harks, T., & Timmermans, V. (2017). Equilibrium computation in atomic splittable singleton congestion games. In *Lecture notes in comput. sci.: vol. 10328, Integer programming and combinatorial optimization* (pp. 442–454). Springer, Cham.
- He, X., & Yesha, Y. (1987). Parallel recognition and decomposition of two terminal series parallel graphs. *Information and Computer*, 75(1), 15–38.
- Hoerl, A. E. (1959). Optimum solution of many variables equations. *Chemical Engineering Progress*, 55, 69–78.
- Hoerl, A. E. (1962). Application of ridge analysis to regression problems. *Chemical Engineering Progress*, 58, 54–59.
- Hoerl, A. E., & Kennard, R. W. (1970). Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1), 55–67.
- Josefsson, M., & Patriksson, M. (2007). Sensitivity analysis of separable traffic equilibrium equilibria with application to bilevel optimization in network design. *Transportation Research Part B*, 41(1), 4–31.
- Klimm, M., & Warode, P. (2022). Parametric computation of minimum-cost flows with piecewise quadratic costs. *Mathematics of Operations Research*, 47(1), 812–846.
- Koçyiğit, Ç., Rujeeraipoon, N., & Kuhn, D. (2022). Robust multidimensional pricing: separation without regret. *Mathematical Programming*, 196(1–2), 841–874.
- Konishi, H. (2004). Uniqueness of user equilibrium in transportation networks with heterogeneous commuters. *Transportation Science*, 38(3), 315–330.
- Landsberger, M., & Meilijson, I. (1994). Co-monotone allocations, Bickel-Lehmann dispersion and the Arrow-Pratt measure of risk aversion. *Annals of Operations Research*, 52, 97–106.
- Ma, Z., Callaway, D. S., & Hiskens, I. A. (2013). Decentralized charging control of large populations of plug-in electric vehicles. *IEEE Transactions on Control Systems Technology*, 21(1), 67–78.
- Mehr, N., & Horowitz, R. (2020). How will the presence of autonomous vehicles affect the equilibrium state of traffic networks? *IEEE Transactions on Control of Network Systems*, 7(1), 96–105.
- Meunier, F., & Pradeau, T. (2014). The uniqueness property for networks with several origin-destination pairs. *European Journal of Operational Research*, 237(1), 245–256.
- Milchtaich, I. (2000). Generic uniqueness of equilibrium in large crowding games. *Mathematics of Operations Research*, 25(3), 349–364.
- Milchtaich, I. (2005). Topological conditions for uniqueness of equilibrium in networks. *Mathematics of Operations Research*, 30(1), 225–244.
- Milchtaich, I. (2006). Network topology and the efficiency of equilibrium. *Games and Economic Behavior*, 57(2), 321–346.
- Milchtaich, I. (2013). Representation of finite games as network congestion games. *International Journal of Game Theory*, 42(4), 1085–1096.
- Nimalsiri, N. L., Mediwaththe, C. P., Ratnam, E. L., Shaw, M., Smith, D. B., & Halgamuge, S. K. (2020). A survey of algorithms for distributed charging control of electric vehicles in smart grid. *IEEE Transactions on Intelligent Transportation Systems*, 21(11), 4497–4515.
- Patriksson, M. (2004). Sensitivity analysis of traffic equilibria. *Transportation Science*, 38(3), 258–281.
- Pradeau, T. (2014). *Congestion games with player-specific cost functions* (Ph.D. thesis), (2014PEST1096), Université Paris-Est.
- Puccetti, G., & Scarsini, M. (2010). Multivariate comonotonicity. *Journal of Multivariate Analysis*, 101(1), 291–304.
- Rossi, T., McNeil, S., & Hendrickson, C. (1989). Entropy model for consistent impact-fee assessment. *Journal of Urban Planning and Development*, 115, 51–63.
- Schmeidler, D. (1982). *Subjective probability without additivity: Technical report*, The Foerder Institute for Economic Research, Tel-Aviv University.
- Schmeidler, D. (1984). *Subjective probability and expected utility without additivity: Technical report*, Institute of Mathematics and its Applications, University of Minnesota.
- Schmeidler, D. (1986). Integral representation without additivity. *Proceedings of the American Mathematical Society*, 97(2), 255–261.
- Schmeidler, D. (1989). Subjective probability and expected utility without additivity. *Econometrica*, 57(3), 571–587.
- Tikhonov, A. N. (1943). On the stability of inverse problems. *C. R. (Doklady) Academy of Science URSS (N.S.)*, 39, 176–179.
- Tikhonov, A. N. (1963). On the solution of ill-posed problems and the method of regularization. *Doklady Akademii Nauk SSSR*, 151, 501–504.
- Tikhonov, A. N., & Arsenin, V. Y. (1977). *Scripta series in mathematics, Solutions of ill-posed problems* (p. xiii+258). V. H. Winston & Sons, Washington, D.C.; John Wiley & Sons, New York-Toronto, Ont.-London.
- Valdes, J., Tarjan, R. E., & Lawler, E. L. (1982). The recognition of series parallel digraphs. *SIAM Journal on Computing*, 11(2), 298–313.
- van de Klundert, J., Cominetti, R., Liu, Y., & Kong, Q. (2023). The interdependence between hospital choice and waiting time – with a case study in urban China. arXiv:2306.16256v1.
- Verbree, J., & Cherukuri, A. (2023). *Wardrop equilibrium and Braess's paradox for varying demand: Technical report*, arXiv preprint arXiv:2310.04256.
- Wan, C. (2016). Strategic decentralization in binary choice composite congestion games. *European Journal of Operational Research*, 250(2), 531–542.
- Wardrop, J. G. (1952). Some theoretical aspects of road traffic research. Vol. I, In *Proceedings of the institute of civil engineers, part II* (pp. 325–378).
- Wilson, R. (1968). The theory of syndicates. *Econometrica*, 36(1), 119–132.
- Yaari, M. E. (1987). The dual theory of choice under risk. *Econometrica*, 55(1), 95–115.