



Phytochemical Study of a Species with Ethnopharmacological Interest: *Sideritis romana* L.

A. Venditti^{1,2*}, C. Frezza², L. Guarcini³, S. Foddai², M. Serafini² and A. Bianco¹

¹Dipartimento di Chimica, Università di Roma "La Sapienza", Piazzale Aldo Moro, 5 - 00185 Roma, Italy.

²Dipartimento di Biologia Ambientale, Università di Roma "La Sapienza", Piazzale Aldo Moro, 5 - 00185 Roma, Italy.

³Dipartimento di Chimica e Tecnologie del Farmaco, Università di Roma "La Sapienza", Piazzale Aldo Moro, 5 - 00185 Roma, Italy.

Authors' contributions

This work was carried out in collaboration between all authors which contributed equally to this study.
All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2016/23809

Editor(s):

(1) Marcello Iriti, Professor of Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

- (1) Arun Kumar, Hindu Post Graduate College, Zamania, India.
(2) Amanda Dias de Araujo Uchoa, University Federal de Pernambuco, Brazil.
(3) Saeed Saeed Alghamdi, Umm al-Qura University, Saudi Arabia.
(4) Daniela Hanganu, Luliu Hațieganu University of Medicine and Pharmacy, Romania.
Complete Peer review History: <http://sciencedomain.org/review-history/13171>

Original Research Article

Received 23rd December 2015
Accepted 15th January 2016
Published 5th February 2016

ABSTRACT

The phytochemical analysis of *Sideritis romana* L., a species largely used as a traditional remedy, led to the isolation and the identification of several acetylated flavonoid glycosides with apigenin, luteolin, hypolaetin and isoscutellarein backbone. Among these, an apigenin derivative (2) was recognized for the first time in *Sideritis* as well as in the *Lamiaceae* family. The iridoidic pattern of this species showed the presence of harpagide, currently considered the main taxonomical maker in this genus, together with 6-deoxyharpagide which is a rare compound since it was previously recognized in a limited number of species of *Lamiaceae*. Other identified iridoids were ajugoside and bartsioside, the latter recognized for the first time in the species as well as in the family. Acetylated flavonoid glycosides and iridoid glucosides are considered of chemotaxonomic relevance in several genera comprised in the Lamioidae subfamily and the occurrence of both of these classes of compounds in *S. romana* was discussed. The co-presence of the acetylated

*Corresponding author: E-mail: alessandro.venditti@uniroma1.it;

apigenin derivative and of 6-deoxyharpagide could be used as a marker at a specific level seen their relative rarity in *Lamiaceae*. Regarding the traditional uses of this species, the isolated compounds may give a rationale from a chemical standpoint.

Keywords: *Sideritis romana*; *Lamiaceae*; *iridoids*; *flavonoids*; *chemotaxonomy*; *traditional medicine*.

1. INTRODUCTION

Sideritis romana L., also known as ironwort, is an annual plant belonging to *Lamiaceae* family (Lamioideae subfamily). It is characterized by an erected stem which is simple or branched at the base. It owns elliptic basal leaves, the upper ones with a lanceolate shape. The flowers are inserted in outdistanced verticillasters and the youngest ones fill the inferior part of the inflorescence. The corolla is yellow with white or pink shades. The fruits are brown trigonal and warty nuculae. It can be found in the Mediterranean area where it widely grows in arid meadows and grazings till the altitude of 1900 m a.s.l. In Italy it is present in almost all the territory except for some regions of northern Italy like Veneto, Trentino Alto Adige and Valle d'Aosta and also in Piedmont where its presence is actually unsure [1]. It can be mistaken for *Stachys ocymastrum* (L.) Briq. but the latter owns leaves with different shape (oval and crenate along the margins excluding the base) and the youngest flowers fill the superior part of the inflorescence [2]. Many species of the genus *Sideritis* (mountain tea) are widely used in traditional medicine in the Mediterranean and Balkan regions as a tonic and for the treatment of several diseases i.e. gastrointestinal complaints, inflammations, and rheumatic disorders [3]. *Sideritis* are used in Turkish folk medicine as beverage in brew form for its high antibacterial, carminative, diuretic and digestive properties [4]. They are also used as anti-inflammatory, anti-ulcer, cytostatic, antimicrobial, vulnerary, astringent and circulation stimulating agents [5]. The genus name *Sideritis* is derived from the Greek word "sideros" (Σίδηρος, iron) and was known to ancient Greeks because of its property to heal wounds caused by iron weapons during battles. In Italy, it is used for its vulnerary properties to heal cuts and wounds of the skin by the shepherds of the Madonie Mountains (Sicily) [6]. Moreover in some Italian regions, the decoy of the exsiccated leaves is used to wash the unconcealed parts of the body in case of fright in order to soothe anxiety [1].

In literature there are little studies on this species focusing mainly on the content of the essential oil

[7,8]. Only particularities of the composition of the polar fraction are reported, i.e the revision of siderin structure [9] and the identification of a new glycosidic flavonoid [10]. There is also in literature a detailed review on *Sideritis* genus and its traditional uses [3] but the species *romana* obviously was not included since a complete phytochemical analysis of the polar fraction was not available. We have then decided to re-study a sample of this species, wanting to build a first more detailed and total molecular pattern on the polar fraction content.

2. MATERIALS AND METHODS

2.1 General

NMR spectra were recorded on a Varian (now Agilent Technologies) Mercury 300 MHz instrument and on a Bruker Avance III 400 MHz instrument, using CDCl₃, CD₃OD or D₂O as deuterated solvents. The chemical shifts are expressed in ppm from TMS in deuteriochloroform; the signal of HDO (s) at 4.78 ppm was used as reference for spectra in D₂O; and the internal solvent signal (*m*5) at 3.31 ppm for spectra recorded in CD₃OD.

MS spectra were performed on a Q-TOF MICRO spectrometer (Waters, Manchester, UK) equipped with an ESI source operating in negative and/or positive ion mode. The flow rate of the sample infusion was 10 µl per min. with 100 acquisitions per spectrum. Data were analysed by the MassLynx software developed by Waters.

Solvents of RPE grade were purchased from Sigma Aldrich (Milan, Italy) or Carlo Erba Reagenti (Milan, Italy) and silica gel 60 (70-230 mesh ASTM) was from Fluka.

2.2 Bidimensional NMR Experiments

Bidimensional spectra were performed on a Bruker Avance III 400 MHz instrument, operating at 9.4 T at 298° K. NOESY experiments were acquired with a spectral width of 15 ppm in both dimensions, a mixing time length of 700 ms, recycle delay of 2 s and a data matrix of 4K x 256 points. HSQC experiments were acquired

with a spectral width of 15 and 250 ppm for the proton and carbon respectively, an average $^1J_{C-H}$ of 145 Hz, recycle delay of 2 s and a data matrix of 4K x 256 points. HMBC experiments were acquired with a spectral width of 15 and 250 ppm for the proton and carbon respectively, long range coupling constant of $^nJ_{C-H}$ of 8 Hz, recycle delay of 2 s and a data matrix of 4K x 256 points.

2.3 Plant Material

A sample of fresh plant material (168.9 g) was harvested at flowering stage at the end of June 2014 in Central Italy on Latium hills at an altitude of 150 m a.s.l. (GPRS coordinates 41.615434, 13.548094). The botanical identification was performed by one of us (A.V.) using available literature [1,2]. The specimen of the studied plant is registered at "Museo Erbario, Dipartimento di Biologia Ambientale, Sapienza Università di Roma" under the following accession number: RO H.A. N°18419.

From these separation procedures twelve compounds were isolated: (*E*)-phytol (**1**) (14.9 mg) [11,12], apigenin 7-*O*-(6''-*O*-acetyl- β -D-glucopyranoside) (**2**) (15.6 mg) [13], 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(3-hydroxy-4-methoxyphenyl)-4*H*-1-benzopyran-4-one (**3**) (39.7 mg), 7-[[2-*O*-(6-*O*-acetyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]oxy]-2-(3,4-dihydroxyphenyl)-5-hydroxy-4*H*-1-benzopyran-4-one (**4**) (11.9 mg), 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one (**5**) (5.0 mg) [14], 7-[[2-*O*- β -D-allopyranosyl- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxy-3-methoxyphenyl)-4*H*-1-benzopyran-4-one (**6**) (4.7 mg) [15], 7-[[6-*O*-acetyl-2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-4*H*-1-benzopyran-4-one (**7**) (5.1 mg), 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one (**8**) (5.5 mg) [16], 6-deoxiharpagide (**9**) (7.0 mg) [17], harpagide (**10**) (10.2 mg) [18], ajugoside (**11**) (3.7 mg) [19] and bartsioside (**12**) (2.1 mg) [20] (Fig. 1).

NMR data of compounds **1** and **9-11** were in accordance with literature data.

2.5 Spectral Data of Isolated Compounds

apigenin 7-*O*-(6''-*O*-acetyl- β -D-glucopyranoside) (2**)**, 1H -NMR (CD₃OD, 400 MHz), δ : 7.86 (2H, d, J = 8.8 Hz, H-2'/H-6'), 6.93 (2H, d, J = 8.8 Hz, H-3'/H-5'), 6.76 (1H, d, J = 2.1 Hz, H-8), 6.64 (1H, s, H-3), 6.49 (1H, d, J = 2.1 Hz, H-6), 5.04 (1H, d, J = 7.3 Hz, H-1''), 4.46 (1H, dd, J = 12.0, 2.2 Hz, H-6''a), 4.23 (1H, dd, J = 12.0, 7.1 Hz, H-6''b), 2.06 (3H, s, CH₃CO).

^{13}C NMR (100 MHz, MeOD) δ : 184.1 (C-4), 172.7 (CH₃CO), 166.7 (C-2), 164.6 (C-7), 162.9 (C-5), 158.9 (C-9), 129.6 (C-2'/C-6'), 123.1 (C-1'), 117.1 (C-3'/C-5'), 104.9 (C-10), 104.2 (C-3), 101.5 (C-1''), 101.1 (C-8), 96.2 (C-6), 77.8 (C-3''), 75.6 (C-5''), 74.7 (C-2''), 71.6 (C-4''), 64.8 (C-6''), 20.8 (CH₃CO). ESI-MS: m/z 496.97 [M+Na]⁺; m/z 472.92 [M-H]⁻.

7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(3-hydroxy-4-methoxyphenyl)-4*H*-1-benzopyran-4-one (3**)**, 1H NMR (400 MHz, DMSO) δ : 7.58 (2H, m, H-2'/H-6'), 6.98 (1H, s, H-3), 6.94 (1H, d, J = 8.9 Hz, H-5'), 6.80 (1H, d, J = 1.8 Hz, H-6), 6.45 (1H, d, J = 1.8 Hz, H-8), 5.21 (1H, d, J = 7.5 Hz, H-1''), 4.79 (1H, d, J = 7.9 Hz, H-1'''), 4.12 (1H, dd, J = 11.7, 1.8 Hz, H-6''a), 4.00 (1H, dd, J = 11.7, 5.4 Hz, H-6''b), 3.89 (3H, s, OCH₃), 1.94 (3H, s, CH₃CO). ESI-MS: m/z 689.11 [M+Na]⁺; m/z 665.02 [M-H]⁻.




Fig. 1. Structures of compounds isolated from *Sideritis romana*

7-[[2-O-(6-O-acetyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]oxy]-2-(3,4-dihydroxyphenyl)-5-hydroxy-4H-1-benzopyran-4-one (4), 1H -NMR (CD₃OD, 300MHz) δ : 7.61 – 7.52 (1H, br s, H-2'), 7.52 (1H, br d, J = 8.5 Hz, H-6'), 6.94 (1H, d, J = 8.4 Hz, H-5'), 6.83 (1H, d, J = 1.9 Hz, H-6), 6.70 (1H, s, H-3), 6.51 (1H, d, J = 1.9 Hz, H-8), 5.24 (1H, d, J = 6.7 Hz, H-1''), 4.98 (1H, d, J = 8.0 Hz, H-1'''), 4.16 (1H, dd, J = 12.2, 1.7 Hz, H-6''a), 2.06 (3H, s, CH₃CO).
ESI-MS: m/z 675.10 [M+Na]⁺; m/z 650.99 [M-H]⁻.

7-[[2-O-(6-O-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxyphenyl)-4H-benzopyran-4-one (5), 1H -NMR (CD₃OD, 300Hz) δ : 7.98 (2H, d, J = 8.2 Hz, H-2'/H-6'), 6.94 (1H, s, H-3), 6.92 (2H, d, J = 8.2 Hz, H-3'/H-5'), 6.73 (1H, d, J = 2.2 Hz, H-6), 6.42 (1H, d, J = 2.1 Hz, H-8), 5.18 (1H, d, J = 7.5 Hz, H-1''), 4.80 (1H, d, J = 7.9 Hz, H-1'''), 4.12 (1H, dd, J = 11.8, 1.9 Hz, H-6''a), 4.00 (1H, dd, J = 11.8, 5.7 Hz, H-6''b), 1.89 (3H, s, CH₃CO).
ESI-MS: m/z 659.12 [M+Na]⁺; m/z 635.02 [M-H]⁻.

7-[[2-O- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxy-3-methoxyphenyl)-4H-1-benzopyran-4-one (6), 1H NMR (400 MHz, MeOD) δ : 7.57 (1H, dd, J = 8.5, 2.1 Hz, H-6'), 7.54 (1H, d, J = 2.1 Hz, H-2'), 6.95 (1H, d, J = 8.5 Hz, H-5'), 6.89 (1H, d, J = 2.2 Hz, H-6), 6.71 (1H, s, H-3), 6.53 (1H, d, J = 2.2 Hz, H-8), 5.27 (1H, d, J = 7.2 Hz, H-1''), 5.01 (1H, d, J = 7.8 Hz, H-1'''), 3.98 (3H, s, 3'-CH₃O), other sugar signals between 3.98 and 3.25 ppm.
ESI-MS: m/z 646.67 [M+Na]⁺; m/z 622.85 [M-H]⁻.

7-[[6-O-acetyl-2-O-(6-O-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-4H-1-benzopyran-4-one (7), 1H NMR (400 MHz, MeOD) δ : 7.56 (1H, br d, J = 8.9 Hz, H-6'), 7.55 (1H, br s, H-2'), 6.96 (1H, d, J = 8.6 Hz, H-5), 6.84 (1H, br s, H-6), 6.53 (1H, s, H-3), 5.25 (1H, d, J = 7.8 Hz, H-1''), 5.00 (1H, d, J = 8.0 Hz, H-1'''), 3.99 (3H, s, 4'-CH₃O), 2.01 (3H, s, CH₃CO), 1.99 (3H, s, CH₃CO).
ESI-MS: m/z 746.97 [M+Na]⁺; m/z 723.58 [M-H]⁻.

7-[[2-O-(6-O-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one (8), 1H NMR (400 MHz, MeOD) δ : 8.00 (2H, d, J = 8.1 Hz, H-2'/H-6'), 6.94 (2H, d, J = 8.3 Hz, H-3'/H-5'), 6.79 (1H, s, H-6), 6.55 (1H, s, H-3), 5.23 (1H, d, J = 7.8 Hz, H-1''), 4.98 (1H, d, J = 8.0 Hz, H-1'''), 1.99 (3H, s, CH₃CO).
ESI-MS: m/z 689.20 [M+Na]⁺; m/z 665.26 [M-H]⁻.

bartsioside (12), 1H NMR (400 MHz, MeOD) δ : 6.20 (1H, dd, J = 6.3, 2.3 Hz, H-3), 5.85 (1H, d, J = 1.7 Hz, H-7), 5.40 (1H, d, J = 5.2 Hz, H-1), 4.98 (1H, d, J = 6.2 Hz, H-4), 4.24 (2H, br d, J = 10.0 Hz, H-10), 3.16 (2H, m, H-5/H-9), 2.84 – 2.70 (1H, m, H-6a), 2.24 – 2.14 (1H, m, H-6b).
ESI-MS: m/z 353.14 [M+Na]⁺

3. RESULTS AND DISCUSSION

The study of the ethanolic extract of the fresh aerial parts of *S. romana* led to the isolation and the identification of twelve compounds namely (*E*)-phytol (**1**), apigenin 7-*O*-(6''-*O*-acetyl- β -D-glucopyranoside) (**2**), 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(3-hydroxy-4-methoxyphenyl)-4*H*-1-benzopyran-4-one (**3**), 7-[[2-*O*-(6-*O*-acetyl- β -D-glucopyranosyl)- β -D-glucopyranosyl]oxy]-2-(3,4-dihydroxyphenyl)-5-hydroxy-4*H*-1-benzopyran-4-one (**4**), 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one (**5**), 7-[[2-*O*- β -D-allopyranosyl- β -D-glucopyranosyl]oxy]-5-hydroxy-2-(4-hydroxy-3-methoxyphenyl)-4*H*-1-benzopyran-4-one (**6**), 7-[[6-*O*-acetyl-2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-4*H*-1-benzopyran-4-one (**7**), 7-[[2-*O*-(6-*O*-acetyl- β -D-allopyranosyl)- β -D-glucopyranosyl]oxy]-5,8-dihydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one (**8**), 6-deoxyharpagide (**9**), harpagide (**10**), ajugoside (**11**) and bartsioside (**12**).

Compound (**1**) is a very common acyclic diterpene alcohol which is part of the chlorophyll structure and is also endowed with interesting biological activities i.e. antiradical, antimicrobial and cytotoxic [21-23]. The presence of (**1**) can explain some of the traditional uses of this species as antibacterial.

Compounds from (**2**) to (**6**) are glycosidic-flavones derivatives of apigenin and luteolin. In particular compounds (**3**), (**4**), (**5**) and (**6**) are diglucosides containing an allose residue in the structure and were already evidenced in some species of the genus such as *S. scardica* Griseb., *S. grandiflora* Salzm. ex Benth., *S. raeseri* Boiss. & Heldr., *S. taurica* Steph. ex Willd., *S. syriaca* L., *S. perfoliata* L., as well as in near species such as *Stachys aegyptiaca* Pers. and *St. anisochila* Vis. & Pancic [14-16,24-27]. These compounds were reported here for the first time in *S. romana*. On the other hand compound (**2**) was evidenced for the first time as a constituent of *Sideritis*, and also in *Lamiaceae* family. Previously, compound (**2**) has been evidenced in few species of Asteraceae such as *Matricaria chamomilla* L. [28] and *Chrysanthemum morifolium* Ramat. [29] and in an Euphorbiaceae species as *Mallotus apelta* (Lour.) Muell.-Arg. [30], thus resulting a quite rare compound. From a taxonomic standpoint it is worth to note that *Lamiaceae* and Asteraceae are families comprised in Asterales order while Euphorbiaceae are comprised in Malpighiales. Acetylated flavonoids derivatives of apigenin and luteolin containing allose have been previously isolated from species comprised in Ajugoideae subfamily such as *Ajuga reptans* and *A. genevensis* [31]. Among flavonoids, isoscutellarein and hypolaetin derivatives of chemotaxonomic relevance (**7** and **8**) were evidenced during this study. In fact, mono and diacetylated isoscutellarein derivatives containing allose have been recently reported for this genus

[32] and are traditionally considered as further chemotaxonomic markers (besides iridoids) since their occurrence seems to be restricted to some related genera in the *Lamiaceae* like *Pogostemon*, *Galeopsis*, *Stachys* and *Sideritis* itself [33].

Considering the ethnomedicinal uses of this species, the presence of a considerable amount of these flavonoid derivatives of apigenin and luteolin, might perhaps justify its traditional use as a herbal tea because apigenin and luteolin glycosides are universally known to exert a strong anxiolytic action [34]. More recently, phenolics of *Sideritis* proved also to possess cytotoxic action toward several cancer lines [35-37] as well as spasmolytic [38] and a potent antioxidant activity [39-41] which resulted comparable to those exerted by *Camellia sinensis* [42]. Moreover, the hypolaetin and isoscutellarein derivatives are endowed with a strong antioxidant and anticholinesterase activity which may result in a neuroprotective effect in the Alzheimer disease [43]. The presence of flavonoids in diglycosidic form improves their solubility in water, the solvent used for infusions, and at the same time their bioavailability. Several analytical studies were conducted to evaluate the phenolic content of mountain teas [5,39,44] and also the urinary excretion of metabolites has been carried out in recent times [45], as a demonstration of the high interest on *Sideritis* species.

Lastly compounds (**9-12**) are iridoids. Iridoids are considered chemotaxonomic markers in *Lamiaceae*, and in particular harpagide and its derivatives. harpagide (**10**) was already evidenced in *Sideritis* [46], while its derivative (**9**) was recognized for the first time as a constituent of *S. romana* as well as in the genus itself. This molecular trait may be characteristic of the studied species since in *Sideritis* were usually

found aucubin derivatives, melittoside or its acylated derivatives: melittoside and 5-alloxy-aucubin were found in *Sideritis italica* (Mill.) Greuter et Burter [32], melittoside, 10-O-(*E*)-feruloylmelittoside, 10-O-(*E*)-*p*-coumaroylmelittoside were evidenced in *S. trojana* Bornm. [47], the latter compound was also recognized in *S. lanata* L. [48]. Harpagide (**10**) is a widespread compound in several species of the *Lamiaceae* while its derivative (**9**) is less common having been evidenced before only in few species such as *Galeopsis tetrahit* L. and *G. pubescens* Besser [17] and *Ajuga iva* (L.) Schreb [49]. It is worth to note that *Galeopsis* genus is comprised in the Lamioideae subfamily of *Lamiaceae* just like *Sideritis* genus, while *Ajuga* is comprised in a different subfamily, Ajugoideae. The presence of (**9**) in *S. romana*, together with the presence of acetylated allosyl flavonoids with apigenin and luteolin aglicones backbone, is a further similarity with the genus *Ajuga*, even if these two genus are comprised in different subfamilies. In fact, there are several reports on isolation of iridoids peculiar of *Ajuga* also in *Sideritis* species: ajugol and/or its derivative ajugoside (**11**), recognized also in the present study, were found in several species of this genus such as *S. montana* L., *S. scardica* Griseb., *S. syriaca* L. [50], *S. perfoliata* L. subsp. *perfoliata* (syn. of *Sideritis perfoliata* L.) [51], *S. libanotica* Labill. subsp. *linearis* (Bentham) Bornm [52] and *S. lycia* Boiss. & Heldr [53]. The presence of common compounds between these two genera, may have chemosystematic implications, revealing a more strict proximity between themselves. It is worth to note that bartsioside (**12**) was found in *Sideritis* and, more in general, in *Lamiaceae* for the first time in this study. The presence of this compound, even if in little amount, is quite unusual since it is mainly recognized from species traditionally comprised in Scrophulariaceae [20,54,55] (the majority of these species have been recently moved in other families [56]) and Plantaginaceae [57]. Nevertheless, this is not unexpected from a biogenetic point of view since bartsioside is formally the 6-deoxyderivative of aucubin and several *Sideritis* showed the presence of iridoids related to aucubin. Moreover, this species showed also the presence of (**9**), an additional 6-deoxylated iridoid. The presence of (**6**) and (**12**) may be an evidence of a metabolic tendency toward the biosynthesis of deoxy-derivative.

The bioactivities of iridoids (antibacterial, anti-inflammatory, and antiviral activities) has been largely studied, in particular the anti-inflammatory

one [58-60]. Also the structure-activity relationship on the basis of the different patterns of substitution (hydroxylation, unsaturation, and acylation) on the iridoid skeleton has been carried out [61]. Accordingly with the last study, the presence of an olefinic bond at C₇-C₈ (aucubin derivative) is one of the most positive character for activity and aucubin derivatives have been found in *Sideritis* [32,47,48]. Among the iridoids identified in *S. romana* bartsioside, presenting a C₇-C₈ unsaturation, may be one of the major responsible of the anti-inflammatory action. Most probably, iridoids are the main responsible of the anti-inflammatory effect also in other *Sideritis* species, even if other active constituents like flavonoids, largely represented in this species, may have obviously a synergistic or modulating action.

4. CONCLUSIONS

In conclusion, during our work on this plant we isolated and identified several different compounds of systematic importance. Among these, four glycosidic apigenin and luteolin derivatives were evidenced besides isoscutellarein and hypolaetin derivatives which characterize the species of this genus. Compound (**2**) resulted a novelty in the studied species as well as in the genus and given its rarity, it may be a suitable marker at a specific level, while compounds from (**3**) to (**8**) were already evidenced in other *Lamiaceae* as *Sideritis* and *Ajuga*. In addition to this, it's worth to note the finding of 6-deoxyharpagide (**9**), harpagide (**10**) and ajugoside (**11**) which, though they are not new in the genus, were reported for the first time in this species, as well as (**12**) never reported before in *Lamiaceae*. The presence of (**9**) is an additional similarity with some *Ajuga* species and, together with the presence of the isolated flavonoids, may be an evidence of a more strict proximity among these genera of *Lamiaceae* even if comprised in two different subfamilies. Moreover, the presence of a huge amount of flavonoids and iridoids, together with (*E*)-phytol, may support the traditional uses of this plant by exerting a wide range of bioactivities.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Pignatti S. Flora d'Italia. Edagricole; 1982. Italian.
- Conti F, Abbate G, Alessandrini A, Blasi C. An annotated checklist of the Italian vascular flora. Rome: Palombi Press; 2005.
- González-Burgos E, Carretero ME, Gómez-Serranillos MP. *Sideritis* spp.: Uses, chemical composition and pharmacological activities — A review. J. Ethnopharmacol. 2011;135(2):209–225.
- Özcan M, Chalchat JC, Akgül A. Essential oil composition of Turkish mountain tea (*Sideritis* spp.). Food Chem. 2001;75:459–463.
- Palomino OM, Gomez-Serranillos P, Carretero E, Villar A. High-performance liquid chromatography of flavonoids from *Sideritis* species. J Chromatography A. 1996;731:103–108.
- Leporatti ML, Impieri M. Ethnobotanical notes about some uses of medicinal plants in Alto Tirreno Cosentino area (Calabria, Southern Italy). J Ethnobiol Ethnomed. 2007;3:34–39.
- Kirimer N, Tabanca N, Özek T, Tümen G, Baser KHC. Essential oils of annual *Sideritis* species growing in Turkey. Pharm Biol. 2000;38(2):106–111.
- Flamini G, Cioni PL, Morelli I, Maccioni S, Tomei PE. Characterization of the volatile fraction of a *Sideritis romana* population from Montemarcello (eastern Liguria). J Essent Oil Res. 1994;6(3):239–242.
- Venturella P, Bellino A, Piozzi F. Revised structure for siderin. Tetrahedron Lett. 1974;12:979–81.
- Venturella P, Bellino A, Papucci A. A new flavone glycoside from *Sideritis romana* L. Heterocycles. 1977;6(3):267–72.
- Bhattacharya AK, Rana KC. Antimicrobial agent, (*E*)-phytol and lauric amide from the plant *Lagascea mollis*. Indian J Chem. 2013;52B:901–903.
- Venditti A, Frezza C, Riccardelli M, Foddai S, Nicoletti M, Serafini M, Bianco A. Secondary metabolites from *Scrophularia canina* L. Nat Prod Res; 2015. DOI: 10.1080/14786419.2015.1122598 (in press).
- Svehlikova V, Bennett RN, Mellon FA, Needs PW, Piacente S, Kroon PA, et al. Isolation, identification and stability of acylated derivatives of apigenin 7-*O*-glucoside from chamomile (*Chamomilla recutita* [L.] Rauschert). Phytochemistry. 2004;65(16):2323–2332.
- El-Ansari MA, Barron D, Abdalla MF, Saleh NAM, Le Quere JL. Flavonoid constituents of *Stachys aegyptiaca*. Phytochemistry. 1991;30(4):1169–1173.
- Rabanal RM, Valverde S, Martin-Lomas M, Rodriguez B, Chari VM. Chrysoeriol 7-(2"-*O*- β -D-allopyranosyl)- β -D-glucopyranoside from *Sideritis grandiflora*. Phytochemistry. 1982;21(7):1830–1832.
- Lenherr A, Mabry TJ. Acetylated allose-containing flavonoid glucosides from *Stachys anisochila*. Phytochemistry. 1987;26(4):1185–1188.
- Sticher O, Rogenmoser E, Weisflog A. New iridoid glucosides from *Galeopsis tetrahit* and *Galeopsis pubescens*. Tetrahedron Lett. 1975;5:291–294.
- Bobbitt JM, Segebarth KP. The iridoid glycosides and similar substances. In: Taylor WI, Battersby AR, editors. Cyclopentanoid terpene derivatives. New York: Marcel Dekker Inc; 1969.
- Guiso M, Marini-Bettolo R, Agostini A. Iridoids. XIII. Ajugoside and ajugol. Structure and configuration. Gazz Chim Ital. 1974;104(1-2):25–33.
- Bianco A, Guiso M, Iavarone C, Trogolo C. Iridoids. XX. Bartsioside, structure and configuration. Gazz Chim Ital. 1976; 106(7-8):725–732.
- Pejin B, Savic A, Sokovic M, Glamoclija J, Ciric A, Nikolic M, et al. Further in vitro evaluation of antiradical and antimicrobial activities of phytol. Nat Prod Res. 2014a;28(6):372–376.
- Pejin B, Kojic V, Bogdanovic G. An insight into the cytotoxic activity of phytol at *in vitro* conditions. Nat Prod Res. 2014b;28(22):2053–2056.
- Pejin B, Ciric A, Glamoclija J, Nikolic M, Sokovic M. *In vitro* anti-quorum sensing activity of phytol. Nat Prod Res. 2015;29(4):374–377.
- Petreska JS, Stefkov G, Kulevanova S, Alipieva K, Bankova V, Stefova M. Phenolic compounds of mountain tea from the Balkans: LC/DAD/ESI/MSn profile and content. Nat Prod Commu. 2011;6(1):21–30.

25. Petreska SJ, Stefova M. Evaluation of the ion trap MS performance for quantification of flavonoids and comparison to UV detection. *J Mass Spect.* 2012;47(11): 1395-1406.
26. Petreska SJ, Stefova M. Assay of urinary excretion of polyphenols after ingestion of a cup of mountain tea (*Sideritis scardica*) measured by HPLC-DAD-ESI-MS/MS. *J Agric Food Chem.* 2013;61(44):10488-10497.
27. Faiella L, Dal Piaz F, Bader A, Braca A. Diterpenes and phenolic compounds from *Sideritis pullulans*. *Phytochemistry.* 2014; 106:164-170.
28. Ganzera M, Guggenberger M, Stuppner H, Zidorn C. Altitudinal variation of secondary metabolite profiles in flowering heads of *Matricaria chamomilla* cv. BONA. *Planta Med.* 2008;74(4):453-457.
29. Shu J, Huang G, Mao J. Isolation and identification of flavonoids and phenolic acid in 95% ethanol extraction of Chuju. *Zhongguo Shipin Xuebao.* 2013;13(4):207-213.
30. Wei S, Qu L, Zheng Z-W. Flavonoids in *Mallotus apelta* (Lour.) Muell.-Arg. *Zhongchengyao.* 2012;34(4):691-693.
31. Nikolova M, Asenov A. Surface flavonoid aglycones in newly studied plant species. *Nat Prod Res.* 2006;20(1):103–106.
32. Venditti A, Bianco A, Maggi F, Nicoletti M. Polar constituents composition of endemic *Sideritis italica* (MILL.) GREUTER et BURTER from Central Italy. *Nat Prod Res.* 2013a;27(15):1408-1412.
33. Tomás-Barberan F, Gil MI, Ferreres F, Tomás-Lorente F. Flavonoid p-coumaroylglucosides and 8-hydroxyflavone allosylglucosides in some Labiatae. *Phytochemistry.* 1992;31(9): 3097–3102.
34. McKay DL, Blumberg JB. A review of the bioactivity and potential health benefits of chamomile tea (*Matricaria recutita* L.). *Phytother Res.* 2006;20:519–530.
35. Tadic VM, Jeremic I, Dobric S, Isakovic A, Markovic I, Trajkovic V, et al. Anti-inflammatory, gastroprotective, and cytotoxic effects of *Sideritis scardica* extracts. *Planta Med.* 2012;78(5):415-427.
36. Jeremic I, Tadic V, Isakovic A, Trajkovic V, Markovic I, Redzic Z, et al. The mechanisms of *in vitro* cytotoxicity of mountain tea, *Sideritis scardica*, against the C6 glioma cell line. *Planta Med.* 2013;79(16):1516-1524.
37. Kogiannou DAA, Kalogeropoulos N, Kefalas P, Polissiou MG, Kaliora AC. Herbal infusions; Their phenolic profile, antioxidant and anti-inflammatory effects in HT29 and PC3 cells. *Food Chem Toxicol.* 2013;61:152-159.
38. Brankovic S, Kitic D, Radenkovic M, Veljkovic S, Jankovic T, Savikin K, et al. Spasmolytic activity of the ethanol extract of *Sideritis raeseri* spp. raeseri Boiss. & Heldr. on the isolated rat ileum contractions. *J Med Food.* 2011;14(5):495-498.
39. Petreska J, Stefova M, Ferreres F, Moreno DA, Tomas-Barberan FA, Stefkov G, et al. Potential bioactive phenolics of Macedonian *Sideritis* species used for medicinal "Mountain Tea". *Food Chem.* 2011a;125(1):13-20.
40. Zacharoula I, Linardaki CG, Vasilopoulou CC, Gregoris Iatrou FNL, Marigoula M. Differential antioxidant effects of consuming tea from *Sideritis clandestina* subsp. peloponnesiaca on cerebral regions of adult mice. *J Med Food.* 2011; 14(9):1060-1064.
DOI: 10.1089/jmf.2010.0187
41. Goulas V, Exarchou V, Kanetis L, Gerathanassis IP. Evaluation of the phytochemical content, antioxidant activity and antimicrobial properties of mountain tea (*Sideritis syriaca*) decoction. *J Funct Foods.* 2014;6:248-258.
42. Danesi F, Saha S, Kroon PA, Glibetic M, Konic-Ristic A, D'Antuono LF, et al. Bioactive-rich *Sideritis scardica* (mountain tea) is as potent as Camellia sinensis tea at inducing cellular antioxidant defences and preventing oxidative stress. *J Sci Food Agric.* 2013;93(14):3558-3564.
43. Uriarte-Pueyo I, Calvo MI. Structure-activity relationships of acetylated flavone glycosides from *Galeopsis ladanum* L. (*Lamiaceae*). *Food Chem.* 2010;120(3): 679-683.
44. Petreska J, Stefkov G, Kulevanova S, Alipieva K, Bankova V, Stefova M. Phenolic compounds of mountain tea from the Balkans: LC/DAD/ESI/MSn profile and content. *Nat Prod Commun.* 2011b;6(1): 21-30.
45. Petreska SJ, Stefova M. Assay of urinary excretion of polyphenols after ingestion of a cup of mountain tea (*Sideritis scardica*) measured by HPLC-DAD-ESI-MS/MS. *J Agric Food Chem.* 2013;61(44):10488-10497.

46. Gritsenko EN, Kostyuchenko OI, Fefer IM, Gritsenko EN, Kobzar AY. Iridoids of some genera of the Labiatae family. Mater. S'ezda Farm. B. SSR, 3rd. 1977;164-166, Ed Urvantsev IF; Minsk. Gos. Med. Inst., Minsk, USSR. Russian.
47. Kirmizibekmez H, Ariburnu E, Masullo M, Festa M, Capasso A, Yesilada E, et al., Iridoid, phenylethanoid and flavonoid glycosides from *Sideritis trojana*. Fitoterapia. 2012;83(1):130-136.
48. Alipieva KI, Kostadinova EP, Evstatieva LN, Stefova M, Bankova VS. An iridoid and a flavonoid from *Sideritis lanata* L. Fitoterapia. 2009;80(1):51-53.
49. Assaad AM, Lahloub MF. Iridoid glucosides of *Ajuga iva* (L.) Schreb (*Lamiaceae*). Alex J Pharm Sci. 1988;2(2): 132-5.
50. Koleva II, Handjieva NV, Study of the iridoid glycosides in *Sideritis (Lamiaceae)*. Nauchni Trudove - Vissh Institut po Khranitelna i Vkusova Promishlenost, Plovdiv. 1997;42:75-79. Bulgarian.
51. Charami M-T, Lazari D, Karioti A, Skaltsa H, Hadjipavlou-Litina D, Souleles C. Antioxidant and antiinflammatory activities of *Sideritis perfoliata* subsp. *perfoliata (Lamiaceae)*. Phytother Res. 2008;22(4): 450-454.
52. Ezer N, Akcos Y, Rodriguez B, Abbasoglu U. An iridoid glucoside from *Sideritis libanotica* Labill. subsp. *linearis* (Benth) Bornm., and its antimicrobial activity. Hacettepe Universitesi Eczacilik Fakultesi Dergisi. 1995;15(1):15-21. Turkish.
53. Akcos Y, Ezer N, Ozcelik B, Abbasoglu U. Iridoid glucosides from *Sideritis lycia* Boiss. & Heldr. and its antimicrobial activities. FABAD Farmasotik Bilimler Dergisi. 1998;23(3):99-103.
54. Bianco A, Passacantilli P, Righi G, Nicoletti M. Iridoids in the flora of Italy. Part 10. Iridoid glucosides from *Parentucellia viscosa*. Phytochemistry. 1985;24(8):1843-1845.
55. Venditti A, Serrilli AM, Bianco A. Iridoids from *Bellardia trixago* (L.) All. Nat Prod Res. 2013b;27(15):1413-1416.
56. Olmstead RG, De Pamphilis CW, Wolfe AD, Young ND, Elisons WJ, Reeves PA. Disintegration of the *Scrophulariaceae*. Am J Bot. 2001;88(2):348-361.
57. Ronsted N, Franzzyk H, Molgaard P, Jaroszewski JW, Jensen SR. Chemotaxonomy and evolution of *Plantago* L. Plant Syst Evol. 2003;242(1-4):63-82.
58. Zhang L, Feng L, Jia Q, Xu J, Wang R, Wang Z, et al. Effects of β -glucosidase hydrolyzed products of harpagide and harpagoside on cyclooxygenase-2 (COX-2) *in vitro*. Bioorg Med Chem. 2011; 19:4882–4886.
59. Georgiev M, Pastore S, Lulli D, Alipieva K, Kostyuk V, Potapovich A, et al. *Verbascum xanthophoeniceum*-derived phenylethanoid glycosides are potent inhibitors of inflammatory chemokines in dormant and interferon-gamma-stimulated human keratinocytes. J Ethnopharmacol. 2012;144(3):754-760.
60. Wen B, He R, Li P, Xu Q, Lu Y, Peng B, et al. Pharmacokinetics of 8-O-acetylharpagide and harpagide after oral administration of *Ajuga decumbens* Thunb extract in rats. J Ethnopharmacol. 2013;147(2):503-508.
61. Recio MC, Giner RM, Mánez S, Ríos JL. Structural consideration on the iridoids as anti-inflammatory agents. Planta Med. 1994;60:232–234.

© 2016 Venditti et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/13171>