



Evaluating the Role of Seagrass in Cenozoic CO₂ Variations

Marco Brandano^{1, 2*}, Marco Cuffaro², Giovanni Gaglianone¹, Patrizio Petricca³, Vincenzo Stagno¹ and Guillem Mateu-Vicens^{4, 5}

¹ Dipartimento Scienze della Terra, Università Roma La Sapienza, Rome, Italy, ² Istituto di Geologia Ambientale e Geoingegneria, CNR, Rome, Italy, ³ GFZ German Research Centre for Geosciences, Telegrafenberg, Potsdam, Germany, ⁴ Càtedra Guillem Colom Casasnovas, Universitat de les Illes Balears, Palma de Mallorca, Spain, ⁵ Laboratorio de Zoología, Departament de Biologia, Universitat de les Illes Balears, Palma de Mallorca, Spain

Marine seagrass angiosperms play an important role in carbon sequestration, removing carbon dioxide from the atmosphere and binding it as organic matter. Carbon is stored in the plants themselves, but also in the sediments both in inorganic and organic forms. The inorganic component is represented by carbonates produced by calcareous organisms living as epiphytes on seagrass leaves and rhizomes. In this paper, we find that the rate of seagrass epiphyte production (leaves and rhizomes) averages 400 g m⁻² yr⁻¹, as result of seagrass sampling at seven localities along the Mediterranean coasts, and related laboratory analysis. Seagrasses have appeared in the Late Cretaceous becoming a place of remarkable carbonate production and C sequestration during the whole Cenozoic era. Here, we explore the potential contribution of seagrass as C sink on the atmospheric CO₂ decrease by measuring changes in seagrass extent, which is directly associated with variations in the global coastal length associated with plate tectonics. We claim that global seagrass distribution significantly affected the atmospheric composition, particularly at the Eocene-Oligocene boundary, when the CO₂ concentration fell to 400 ppm, i.e., the approximate value of current atmospheric CO₂.

OPEN ACCESS

Edited by:

Cody Springer Sheik, University of Minnesota Duluth, USA

Reviewed by:

John Senko, University of Akron, USA Alberto Saez, University of Barcelona, Spain

*Correspondence: Marco Brandano marco.brandano@uniroma1.it

Specialty section:

This article was submitted to Microbiological Chemistry and Geomicrobiology, a section of the journal Frontiers in Environmental Science

Received: 13 July 2016 Accepted: 24 October 2016 Published: 11 November 2016

Citation:

Brandano M, Cuffaro M, Gaglianone G, Petricca P, Stagno V and Mateu-Vicens G (2016) Evaluating the Role of Seagrass in Cenozoic CO₂ Variations. Front. Environ. Sci. 4:72. doi: 10.3389/fenvs.2016.00072 Keywords: seagrass, atmospheric CO_2 , cenozoic, plate reconstructions, coast

INTRODUCTION

Seagrasses are marine angiosperms that form extensive submarine meadows in the photic zone along temperate to tropical coastlines worldwide (Short and Wyllie-Echeverria, 1996; Short et al., 2007). They first *(Posidonia cretacea)* appeared during the Late Cretaceous (Brasier, 1975; den Hartog and Polderman, 1975; Eva, 1980; Ivany et al., 1990) in the Tethys Ocean. Since the early Eocene this ecosystem became well established and spread throughout the Tethys (Brasier, 1975; Óosović et al., 2004; Zamagni et al., 2008; Tomás et al., 2016; Tomassetti et al., 2016), the Western Atlantic-Caribbean (Vélez-Juarbe, 2014) and in the Indo-Pacific realm when they colonized the euphotic zone of coastal environments as well as carbonate platforms.

Until the 2000s only few works focus on geological record of seagrass (Brasier, 1975; Eva, 1980; Wanless, 1981; Ivany et al., 1990). This is likely a consequence of the scarcity of fossil remains of seagrasses, due to the low potential of preservation of these plants (Brasier, 1975; Reich et al., 2015 and references therein). On the contrary, in these last years many works have focused on the identification of paleo-seagrasses through the recognition of indirect sedimentological and biological indicators, by comparison with modern seagrass habitats (e.g.,

1

Mateu-Vicens et al., 2008, 2012; Reich et al., 2015; Darroch et al., 2016). These works well evidence the extensive colonization by seagrass in the photic zone of coastal environment and carbonate platforms at global scale (Reich et al., 2015 and reference therein).

Seagrasses have two clearly distinguishable levels, rhizomes and blades. Rhizomes run in sediments beneath the seabed, and have regularly spaced nodes, each bearing roots below and an erect stem or shoot with the blades above. Each level has characteristic associated communities living attached to the plant (Langer, 1988; Mateu-Vicens et al., 2008). A diverse array of biota dwells in seagrass meadows as epiphytic (foraminifers, bryozoans, serpulids and encrusting coralline algae) or infaunal forms (echinoids, mollusks) (**Figures 1A,B**). Many of these organisms secrete calcareous skeletons (Perry and Beavington-Penney, 2005; Brandano et al., 2009; Mateu-Vicens et al., 2010, 2012), consequently seagrasses hosted a significant carbonate production.

The distribution of seagrass species is the result of combined plant sexual reproduction and clonal growth (**Figure 1C**) influenced by dispersal and environmental limitations (Spalding et al., 2003). All seagrass species are capable of asexual reproduction, giving rise to modular units known as ramets, through horizontal growth of the rhizome genetically identical to the parent plant, the genet. The reproductive strategy, which involves clonal growth and production of long-lived, locally dispersed seeds may provide an evolutionary advantage to plants growing in environments subject to temporally unpredictable major disturbances (Rasheed, 2004).

Lastly, seagrasses play an important role in carbon sequestration as they remove carbon dioxide from the atmosphere and bind it as organic matter (Fourgurean et al., 2012). The carbon (C) sequestered in vegetated coastal ecosystems, specifically mangrove forests, seagrass beds and salt marshes, has been termed "blue carbon" (Nellemann et al., 2009). Blue carbon is sequestered over the short term (decennial) in biomass and over longer (millennial) time-scales in sediments (Duarte et al., 2005; Lo Iacono et al., 2008). Carbon is stored above the seabed on plant tissues, underneath the seabed on roots and rhizomes, and in sediments in both organic and inorganic forms, the latter represented by carbonates. Much attention has focused on the quantification of organic C sequestration, and the latest estimates indicate that seagrasses can store 4.2-8.4 Pg C (Fourqurean et al., 2012), and total global C burial by seagrasses may reach 112 Tg C yr⁻¹ (McLeod et al., 2011). However, we point out here that less attention has been paid to C stored by carbonates (inorganic C) produced by epiphytic calcareous organisms that, live on seagrass leaves and rhizomes. This inorganic C, unlike organic C, is locked away..

METHODS

Estimates of Seagrass Carbonate Production

Posidonia oceanica shoots were collected by scuba divers at 5, 11, and 15 mwd (mean water depth) from seagrass meadows in seven locations on the Tyrrhenian coast (Maratea, Ponza, S. Marinella,



FIGURE 1 | (A) Biota dwelling vertical rhizome (*Miniacina miniacea*) and **(B)** leaves as epiphytic forms; **(C)** clonal growth reproductive strategy involves the development of modular units, knowns as ramets, through horizontal growth of the rhizome, genetically identical to the parent plant, the genet; **(D)** Location map of sampling sites for measurement of seagrass carbonate production.

Isola del Giglio, Osalla, Alghero, Crovani see Figure 1D) during spring and autumn 2012 after evaluation of meadow shoot density near the sampling points. The sampling period was chosen taking into account the seagrass growth dependence on seasons (Short et al., 2006). The main morphometric measurements (length, width, and leaf area, leaf length, Leaf Area Index) were performed on 200 leaves from seagrass bundles. The annual production of epiphytic carbonate in the meadows was evaluated independently for the leaves (blades) and its common basal parts (bundle of leaf-sheaths), as well as on the whole leaf bundles. Calcium carbonate content was assessed by geochemical calcimetric analysis on ash from plant parts, previously dried at 105°C for 24 h and weighed, burned in an oven at 550°C for 4 h (LOI₅₅₀ method; (Dean, 1974)), and determination of calcium carbonate content by calcimetric analysis on ash. The epiphytic foliar content of calcium carbonate was converted into epiphytic foliar annual average following Canals and Ballesteros (1997), that is multiplying by 1.88. This process was also extended to the carbonate contribution from basal parts, and then to whole bundles. On whole parts (leaves and basal parts), the annual production of epiphytic carbonate per surface meadow unit of *P. oceanica* was estimated to be about $400 \text{ g m}^{-2} \text{ yr}^{-1}$.

Plate Reconstructions and Coastline Lengths

Absolute plate reconstructions with respect to the Pacific Hotspot Reference frame during the Cenozoic were performed by combining global plate models and finite rotations provided by Seton et al. (2012). The reconstructed positions of continents (Figure S1) were obtained with GPlates software (http://www. gplates.org), and closely coincide with magnetic Chrons, crucial geological events (e.g., the Messinian salinity crisis) and global variation trends of atmospheric CO₂, as reported by Beerling and Royer (2011). The coastline polygons from Seton et al. (2012) were used to compute global coastline lengths at each selected stage. Coastal data were digitalized and simplified due the complexity of several uneven areas (such as Polinesia and Indonesia), underestimating the actual total lengths of the coastlines. Polygon borders presumed to be in contact with seas and oceans were selected and considered as a single boundary according to the reconstruction by Seton et al. (2012). Total coastline lengths and areas of seagrass emplacement were computed at each selected age.

Global CO₂ Estimates

In this work, the Cenozoic CO_2 variations according to Beerling and Royer (2011) are used with one proxy (phytoplankton) to obtain a better comprehension of changes in atmospheric CO_2 concentrations. Our purpose is to quantify the contribution from both inorganic and organic carbon sequestration due to seagrass extent variations during the Cenozoic, and compare them with the reference changes of Beerling and Royer (2011) to evaluate the role of seagrass in CO_2 atmospheric composition. At this aim we consider that,

$$\frac{V_{co_2}}{V_{atm}} = C \tag{1}$$

where V_{CO_2} is the partial volume of the CO₂ in the atmosphere, V_{atm} is the total volume of the atmosphere, and C = 400 ppmv is the current value of the global concentration of carbon dioxide in the atmosphere, expressed in parts per million by volume.

Under the same conditions of temperature T and pressure P, we can write:

$$\frac{PV_{co_2}}{PV_{atm}} = \frac{n_{co_2}RT}{n_{atm}RT'}$$
(2)

where n_{CO_2} and n_{atm} are the CO₂ and the atmosphere number of moles respectively, and *R* is the universal gas constant, so that we have:

$$\frac{V_{co_2}}{V_{atm}} = \frac{M_{co_2}}{m_{co_2}} \frac{m_{atm}}{M_{atm}},\tag{3}$$

where M_{CO_2} is the mass in grams of the carbon dioxide in the atmosphere, $M_{atm} = 5.1480 \times 10^{21}$ g is the total mean mass of the atmosphere (Trenberth and Smith, 2005), $m_{CO2} = 44.00$ g mole⁻¹ and $m_{atm} = 28.97$ g mole⁻¹ are the CO₂ molecular weight and the mean molecular weight of the atmosphere (NASA Earth Fact Sheet, http://nssdc.gsfc.nasa.gov/planetary/factsheet/ earthfact.html), respectively

Using equation (3), we are able to compute the variation of the ratio V_{CO_2}/V_{atm} during the Cenozoic, and evaluate the inorganic and organic carbon sequestration in terms of carbon dioxide mass M_{CO_2} (in grams) in the atmosphere.

The quantity M_{CO_2} depends on the seagrass areal extent, being estimated for the Present using the lower global area of $A_1 = 300,000 \text{ km}^2$ (Charpy-Roubaud and Sournia, 1990), and an upper global area $A_2 = 600,000 \text{ km}^2$ (Duarte et al., 2005). The area depends by the extension of seagrass from the coast, that in turn is under control of the light requirement for photosynthesis Short et al. (2007). The upslope limit of seagrasses is imposed by their necessity of constant immersion in seawater and low hydrodynamic conditions because high energy superficial wave inhibits its growth and development. As seagrass may colonize rocky and sandy substrates in the photic zone (Green and Short, 2003; Mateu-Vicens et al., 2012) the areal extent of seagrass meadows depends by steep gradient of marine seafloor. These requirements suggest that it is not necessary a flat and wide seafloor surface for seagrass meadows colonization, and also steep rocky substrate may host productive seagrasses (Green and Short, 2003). Modern examples of such substrate are represented by the northeastern coast of Philippines or northeastern coast of Haiti in the Caribbean (Green and Short, 2003). On the contrary are well known examples of large continental platforms not colonized by seagrass such as the southwestern Africa or southeastern South America continental platforms.

During the Cenozoic, we expect that seagrass area varied according to plate tectonics. Mantle—reference plate reconstructions with respect to the Pacific hotspot framework during the Cenozoic were performed by combining global plate models and finite rotations provided by Seton et al. (2012). The reconstructed positions of continents (Figure S1) were obtained with GPlates software (http://www.gplates.org), chosen here to

coincide as closely as possible with magnetic chrons, crucial geological events (e.g., the Messinian salinity crisis) and global variation trends of atmospheric CO₂, as reported by Beerling and Royer (2011). The coastline polygons of Seton et al. (2012) are used to compute global coastline lengths *L* at each selected stage in a range of latitude $[-65^{\circ}N, 65^{\circ}N]$ in which seagrasses are presumed to have existed based on paleontological record-IPSIs-, Reich et al. (2015) (see **Figure 1**, Table S1). Offshore distance *d* is assumed constant during the Cenozoic, and is varied to obtain lower and upper global estimates of the seagrass areal extent for the present-day, so that:

$$A_i = L_0 d_i \tag{4}$$

where i = 1, 2, and L_0 is the current total coastline length obtained with coastal polygon model by Seton et al. (2012), at age t = 0 Ma, resulting in $d_1 = 0.889625$ km and $d_2 = 1.779249$ km. Cenozoic coast lengths and seagrass areal extent variations are reported in Table S1.

To compute current inorganic carbon sequestration, laboratory experiments carried out in this study suggest that present-day seagrass coverage produces CaCO₃ carbonate with the rate $a = 400 \text{ g m}^{-2} \text{ yr}^{-1}$, which results in the 44% of CO₂ stored in seagrass carbonates, corresponding to the coefficient b= 176 g m⁻² yr⁻¹. The mass in grams of CO₂ over a year, due to inorganic carbon sequestration, produced by seagrass coverage A_i , with i = 1, 2 is obtained for the Present-day, as it follows:

$$M_{CO_2}^{CaCO_3}(i) = bL_0 d_i \tag{5}$$

Combining Equation (5) with Equation (3), we obtain the carbon dioxide removal (CDR) due to inorganic (CaCO₃) carbon sequestration, during the Cenozoic. At each selected age, results are reported in Table S2 and Table S3, for lower and upper seagrass areal extent A_1 and A_2 , respectively.

To calculate the present-day value of organic carbon sequestration, the estimate procedure of Duarte et al. (2010), Kennedy et al. (2010) and Waycott et al. (2009) were followed. Those authors estimated an organic C sink of $c_1 = 160 \text{ g C m}^{-2} \text{ yr}^{-1}$ and $c_2 = 186 \text{ g C m}^{-2} \text{ yr}^{-1}$ related to the A_1 (300,000 km²) and A_2 (600,000 km²) seagrass areal extent, respectively. The mass in grams of C over a year, due to the organic carbon sequestration, produced by seagrass coverage A_i , with i = 1, 2 is obtained for the present-day, as it follows:

$$M_c^{Org}(i) = c_i L_0 d_i \tag{6}$$

Then, from Rasheed (2004), we know that:

$$M_{\rm CO_2} = M_c \frac{m_{co_2}}{m_c} \tag{7}$$

where $m_{CO2} = 44.00 \text{ g mole}^{-1}$ and $m_C = 12.00 \text{ g mole}^{-1}$ are the CO₂ and C molar mass, respectively.

Combining Equation (7), Equation (6), and Equation (3) we obtain the carbon dioxide removal (CDR) due to organic carbon sequestration during the Cenozoic. At each selected age, results

are reported in Tables S2, S3, for lower and upper seagrass areal extent *A*1 and *A*2, respectively.

Finally, to have an estimate of the global amount of C sink due to both inorganic and organic contributions, we define the cumulative carbon dioxide removal (CDR), as it follows:

$$M_{CO_2}^{cum.}(i) = M_{CO_2}^{CaCO_3}(i) + M_{CO_2}^{Org.}(i),$$
 (8)

where i = 1, 2. Combining equation (8) with equation (3) we obtain the carbon dioxide removal (CDR) due to cumulative effect of carbon sequestration (i.e., inorganic and organic), during the Cenozoic. At each selected age, results are reported in Tables S2, S3, for lower and upper seagrass areal extent A_1 and A_2 , respectively. A third model for the upper seagrass areal extent $A_3 = 100,0000 \text{ km}^2$ with $d_3 = 2.965415 \text{ km}$ can be used to calculate the present-day values of inorganic and organic carbon sequestration taking into account the seagrass loss of 29% worldwide (Waycott et al., 2009), also including the inorganic C sink estimate of $b = 176 \text{ g m}^{-2} \text{ yr}^{-1}$, and the organic C sink estimate of $c_2 = 186 \text{ g C} \text{ m}^{-2} \text{ yr}^{-1}$ respectively. Again, combining Equation (5), Equation (6), Equation (7) with Equation (3), and using Equation (8) we obtain results as in Table S4.

Storage of C by seagrasses may be calculated over 1000-year periods as seagrasses work and live for 1000 of years (Arnaud-Haond et al., 2012). This carbon dioxide removal over 1000 yr (KCDR) is reported in Tables S2–S4, for the inorganic, organic and cumulative contributions of lower and upper seagrass areal extent A_1 , A_2 , and A_3 .

Cenozoic coastal variations, associated with inorganic, organic and cumulative contribution for C sink for Area A_1 and A_3 are reported in Figures S2, S3, respectively.

Linear regressions on CO₂ Cenozoic variations CO₂ and KCDR models (i.e., A_1 , A_2 , and A_3), are reported in Table S5 and Figure S4, and represent the rate of variations, showing a decrease of CO₂ with respect an increase of carbon dioxide removal over 1000 yr.

Partial time interval linear regressions are reported in Table S5 and Figure S5, showing that rates of variations suggest a correspondence at 45% between CO_2 decrease and cumulative KCDR increase during the Cenozoic whereas, if we consider the 41–7 Ma interval, correspondence increases to 66% (Table S5, and Figure S5).

RESULTS AND DISCUSSION

The Mediterranean shelf has classically hosted seagrass sites used to investigate both calcareous production and carbon burial capacity (Canals and Ballesteros, 1997; Duarte et al., 2013). In this paper, carbonate content and (epiphytic) production have been quantified both from seagrass leaves (blades) and from bundles of persistent basal leaf-sheaths resembling "shaving brushes" (Larkum et al., 2006), where the most important seagrass is *Posidonia oceanica* in terms of calcareous epiphyte production (**Figure 1A**). The seagrass biomass exerts a predominant control on the relative abundance of calcareous epiphytes. The most

abundant of these are geniculate (articulated) and non-geniculate (encrusting) coralline algae that, together, comprise up to 50% of epiphytic carbonate. Other significant epiphytes are bryozoans and benthic foraminifers, which contribute approximately in equal amounts (\sim 25% each) to the carbonate production. Calcareous epiphyte abundance peaks at water depths of ~ 10 meters. The rate of epiphyte production (leaves and rhizomes) averages 400 g m⁻² yr⁻¹. Today, seagrass meadows are widespread in many tropical and temperate, shallow marine, and estuarine environments. Data from leaf carbonate production are available from both tropical and temperate environments (Nelsen and Ginsburg, 1986; Bosence, 1989; James et al., 2012), showing that the total epiphytic production for seagrass is within the range of the Tyrrhenian seagrass. Throughout the world, seagrasses are estimated to cover 0.6×10^6 km², equivalent to 10% of the coastal ocean surface, an area comparable to that covered by coral reefs and mangroves (Charpy-Roubaud and Sournia, 1990; Hemminga and Duarte, 2000). Consequently, with an average carbonate production of $400 \text{ g m}^{-2} \text{ yr}^{-1}$, the CO₂ stored in seagrass carbonates may be estimated at up to 105 Tg yr^{-1} (10.5 × 10⁷ t).

Seagrasses have been integral components of shallow marine ecosystems since their appearance in the Late Cretaceous (den Hartog, 1970; Brasier, 1975; Ivany et al., 1990). Because the calcareous skeletons of many seagrass dwellers contribute significantly to the sediment of photic shallow water environments, seagrass meadows have become host of substantial carbonate production and C (organic and inorganic) sequestration since the Cenozoic era.

Reconstructions of atmospheric CO2 concentrations for the Cenozoic have improved greatly in the last few decades (e.g., Beerling and Royer, 2011), although many uncertainties still exist, with CO₂ estimates varying greatly within the same time interval used as a proxy (Royer et al., 2012). However, the review of CO₂ concentration estimates by Beerling and Royer (2011) documented a consistent pattern of CO₂ change. According to the above authors' compilation in the early Paleogene atmospheric CO₂ was about 400 ppm, followed by a first peak to nearly 1900 ppm at 50 Ma and two others above 1200 ppm until the Eocene-Oligocene boundary when a decreasing trend began, with values around 400 ppm throughout the Oligocene. During the Miocene a few data points exceed 400 ppm, but most fall in the range 200–350 ppm. The bulk of the data from the Pliocene are in the range 250-400 ppm, followed by a general decrease during the Pleistocene.

The causes of the reduction in atmospheric CO_2 levels at the Eocene-Oligocene boundary are still debated (Willenbring and von Blanckenburg, 2010). According to Allen and Armstrong (2008), the decrease in atmospheric CO_2 was the result of complementary mechanisms, such as waning pre-collision arc magmatism, increased weathering of silicate, storage of organic carbon in Paratethyan basins, and oceanographic changes producing upwelling.

Changes in the atmospheric CO_2 concentrations, however, may have had severe consequences for marine ecosystems. Elevated atmospheric CO_2 phases in the Earth's history had induced important crisis and extinctions of marine calcifiers (Weissert and Erba, 2004; Erba et al., 2010; Črne et al., 2011; Hönisch et al., 2012). However, with the exception of PETM (Paleocene Eocene Thermal Maximum) event, the Cenozoic interval was not affected by important changes of CaCO₃ saturation (Hönisch et al., 2012).

The role of modern seagrasses as global C sinks was demonstrated and quantified once the total area of Earth covered by seagrass meadows was estimated. Seagrasses grow over a large range of latitudes, from the tropics to close to Arctic regions (N70°, Norway) in coastal waters, with a seafloor illuminated by 1.0–29% of surface irradiance, in water depths of up to -60 m, average -25 m (Green and Short, 2003). On the other hand, the global extent of seagrasses is directly linked to global coastal length. As a consequence, changes in the latter over time may influence global seagrass coverage and its contribution to C storage.

The potential contribution of seagrasses in reducing atmospheric CO_2 during the Cenozoic can be evaluated by measuring changes in seagrass extent, which is directly associated with variations in global coastal length due to plate tectonics. Estimates of coastal length changes are provided here by means of plate tectonic reconstructions. During the Cenozoic the position and shape of the continents were modified by the movement of lithospheric plates with respect to the underlying mantle (**Figure 2**).

Using the rotation vector, plate polygon and coastline data proposed by Seton et al. (2012), we computed global plate reconstructions at several key-time points (Figure S1)



FIGURE 2 | Present-day (0 Ma) and reconstructed (65 Ma) positions of continents using models provided by Seton et al. (2012) (red polygons). Black lines represent digitalized, and simplified data, estimated in a range of latitude [-65°N, 65°N] in which seagrasses are presumed to have existed, and utilized to compute global coastal lengths at several key-time points.

overlapping possible magnetic Chrons, crucial geological events (e.g., the Messinian salinity crisis) and trends of variations in atmospheric CO₂, as reported by Beerling and Royer (2011). Then, with the reconstructed positions of continents, we estimated the total length of the global coastlines in a range of latitude $[-65^{\circ}N, 65^{\circ}N]$ in which seagrasses are presumed to have existed over the entire investigated period of time (see Methods paragraph).

From 65 to 41 Ma we observe a general increase in the coastline length (Figure 3), mostly due to the southward movement of North America and Eurasia over the latitude hypothesized for seagrass growth, at 65°N. The appreciable decrease in coastlines at 34 Ma can be attributed to the evolution of the India-Eurasia continental collision: a relevant increase in coastline lengths between 34 and 28 Ma is observed, probably due to the opening of the Gulf of Aden and the beginning of continental rifting in the Red Sea between the Africa and Arabia plates. Lastly, a steady decrease in length is noted after 14 Ma, mainly due to the continental collision of the Arabia and Eurasia plates, with the exception of a singular increase at 5 Ma, ascribed to the closure of North America and South America. The curves of CaCO₃ carbonate production and organic contribution to carbon dioxide removal (CDR) and their cumulative curve (i.e., CaCO₃ plus organic C) during the Cenozoic are shown in Figure 3 and compared with variations in coastline length (31.0 $\times 10^4$ to 36.7 $\times 10^4$ km). Here, we present 3 models (A₁₋₃) that take into account the dependence of CO₂ sequestration on the seagrass areal extent, being evaluated for the Present-day with a lower global estimate of $A_1 = 300,000 \text{ km}^2$ (29), an upper global estimate of $A_2 = 600,000 \text{ km}^2$ (19) and an overestimation of A_3 = 1,000,000 km². We report the result of calculation for the A_2 model, for A_1 and A_2 see Figures 3, 4.

The contribution of CaCO₃ to CO₂ sequestration in the A₂ was found to vary from 12.45 × 10⁻³ to 14.72 × 10⁻³ ppm yr⁻¹. Higher values were observed for the organic component, which ranged from 48.26 × 10⁻³ to 57.52 × 10⁻³ ppm yr⁻¹. The cumulative contribution of the two components was in the range 60.71×10^{-3} to 71.79×10^{-3} ppm yr⁻¹.

Storage of C by seagrasses may be calculated over 1000year periods, because seagrasses work and live for 1000 of years (Arnaud-Haond et al., 2012). This cumulative carbon dioxide removal over 1000 years (KCDR) (80.43-95.11 ppm) throughout the Cenozoic is shown in **Figure 4** and compared with the value of atmospheric CO₂ reported by Beerling and Royer (2011), including values for the key time used to calculate plate reconstructions and coastline lengths. **Figure 3A** shows the general match between increased C sequestration and increased coastal length. Linear regressions on both curves, i.e., cumulative KCDR and atmospheric CO₂, show a general increase in seagrass contributions with respect to a global decrease in atmospheric CO₂ during the Cenozoic (Figure S4).

Nowadays, the role of the oceans as carbon sinks is widely documented (Duarte et al., 2005; McLeod et al., 2011; Fourqurean et al., 2012; Karl et al., 2012). Phytoplankton photosynthesis is one of the mechanism leading to C sequestration through the deposition and burial of the particulate organic material at the ocean floor formed by phytoplankton (Raven and Falkowski,



FIGURE 3 | (A) Cenozoic cumulative carbon dioxide removal (CDR) for model A₂ (red line, open squares, compared with variations of coastline length (black line open triangles). (B) Cenozoic inorganic (CaCO₃) carbon dioxide removal (CDR) for model A₂. (C) Cenozoic organic carbon dioxide removal (CDR) for model A₂. Dashed pattern represents the Eocene-Oligocene time interval. Data are reported in Table S3.

1999). However, the carbon burial capacity of coastal vegetated habitats (seagrass, mangroves and salt marshes) is 180 times greater than the average burial rate in the open ocean (Duarte et al., 2005), reaching 233 Tg C yr⁻¹ (McLeod et al., 2011), values comparable with the C burial in lakes and peatlands that is about 300 Tg C yr⁻¹ (Dean and Gorham, 1998). Consequently, it is out of question that blue carbon sinks play a key role in the oceanic carbon cycle (Duarte et al., 2005; McLeod et al., 2011; Fourqurean et al., 2012). On the contrary, it is more difficult to prove the role of blue carbon and, more in general, of the ocean as C sink in the fossil record. The obtained data for seagrass are in agreement with the present day role of ocean as natural sink. The obtained values of carbon dioxide sequestration are in the order of atmospheric CO₂ fluctuations during glacial cycles (Kohfeld et al., 2005).

By looking at the specific time-points and analyzing each rate of the two curves compared (i.e., cumulative KCDR increase and CO₂ decrease, Figure S5) with linear regressions for each time interval, the correspondence decreases to 45%, probably due to the lack of accuracy of CO₂ data from 65 to 41–34 Ma. In addition, after the Late Miocene (5 Ma), the lack of correspondence may be due to the development of the Arctic ice cap, marked by decreasing temperature and consequent decreasing atmospheric CO₂. The resulting increase of CO₂ solubility in ocean seawater, together with the



Mediterranean Messinian crisis, probably reduced the seagrass extent.

It should also be noted that the Red Sea syn-rift Miocene evaporatic event and the Mediterranean Messinian salinity crisis were included in our computation of coastlines, although a decreased seagrass productivity is expected in these areas during those specific periods. This means that 4000 km² of seagrass production in the Red Sea should be neglected in our record, almost from 28 to 5 Ma. In addition, at 5 Ma, we have an over-estimated length of coastlines producing seagrasses in the Mediterranean, so that non-accurate comparisons can be made for the 7-5 and 5-1 time intervals. Focusing on those keytime intervals, for which CO2 data are well constrained (e.g., from 41 to 7 Ma) the global correspondence rises up to 66% except for the time ranges 23-20 and 20-14 Ma. The most evident shift occurs with the Eocene-Oligocene CO2 decline (Figure 4). Both curves, i.e., increasing cumulative KCDR and decreasing CO₂ (Figure 4 and Figure S5), plot on higher rates of the considered time intervals, estimated with linear regressions to be 5.32 ppm Ma⁻¹ and -87.00 ppm Ma⁻¹, respectively (Figure S5). In conclusion, we provide an initial estimate of the contribution of seagrass meadows as C sink since their occurrence at the end of the Cretaceous. We admit that this result might be underestimated due to the simplified models used for coastal length computations, and it would need of more specific regional tectonic investigations and ecological constraints associated with eustatic processes in order to improve the accuracy of seagrass involvement to CO₂ decline. However, we are confident that this study represents a valid quantitative and global estimate of C sequestration by seagrass in the geological record since Cretaceous time, to date missing in literature.

AUTHOR CONTRIBUTIONS

MB, MC, VS, and GM developed the ideas and the methods for this study. GG carried out sampling and laboratory analyses. MC and PP carried out plate tectonic reconstructions and coastline length computations. MB and MC wrote jointly the paper. All the authors discussed extensively the results and the interpretations

FUNDING

Funding by Sapienza University of Rome and IGAG-CNR.

ACKNOWLEDGMENTS

Discussions with Marco Ligi and Fabio Trippetta were appreciated. Many of the figures were made with the Generic Mapping Tools of Wessel and Smith (1995) (http://gmt.soest.hawaii.edu/). Criticisms by two reviewers and A. Editor Cody Sheik greatly improved the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: http://journal.frontiersin.org/article/10.3389/fenvs. 2016.00072/full#supplementary-material

REFERENCES

- Allen, M. B., and Armstrong, H. A. (2008). Arabia-Eurasia collision and the forcing of mid Cenozoic global cooling. *Paleoecol. Palaeogeogr. Palaeocl.* 265, 52–58. doi: 10.1016/j.palaeo.2008.04.021
- Arnaud-Haond, S., Duarte, C. M., Diaz-Almela, E., Marbà, N., Sintes, T., and Serrão, E. A. (2012). Implications of extreme life span in clonal organisms: millenary clones in meadows of the Threatened Seagrass: *Posidonia oceanica. PLoS ONE* 7:e30454. doi: 10.1371/journal.pone.00 30454
- Beerling, D. J., and Royer, D. L. (2011). Convergent cenozoic CO₂ history. Nat. Geosci. 4, 418–420. doi: 10.1038/ngeo1186
- Bosence, D. (1989). Biogenic carbonate production in florida bay. *Bull. Mar. Sci.* 44, 419–433.
- Brandano, M., Frezza, V., Tomassetti, L., Pedley, M., and Matteucci, R. (2009). Facies analysis and palaeoenvironmental interpretation of the Late Oligocene Attard Member (Lower Coralline Limestone Formation), Malta. *Sedimentology* 56, 1138–1158. doi: 10.1111/j.1365-3091.2008.01023.x
- Brasier, M. D. (1975). An outline history of seagrass communities. *Palaeontology* 18, 681–702.
- Canals, M., and Ballesteros, E. (1997). Production of carbonate particles by phytobenthic communities on the Mallorca-Menorca shelf, northwestern mediterranean sea. *Deep Sea Res. II*, 44, 611–629. doi: 10.1016/s0967-0645(96)00095-1
- Charpy-Roubaud, C., and Sournia, A. (1990). "The comparative estimation of phytoplanktonic, microphytobenthic and macrophytobenthic primary production in the oceans," in *Microalgal Growth: Inputs and Losses, Practical Approaches: Marine Microbial Food Webs*, eds S. Y. Maestrini and F, Rassoulzadegan (Rome: Sapienza Press), 31–57.
- Ćosović, V., Drobne, K., and Moro, A. (2004). Paleoenvironmental model for Eocene foraminiferal limestones of the Adriatic carbonate platform (Istrian Peninsula). *Facies* 50, 61–75. doi: 10.1007/s10347-004-0006-9
- Črne, A. E., Weissert, H., Goričan, Š., and Bernasconi, S. M. (2011). A biocalcification crisis at the Triassic-Jurassic boundary recorded in the Budva Basin (Dinarides, Montenegro). *Geol. Soc. Am. Bull.* 123, 40–50. doi: 10.1130/B30157.1
- Darroch, S. A., Locatelli, E. R., McCoy, V. E., Clark, E. G., Anderson, R. P., Tarhan, L. G., et al. (2016). Taphonomic disparity in foraminifera as a paleo-indicator for seagrass. *Palaios* 31, 242–258. doi: 10.2110/palo.2015.046
- Dean, W. E. (1974). Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: comparison with other methods. J. Sed. Petrol. 44, 242–248.
- Dean, W. E., and Gorham, E., (1998). Magnitude and significance of carbon burial in lakes, reservoirs, and peatlands. *Geology* 26, 535–538.
- den Hartog, C. (1970). The sea-grasses of the World. North-Holland Publ. *Co: Verh kon Ned Akad Wet Afd Natuurkundem* 59, 275.
- den Hartog, C., and Polderman, P. J. G. (1975). Changes in the seagrass populations of the Dutch Wadden Zee. *Aquat. Bot.* 1, 141–147.
- Duarte, C. M., Kennedy, H., Marbà, N., and Hendriks, I. (2013). Assessing the capacity of seagrass meadows for carbon burial: current limitations and future strategies. *Ocean Costal Manage* 83, 32–38. doi: 10.1016/j.ocecoaman.2011.09.001
- Duarte, C. M., Marbà, N., Gacia, E., Fourqurean, J. W., Beggins, J., Barròn, C., et al. (2010). Seagrass community metabolism: assessing the carbon sink capacity of seagrass meadows. *Glob. Biogeochem. Cycles* 24:GB4032. doi: 10.1029/2010GB0 03793
- Duarte, C. M., Middelburg, J. J., and Caraco, N. (2005). Major role of marine vegetation on the oceanic carbon cycle. *Biogeoscences* 2, 1–8. doi: 10.5194/bg-2-1-2005
- Erba, E., Bottini, C., Weissert, H., and Keller, C. E. (2010). Calcareous nannoplankton response to surface water acidification around Oceanic Anoxic Event 1a. *Science* 329, 428–432. doi: 10.1126/science.1188886
- Eva, A. N. (1980). Pre-miocene seagrass communities in the Caribbean. *Palaeontology* 23, 231–236.
- Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., et al. (2012). Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.* 5, 505–509. doi: 10.1038/ngeo1477

- Green, E. P., and Short, F. T. (2003). World Atlas of Seagrasses. Prepared by the UIMEP World Conservation Monitoring Centre. Berkeley, CA: University of California Press.
- Hemminga, M., and Duarte, C. M. (2000). Seagrass Ecology. Cambridge: Cambridge University Press.
- Hönisch, B., Ridgwell, A., Schmidt, D. N., Thomas, E., Gibbs, S. J., Sluijs, A., et al. (2012). The geological record of ocean acidification. *Science* 335, 1058–1063. doi: 10.1126/science.1208277
- Ivany, L. C., Portell, R. W., and Jones, D. S. (1990). Animal-plant relationships and paleobiogeography of an Eocene seagrass community from Florida. *Palaios* 5, 244–258. doi: 10.2307/3514943
- James, N. P., Bone, Y., Brown, K. M., and Cheshire, A. (2012). "Calcareous epiphyte production in cool-water carbonate seagrass depositional environments; Southern Australia," in *Perspectives in Carbonate Geology: A Tribute to the Career of Robert Nathan Ginsburg* (London: Special Publication 41 of the IAS), 123–148.
- Karl, D. M., Church, M. J., Dore, J. E., Letelier, R. M., and Mahaffey, C., (2012). Predictable and efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen fixation. *Proc. Natl. Acad. Sci. U.S.A.* 109, 1842–1849. doi: 10.1073/pnas.1120312109
- Kennedy, H., Beggins, J., Duarte, C. M., Fourqurean, J. W., Holmer, M., Marbà, N., et al. (2010). Seagrass sediments as a global carbon sink: isotopic constraints. *Glob. Biogeochem. Cycles* 24:GB4026. doi: 10.1029/2010GB 003848
- Kohfeld, K. E., Le Quéré, C., Harrison, S. P., and Anderson, R. F., (2005). Role of marine biology in glacial-interglacial CO₂ cycles. *Science* 308, 74–78. doi: 10.1126/science.1105375
- Langer, M. R. (1988). Recent epiphytic foraminifera from Vulcano (Mediterranean Sea). Rev. Paléobiol. BENTHOS'86 Special 2, 827–832.
- Larkum, A. W. D., Orth, R. J., and Duarte, C. M. (2006). Seagrasses: Biology, Ecology and Conservation. Dordrecht: Springer, 691.
- Lo Iacono, C., Mateo, M. A., Gràcia, E., Guasch, C. R., Serrano, L., and Dañobeitia, J. (2008). Very high-resolution seismo-acoustic imaging of seagrass meadows (Mediterranean Sea): implications for carbon sink estimates. *Geophys. Res. Lett.* 35:L18601. doi: 10.1029/2008GL034773
- Mateu-Vicens, G., Box, A., Deudero, S., and Rodríguez, B. (2010). Comparative analysis of epiphytic foraminifera in sediments colonized by seagrass Posidonia oceanica and invasive macroalgae *Caulerpa* spp. J. Foramin. Res. 40, 134–147. doi: 10.2113/gsjfr.40.2.134
- Mateu-Vicens, G., Brandano, M., Gaglianone, G., and Baldassarre, A. (2012). Seagrass-meadow sedimentary facies in a mixed siliciclasticcarbonate temperate system in the Tyrrhenian Sea (Pontinian Islands, Western Mediterranean). J. Sediment Res. 82, 451–463. doi: 10.2110/jsr. 2012.42
- Mateu-Vicens, M., Hallock, P., and Brandano, M. (2008). A depositional model and paleoecological reconstruction of the lower Tortonian distally steepened ramp of Menorca (Balearic Islands, Spain). *Palaios* 23, 465–481. doi: 10.2110/palo.2007.p07-061r
- McLeod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., et al. (2011). A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front. Ecol. Environ.* 9, 552–560. doi: 10.1890/110004
- Nellemann, C., Corcoran, E., Duarte, C. M., Valdés, L., De Young, C., Fonseca, L., et al. (2009). *Blue Carbon. A Rapid Response Assessment*. Arendal: GRID-Arendal: United Nations Environment Programme.
- Nelsen, J. E., and Ginsburg, R. N. (1986). Calcium carbonate production by epibionts on thalassia in florida bay. J. Sediment. Petrol. 56, 622–628.
- Perry, C. T., and Beavington-Penney, S. J. (2005). Epiphytic calcium carbonate production and facies development within sub-tropical seagrass beds, Inhaca Island, Mozambique. *Sediment Geol.* 174, 161–176. doi: 10.1016/j.sedgeo.2004.12.003
- Rasheed, M. A. (2004). Recovery and succession in a multi-species tropical seagrass meadow following experimental disturbance: the role of sexual and asexual reproduction. *J. Exp. Mar. Biol. Ecol.* 310, 13–45. doi: 10.1016/j.jembe.2004.03.022
- Raven, J. A., and Falkowski, P. G. (1999). Oceanic sinks for atmospheric CO₂. *Plant Cell Environ*. 22, 741–755.

- Reich, S., Di Martino, E., Todd, J. A., Wesselingh, F. P., and Renema, W. (2015). Indirect paleo-seagrass indicators (IPSIs): a review. *Earth Sci. Rev.* 143, 161–186. doi: 10.1016/j.earscirev.2015.01.009
- Royer, D. L., Pagani, M., and Beerling, D. J. (2012). Geobiological constraints on Earth system sensitivity to CO₂ during the Cretaceous and Cenozoic. *Geobiology* 10, 298–310. doi: 10.1111/j.1472-4669.2012. 00320.x
- Seton, M., Müller, R. D., Zahirovic, S., Gaina, D., Torsvik, T., Shepard, G., et al. (2012). Global continental and ocean basin reconstructions since 200 Ma. *Earth Sci. Rev.* 113, 212–270. doi: 10.1016/j.earscirev.2012.03.002
- Short, F. T., Carruthers, T., Dennison, W., and Waycott, M. (2007). Global seagrass distribution and diversity: a bioregional model. J. Exp. Mar. Biol. Ecol. 350, 3–20. doi: 10.1016/j.jembe.2007.06.012
- Short, F. T., Koch, E., Creed, J. C., Magalhaes, K. M., Fernandez, E., and Gaeckle, J. L. (2006). SeagrassNet monitoring across the Americas: case studies of seagrass decline. *Mar. Ecol.* 27, 277–289. doi: 10.1111/j.1439-0485.2006. 00095.x
- Short, F. T., and Wyllie-Echeverria, S. (1996). Natural and human-induced disturbances of seagrass. *Environ. Conserv.* 23, 17–27.
- Spalding, M., Taylor, M., Ravilious, C., Short, F., and Green, E. (2003). "Global overview: the distribution and status of seagrasses," in *World Atlas of Seagrasses*, eds E. P Green and F. T. Short (Berkeley, CA: University of California Press), 5–26.
- Tomassetti, L., Benedetti, A., and Brandano, M. (2016). Middle Eocene seagrass facies from Apennine carbonate platforms (Italy). Sediment. Geol. 335, 136–149. doi: 10.1016/j.sedgeo.2016. 02.002
- Tomás, S., Frijia, G., Bömelburg, E., Zamagni, J., Perrin, C., and Mutti, M. (2016). Evidence for seagrass meadows and their response to paleoenvironmental changes in the early Eocene (Jafnayn Formation, Wadi Bani Khalid, N Oman). Sediment. Geol. 341, 189–202. doi: 10.1016/j.sedgeo.2016. 05.016
- Trenberth, K. E., and Smith, L. (2005). The mass of the atmosphere: a constraint on global analyses. J. Clim. 18, 864–875. doi: 10.1175/jcli-3299.1

- Vélez-Juarbe, J. (2014). Ghost of seagrasses past: using sirenians as a proxy for historical distribution of seagrasses. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 400, 41–49.
- Wanless, H. R. (1981). Fining-upwards sedimentary sequences generated in seagrass beds. J. Sediment. Res. 51, 445–454.
- Waycott, M., Duarte, C. M., Carruthers, T. J. B., Orth, R. J., Denninson, W. C., Olyarnik, S., et al. (2009). Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. Natl. Acad. Sci. U.S.A.* 106, 12377–12381. doi: 10.1073/pnas.0905620106
- Weissert, H., and Erba, E. (2004). Volcanism, CO₂ and palaeoclimate: a Late Jurassic–Early Cretaceous carbon and oxygen isotope record. J. Geol. Soc. 161, 695–702.
- Wessel, P., and Smith, W. H. (1995). The Generic Mapping Tools, GMT, Version 3: Technical Reference and Cookbook. Manoa: School of Ocean and Earth Science and Technology, University of Hawaii at Manoa.
- Willenbring, J. K., and von Blanckenburg, F. (2010). Long-term stability of global erosion rates and weathering during late-Cenozoic cooling. *Nature* 465, 211–214. doi: 10.1038/nature09044
- Zamagni, J., Mutti, M., and Košir, A. (2008). Evolution of shallow benthic communities during the late Palaeocene–earliest Eocene transition in the Northern Tethys (SW Slovenia). *Facies* 54, 25–43. doi: 10.1007/s10347-007-0123-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Brandano, Cuffaro, Gaglianone, Petricca, Stagno and Mateu-Vicens. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.