

## New insight on the photofragmentation of $\text{CH}_2\text{I}_2$

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 112064

(<http://iopscience.iop.org/1742-6596/635/11/112064>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

### Download details:

IP Address: 150.146.205.185

This content was downloaded on 24/11/2016 at 09:29

Please note that [terms and conditions apply](#).

You may also be interested in:

[Selectivity in the photofragmentation of halo-pyrimidines](#)

P Bolognesi, A Kettunen, A Cartoni et al.

[Photofragmentation of halogenated pyrimidine molecules in the VUV range](#)

M C Castrovilli, P Bolognesi, D Catone et al.

[Photofragmentation of colloidal solutions of gold nanoparticles under femtosecond laser pulses in IR and visible ranges](#)

P A Danilov, D A Zayarnyi, A A Ionin et al.

[Photofragmentation and vibrational relaxation of size-selected clusters ions : Non-adiabatic molecular dynamics simulations](#)

J Douady, B Gervais, E Jacquet et al.

[Photofragmentation of a DNA nucleoside thymidine; valence- vs. core ionization](#)

E Itälä, K Kooser, T Hägerth et al.

[XUV photofragmentation of small water cluster cations at FLASH](#)

C Domesle, L Lammich, B Jordon-Thaden et al.

## New insight on the photofragmentation of CH<sub>2</sub>I<sub>2</sub>

A. R. Casavola\*, A. Cartoni<sup>†1</sup>, P. Bolognesi <sup>\*2</sup>, S. Borocci<sup>#</sup>, L. Avaldi\*

\*Istituto di Struttura della Materia, Area della Ricerca Roma 1, 00016, Monterotondo Scalo, Italy


<sup>†</sup> Dip. di Chimica, Università di Roma Sapienza, p.le Aldo Moro 5, 00185 Roma, Italy

<sup>#</sup>Dip. per l' Innovazione nei sistemi Biologici, Agroalimentari e Forestali (DIBAF), Università della Tuscia, L.go dell'Università, s.n.c., 01100 Viterbo, Italy

**Synopsis** An experimental and theoretical investigation of the VUV photofragmentation of CH<sub>2</sub>I<sub>2</sub> molecule is reported. The work focuses on the I-loss channel. The experimental results demonstrate that at energies lower than 12 eV diiodomethane radical cation dissociates only into the CH<sub>2</sub>I<sup>+</sup> ion and an I atom while the theoretical calculations show that isomerisation of [CH<sub>2</sub>I<sub>2</sub>]<sup>+</sup> into iso-diiodomethane [CH<sub>2</sub>I-I]<sup>+</sup> intermediates can occur before the loss of iodine atom.

Diiodomethane is a halomethane widely studied for its role played in the terrestrial atmosphere as well as in solution chemistry. In the environment CH<sub>2</sub>I<sub>2</sub> is a source of iodine atom which affects the ozone chemistry and contributes to the formation of the ultrafine marine aerosol particles, while in photoreaction chemistry this molecule is used for cyclopropanation of alkene. Different studies [1] have demonstrated that in condensed phase diiodomethane molecule under photon irradiation isomerizes into iso-diiodomethane and becomes a source of “CH<sub>2</sub>” species. Moreover in the recent literature it has been reported that in the gas phase halomethane can isomerize into isohalomethane before losing the halogen molecule [2]. In a previous study [3] we have investigated the photofragmentation of dihalomethanes (CH<sub>2</sub>X<sub>2</sub>, X=F, Cl, Br, I) and in particular we found that diiodomethane generates different fragment ions and the I-loss channel is the one opening at the lowest energy. In this work we report an experimental and theoretical investigation of VUV photofragmentation of CH<sub>2</sub>I<sub>2</sub> molecule with the main focus on I-loss channel, which occurs from the lower electronic ionic states as showed in our PEPICO experiments.

Theoretical calculations have been performed at the MP2/CCSD(T) level of theory to obtain the structures and the energy of each stationary point found for the [CH<sub>2</sub>I<sub>2</sub>]<sup>+</sup> ion. The results show that the isomerization of diiodomethane radical cation [CH<sub>2</sub>I<sub>2</sub>]<sup>+</sup> into iso-diiodomethane intermediates [CH<sub>2</sub>I-I]<sup>+</sup> is an energetically competing channel with respect to the direct C-I bond breaking (Figure 1).



**Figure 1.** Schematic representation of energetic paths to I atom and CH<sub>2</sub>I<sup>+</sup> ion from [CH<sub>2</sub>I<sub>2</sub>]<sup>+</sup> through [CH<sub>2</sub>I-I]<sup>+</sup> intermediates (black pattern on the right).

### Acknowledgment

This work is supported by the MIUR FIRB RBFR10SQZI project 2010.

### References

- [1] Phillips, D. L.; Fang, W.-H.; Zheng, X. J. *Am. Chem. Soc.* 2001, 123, 4197-4203.
- [2] Kalume, A.; George, L.; Reid, S. A. *J. Phys. Chem. Lett.* 2010, 1, 3090-3095.
- [3] Cartoni, A.; Bolognesi, P.; Fainelli, E.; Avaldi L. *J. Chem. Phys.* 2014, 140, 184307-1--13

<sup>1</sup>E-mail: [antonella.cartoni@uniroma1.it](mailto:antonella.cartoni@uniroma1.it)

<sup>2</sup>E-mail: [paola.bolognesi@cnr.it](mailto:paola.bolognesi@cnr.it)

