

# Characterization of the transcription factor encoding gene, *KIADR1*: metabolic role in *Kluyveromyces lactis* and expression in *Saccharomyces cerevisiae*

Silvia Cardarelli,<sup>1</sup>† Sirio D'Amici,<sup>1</sup>† Paola Tassone,<sup>1</sup> Angela Tramonti,<sup>2</sup> Daniela Uccelletti,<sup>1</sup> Patrizia Mancini<sup>3</sup> and Michele Saliola<sup>1</sup>

## Correspondence

Michele Saliola  
michele.saliola@uniroma1.it

<sup>1</sup>Department of Biology and Biotechnology 'C. Darwin', Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy

<sup>2</sup>CNR Department of Biochemical Sciences 'Rossi Fanelli', Istituto di Biologia e Patologia Molecolari, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy

<sup>3</sup>Department of Experimental Medicine, Sapienza University of Rome, Viale Regina Elena 324, 00161 Rome, Italy

In *Saccharomyces cerevisiae*, Adr1 is a zinc-finger transcription factor involved in the transcriptional activation of *ADH2*. Deletion of *KIADR1*, its putative ortholog in *Kluyveromyces lactis*, led to reduced growth in glycerol, oleate and yeast extract-peptone medium suggesting, as in *S. cerevisiae*, its requirement for glycerol, fatty acid and nitrogen utilization. Moreover, growth comparison on yeast extract and peptone plates showed in *K. lactis* a KIAdr1-dependent growth trait not present in *S. cerevisiae*, indicating different metabolic roles of the two factors in their environmental niches. *KIADR1* is required for growth under respiratory and fermentative conditions like *KIADH*, alcohol dehydrogenase genes necessary for metabolic adaptation during the growth transition. Using in-gel native alcohol dehydrogenase assay, we showed that this factor affected the Adh pattern by altering the balance between these activities. Since the activity most affected by KIAdr1 is KIAdh3, a deletion analysis of the *KIADH3* promoter allowed the isolation of a DNA fragment through which KIAdr1 modulated its expression. The expression of the *KIADR1-GFP* gene allowed the intracellular localization of the factor in *K. lactis* and *S. cerevisiae*, suggesting in the two yeasts a common mechanism of KIAdr1 translocation under fermentative and respiratory conditions. Finally, the chimeric *KI/ScADR1* gene encoding the zinc-finger domains of KIAdr1 fused to the transactivating domains of the *S. cerevisiae* factor activated in *Scadr1Δ* the transcription of *ADH2* in a ScAdr1-dependent fashion.

Received 3 August 2016

Accepted 14 September 2016

## INTRODUCTION

Four major alcohol dehydrogenase (*ADH*) genes are present in the yeast *Saccharomyces cerevisiae*. *ADH1* encodes the fermentative enzyme involved in the production of ethanol (Lutstorf & Megnet, 1968; Bennetzen & Hall, 1982). *ADH2* codes for a glucose-repressible enzyme involved in the oxidation of the ethanol produced during fermentation (Ciriacy, 1975, 1979; Denis *et al.*, 1981). *ADH3* codes for a

mitochondrial enzyme, a component of the ethanol/acetaldehyde shuttle (Young & Paquin, 1985; Bakker *et al.*, 2000), while *ADH5* codes for a minor activity whose physiological role has not yet been defined (Dickinson *et al.*, 2003).

Genetic analysis of the *ADH2* locus led to the identification and isolation of *ADR1* (Ciriacy, 1979; Denis & Young, 1983), a gene coding for a carbon source-responsive zinc-finger transcription factor, the major regulatory protein involved in the transcriptional activation of *ADH2*. This factor is also required for the induction of genes that are regulated by glucose repression (Young *et al.*, 2003), for the biogenesis of peroxisomes and for ethanol, glycerol and fatty acid utilization (Simon *et al.*, 1991; Young *et al.*, 2003; Tachibana *et al.*, 2005).

†These authors contributed equally to this work.

Abbreviations: ADH, alcohol dehydrogenase;  $\beta$ -gal,  $\beta$ -galactosidase; RE, repression ethanol; YE, yeast extract; YNB, yeast nitrogen base; YP, yeast extract-peptone.

*S. cerevisiae* is a Crabtree-positive yeast in which respiration is completely repressed by glucose even under aerobic conditions, growth being supported solely by fermentation. In contrast, in the Crabtree-negative *Kluyveromyces lactis*, both respiratory and fermentative pathways coexist during growth on glucose (De Deken, 1966), although respiration appears dispensable since antimycin A does not inhibit growth on glucose (Rag<sup>+</sup> phenotype) (Goffrini *et al.*, 1989; Wésolowski-Louvel *et al.*, 1992).

In *K. lactis*, four *ADH* genes have also been identified (*ADH1–ADH4*) (Saliola *et al.*, 1990, 1991; Shain *et al.*, 1992). *KIADH1* and *KIADH2* encode two enzymes preferentially expressed on fermentative carbon sources and *KIADH3* and *KIADH4* code for two mitochondrial enzymes, which are probably components of the ethanol/acetaldehyde shuttle (Saliola & Falcone, 1995; Bakker *et al.*, 2000; Overkamp *et al.*, 2002; Saliola *et al.*, 2008). The latter two genes are regulated at the transcriptional level by carbon source-dependent growth conditions and their expression is mutually exclusive (Saliola & Falcone, 1995). In fact, *KIADH4* is specifically induced by ethanol added to the medium or produced by fermentation. On the other hand, this compound specifically represses the expression of *KIADH3*, which, differently from *KIADH4*, is highly expressed on all respiratory carbon sources with the exception of ethanol (Saliola & Falcone, 1995). The corresponding activities have been identified by genetic analysis in native PAGE by an ADH-specific staining assay and characterized at the biochemical level (Lutstorf & Megnet, 1968; Mazzoni *et al.*, 1992; Saliola & Falcone, 1995; Bozzi *et al.*, 1997). Although a deletion analysis of the *KIADH4* and *KIADH3* promoters led to the identification of regulatory elements, no transcription factors directly involved in their activation have yet been found (Saliola *et al.*, 1999, 2007; Mazzoni *et al.*, 2000). The identification of a putative ortholog of *ScADR1* in the genome of *K. lactis* (KLLA0F13046g) (Dujon *et al.*, 2004; Bussereau *et al.*, 2006) led us to investigate its role in the regulation of *ADH* genes and in the metabolism of this yeast.

*KIADR1* is required for growth in glycerol, oleate and yeast extract-peptone (YP) medium. Measurements of glycerol, ethanol and oxygen reoxidation rate showed that *KIADR1* regulates both respiratory and fermentative metabolisms displaying an Adh pattern with an altered balance between the four

different Adh activities. Since the activity mostly affected by *KIADR1* was *KIADH3*, a deletion analysis of the *KIADH3* promoter allowed the isolation of a DNA fragment through which *KIADR1* modulates the expression of *KIADH3* and, indirectly, that of *KIADH4*.

In addition, the *KIADR1-GFP* gene allowed the intracellular localization of the fusion protein under different growth conditions in both yeasts.

Finally, a chimeric *K. lactis/S. cerevisiae* *ADR1* containing the zinc-finger domains of *K. lactis* activated in *Scadr1Δ* the transcription of *ADH2* under de-repressing conditions like the endogenous factor.

## METHODS

**Strains, media and culture conditions.** The strains used in this work are reported in Table 1.

Cultures were grown under shaking conditions at 28 °C in YP [1 % Difco yeast extract (YE), 2 % Difco Bacto-peptone] or in minimal yeast nitrogen base (YNB) medium (6.7 g l<sup>-1</sup> Difco YNB) supplemented by different carbon sources at concentrations specified below, while *S. cerevisiae* cultures were also supplemented by Bacto Casamino Acids (Difco) at a concentration of 2.5 mg ml<sup>-1</sup>. Oleate medium contained 0.1 % oleic acid, 0.02 % Tween 40, 0.1 % YE and 0.67 % YNB. Ergosterol was added to YPD plated at 20 µg ml<sup>-1</sup>. Antimycin A was supplemented at a concentration of 5 µM in YP plates containing 5 % glucose. Geneticin (G418) concentration in the medium was 100 µg ml<sup>-1</sup> (*K. lactis*) or 200 µg ml<sup>-1</sup> (*S. cerevisiae*). Minimal medium was supplemented with auxotrophic requirements at the final concentration of 10 µg ml<sup>-1</sup>. *Escherichia coli* strain DH5α was used for the propagation of plasmid DNA. Cultures were grown at 37 °C on LB medium (0.5 % YE, 1 % Difco tryptone, 0.5 % NaCl) supplemented with 100 µg ml<sup>-1</sup> ampicillin.

**In-gel ADH staining activity and β-galactosidase assay.** Preparation of cellular extracts, clear native PAGE, electrophoresis conditions and ADH staining assay were carried out as previously described (Saliola & Falcone, 1995).

The construction of the *lacZ/KIADH3* fusion vectors used in this study for the deletion analysis of the *KIADH3* promoter (pCG1, pCG17, pCG13, pCG11, pCG7, pCG12, pCG29, pCG30, pCG31 and pCG32, respectively) was as previously described (Saliola *et al.*, 2007). The vector pCG33 was obtained from pCG32 following *SacI* digestion and religation. This plasmid contains the first 529 bp of the promoter fused to the *KIADH3* gene.

Yeast transformants, used for β-galactosidase (β-gal) analysis, were selected for leucine prototrophy. Colonies were collected after 3 days of growth, inoculated into 2 ml of YNB containing 0.2 % glucose and

**Table 1.** Yeast strains used in this study

Strains	Genotype	Reference
MW179-1D	<i>MATα metA1 ade-T600 leu2 trpA1 ura3 lac4</i>	Saliola <i>et al.</i> (2008)
MW179-1D/ <i>KIADR1Δ</i>	<i>KIADR1 :: KanMX4</i>	This study
MW179-1D/ <i>KIADH3Δ</i>	<i>KIADH3 :: URA3</i>	This study
MW179-1D/ <i>KIADR1Δ KIADH3Δ</i>	<i>KIADH3 :: URA3</i>	This study
MW179-1D/ <i>KIADH3Δ</i>	<i>KIADR1 :: KanMX4</i>	Saliola <i>et al.</i> (2008)
BY4741	<i>MATα his3Δ0 leu2Δ0 met15Δ0 ura3Δ0</i>	Euroscarf coll
BY4741/ <i>ADR1Δ</i>	<i>ADR1 :: KanMX4</i>	Euroscarf coll

grown for a further 2 days. Finally, 200 µl of these cultures was inoculated in 10 ml of YNB containing 2 % ethanol or 2 % glycerol plus 0.1 % glucose and grown for a further 60 h ready for β-gal determination. Parental and deleted strain cultures grown on ethanol showed a higher biomass yield (roughly  $3 \times 10^8$  cells ml<sup>-1</sup>) than those grown in 2 % glycerol plus 0.1 % glucose (roughly  $2 \times 10^8$  cells ml<sup>-1</sup>). Four independent clones were inoculated for each construct and each β-gal value represents the media from at least three independent determinations. β-Gal values with a standard deviation >20 % were excluded from the determinations. β-Gal activity (nmol min mg<sup>-1</sup> protein) was measured in crude extract prepared by breakage of yeast cells with glass beads according to Kaiser *et al.* (1994).

**Assay methods.** Glucose, glycerol and ethanol concentrations in culture supernatants were measured using commercial kits from R-Biopharm according to the manufacturer's instructions.

**Amplification of *KIADR1* and construction of centromeric and multicopy plasmids.** The *KIADR1* gene was amplified as *XbaI* fragments from the *K. lactis* genome with the following primers: forward: 5'-GGGTCTAGacctaccagcaagtgcagcttc-3' and reverse: 5'-GGGTCTagatagataatcgccacaattccac-3' (in capital letters are sequences introduced to produce a *XbaI* site). The primer sequences were located at -1216 bp upstream of the ATG and +561 bp downstream of the stop codon. The amplified *KIADR1* was cloned as a 5.9 kbp *XbaI* fragment into the *XbaI* site of the pTZ19 vector (Pharmacia) to harbour pTZ19/*KIADR1*. The 5.9 kbp *KIADR1 XbaI* fragment was also cloned in the *XbaI* site of the *K. lactis* centromeric KCplac13 to harbour KCp-*KIADR1*, in the *SpeI* site of the multi-copy pKL plasmid (Saliola *et al.*, 2008) to harbour p*KIADR1* and in the *XbaI* site of the *S. cerevisiae* centromeric pRS416 plasmid (pRS-*KIADR1*).

**Construction of plasmids with the *KIADR1* gene fused in frame with the GFP gene.** pTZ19/*KIADR1* was digested with *HpaI* (site located at the 3' end terminal part of the gene 66 nucleotides before the stop codon) and ligated with the 0.8 kbp *EcoI*CRI-digested GFP gene fragment. Analysis of *E. coli* clones transformed with the ligated fragments allowed the isolation of pTZ18/*KIADR1*-GFP. This plasmid contains the *KIADR1* gene fused in frame with the GFP at the *HpaI* site. This gene codes for a chimeric protein in which KlAdr1 is devoid of the last 22 amino acids as compared to the WT protein. pTZ18/*KIADR1*-GFP was digested with *XbaI*, and the purified *KIADR1*-GFP 6.8 kbp fragment was cloned in the *XbaI* and *SpeI* sites of the centromeric (KCplac13) and multi-copy (pKL) *K. lactis* plasmids, and into *S. cerevisiae* pRS416 to harbour KCp-*KIADR1*-GFP, p*KIADR1*-GFP and pRS-*KIADR1*-GFP, respectively.

Construction of the centromeric KCplac13 vector containing the chimeric *KIGUT2*-GFP gene has already been described (Saliola *et al.*, 2008).

**Construction of the *KIADR1* deletion cassette.** The deletion cassette was constructed by digesting pTZ19/*KIADR1* with *StuI*-*BglII* and cloning the purified KanMX4 *EcoRV*-*BglII*-digested fragment obtained from the pFA6a vector at these sites. The final plasmid isolated (pTZ19/*KIADR1*Δ) contained a deletion of 95 % of *KIADR1*. The cassette was linearized by *XbaI* digestion and transformed into strain MW179-1D for gene replacement. Transformants were selected on plates containing G418. Resistant clones were first analysed by Southern blot: DNA prepared from putative deleted strains was transferred to filters and hybridized concomitantly with <sup>32</sup>P DNA probes containing the deleted portions of the *KIADR1* gene and the KanMX4 gene. The correct integration of the cassette into the *KIADR1* locus was finally confirmed by PCR using the primers previously used for its amplification and the KanMX4 K1: 5'-ggatgatggcctaagtacg-3' and K2: 5'-gttcattgatgctcgatgag-3' primers. One of these deletants, named MW179-1D/*KIADR1*Δ, was further characterized.

### Construction of strains with the *KIADR1* and *KIADH3* deletion.

The strains MW179-1D/*KIADH3*Δ and MW179-1D/*KIADR1*Δ *KIADH3*Δ have been constructed transforming the parental strain with the linearized *KIADH3*::URA3 DNA deletion cassette (Mazzoni *et al.*, 1992). Putative deleted clones selected for uracil prototrophy were analysed for *KIADH3* activity on native gel. Strains unable to express this activity were further verified by PCR for *KIADH3* deletion with the primers (forward) 5'-ggtgaaaacgtcaagggtcg-3', (reverse) 5'-agatggcaaccagtcaccag-3' and URA3 (forward) 5'-gcagtactctgcccgtgttatac-3'.

### Construction of centromeric and multi-copy *ScADR1* plasmids.

A pRS314 plasmid containing the *ScADR1* gene with its own promoter and terminator (Young *et al.*, 1998) was digested with *SpeI*-*SalI*, and the 6.2 kbp purified fragment was cloned in the *K. lactis* KCplac13 centromeric and pKL multicopy vectors (Saliola *et al.*, 2008), digested with the same enzymes, to harbour KCp-*ScADR1* and pKL-*ScADR1*, respectively.

### Construction of the chimeric *K. lactis/S. cerevisiae* ADR1 plasmid.

Construction of the *K. lactis/S. cerevisiae* ADR1 gene was performed in two steps. First, the 5' end of *KIADR1* (containing 980 bp of the promoter and the sequence coding for the first 184 amino acids, comprising the zinc-finger domains of the protein) was amplified by PCR using the following primers: forward 5'-tgctcagtcctccacctct-3' and reverse-5' GGGTCTAGAgatataatcgccacaattccac-3'. The amplified fragment was digested with *PmlI*-*XbaI* and cloned in the *EcoI*CR-*XbaI*-digested sites of pRS416 to harbour pRS-5'*KIADR1*. In the second step, the 4.1 kbp fragment of the *ScADR1* gene, containing the DNA sequence that codes for the portion of the protein comprised from amino acid 164 to amino acid 1323 plus 600 bp of the terminator, was amplified by PCR with the following primers: forward 5'-gggactag-taccaagaaagtgtcgagaact-3' and reverse 5'-gggtctagatataatcgccacaattccac-3'. The PCR product was digested with *SpeI*-*XhoI*, purified and ligated with the pRS-5'*KIADR1* plasmid digested with *XbaI*-*XhoI*. The final plasmid was called pRS-KI/*ScADR1* and contained the *K. lactis/S. cerevisiae* chimeric gene.

**DASPMI and GFP fluorescence microscopy.** DASPMI (2-[4dimethylaminostryl]-1-methylpyridinium iodide) mitochondria-specific staining of live cells was carried out according to Yaffe (1995).

Cells transformed with GFP plasmids were grown under selective conditions to the exponential phase and then shifted for 6 h in YP containing carbon sources at concentrations specified in the text. To assess the fluorescent signal, cells were analysed with an Axio Observer inverted microscope (Zeiss) then scanned in a series of 0.3 µm sequential sections with AxioVision software (Zeiss) that allows 2D reconstruction.

**General methods.** DNA manipulation, plasmid engineering and other techniques were performed according to standard procedures. Yeast transformation was performed by electroporation with a Biorad Gene Pulser apparatus. Putative KlAdr1-binding sites were retrieved using the FIMO tool of the MEME suite (Bailey *et al.*, 2009).

Respiratory rates were measured at 30 °C using a Clark-type electrode according to the method of Ferrero *et al.* (1981).

## RESULTS

### Carbon source utilization in *KIADR1*Δ

The *K. lactis* ADR1 gene was amplified by PCR using primers designed on the basis of the reported Génolevures genome sequence (Dujon *et al.*, 2004; Bussereau *et al.*, 2006). The role of this gene was studied by deleting *KIADR1* with the KanMX4 gene and one of the deleted strains, named MW179-1D/*KIADR1*Δ, was further characterized.

Growth tests on *Kladr1Δ* were performed in rich YP medium and minimal medium (YNB) in the presence of respiratory and fermentative carbon sources. The growth of the parental and *Kladr1Δ* strains showed it to be similar on YNB plates containing glucose, lactate or ethanol, whereas in glycerol the mutant displayed a highly reduced growth (Fig. 1a). Unexpectedly, *Kladr1Δ* showed reduced growth on YP medium that was not ameliorated by the addition of glycerol (YPG) (Fig. 1a), suggesting the requirement of KlADR1 for the utilization of glycerol and for the mobilization of nutrient sources contained in YP medium.


The two strains showed a comparable growth on YPD, while the addition of antimycin A, an inhibitor of the respiratory transport chain, reduced the growth of the mutant (Fig. 1a). The addition of ergosterol, a structural component required for membrane fluidity and permeability (Parks *et al.*, 1995), to plates containing YPD-antimycin A recovered the growth of *Kladr1Δ* to nearly parental levels (Fig. 1a). Finally, *Kladr1Δ* was unable to grow on plates containing oleate as a sole carbon source (Fig. 1a) like its *S. cerevisiae* counterpart (Simon *et al.*, 1991).

The growth of the *Kladr1Δ* in YP medium (Fig. 1a) was compared to that of *S. cerevisiae adr1Δ* and was extended to

plates containing only YE or peptone. *Kladr1Δ* showed reduced growth on YP and severely reduced growth on YE and peptone, revealing that these phenotypes are specific to *Kladr1Δ* (Fig. 1b). In fact, the WT, the *Kladr1Δ* complemented with KCp-KIADR1 and also the *Klglut2Δ* strain, unable to grow under glycerol conditions like *Kladr1Δ* (Fig. 1b) (Saliola *et al.*, 2008), showed comparable growth ability on these media (Fig. 1b). In contrast, *Scadr1Δ* and its parental strain were unable to grow on these nutrient sources, suggesting a different role of these two factors in the mobilization of complex resources like peptone and YE (Fig. 1b).

The decreased fermentative capability of *Kladr1Δ*, deduced from sensitivity to antimycin A (Goffrini *et al.*, 1989), were associated with reduced respiratory capability. In fact, *Kladr1Δ* showed a reduction of 50% O<sub>2</sub> respiration rate in the presence of glucose and ethanol, as compared to its parental strain (glucose: 45±6 vs 22±3; ethanol: 51±6 vs 27±3 μl O<sub>2</sub> per hour per milligram dry weight), while in lactate, the oxygen consumption was reduced to 30% (41±3 vs 14±1).

*K. lactis* mainly grows under aerobic conditions (De Deken, 1966) as opposed to *S. cerevisiae*, where respiration



**Fig. 1.** (a) Growth testing of *K. lactis* MW179-1D (WT) and *Kladr1Δ*. (b) Growth test comparison of *K. lactis* and *S. cerevisiae* parental and *adr1Δ* strains. *KlGlut2Δ* and *KlADR1Δ* harbouring the centromeric KCp-KIADR1 were used as controls for growth on YNB plates supplemented by glycerol and peptone, and YE and YP plates. Cells were grown on plates containing YNB or YP medium supplemented by glucose (YNB+D, YPD), ethanol (YNB+E), glycerol (YNB+G, YPG) and lactate (YNB+L) at a concentration of 2%; in oleate medium; or on plates containing only peptone, YE or YP medium (YP). YPD plates were also supplemented by antimycin A (YPD+AA) or antimycin A plus ergosterol (YPD+AA Erg). *K. lactis* cultures were pre-grown in YNB containing 1% glucose, and *S. cerevisiae* cultures were pre-grown in YNB containing 1% glucose supplemented with casamino acids. Cultures were adjusted to the same density, and 5 μl of serial 10-fold dilutions was spotted onto the medium indicate. The initial concentration was  $1 \times 10^7$  cells ml<sup>-1</sup>.

is completely repressed by glucose. However, when the concentration of oxygen in the culture was reduced to a critical point, as compared to the cellular biomass and glucose concentration, cells began to ferment, accumulating ethanol in the medium (Lemaire & Wésolowski-Louvel, 2004). Accordingly, determination of the main parameters describing the growth behaviour of *Kladr1Δ* can provide insight into the respiratory-fermentative capabilities of the mutant. These parameters, reported in the curves of Fig. 2, display (at time intervals) cell growth, glucose consumption and the production of ethanol and glycerol. The growth curve of the mutant showed a profile not very different from that of WT (Fig. 2). Glucose consumption was also very similar in the two strains, at least over the first 18 h, and was followed by a slightly delayed consumption of glucose and higher accumulation of ethanol (2.4 vs 1.9 g l<sup>-1</sup>) in *Kladr1Δ* (Fig. 2). Glycerol is one of the end products of fermentation; its determination in the supernatant of the two strains showed a significantly lower amount, and absence of its reoxidation, in *Kladr1Δ* as compared to WT (Fig. 2).


### In-gel native PAGE analysis of Adh pattern in *Kladr1Δ*

The four *ADH* genes identified in *K. lactis* (*KIADH1–KIADH4*) are regulated at the transcriptional level by carbon source-dependent growth conditions (Saliola & Falcone, 1995). The corresponding activities have been identified, in native gel from *adh* mutant extracts, by ADH staining assay (Mazzoni *et al.*, 1992; Saliola & Falcone, 1995) that allows dissection of the *S. cerevisiae* ADH system (Lutstorf & Megnet, 1968; Ciriacy, 1975, 1979).

ScAdr1 is the transcription factor responsible for the activation of the *S. cerevisiae* *ADH2* gene (Ciriacy 1979; Denis & Young, 1983). To test whether KlAdr1 is involved in the activation of *KIADH* genes, we analysed the *K. lactis* Adh pattern. Cellular extracts from respiratory and fermentative-grown cultures of the two strains were fractionated on native PAGE and stained for ADH activity. The cultures grown in 7% and 1% glucose-containing medium represent conditions completely fermentative or as mixed respiratory-fermentative (Saliola & Falcone, 1995). These differences were confirmed by the Adh patterns (Fig. 3a, lanes 1 and 2). Comparison of the two Adh patterns (Fig. 3a) showed reduced levels of both cytosolic (Adh1 and Adh2) and mitochondrial activity (Adh3 and Adh4) in *Kladr1Δ* (lanes 5–8 vs lanes 1–4), indicating that KlAdr1 can regulate the expression of these genes.


To better assess its role in the expression of *KIADH* genes, both *KIADR1* and *ScADR1* were over-expressed from centromeric and multi-copy plasmids and their expression was compared, on cultures grown in different carbon sources, by Adh pattern. The strain harbouring the centromeric KCp-KIADR1 displayed in glucose a reduced amount of KlAdh4 (Fig. 3b, lanes 1 and 2 vs Fig. 3a, lanes 1 and 2), an activity that increases during fermentation or growth in ethanol (Fig. 3a and b, lane 4) (Saliola & Falcone, 1995). The strain transformed with KCp-ScADR1 showed an Adh pattern almost identical to that of the untransformed strain (Fig. 3b lanes 5–8).

In contrast, the strains over-expressing either factor displayed highly increased levels of KlAdh3 (Fig. 3c), even in ethanol (lanes 4 and 8), a compound that specifically represses the expression of this gene (Saliola & Falcone,



**Fig. 2.** Growth curves and metabolite determination of WT and *Kladr1Δ*. Cultures were grown in YP medium containing 1% glucose. Glycerol (G), glucose (D) and ethanol (E) were determined at time intervals in culture supernatants. *adr1Δ*=*Kladr1Δ*. Each value is the average of three independent determinations.





**Fig. 3.** In-gel native Adh pattern from extracts of *K. lactis* cultures grown in fermentative/respiratory carbon sources. Adh-stained activities from extracts of (a) WT and *Kladr1*Δ. (b) WT transformed with centromeric KCP-KIADR1 (WT+cKIADR1) or KCP-ScADR1 (WT+cScADR1) plasmids. (c) WT transformed with the multi-copy pKIADR1 (WT+pKIADR1) or pKL-ScADR1 (WT+pScADR1) plasmids. Cellular extracts were prepared from cultures grown on YP containing 7% (D7) or 1% (D1) glucose, 2% glycerol (G) or 2% ethanol (E). The migrating position of each activity is also reported. The arrows indicate the hetero-tetramers formed by KIAdh3 and KIAdh4 activity.

1995; Saliola *et al.*, 2007). Since the regulation of *KIADH3* and *KIADH4* genes is mutually exclusive (Saliola & Falcone, 1995), the increased Adr1-dependent expression of *KIADH3* alters the balance between the two genes leading to the formation of hetero-tetramers between the activity of KIAdh3 and KIAdh4 (see arrows in Fig. 3c; and the next section) (Saliola *et al.*, 2007). Indeed, strains over-expressing *ADR1* displayed very high levels of KIAdh3, but lower levels of KIAdh4. However, the over-expression of *ScADR1* was unable to complement the growth-deficient glycerol phenotype of *Kladr1*Δ (data not shown).

### Deletion analysis of the *KIADH3* promoter in *Kladr1*Δ


An extensive deletion analysis of the *KIADH3* promoter fused to the *lacZ* gene reporter allowed the functional

identification of positive and negative elements. Suggesting very complex regulation of this gene (Saliola *et al.*, 2007). The positive (+) and the short *cis*-acting repression ethanol (RE) element previously identified are reported in Fig. 4(a).

To test whether KIAdr1 is involved in the control of *KIADH3*, deleted fragments of its promoter fused to *lacZ* were transformed in parental and *Kladr1*Δ strains (Fig. 4a).  $\beta$ -Gal values were determined under ethanol-repressive and glycerol plus 0.1% glucose-inducing conditions (see Fig. 3a, lanes 4 and 3, respectively) to bypass the inability of *Kladr1*Δ inability to grow in glycerol. pCG1, pCG17, pCG13, pCG11 and pCG12 containing promoter fragments of increased deleted length showed comparable levels of activity in both parental and mutant strains (Fig. 4a). In contrast, the *Kladr1*Δ strain transformed with pCG7 (first 529 bp of the promoter) showed comparable levels of  $\beta$ -gal activity under glycerol and ethanol conditions (323 and 371 nmol min<sup>-1</sup> protein, respectively). These values are very different to those measured in the parental strain (445 and 83, respectively) (Fig. 4a), suggesting the presence on the DNA<sub>529-297</sub> fragment of an element conferring constitutive  $\beta$ -gal activity. In this process, KIAdr1 seems to modulate the positive (+) factor element conferring a role of repressor in ethanol and one of activator in glycerol. However, the  $\beta$ -gal values shown by the two pCG11-containing strains suggested that the KIAdr1-dependent role noted for pCG7 is completely abolished by the presence of another element(s) located in the DNA<sub>660-529</sub> fragment (Fig. 4a). Analysis of pCG8 ( $\Delta$ 660–529) confirmed the role of the two elements (Fig. 4a). In fact, the repressive role of KIAdr1 (Adr1<sub>RE</sub>) and the presence of an antagonistic dominant activator partner on the DNA<sub>660-529</sub> fragment are in agreement with  $\beta$ -gal measured in the parental strain and its *Kladr1*Δ strain (79 and 18 vs 616 and 150) (Fig. 4a). The low  $\beta$ -gal value measured under the two conditions in the parental strain with pCG8 (79 and 18), as compared to *Kladr1*Δ (616 and 150), is probably due to the repressive role of Adr1<sub>RE</sub> in the absence of the dominant DNA<sub>660-529</sub>-positive element. Conversely, the different  $\beta$ -gal values measured under ethanol conditions (31 vs 59) in the two pCG1-containing strains can be interpreted in the same way.

To confirm the interfering role of KIAdr1 in the regulation of *KIADH3* and *KIADH4* (see Fig. 3c), we transformed *Kladr1*Δ, *Kladh3*Δ and *Kladh3*Δ with promoter constructs containing different combinations of positive and negative elements fused to *KIADH3*, used as a gene reporter (Fig. 4b–d) (Saliola *et al.*, 2007). The extracts from the ethanol- and glycerol-grown cultures of the two transformed strains were fractionated in native gel and their Adh staining activity compared (Fig. 4b).

The patterns were similar in the two strains transformed with pCG29, pCG30, pCG31 and pCG32 (Fig. 4b) (Saliola *et al.*, 2007), whereas the activity of pCG33 extracts (see scheme of Fig. 4c) showed interesting results. The expression of this plasmid, together with pCG32 used as a control, confirmed the role of KIAdr1 in this gene (Fig. 4d). In fact, the double



**Fig. 4.** Deletion analysis of the *KIADH3* promoter fused to the *lacZ* and *KIADH3* gene reporters. (a)  $\beta$ -Gal activity (nmol min $^{-1}$  protein) measured in WT and *Kladr1 $\Delta$*  cells transformed with various promoter fragment constructs fused to *lacZ*. (b) Adh staining activity of extracts of *Kladr1 $\Delta$* , *Kladh3 $\Delta$*  (*adr1 $\Delta$ adh3 $\Delta$* ) and *Kladh3 $\Delta$*  (*adh3 $\Delta$* ) cells transformed with various promoter fragment constructs fused to *KIADH3*. (c) Centromeric plasmids carrying *KIADH3* under different promoter forms used in this analysis. (d) Adh staining activity of extracts of *Kladr1 $\Delta$* , *Kladh3 $\Delta$*  and *Kladh3 $\Delta$*  cells transformed with pCG32 and pCG33. A scheme reports the homo-/hetero-tetrameric composition of each Adh band in the figure. Cells were grown for 60 h in YNB medium under repressing (E=2% ethanol) and de-repressing (2% glycerol and 0.1% glucose) conditions. Each  $\beta$ -gal value is the average of at least three independent clones analysed. The Adh pattern was analysed in three independent clones for each construct. The position on the promoter of positive (+) and negative (RE and Adr1<sub>RE</sub>) elements is also reported.

mutant harbouring pCG33 showed the presence of KlAdh3 in both substrates (Fig. 4d, lanes 4 and 9), as compared to *Kladh3 $\Delta$*  (lanes 2 and 7). However, the highly reduced levels of KlAdh3 in ethanol (Fig. 4d, lanes 4), as compared to glycerol (Fig. 4d, lane 9) and to the  $\beta$ -gal values in Fig. 4(a) (371 vs 323 nmol min $^{-1}$  protein), were interpreted according to the scheme reported on the left of Fig. 4(d). The absence of KlAdr1 leads to the contemporary expression of the two genes, as shown by the assembly of both KlAdh3 and KlAdh4 homo- and hetero-tetramers (Fig. 4d, lanes 4, 8 and 9 and Fig. 4b, lanes 2 and 3) in a manner similar to the over-expression of *KIADR1* in WT (see Fig. 3c).

### Cellular localization of KlAdr1


In *S. cerevisiae*, expression of the *ScADR1-GFP* fusion gene allowed the intracellular localization of ScAdr1 under repressed and de-repressed conditions (Sloan *et al.*, 1999; Huh *et al.*, 2003; Moreno-Cermeno *et al.*, 2013).

In order to visualize the localization of KlAdr1 in *K. lactis*, the centromeric Kcp-KIADR1-GFP plasmid was inserted into the parental and *Kladr1 $\Delta$*  strains. Growth of the transformed *Kladr1 $\Delta$*  strain on glycerol-containing plates

confirmed the functionality of the *KIADR1-GFP* gene (data not shown). To visualize the fusion protein, WT-transformed cells were grown to the mid-exponential phase in YNB containing 1% glucose and shifted for 6 h on fermentative (glucose 7%) under respiratory conditions (glycerol 2% and ethanol 2%). As shown in Fig. 5(a), cells grown in glucose 7% displayed a diffuse fluorescence suggesting a mainly cytoplasmic localization of KlAdr1-GFP. In contrast, cells shifted in glycerol or ethanol showed a weak, localized fluorescent signal in glycerol and a stronger one in ethanol, indicating that a substantial amount of the chimeric protein was localized within the nucleus (Fig. 5a). These results suggested a common mechanism of gene activation in the two yeasts under fermentative/repressive and respiratory/derepressive conditions.

### Cellular localization of KIGut2 in *Kladr1 $\Delta$*

Since *KIGUT2* is required for growth in glycerol (Saliola *et al.*, 2008) and, in *S. cerevisiae*, this gene is under the control of Adr1 (Tachibana *et al.*, 2005), we analysed the localization of KIGut2 in *Kladr1 $\Delta$*  by means of its fusion with GFP.



**Fig. 5.** Fluorescence analysis of WT and *Kladr1Δ* cells transformed with the centromeric KCp-KIADR1-GFP or KCp-KIGUT2-GFP plasmids or treated with DASPMI. Micrographs show (a) the intracellular localization of KIADR1-GFP in the WT under fermentative and respiratory conditions; (b) the intracellular localization of KIGut2-GFP in the WT and in the *Kladr1Δ*; and (c) the mitochondrial morphology of WT and *Kladr1Δ* cells stained with DASPMI. Cells were grown to late exponential phase in YNB supplemented by 1% glucose. Cells were harvested, then shifted for 6 h on YP containing 7% (YPD7) or 1% glucose (YPD), 2% glycerol (YPG) or 2% ethanol (YPE) with/without DASPMI.

In the parental strain, the expression of the *KIGUT2-GFP* gene showed mitochondrial localization of the chimeric protein under glucose conditions (Fig. 5b) (Saliola *et al.*, 2010), while fluorescence distribution was reduced in intensity and slightly more diffuse in *Kladr1Δ*. On the contrary, under glycerol and ethanol conditions, the mutant displayed a reduced fluorescence signal, altered in definition and slightly dispersed throughout the cell (Fig. 5b).

In *S. cerevisiae*, Adr1 coordinates the pathways that generate acetyl-CoA and NADH from non-fermentable substrates (Young *et al.*, 2003), playing a role in the mitochondrial network dynamic (Cherry & Denis, 1989). Similarly, *Kladr1Δ* showed highly reduced oxygen reoxidation rates, the inability to utilize respiratory substrates like glycerol, and also that the intracellular distribution of KIGut2 is slightly more diffuse. Therefore, cells were analysed by the DASPMI vital dye assay to test the effect of deletion of *KIADR1* on the mitochondrial network. Under glycerol conditions, this assay visualized a mitochondrial network frame with a punctuated shape slightly more accentuated in the mutant, as compared to the WT (Fig. 5c).

### Expression of *KIADR1* in *S. cerevisiae*


The over-expression of *ScADR1* in *K. lactis* influenced the Adh pattern like the endogenous gene (Fig. 3c, lanes 5–8 vs lanes 1–4). Thus, a complementation test was performed to see whether KIAdr1 is capable of activating transcription of the *S. cerevisiae ADH2* gene. To this end, pRS-KIADR1-GFP was transformed into the *Scadr1Δ* strain and the extracts were tested on native PAGE for Adh2 activity. Indeed, as can be seen from Fig. 6(a), the extracts of four *KIADR1-GFP*-transformed clones showed the presence of Adh2 only under derepressing conditions (lanes 3–6 vs lanes 9–12), as compared to the parental (lanes 1 and 2) and *Scadr1Δ* (lanes 7 and 8)

strains. However, this activity (Fig. 6a, lanes 3–6) was not displayed at levels comparable to those of WT (lane 2). The presence of the KIAdr1-GFP protein in the transformants was also analysed under fluorescent microscopy. The weak fluorescent signal visualized under repressing and de-repressing conditions indicated that KIAdr1 is expressed in *S. cerevisiae* (Fig. 6b). The protein would appear to be distributed in the cytoplasm under repressing conditions, while in glucose 0.2%, fluorescence distribution suggested its nuclear localization (Fig. 6b).

To discover why the *S. cerevisiae* strain expressing *KIADR1-GFP* displayed reduced levels of KIAdh2, we focused on the zinc-finger DNA-binding domains of this protein being more highly conserved between the two factors (Fig. 6c) (Parua *et al.*, 2010). The scheme of Fig. 6c also shows the identity between the domains of the two proteins, the phosphorylated serine of Adr1 (\*S) (Denis *et al.*, 1992) and the putative one of KIAdr1 (<sup>S</sup>S). The location of this serine (<sup>S</sup>S<sub>312</sub>) suggested the presence of an extra domain of roughly 50 amino acids in *K. lactis*, as compared with that of ScAdr1 (\*S<sub>230</sub>) (Fig. 6c, – aa insertion).

A chimeric *ADR1* gene was constructed to test whether the ability of KIAdr1 to activate ScAdh2 depends on its DNA-binding domain. The chimeric gene (*KI/ScADR1*) contains the *K. lactis* promoter and the sequence coding for the zinc-finger domains in frame with the *ScADR1* sequence encoding the transactivating domains (Bemis & Denis, 1988) (scheme of Fig. 6c). This chimera, when transformed into the *Scadr1Δ* strain under derepressed conditions, allowed the appearance of the native ScAdh2 activity like the endogenous factor (Fig. 6d, lanes 7 and 8 vs lanes 1 and 2). These results indicated that the zinc-finger domains of KIAdr1 can recognize UAS1 or other UAS1-like elements of its promoter activating the transcription of *ScADH2* in a ScAdr1-dependent manner (Shuster *et al.*, 1986).





**Fig. 6.** Expression analysis of *KIADR1-GFP* in *S. cerevisiae* by in-gel native Adh pattern and GFP localization. (a) In-gel native Adh pattern from extracts of BY4741 (WT), *Scadr1Δ* (*Scadr1Δ*) and *Scadr1Δ* transformed with KcP-*KIADR1-GFP*. (b) Micrographs showing the intracellular localization of *KIAdr1* in *Scadr1Δ* cells transformed with KcP-*KIADR1-GFP*. (c) A scheme showing the identities between *K. lactis* and *S. cerevisiae* Adr1 domains and the construction strategy for the chimeric fusion *KI/ScAdr1* factor. (d) Expression analysis of the *KI/ScADR1* chimeric gene in the *Scadr1Δ* by in-gel native Adh pattern from extracts of BY4741 (WT), *Scadr1Δ* (*Scadr1Δ*) and *Scadr1Δ* transformed with pRS-*KI/ScADR1* (*Scadr1Δ*+chim). Cellular extracts were prepared from YNB-casamino acid-supplemented cultures containing 2% (D2) or 0.2% (D.2) glucose, ethanol 2% (E2) or glycerol 2% (G2). Cells transformed with KcP-*KIADR1-GFP* were pre-grown to the early exponential phase in 2% glucose, shifted in repressing (D2) and de-repressing (D.2) glucose medium and grown for a further 6 h.

## DISCUSSION

### Role of *KIAdr1* in the metabolism of *K. lactis*

In this study, we report the characterization of the *K. lactis* *ADR1* gene. In *S. cerevisiae*, *Adr1* is a zinc-finger transcription factor that regulates the transcription of *ADH2* and other genes required for the utilization of respiratory carbon sources, peroxisomal biogenesis, fatty acid metabolism and

other cellular functions (Denis & Young, 1983; Simon *et al.*, 1991; Young *et al.*, 2003; Tachibana *et al.*, 2005).

In *K. lactis*, *KIAdr1* is required for growth in glycerol, oleate and YP medium (Fig. 1a). Moreover, the sensitivity of *Kladr1Δ* to antimycin A, its reduced oxygen reoxidation rate and the different accumulation of ethanol and glycerol in glucose-grown culture indicated a role for *KIAdr1* under respiratory and fermentative conditions (Figs 1 and 2).

Yeast cells produce and accumulate glycerol from the intermediate of glycolysis dihydroxyacetone phosphate as the main compatible solute for redox balance and osmoregulation (Hohmann, 2002). It follows that the reduced glycolytic flux of *Kladr1Δ* is unable to support the accumulation of glycerol for the cytoplasmic reoxidation of excess NAD (P)H to maintain redox balance (Fig. 2) (Cialfi *et al.*, 2011). As a consequence, the metabolism of these cells becomes respiro-fermentative (González-Siso *et al.*, 2000) and the respiratory chain intervenes to neutralize excess redox (Overkamp *et al.*, 2002). Therefore, blocking of the respiratory chain by antimycin A inhibits the growth of this mutant, while the addition of ergosterol, a major structural component required for the control of membrane fluidity (Park *et al.*, 1995), can recover the growth deficiency (Fig. 1). These data and the growth-deficient oleate phenotypes (Fig. 1) indicated that the altered redox balance of *Kladr1Δ* is the limiting factor in the adaptation of plasma membrane fatty acid composition during the transition from respiratory to fermentative conditions (Heipieper *et al.*, 2000; Cialfi *et al.*, 2011).

### Role of *KIAdr1* in the ADH system of *K. lactis*

Under fermentative and respiratory conditions, Adhs are required for the maintenance of intracellular redox balance (Bakker *et al.*, 2000; Overkamp *et al.*, 2002; Saliola *et al.*, 2012). Since *KIAdr1* intervenes in the regulation of these physiological conditions (Fig. 2), we showed, by in-gel native assays, that the deletion and over-expression of this factor altered the Adh pattern in all carbon sources. In particular, *KIAdh3* was the activity most affected by its over-expression (Fig. 3). Moreover, extensive deletion analysis allowed the isolation of a *KIADH3* promoter fragment (DNA<sub>529-297</sub>) in which *KIAdr1* exerted its role in modulating the expression of this gene (Fig. 4).

In this process, the role of *KIAdr1* is to tune *KIADH3*, repressing its expression in ethanol and inducing it in glycerol. We propose that the modulatory role of *KIAdr1* in this gene is to avoid the contemporary expression of *KIADH3* and *KIADH4* and the assembly of hetero-tetramers of the two activities (Fig. 4d). In fact, the two Adhs displayed different biochemical properties (Bozzi *et al.*, 1997), in agreement with their specific activation during growth under mainly fermentative (*KIADH4*) and respiratory (*KIADH3*) conditions (Saliola & Falcone, 1995; Saliola *et al.*, 2006).

Using the FIMO tool of the MEME suite (Bailey *et al.*, 2009), we were able to identify on this DNA fragment a

putative UAS<sub>509-490</sub> element showing a significant identity with the UAS1 of *ScADH2* (Shuster *et al.*, 1986). Since *KIADH3* seems to be regulated by an intricate and concerted balance of many positive and negative factors (Fig. 4) (Saliola *et al.*, 2007), further research will be necessary to confirm the binding of KlAdr1 to this element, the roles and the identities of the other factors and their relation with KlAdr1.

### Intracellular localization of KlAdr1 and KlGut2

Minor differences have been reported in the localization of Adr1 in *S. cerevisiae*, probably due to the diverse genetic context of the strains used (Sloan *et al.*, 1999; Huh *et al.*, 2003; Moreno-Cermeño *et al.*, 2013). In *K. lactis*, the diffuse fluorescence displayed under fermentative conditions suggested a mainly cytoplasmic localization of the KlAdr1-GFP. In contrast, in glycerol- and ethanol-grown cells, the fluorescent signal suggested the localization of the protein within the nucleus (Fig. 5a). We may emphasize that the nuclear localization of KlAdr1 is in agreement with the regulation of *KIADH3* in glycerol-inducing and ethanol-repressive media, as suggested by the deletion data (see Fig. 4a and d).

The localization of KlGut2-GFP was also performed, to provide insights into the role of KlAdr1 on *KIGUT2*, a gene that, in *S. cerevisiae*, is regulated by Adr1 (Tachibana *et al.*, 2005).

In *K. lactis*, KlGut2-GFP displayed mitochondrial localization (Saliola *et al.*, 2010) while fluorescence distribution was reduced in intensity in *Kladr1Δ* as compared to WT (Fig. 5b). However, under glycerol and ethanol conditions, the mutant displayed a stronger fluorescence signal and reduced intensity, definition and dispersal throughout the cell than in glucose (Fig. 5b). These results indicated that KlAdr1 is not directly involved in its regulation and, as for *KIADH3*, seems to modulate its expression; or it may be necessary to addressing KlGut2 into the mitochondrial compartment. In fact, as shown by DASPMI vital dye assay, *Kladr1Δ* cells displayed a mitochondrial network distribution only slightly punctuated as compared to WT. Although this result suggested a role for KlAdr1 in the maintenance of the mitochondrial network, like its *S. cerevisiae* counterpart (Cherry & Denis, 1989), the effect of its deletion seems more marked on the localization and distribution of KlGut2.

### Expression of *KIADR1* in *S. cerevisiae*

By cross-complementation experiments, we were able to define the extent of KlAdr1-activating expression. *KIADR1* was shown to be expressed in *Scadr1Δ* and its product localized in the cytoplasm in glucose-repressing medium, and in the nucleus under de-repressing conditions (Fig. 6b). Moreover, KlAdr1 induced the expression of *ScADH2* under de-repressing conditions, although with reduced levels of ScAdh2 as compared to WT (Fig. 6a). This result suggested

a limited functionality of KlAdr1 in the activation of *ScADH2* expression, while the introduction of the chimeric *KI/ScADR1* gene in *Scadr1Δ* led to levels of ScAdh2 similar to those of the WT. It follows that the zinc-finger domains of KlAdr1 must bind to a *ScADH2* promoter element(s) regulating its expression in a ScAdr1-dependent fashion (Fig. 6d). We suggest that this domain of KlAdr1 binds to the same sequence of ScAdr1, while the transactivator domains have diverged between the two yeasts and cannot be fully recognized by the evolutionarily diverged *S. cerevisiae* apparatus (Bussereau *et al.*, 2006).

Finally, growth test comparison of the two *adr1Δ* yeast strains showed the inability of the *S. cerevisiae* parental and mutant strains to grow in the presence of rich nutrient sources without the addition of carbohydrate (YE, peptone or YP) (Fig. 1b). Since *K. lactis* has been isolated from mammalian fluids (milk) particularly rich in nutrients (Rodicio & Heinisch, 2013), these data suggest that the adaptation to this habitat and/or the intracellular mobilization of nutrients as sources of carbon, nitrogen and co-factors could be a specific KlAdr1-dependent trait.

In conclusion, these factors have evolved and contribute to the metabolic adaptation of the two yeasts to their specific environmental niches.

### ACKNOWLEDGEMENTS

This work was funded by 'Ateneo 2012' – Sapienza University of Rome. SC is a recipient of a postdoctoral fellowship of 'Ateneo 2014'. We would like to thank C. Falcone (Sapienza University of Rome) for support during the execution of this project and E. Zanni, C. Lanini and M. L. Sponziello (Sapienza University of Rome) for technical help. We would also like to thank T. Lodi of the University of Parma, Italy, for data on cell respiration and E. T. Young of the University of Washington, USA, for suggestions and for having provided the plasmid with the ScADR1 gene.

### REFERENCES

- Bailey, T. L., Boden, M., Buske, F. A., Frith, M., Grant, C. E., Clementi, L., Ren, J., Li, W. W. & Noble, W. S. (2009). MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**, W202–W208.
- Bakker, B. M., Bro, C., Kötter, P., Luttk, M. A., van Dijken, J. P. & Pronk, J. T. (2000). The mitochondrial alcohol dehydrogenase Adh3p is involved in a redox shuttle in *Saccharomyces cerevisiae*. *J Bacteriol* **182**, 4730–4737.
- Bemis, L. T. & Denis, C. L. (1988). Identification of functional regions in the yeast transcriptional activator *ADR1*. *Mol Cell Biol* **8**, 2125–2131.
- Bennetzen, J. L. & Hall, B. D. (1982). The primary structure of the *Saccharomyces cerevisiae* gene for alcohol dehydrogenase. *J Biol Chem* **257**, 3018–3025.
- Bozzi, A., Saliola, M., Falcone, C., Bossa, F. & Martini, F. (1997). Structural and biochemical studies of alcohol dehydrogenase isozymes from *Kluyveromyces lactis*. *Biochim Biophys Acta* **1339**, 133–142.
- Bussereau, F., Casaregola, S., Lafay, J. F. & Bolotin-Fukuhara, M. (2006). The *Kluyveromyces lactis* repertoire of transcriptional regulators. *FEMS Yeast Res* **6**, 325–335.

- Cherry, J. R. & Denis, C. L. (1989). Overexpression of the yeast transcriptional activator *ADR1* induces mutation of the mitochondrial genome. *Curr Genet* **15**, 311–317.
- Cialfi, S., Uccelletti, D., Carducci, A., Wésolowski-Louvel, M., Mancini, P., Heipieper, H. J. & Saliola, M. (2011). KIHs1 is a component of glycerol response pathways in the milk yeast *Kluyveromyces lactis*. *Microbiology* **157**, 1509–1518.
- Ciriacy, M. (1975). Genetics of alcohol dehydrogenase in *Saccharomyces cerevisiae*. II. Two loci controlling synthesis of the glucose-repressible ADH II. *Mol Gen Genet* **138**, 157–164.
- Ciriacy, M. (1979). Isolation and characterization of further *cis*- and *trans*-acting regulatory elements involved in the synthesis of glucose-repressible alcohol dehydrogenase (ADHIII) in *Saccharomyces cerevisiae*. *Mol Gen Genet* **176**, 427–431.
- De Deken, R. H. (1966). The Crabtree effect: a regulatory system in yeast. *J Gen Microbiol* **44**, 149–156.
- Denis, C. L. & Young, E. T. (1983). Isolation and characterization of the positive regulatory gene *ADR1* from *Saccharomyces cerevisiae*. *Mol Cell Biol* **3**, 360–370.
- Denis, C. L., Ciriacy, M. & Young, E. T. (1981). A positive regulatory gene is required for accumulation of the functional messenger RNA for the glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*. *J Mol Biol* **148**, 355–368.
- Denis, C. L., Fontaine, S. C., Chase, D., Kemp, B. E. & Bemis, L. T. (1992). *ADR1c* mutations enhance the ability of *ADR1* to activate transcription by a mechanism that is independent of effects on cyclic AMP-dependent protein kinase phosphorylation of Ser-230. *Mol Cell Biol* **12**, 1507–1514.
- Dickinson, J. R., Salgado, L. E. & Hewlins, M. J. (2003). The catabolism of amino acids to long chain and complex alcohols in *Saccharomyces cerevisiae*. *J Biol Chem* **278**, 8028–8034.
- Dujon, B., Sherman, D., Fischer, G., Durrens, P., Casaregola, S., Lafontaine, I., De Montigny, J., Marck, C., Neuvéglise, C. & other authors (2004). Genome evolution in yeasts. *Nature* **430**, 35–44.
- Ferrero, I., Viola, A. M. & Goffeau, A. (1981). Induction by glucose of an antimycin-insensitive, azide-sensitive respiration in the yeast *Kluyveromyces lactis*. *Antonie Van Leeuwenhoek* **47**, 11–24.
- Goffrini, P., Algeri, A. A., Donnini, C., Wesolowski-Louvel, M. & Ferrero, I. (1989). *RAG1* and *RAG2*: nuclear genes involved in the dependence/independence on mitochondrial respiratory function for the growth on sugars. *Yeast* **5**, 99–106.
- González-Siso, M. I., Freire-Picos, M. A., Ramil, E., González-Domínguez, M., Rodríguez Torres, A. & Cerdán, M. E. (2000). Respiratory fermentative metabolism in *Kluyveromyces lactis*: insights and perspectives. *Enzyme Microb Technol* **26**, 699–705.
- Heipieper, H. J., Isken, S. & Saliola, M. (2000). Ethanol tolerance and membrane fatty acid adaptation in *adh* multiple and null mutants of *Kluyveromyces lactis*. *Res Microbiol* **151**, 777–784.
- Hohmann, S. (2002). Osmotic stress signaling and osmoadaptation in yeasts. *Microbiol Mol Biol Rev* **66**, 300–372.
- Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S. & O'Shea, E. K. (2003). Global analysis of protein localization in budding yeast. *Nature* **425**, 686–691.
- Kaiser, C., Michaelis, S. & Mitchell, A. (1994). *Methods in Yeast Genetics*. New York: Cold Spring Harbour Laboratory Press.
- Lemaire, M. & Wésolowski-Louvel, M. (2004). Enolase and glycolytic flux play a role in the regulation of the glucose permease gene *RAG1* of *Kluyveromyces lactis*. *Genetics* **168**, 723–731.
- Lutstorf, U. & Megnet, R. (1968). Multiple forms of alcohol dehydrogenase in *Saccharomyces cerevisiae*. I. Physiological control of ADH-2 and properties of ADH-2 and ADH-4. *Arch Biochem Biophys* **126**, 933–944.
- Mazzoni, C., Saliola, M. & Falcone, C. (1992). Ethanol-induced and glucose-insensitive alcohol dehydrogenase activity in the yeast *Kluyveromyces lactis*. *Mol Microbiol* **6**, 2279–2286.
- Mazzoni, C., Santori, F., Saliola, M. & Falcone, C. (2000). Molecular analysis of  $UAS_E$ , a *cis* element containing stress response elements responsible for ethanol induction of the *KIADH4* gene of *Kluyveromyces lactis*. *Res Microbiol* **151**, 19–28.
- Moreno-Cermeño, A., Alsina, D., Cabisco, E., Tamarit, J. & Ros, J. (2013). Metabolic remodeling in frataxin-deficient yeast is mediated by Cth2 and Adr1. *Biochim Biophys Acta* **1833**, 3326–3337.
- Overkamp, K. M., Bakker, B. M., Steensma, H. Y., van Dijken, J. P. & Pronk, J. T. (2002). Two mechanisms for oxidation of cytosolic NADPH by *Kluyveromyces lactis* mitochondria. *Yeast* **19**, 813–824.
- Parks, L. W., Smith, S. J. & Crowley, J. H. (1995). Biochemical and physiological effects of sterol alterations in yeast – a review. *Lipids* **30**, 227–230.
- Parua, P. K., Ratnakumar, S., Braun, K. A., Dombek, K. M., Arms, E., Ryan, P. M. & Young, E. T. (2010). 14-3-3 (Bmh) proteins inhibit transcription activation by Adr1 through direct binding to its regulatory domain. *Mol Cell Biol* **30**, 5273–5283.
- Rodicio, R. & Heinisch, J. J. (2013). Yeast on the milky way: genetics, physiology and biotechnology of *Kluyveromyces lactis*. *Yeast* **30**, 165–177.
- Saliola, M. & Falcone, C. (1995). Two mitochondrial alcohol dehydrogenase activities of *Kluyveromyces lactis* are differently expressed during respiration and fermentation. *Mol Gen Genet* **249**, 665–672.
- Saliola, M., Shuster, J. R. & Falcone, C. (1990). The alcohol dehydrogenase system in the yeast, *Kluyveromyces lactis*. *Yeast* **6**, 193–204.
- Saliola, M., Gonnella, R., Mazzoni, C. & Falcone, C. (1991). Two genes encoding putative mitochondrial alcohol dehydrogenases are present in the yeast *Kluyveromyces lactis*. *Yeast* **7**, 391–401.
- Saliola, M., Mazzoni, C., Solimando, N., Crisà, A., Falcone, C., Jung, G. & Fleer, R. (1999). Use of the *KIADH4* promoter for ethanol-dependent production of recombinant human serum albumin in *Kluyveromyces lactis*. *Appl Environ Microbiol* **65**, 53–60.
- Saliola, M., De Maria, I., Lodi, T., Fiori, A. & Falcone, C. (2006). *KIADH3*, a gene encoding a mitochondrial alcohol dehydrogenase, affects respiratory metabolism and cytochrome content in *Kluyveromyces lactis*. *FEMS Yeast Res* **6**, 1184–1192.
- Saliola, M., Getuli, C., Mazzoni, C., Fantozzi, I. & Falcone, C. (2007). A new regulatory element mediates ethanol repression of *KIADH3*, a *Kluyveromyces lactis* gene coding for a mitochondrial alcohol dehydrogenase. *FEMS Yeast Res* **7**, 693–701.
- Saliola, M., Sponziello, M., D'Amici, S., Lodi, T. & Falcone, C. (2008). Characterization of *KIGUT2*, a gene of the glycerol-3-phosphate shuttle, in *Kluyveromyces lactis*. *FEMS Yeast Res* **8**, 697–705.
- Saliola, M., D'Amici, S., Sponziello, M., Mancini, P., Tassone, P. & Falcone, C. (2010). The transdehydrogenase genes *KINDE1* and *KINDII* regulate the expression of *KIGUT2* in the yeast *Kluyveromyces lactis*. *FEMS Yeast Res* **10**, 518–526.
- Saliola, M., Tramonti, A., Lanini, C., Cialfi, S., De Biase, D. & Falcone, C. (2012). Intracellular NADPH levels affect the oligomeric state of the glucose 6-phosphate dehydrogenase, *Eukaryot. Cell* **11**, 1503–1511.
- Shain, D., Salvatore, C. & Denis, C. (1992). Evolution of the alcohol dehydrogenase (ADH) genes in yeast: characterization of a fourth ADH in *Kluyveromyces lactis*. *Mol Gen Genet* **232**, 478–488.
- Shuster, J., Yu, J., Cox, D., Chan, R. V., Smith, M. & Young, E. (1986). *ADR1*-mediated regulation of *ADH2* requires an inverted repeat sequence. *Mol Cell Biol* **6**, 1894–1902.
- Simon, M., Adam, G., Rapatz, W., Spevak, W. & Ruis, H. (1991). The *Saccharomyces cerevisiae* *ADR1* gene is a positive regulator of transcription of genes encoding peroxisomal proteins. *Mol Cell Biol* **11**, 699–704.

**Sloan, J. S., Dombek, K. M. & Young, E. T. (1999).** Post-translational regulation of Adr1 activity is mediated by its DNA binding domain. *J Biol Chem* **274**, 37575–37582.

**Tachibana, C., Yoo, J. Y., Tagne, J. B., Kacherovsky, N., Lee, T. I. & Young, E. T. (2005).** Combined global localization analysis and transcriptome data identify genes that are directly coregulated by Adr1 and Cat8. *Mol Cell Biol* **25**, 2138–2146.

**Wésolowski-Louvel, M., Prior, C., Bornecque, D. & Fukuhara, H. (1992).**  $Rag^-$  mutations involved in glucose metabolism in yeast: isolation and genetic characterization. *Yeast* **8**, 711–719.

**Yaffe, M. P. (1995).** Isolation and analysis of mitochondrial inheritance mutants from *Saccharomyces cerevisiae*. *Methods Enzymol* **260**, 447–453.

**Young, E. T. & Paquin, D. (1985).** Isolation and DNA sequence of *ADH3*, a nuclear gene encoding the mitochondrial isozyme of alcohol dehydrogenase in *Saccharomyces cerevisiae*. *Mol Cell Biol* **5**, 3024–3034.

**Young, E. T., Saario, J., Kacherovsky, N., Chao, A., Sloan, J. S. & Dombek, K. M. (1998).** Characterization of a p53-related activation domain in Adr1p that is sufficient for ADR1-dependent gene expression. *J Biol Chem* **273**, 32080–32087.

**Young, E. T., Dombek, K. M., Tachibana, C. & Ideker, T. (2003).** Multiple pathways are co-regulated by the protein kinase Snf1 and the transcription factors Adr1 and Cat8. *J Biol Chem* **278**, 26146–26158.

---

Edited by: V. J. Cid and H. Sychrova