



Available online at:

<http://www.italian-journal-of-mammalogy.it/article/view/12101/pdf>

doi:10.4404/hystrix-27.1-12101

Commentary

## Species Distribution Models: exploring patterns and processes in mammal ecology

Danilo RUSSO<sup>1,2,\*</sup>, Luigi MAIORANO<sup>3</sup>, Hugo REBELO<sup>4,5,2</sup>, Damiano G. PREATONI<sup>6</sup>

<sup>1</sup>Wildlife Research Unit, Dipartimento di Agraria, Università degli Studi di Napoli Federico II, via Università 100, 80055 Portici (Napoli), Italy

<sup>2</sup>School of Biological Sciences, University of Bristol, Life Sciences Building, Bristol BS8 1TQ, UK

<sup>3</sup>Department of Biology and Biotechnologies "Charles Darwin", Università di Roma "La Sapienza", viale dell'Università 32, 00185 Roma (RM), Italy

<sup>4</sup>CIBIO/InBIO, University of Porto, Campus Agrário Vairão, 4485-661 Vairão, Portugal

<sup>5</sup>CEABN/InBIO, Universidade de Lisboa, Tapada da Ajuda, 1349-017 Lisboa, Portugal

<sup>6</sup>Unità di Analisi e Gestione delle Risorse Ambientali – Guido Tosi Research Group, Dipartimento di Scienze Teoriche e Applicate, Università degli Studi dell'Insubria, via J.H. Dunant 3, 21100 Varese, Italy

### Abstract

Modelling species distribution is one of the most widely used approaches used by modern ecologists to predict species occurrence as well as to explore the environmental variables that determine the presence of a species in an habitat.

Given the outstanding importance of mammals in the context of conservation or management actions, it is not surprising that an ever increasing number of mammalogists have successfully employed species distribution models over the last few years. As a scientific journal devoted to mammal natural history, Hystrix, the Italian Journal of Mammalogy is pleased to present a collection of articles forming a special issue that deals with such an important and timely matter.

### Article history:

Received: 26 June 2016

Accepted: 30 June 2016

### Acknowledgements

As co-editors of this issue, the Authors would like to warmly thank all the authors that contributed with their work, for their enthusiastic support and for the quality of their contributions.

Modelling species distribution is one of the most widespread approaches used by modern ecologists to detect which environmental variables influence current species occurrence, and may also be successfully applied to infer past or future distribution patterns. This method has a wide array of applications for ecological and conservation studies, making it possible to provide answers to many questions, the most basic of which is to identify areas where a given species is likely to occur. More sophisticated theoretical goals may also be tackled, from assessing the role of factors that shape biodiversity patterns in time and space to exploring niche segregation in sympatric species (Russo et al., 2014, 2015). Undoubtedly, what has made such techniques especially attractive is the relevance they have for conservation or management purposes, e.g. they may: help to focus survey work on areas where a target taxon is more probably present, making field operations more cost-effective (Rebello and Jones, 2010); detect regions of high habitat suitability in need of protection, or those in need of appropriate restoration (Razgour et al., 2011); predict invasion patterns by alien species (Di Febbraro et al., 2016, this issue); inform managers about the best places to carry out reintroduction or restocking (Wilson et al., 2011); help carry out effective gap analysis (Bosso et al., 2013).

Given the outstanding importance of mammals as targets of conservation or management actions, it is not surprising that an ever increasing number of mammalogists have successfully employed species distribution models (SDM) over the last few years. As a scientific journal devoted to mammal natural history, Hystrix, the Italian Journal of Mammalogy is pleased to present a collection of articles forming a special issue that deals with such an important and timely matter.

The collection of papers provides a broad overview spanning across different subjects, time and spatial scales. A community-scale assessment is provided by Gomes Vale et al. (2016, this issue), who employed GIS and ecological niche analysis for a 22-species strong mammal assemblage in West Sahara-Sahel to identify the main areas of occur-

rence, explore biogeographic relationships, and detect species richness hotspots. A similar objective is pursued by Cooper-Bohannon et al. (2016, this issue), but their distributional analysis has concerned 58 bat species from South Africa, showing how species of high conservation value may occur in low-species richness areas, which should therefore not be neglected by conservationists. In another study, Razgour et al. (2016, this issue) offer a review of SDM studies on bats remarking that despite the increased popularity of this approach among bat specialists, there is still much space for enhancing scopes and applications by including factors such as dispersal, landscape connectivity and biotic interactions in bat models.

Other articles focus on single species of management importance and show how spatial and time scales influence model outputs with clear conservation implications. Puddu and Maiorano (2016, this issue) present a novel approach to map not only high suitability patches, but also the potential corridors connecting them for the mouflon (*Ovis aries*) in Sardinia, whereas Jones et al. (2016a, this issue) predictions on how different forest management options may affect survival of European red squirrels (*Sciurus vulgaris*) at two UK sites make it possible to identify felling and restocking adjustments that may lead to a dramatic survival increase.

One of the possibilities offered by SDM is to explore how interactions between factors may ultimately shape species distributions. An interesting example is given by Jones et al. (2016b, this issue), whose prediction of the potential distribution of African wild dogs *Lycaon pictus* under future climate and land use changes also incorporates the influence of its potential competitor, the leopard *Panthera pardus*. Time is certainly a crucial variable that may challenge model reliability, yet few studies have dealt with this important aspect, even since the start of the research with SDM. We present one exception by Areias-Guerreiro et al. (2016, this issue) who tested the current validity of past models of otter (*Lutra lutra*) distribution in Spain generated from 1998 records using a new dataset, coming up with the encouraging conclusion that otters expanded towards the predicted areas, thus remarking the usefulness of medium term models. Tackling the species scale from a differ-

\* Corresponding author

Email address: danrusso@unina.it (Danilo Russo)

ent perspective, Di Marco et al. (2016, this issue) examine how to scale population persistence targets from populations up to species using the European ground squirrel (*Spermophilus citellus*) as a case-study.

Given the ever growing concern caused by biological invasions and the threat posed by alien squirrels, two papers refer to this topic. Di Febbraro et al. (2016, this issue) offer a global picture of predicted current and 2070 potential distributions of eight squirrel species, identifying current potential hotspots of invasion and assessing climate change influences by 2070, whereas White et al. (2016, this issue) model how squirrelpox may spread in the well known red and grey squirrel system. Their model includes a realistic representation of habitat in Southern Scotland and provides important management insights.

Overall, although this special issue has inevitably missed some aspects of the vast universe of applications of SDMs to mammal studies, it showcases an attractive selection of articles highlighting the latest trends in SDM research while also providing much needed cautionary notes on the use of this technique.

Overall, we are confident that our readers will enjoy the articles and that this special issue may boost further developments on this theme. ☞☞

## References

Areias-Guerreiro J., Mira A., Barbosa A., 2016. How well can models predict changes in species distributions? A 13-year-old otter model revisited. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11867

Bosso L., Rebelo H., Garonna A.P., Russo D., 2013. Modelling geographic distribution and detecting conservation gaps in Italy for the threatened beetle *Rosalia alpina*. *Journal for Nature Conservation* 21: 72–80.

Cooper-Bohannon R., Rebelo H., Jones G., Cotterill F., Monadiem A., Schoeman M., Taylor P., Park, K., 2016. Predicting bat distributions and diversity hotspots in southern Africa. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11722

Di Febbraro M., Martinoli A., Russo D., Preatoni D., Bertolino S., 2016. Modelling the effects of climate change on the risk of invasion by alien squirrels. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11776

Di Marco M., Santini L., Visconti P., Mortelliti A., Boitani L., Rondinini C., 2016. Using habitat suitability models to scale up population persistence targets. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11660

Gomes Vale C., Campos J., Silva T., Gonçalves D., Sow A., Martínez-Freiría F., Boratyński Z., Brito, J., 2016. Biogeography and conservation of mammals from the West Sahara-Sahel: an application of ecological niche-based models and GIS. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11659

Jones M., Bertola L., Razgour O., 2016. Predicting the effect of interspecific competition on habitat suitability for the endangered African wild dog under future climate and land cover changes. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11678

Jones H., White A., Geddes N., Clavey P., Farries J., Dearnley T., Boots M., Lurz P., 2016. Modelling the impact of forest design plans on an endangered mammal species: the Eurasian red squirrel. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11673

Puddu G., Maiorano L., 2016. Combining multiple tools to provide realistic potential distributions for the mouflon in Sardinia: species distribution models, spatial pattern analysis and circuit theory. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11695

Razgour O., Hanmer J., Jones G., 2011. Using multi-scale modelling to predict habitat suitability for species of conservation concern: The grey long-eared bat as a case study. *Biological Conservation* 144: 2922–2930.

Razgour O., Rebelo H., Di Febbraro M., Russo D., 2016. Painting maps with bats: species distribution modelling in bat research and conservation. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11753

Rebelo H., Jones G., 2010. Ground validation of presence-only modelling with rare species: a case study on *Barbastella barbastellus* (Chiroptera: Vespertilionidae). *Journal of Applied Ecology* 47: 410–420.

Russo D., Di Febbraro M., Rebelo H., Mucedda M., Cistrone L., Agnelli P., De Pasquale P.P., Martinoli A., Scaravelli D., Spilinga C., Bosso L., 2014. What story does geographic separation of insular bats tell? A case study on Sardinian rhinolophids. *PLoS ONE* 9(10): e110894. doi:10.1371/journal.pone.0110894

Russo D., Di Febbraro M., Cistrone L., Jones G., Smeraldo S., Garonna A.P., Bosso L., 2015. Protecting one, protecting both? Scale-dependent ecological differences in two species using dead trees, the rosalia longicorn beetle and the barbastelle bat. *Journal of Zoology, London* 297: 165–175.

White A., Lurz P., Bryce J., Tonkin M., Ramoo K., Bamforth L., Jarrott A., Boots M., 2016. Modelling disease spread in real landscapes: Squirrelpox spread in Southern Scotland as a case study. *Hystrix* 27(1). doi:10.4404/hystrix-27.1-11657

Wilson C.D., Roberts D., Reid, N., 2011. Applying species distribution modelling to identify areas of high conservation value for endangered species: A case study using *Margaritifera margaritifera* (L.). *Biological Conservation* 144: 821–829.