

One Gene and Two Proteins: a Leaderless mRNA Supports the Translation of a Shorter Form of the *Shigella* VirF Regulator

Maria Letizia Di Martino,^a Cédric Romilly,^b E. Gerhart H. Wagner,^b Bianca Colonna,^a Gianni Prosseda^a

Istituto Pasteur Italia-Fondazione Cenci Bolognetti, Department of Biology and Biotechnology C. Darwin, Sapienza Università di Roma, Rome, Italy^a; Department of Cell and Molecular Biology, Biomedical Center, Uppsala University, Uppsala, Sweden^b

ABSTRACT VirF, an AraC-like activator, is required to trigger a regulatory cascade that initiates the invasive program of *Shigella* spp., the etiological agents of bacillary dysentery in humans. VirF expression is activated upon entry into the host and depends on many environmental signals. Here, we show that the *virF* mRNA is translated into two proteins, the major form, VirF₃₀ (30 kDa), and the shorter VirF₂₁ (21 kDa), lacking the N-terminal segment. By site-specific mutagenesis and toeprint analysis, we identified the translation start sites of VirF₃₀ and VirF₂₁ and showed that the two different forms of VirF arise from differential translation. Interestingly, *in vitro* and *in vivo* translation experiments showed that VirF₂₁ is also translated from a leaderless mRNA (llmRNA) whose 5' end is at position +309/+310, only 1 or 2 nucleotides upstream of the ATG84 start codon of VirF₂₁. The llmRNA is transcribed from a gene-internal promoter, which we identified here. Functional analysis revealed that while VirF₃₀ is responsible for activation of the virulence system, VirF₂₁ negatively autoregulates *virF* expression itself. Since VirF₂₁ modulates the intracellular VirF levels, this suggests that transcription of the llmRNA might occur when the onset of the virulence program is not required. We speculate that environmental cues, like stress conditions, may promote changes in *virF* mRNA transcription and preferential translation of llmRNA.

IMPORTANCE Shigella spp. are a major cause of dysentery in humans. In bacteria of this genus, the activation of the invasive program involves a multitude of signals that act on all layers of the gene regulatory hierarchy. By controlling the essential genes for host cell invasion, VirF is the key regulator of the switch from the noninvasive to the invasive phenotype. Here, we show that the *Shigella virF* gene encodes two proteins of different sizes, $VirF_{30}$ and $VirF_{21}$, that are functionally distinct. The major form, $VirF_{30}$, activates the genes necessary for virulence, whereas the minor $VirF_{21}$, which shares the C-terminal two-thirds of $VirF_{30}$, negatively autoregulates *virF* expression itself. $VirF_{21}$ is transcribed from a newly identified gene-internal promoter and, moreover, is translated from an unusual leaderless mRNA. The identification of a new player in regulation adds complexity to the regulation of the *Shigella* invasive process and may help development of new therapies for shigellosis.

Received 8 October 2016 Accepted 11 October 2016 Published 8 November 2016

Citation Di Martino ML, Romilly C, Wagner EGH, Colonna B, Prosseda G. 2016. One gene and two proteins: a leaderless mRNA supports the translation of a shorter form of the Shigella VirF regulator. mBio 7(6):e01860-16. doi:10.1128/mBio.01860-16.

Editor Bonnie Bassler, Princeton University

Copyright © 2016 Di Martino et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Gianni Prosseda, gianni.prosseda@uniroma1.it, or E. Gerhart H. Wagner, gerhart.wagner@icm.uu.se.

This article is a direct contribution from a Fellow of the American Academy of Microbiology. External solicited reviewers: Carmen Buchrieser, Pasteur Institute, Paris, France; Charles Dorman, University of Dublin, Trinity College.

higella spp. are highly adapted human pathogens that cause Dacillary dysentery (1). The sophisticated infectious strategy of Shigella depends on the capacity to invade, disrupt, and cause inflammatory destruction of the intestinal epithelial barrier (2, 3). Activation of the invasive program is exceptionally complex and involves many signals affecting gene regulation at different levels. A key factor is VirF, an AraC-like transcription factor (TF) whose expression is activated as Shigella bacteria sense entry into the host environment (4, 5). In a cascade model, VirF triggers activation of the virB and icsA genes. IcsA affects bacterial intracellular spreading, and VirB promotes expression of several virulence genes, including those encoding a type III secretion system (T3SS), its effectors, and the last regulator of the cascade, MxiE (6, 7). Interestingly, MxiE, another AraC-like TF, appears to rely on high-level transcriptional slippage to generate its reading frame from two separate open

reading frames (ORFs) (8). The genes virF, icsA, virB, and those controlled by VirB are located on a virulence plasmid (pINV) and are silenced outside the human host (9). At low temperatures, the nucleoid-associated protein H-NS represses transcription of the virulence genes (5). In a temperaturedependent manner, H-NS interacts with two sites within the *virF* promoter, spaced by an intrinsically curved DNA region, to prevent access of RNA polymerase (5, 10, 11). At a permissive temperature (37°C), reduced DNA curvature counteracts H-NS binding (4, 12) and unmasks a binding site for the nucleoid protein FIS to activate *virF* transcription (13). VirF then relieves H-NS-mediated repression of virB and icsA and directly stimulates transcription (14, 15). By binding upstream of the virF promoter between two H-NS sites, VirB also counteracts H-NS-dependent repression of virF transcription (16). Expression of *virF* is further modulated by integration host factor (IHF) (17) and other environmental factors, such as pH and osmolarity (7), and is affected by tRNA modifications (18).

The relevance of *virF* activation for the invasive program is further supported by posttranscriptional regulation of *icsA*. RnaG is a *cis*-encoded antisense RNA that promotes premature termination of the *icsA* mRNA (19). VirF binds the RnaG promoter and decreases *rnaG* expression (14). Thus, VirF plays a dual role: (i) it relieves H-NS-mediated repression to activate *icsA* transcription, and (ii) it represses RnaG transcription, thus increasing the level of *icsA* mRNA (14). VirF also globally activates the expression of chromosomal genes in both *Shigella* and *Escherichia coli*. In particular, VirF appears to play a role in shaping the *Shigella* transcriptional program to better match the requirements of an effective intracellular life (20–22).

Like other members of the AraC family of transcriptional regulators, VirF has a conserved, carboxy-terminal DNA-binding domain with two helix-turn-helix (HTH) motifs. AraC-like proteins are typically insoluble and, accordingly, problems with VirF purification have hampered biochemical studies (23). Mutagenesis experiments indicated that the N-terminal domain of VirF promotes dimerization while C-terminal HTH2 motif mutants are nonfunctional (24).

While attempting a thorough characterization of VirF, we found that the *virF* mRNA (R1) is subject to differential translation, giving rise to two forms of VirF. VirF₃₀ activates the virulence system and some chromosomal genes, whereas VirF₂₁ exerts negative feedback control on *virF* expression itself.

Moreover, we identified a second *virF* mRNA species (R2) with a 5' end at position nucleotide (nt) +309/+310. This leaderless yet translation-competent mRNA is transcribed from a geneinternal promoter. Possible implications in an interplay between environmental sensing and virulence gene expression are discussed.

RESULTS

The virF gene encodes two independently translated proteins, VirF₃₀ and VirF₂₁. Earlier experiments on *E. coli* minicells carrying the virF gene on recombinant plasmids from *Shigella flexneri* and *Shigella sonnei* indicated two main VirF protein forms of about 30 and 21 kDa and a minor form of 27 kDa (25, 26). The significance of the 27- and 21-kDa forms remained unclear, and it seemed possible that they were degradation products of fulllength VirF (27). To analyze which VirF forms are present in *Shigella*, a 3×FLAG tag sequence was inserted at the 3' end of the *S. flexneri* M90T virF ORF. Western blot analysis (Fig. 1A) confirmed that two VirF proteins, VirF₃₀ (30 kDa) and VirF₂₁ (21 kDa), are expressed by *S. flexneri*. The 27-kDa form was not detected.

The sequence of the *virF* gene contains three putative start codons, all in the same frame, for $VirF_{30}$ and an internal ATG codon, consistent with independent translation of $VirF_{21}$ (25). Thus, we determined at which ATG codons $VirF_{30}$ and $VirF_{21}$ translation initiates. In the absence of a recognizable Shine-Dalgarno (SD) sequence upstream, prediction of the ATG encoding the N-terminal methionine of $VirF_{30}$ was difficult. Thus, each of the ATG codons (codons 1, 2, and 4; codon 3 encodes Asp) (Fig. 1B) was tested for translation initiation activity by using plasmids carrying the *virF* promoter followed by a *virF-lacZ* translational fusion. Plasmid pFL-4A is fused in frame after the fourth *virF* codon (third Met codon), and pFL-1A is fused after the first



FIG 1 Detection of two VirF protein variants. (A) Western blot analysis with anti-FLAG antibody of whole-cell extracts of *S. flexneri* M90T carrying *virF*- $3 \times$ FLAG. Two forms are indicated, VirF₃₀ and VirF₂₁. (B) Schematic organization of the *virF* gene of *Shigella*. The methionine (M) codons relevant for this study are highlighted. The transcription start site (+1) was identified previously (5).

ATG (Fig. 2A). β -Galactosidase activities indicated that the construct with all three ATGs has \approx 5-fold-higher activity than the one fused after ATG₁. Thus, ATG₂ and/or ATG₄ appear to be required for high translation of VirF₃₀, and ATG₁ gives a minor contribution. ATG₄, which has a short upstream SD-like (GAA) sequence, was tested by introducing an $ATG_4 \rightarrow GGG$ (Gly) mutation into pFL-4A. This plasmid, pFL-M4G, in which ATG₁ and ATG₂ are still present, gave very low reporter gene activity (Fig. 2A), suggesting ATG₄ as the main VirF₃₀ start codon. Western blot analysis supported this. VirF₃₀ was produced only from the wild-type (wt) virF gene, but not when ATG₄ had mutated (Fig. 2B, cf. pMYSH6504 and pF-M4G). To corroborate this finding in vitro, we used a toeprinting assay to analyze the formation of ribosomal initiation complexes on virF mRNA (28). A predominant toeprint was seen 17 nt downstream of AUG₄ and a minor one 16 nt downstream of AUG₁ (Fig. 2C), in line with our in vivo results (Fig. 2A and B). Additional bands downstream of position +17 of AUG₄ implicated possible 30S binding-driven structure changes resulting in reverse transcription pauses. In conclusion, translation of VirF₃₀ initiates predominantly at ATG₄. Throughout the remainder of this paper, codon positions are accordingly renumbered, with ATG₄ as codon 1.

While searching for a VirF₂₁ translation start site, we noticed an in-frame ATG codon within virF at position 311 to 313 (relative to +1 of *virF*) (Fig. 1B), consistent with translation of the minor form of VirF. To validate ATG₈₁ (formerly ATG₈₄) as the start codon for VirF21, two mutations were introduced into virF, generating a codon change and a frameshift, respectively. To mutate ATG₈₁ to a different codon that would retain VirF₃₀ function, we changed the ATG (mRNA position 311 to 313) to CTG (Met to Leu; pF-M81L) (Fig. 3A) or to ATC (Met to Ile; pF-M81I). Neither mutation should affect VirF₃₀ translation but should abolish independent translation of VirF₂₁. Both mutant VirF₃₀ proteins were tested for activated expression of *virB* in a *virF*-defective S. flexneri strain (M90TFd) (see Table S1 in the supplemental material) carrying plasmids expressing wt VirF, VirF_{M811}, or VirF_{M81I}. VirF_{M81L} but not VirF_{M81I} activated virB to a level comparable to wt (see Fig. S1 in the supplemental material). Thus, the substitution in VirF_{M811} impairs VirF₃₀ functionality, and therefore only pF-M81L was used in subsequent experiments. Moreover, the exclusive expression of VirF₃₀ upon Met \rightarrow Leu substitution (Fig. 3B) identified ATG_{81} as the start codon for $VirF_{21}$.

To uncouple the translation of VirF₃₀ and VirF₂₁, we inserted a



FIG 2 Identification of the translation start codon of VirF₃₀. (A, left) Schematic of the pFL plasmids used. Plasmids pFL-1A and pFL-4A carry a translational fusion between the 5'-UTR of *virF* mRNA after Met1 (pFL-1A) or Met4 (pFL-4A), in frame with *lacZ*. (Right) pFL-M4G, the Met4 codon, was replaced by Gly (ATG to GGG). β -Galactosidase activities of *E. coli* strain DH10b carrying the same *virF-lacZ* plasmids are shown. Strains harboring the pRS414 vector showed very low β -galactosidase activity levels (2 to 4 Miller units) relative to the values obtained. Values are averages of three experiments, and standard deviations are indicated. (B) Western blot with VirF antibodies on extracts from MG1655 harboring pMYSH6504, a plasmid carrying the wt *virF* gene of *S. flexneri*, or pF-M4G (pMYSH6504 with the M4G substitution). The asterisk indicates unspecific cross-hybridization with a protein in the extract. (C) Toeprint assay results on the +1 *virF* mRNA (see Materials and Methods). The mRNA was incubated alone (lane 1), with 30S (lane 2), and with 30S and tRNA^{fMet} (lane 3). Toeprints at position +16 from ATG1 and at +17 from ATG4 are indicated by a black circle and asterisk, respectively. Sequencing ladders were generated with the same 5'-end-labeled primer.

single G between positions +207 and +208 of *virF* to create a frameshift (pF-FS) into two stop codons (UGA and UAG, +243 to +249), upstream of ATG₈₁ of VirF₂₁ (Fig. 3A). The wt *virF* gene and its M81L (substitution) and frameshift mutant variants were FLAG tagged, resulting in plasmids pGEM-6504-FT, pGEM-M81L-FT, and pGEM-FS-FT. Western blot analysis of total protein from *E. coli* cells showed that only VirF₃₀ is translated from pGEM-M81L-FT and only VirF₂₁ is translated from pGEM-FS-FT (Fig. 3B). Western blotting assays with cells with untagged plasmids confirmed this result (see Fig. S2 in the supplemental material). Since premature termination of frameshifted *virF* only abolished the synthesis of VirF₃₀ and not that of VirF₂₁, both proteins are independently translated.

The relative expression of the two VirF forms was further analyzed by translational *lacZ* fusions. Four *virF-lacZ* fusions enabled us to monitor translation of VirF₂₁ (pRS-M4G and pRS-FS) and VirF₃₀ (pRS-M81L), in comparison to total wt *virF* mRNA translation (pRS-6504). The β -galactosidase levels from pRS-M4G and pRS-FS were about 3-fold lower than those from pRS-6504 and pRS-M81L, which is congruent with the Western blot results shown in Fig. 3B; under these experimental conditions, VirF₂₁ is a minor fraction of the total VirF protein pool. **VirF**₂₁ **negatively autoregulates the** *virF* **gene**. What is the function of the independently translated VirF₂₁? To test whether VirF₂₁ can activate virulence genes, the promoters of *virB* and *icsA* were transcriptionally fused to *lacZ* and transferred to the chromosome of the *E. coli* K-12 strain P90C, generating P90C λ B and P90C λ A, respectively (see Materials and Methods). Activation by wt VirF₃₀ and VirF₂₁ (pMYSH6504), VirF₃₀ (pF-M81L), and VirF₂₁ (pF-FS) was monitored in strain P90C λ B or P90C λ A.

Figure 4A shows that VirF₃₀ alone (pF-M81L) induced the expression of both *lacZ* fusions to a level similar to that in the presence of both VirF₃₀ and VirF₂₁ (pMYSH6504). VirF₂₁ alone (pF-FS) failed to activate (Fig. 4A). Quantitative reverse transcription-PCR (qRT-PCR) results with the *S. flexneri* strain M90T Fd (*virF* defective) carrying the same three plasmids supported this conclusion (Fig. 4B). Thus, a role for VirF₂₁ in the activation of the virulence cascade of *Shigella* is ruled out. A qRT-PCR experiment also confirmed that the previously shown VirF-dependent activation of some chromosomal heat shock genes (20) cannot be carried out by VirF₂₁ (see Fig. S3 in the supplemental material).

To address putative functions of $VirF_{21}$, we investigated its role in positive or negative autoregulation of the *virF* gene. An *E. coli* K-12 strain harboring a P_{virF} -lacZ fusion (DH10b pvirF-lacZ) was



FIG 3 Differential translation of VirF₃₀ and VirF₂₁. (A) Schematic representation of the constructs used to exclusively produce VirF₃₀ or VirF₂₁. Sequences relevant for the construction of plasmid pF-M81L (M81L substitution) or plasmid pF-FS (insertion of G at position +208) are highlighted. Plasmids are derivatives of pMYSH6504. (B) Western blot analysis (with anti-FLAG antibody) of extracts of *E. coli* DH10b carrying pGEM-6504-FT (VirF₃₀ and VirF₂₁), pGEM-M81L-FT (only VirF₃₀), or pGEM-FS-FT (only VirF₂₁). (C) β-Galactosidase activity levels of the *virF-lacZ* plasmids obtained by fusing a fragment (-289 to +405) of the *virF* gene of pMYSH6504 (pRS-6504), pF-M81L (pRS-M81L), pF-M4G (pRS-M4G), pF-FS (pRS-FS), or pF-FS-M81L (pRS-FS-M81L) as the control, to the promoterless *lacZ* gene of pRS414. Values are averages of three experiments, and standard deviations are indicated.

transformed with plasmids that expressed either Ptac promoterdriven VirF₃₀ (pAC-30) or VirF₂₁ (pAC-21). Figure 5A clearly shows that VirF₂₁, but not VirF₃₀, strongly repressed virF expression, and qRT-PCR on the same samples showed corresponding decreases in *lacZ* mRNA levels in the presence of $VirF_{21}$ (Fig. 5B). To validate VirF₂₁-mediated repression of virF transcription in Shigella, we asked whether increased VirF₂₁ levels would reduce the expression of the VirF-activated virB gene. qRT-PCR experiments in the virF-defective strain M90T Fd expressing VirF₃₀ from pF-M81L confirmed severely reduced virB transcription upon induction (isopropyl- β -D-thiogalactopyranoside [IPTG]) of VirF₂₁ (pAC-21) (Fig. 5C). To monitor the VirF₂₁ induction-dependent effect on the VirF protein level, we introduced pAC-21 in the S. flexneri strain that contained the 3×FLAG virF gene (M90T F3xFT; see above). This setup permitted us to assess the levels of VirF21 and VirF30 encoded by pINV by FLAG-tagged antibodies as a function of increasing levels of untagged VirF₂₁ expressed from pAC-21 (monitored via a halon anti-VirF antibody). Figure 5D shows that increasing the VirF₂₁ concentration resulted in a decrease in VirF₃₀, confirming that VirF₂₁ negatively autoregulates *virF* expression.

In addition, we performed DNase I footprinting by *in vitro*translated VirF₂₁ on both strands of the *virF* promoter region. VirF₂₁ was translated in an *in vitro* system (PureSystem) (see Materials and Methods), using a PCR-generated DNA template for $virF_{21}$ -only transcription and translation. VirF₂₁ translation was verified by Western blotting (see Fig. S4 in the supplemental material). Figure 5E indicates that $VirF_{21}$ binding conferred protection of the *virF* promoter region between positions -90 and -20 on the plus strand and approximate positions -60 to -10 on the minus strand and enhanced minus-strand cleavage from about positions -70 to -90. This result, together with data from *in vivo* experiments (Fig. 5A and B), strongly suggests that the transcriptional repression of *virF* by $VirF_{21}$ depends on its direct binding to the consensus *virF* promoter elements.

Identification of a VirF₂₁-encoding leaderless mRNA. The above results showed that two VirF proteins are independently translated. Whether both are translated from the same mRNA, or different versions of virF transcripts, was unknown. The possibility of different mRNAs was suggested by two virF mRNA variants detected in a Northern blot assay performed with total RNA from strain M90T Fd complemented with the virFencoding pMYSH6504 and with plasmid-free M90T Fd (Fig. 6A). An ≈960-nt band (full-length virF mRNA; R1) and an ≈680-nt mRNA that might support translation of $VirF_{21}$ (R2) were visible. To test whether R2 virF mRNA is transcribed from a virF internal promoter or generated by processing, *virF-lacZ* transcriptional fusions and primer extension (PE) analyses were used. We constructed four *virF-lacZ* fusions starting at positions +70, +145, +205, and +305; all were fused at +405. The β -galactosidase activities clearly indicated the presence of a promoter between +205 and +305; truncation up to position +305 produced back-





FIG 4 Functional analysis of VirF₃₀ and VirF₂₁. (A) β-Galactosidase activity of *E. coli* P90C carrying *virB-lacZ* and *icsA-lacZ* transcriptional fusions, transformed with a plasmid expressing VirF₃₀ and VirF₂₁ (pMYSH6504), only VirF₃₀ (pF-M81L), or only VirF₂₁ (pF-FS). Values are averages of three experiments, with standard deviations indicated. (B) *In vivo* levels of *virB* and *icsA* mRNA as a function of the two VirF forms, monitored by qRT-PCR in a Δ*virF S. flexneri* strain (M90T Fd) transformed with pMYSH6504 (VirF₃₀ and VirF₂₁), pF-M81L (VirF₃₀), or pF-FS (VirF₂₁). Expression levels were monitored in M90T or M90T Fd as controls. Samples were run in triplicate, and error bars show the calculated maximum (RQMax) and minimum (RQMin) levels that represent the standard error of the mean expression level (RQ value).

ground values (Fig. 6B). A promoter was indeed predicted by PromoterHunt (29), with consensus -10 (CATTAT; +298 to +303) and -35 elements (TTGACA; +276 to +289) (Fig. 6C). After mutagenesis of the -10 box [CATTAT to CGTTAT; pRS-F(+205 -10mut)], we observed a severe reduction (\approx 7-fold) in the β -galactosidase level. This new promoter was further delineated by PE analysis on RNA extracted from *E. coli* cells harboring the different *virF-lacZ* plasmids. This showed 5' ends at positions +309, +310 (major band), and +311. All three bands were absent in the PE on pRS-F(+305), while with pRS-F(+205 -10mut) the +309/310 bands were not detected. The weaker band at +311 is consistent with a shifted -10 box (data not shown). Thus, the R2 *virF* mRNA is transcribed from a second *virF* promoter, with a transcription start site at position +309/+310. The 5' ends at +309 to 311 and the start codon at +311 to 313 imply that the R2 mRNA is leaderless (Fig. 6C). To test whether the llmRNA is VirF₂₁ translation competent, we cloned the sequence corresponding to R2 mRNA, and also the entire R1 mRNA as a control, downstream of a T7 promoter. To ensure a correct 5' end of the R2 mRNA *in vivo* (5' U₊₃₀₉ as +1) (Fig. 6C), a hammerhead ribozyme sequence downstream of the T7 promoter (see Fig. S5 in the supplemental material) was introduced to generate an R2 mRNA starting at position +309. The plasmids carrying the R1 or R2 transcripts, pAC-T730-FT (R1; virF +1 to +888) and pAC-T7-HH-21-FT (R2; virF +309 to +888) also harbored 3' FLAG tags in *virF*. Upon IPTG induction, *virF* mRNA transcription from the T7 promoter was induced in *E. coli* BL21(D3) harboring either plasmid. PE analysis verified the expected 5' ends of both transcripts (Fig. 7A).

VirF21 translation from the leaderless R2 mRNA was tested by immunoblot analysis on protein extracts after induction. VirF21 was detected in cells carrying pACT7-HH-21-FT, confirming that R2 is a leaderless translation-proficient mRNA (Fig. 7B, right panel). Translation of both VirF forms was observed in cells harboring pAC-T730-FT (Fig. 7B, left panel). In vitro translation in the PureSystem (30) was tested on R1 and R2 mRNAs carrying FLAG tag sequences. Translation products were analyzed with anti-FLAG antibodies. In agreement with the in vivo results, R1 mRNA supported translation of both VirF forms, whereas the leaderless R2 transcript only produced VirF₂₁ (Fig. 7C). Furthermore, toeprint experiments on in vitro-transcribed virF R1 mRNA (start, +1) showed a strong RT stop near the 5' end, indicating initiation complex formation at AUG₄ (compare with Fig. 2C). In contrast, a specific toeprint was observed at position +326 for the llmRNA R2 (start, +309) (Fig. 7D). This toeprint was absent on R1 mRNA, indicating a strong preference for VirF₃₀ translation from the full-length mRNA. Together, these results suggest that a new virF promoter generates a llmRNA variant (R2 mRNA) dedicated to the exclusive translation of VirF21.

DISCUSSION

The complex regulatory cascade for activation of the Shigella virulence genes depends on the VirF protein (7). VirF is at the heart of the switch from the noninvasive to the invasive phenotype. Thus, it is not surprising that its expression is triggered by many environmental signals and that it is controlled at several levels (2, 4, 10, 17). Since its discovery, VirF was known to be present in three forms that differ in size: 30, 27, and 21 kDa (25). The smaller forms were ignored as likely degradation products. Here, we report that the VirF 21-kDa form is translated as an independent polypeptide. Our results address how the VirF21 variant is produced and suggest an autoregulatory role in virF expression. As a first step, we identified the translation start sites of VirF₃₀ and VirF₂₁. Of the three Met codons among the first four codons of the predicted virF ORF, only ATG4 was essential for VirF₃₀ translation (Fig. 1 and 2). Replacement with GGG drastically reduced VirF, as monitored by Western blotting or β -galactosidase activity of virF-lacZ translational fusions (Fig. 2A and B). The identification of ATG4 as a start codon was further supported by toeprint analysis (Fig. 2C). The start codon consistent with the size of VirF₂₁ is ATG81; accordingly, replacement with CTG blocks VirF₂₁ production (Fig. 3B).

Interestingly, while the wt *virF* mRNA is translated into both $VirF_{30}$ and $VirF_{21}$ *in vivo*, a frameshift mutation upstream of



FIG 5 VirF₂₁ autoregulates *virF* expression. (A) β-Galactosidase activity of *virF-lacZ* fusions in response to increased levels of VirF₂₁ or VirF₃₀. Ectopic expression of VirF₂₁ or VirF₃₀ was obtained in *E. coli* p_{*virF}-lacZ* strains carrying pAC-21 or pAC-30, respectively. pGIP7, empty vector. Values are averages of three experiments, and standard deviations indicated. (B) *In vivo* levels of *lacZ* mRNA were monitored in the same samples used in the β-galactosidase assay summarized in panel A. Triplicate samples were evaluated, and error bars indicate standard errors of the mean expression levels (RQ values). (C) *In vivo* levels of *virB* mRNA were monitored in the Δ*virF S*. *flexneri* strain (M90T Fd) carrying pF-M81L (VirF₃₀) or ectopically expressing VirF₂₁ under IPTG control (pAC-21). Triplicate samples were evaluated; error bars show standard errors of the mean expression of VirF₃₀ was monitored with an anti-FLAG antibody. VirF₂₁ induction was monitored with an anti-VirF antibody. Asterisks indicate unspecific cross-hybridization with an unknown protein in the extract. (2) *U*µl of *in vitro*-translated VirF₂₁. The samples were generated with the same 5'-end-labeled plus- or minus-strand-specific primers. The VirF₂₁-protected site is indicated by vertical black lines and shown by shading on both strands of the *virF* promoter sequence.</sub>



FIG 6 *In vivo* analysis of *virF* transcripts. (A) Northern hybridization of 10 μ g of total RNA from *S. flexneri* strain M90TFd, with or without pmysh6504 (*virF* wt) and a *virF*-specific ³²P-labeled probe indicated two major mRNA variants. (B) Schematic representation of *virF-lacZ* transcriptional fusions carrying truncations of the region upstream of the translational start site of VirF₂₁. The ATG for VirF₂₁ is indicated as a reference, and the putative promoter is depicted by an arrow. The β -galactosidase activities of the *virF-lacZ* fusion strains were determined. Values reported are in Miller units and represent the averages \pm standard deviations of five independent experiments. (C) Schematic representation of the new *virF* promoter. The positions of the -35 and -10 elements are indicated, and the mutated -10 box (-10mut) is shown above. PE analysis results are shown for total RNA extracted from *E. coli* cells carrying the indicated plasmids. Three 5' ends at position +309, +310, and +311 are indicated by asterisks. The -10 mutation shows only a 5' end at +311.

ATG81 affects only the production of VirF₃₀, and not that of VirF₂₁. Thus, the two forms are independently translated (Fig. 3B); consequently, a derivative with either the FS mutation or the M81L substitution gives only VirF₂₁ or only VirF₃₀, respectively. β -Galactosidase fusion and immunoblot analyses (Fig. 3C)

showed that the expression level of $VirF_{21}$ under our experimental conditions is generally lower than that of $VirF_{30}$.

 $VirF_{21}$ is clearly not functionally redundant with $VirF_{30}$. Unlike $VirF_{30}$, it does not restore the expression of VirF-regulated genes in a *virF*-defective *S. flexneri* mutant (Fig. 4). Instead, overexpress-



FIG 7 Analysis of VirF₂₁ translation from the *virF* R2 transcript. (A) Primer extension analysis of total RNA from BL21(DE3) cells carrying pAC-T730FT or pACT7-HH-21-FT with or without induction using IPTG. The arrowhead indicates the position of hammerhead cleavage. (B) Western blot analysis of total protein extracts from strain BL21(DE3) cells carrying pAC-T730FT or pACT7-HH-21-FT, with or without induction by IPTG. Shown is translation of both VirF₃₀ and VirF₂₁ in the presence of pACT7-HH-21-FT or of only VirF₂₁ in the presence of pACT7-HH-21-FT. (C) *In vitro* translation of *virF* R1-3XFT (start, + 1) and R2-3XFT (start, + 309) mRNAs. Both VirF₃₀ and VirF₂₁ were translated from *virF* R1-3XFT, but only VirF₂₁ was translated from *virF* R2-3XFT mRNA. Asterisk, unspecific cross-hybridization with a protein in the extract. In the blot on the right, we included *ompF* mRNA as an internal canonical, SD-dependent translation control. (D) Toeprint assay results with +1 (full-length) and +309 (leaderless) *virF* mRNAs. The mRNAs were incubated alone (control; lanes 1 and 4), with 30S (lanes 2 and 5), or with 30S and tRNA^{-fMet} (lanes 3 and 6). A specific toeprint was observed on full-length mRNA (+1) near the 5' end (black circle) (compare with Fig. 2C). A second toeprint, specific to the llmRNA, is at position +326 (black asterisk). Sequencing ladders were generated with the same 5'-end-labeled primer.

sion of VirF₂₁ negatively autoregulates *virF* expression, reducing intracellular levels of VirF₃₀ and causing reduced *virB* expression (Fig. 5B). This negative autoregulation is likely due to VirF₂₁ binding to the *virF* promoter, as indicated by the position of a DNase I footprint (Fig. 5E) which is predicted to interfere with RNA polymerase access.

An arrangement based on a smaller protein controlling a larger one, with both of them encoded by the same gene, applies to Tn5 transposase (31). The form of Tn5 transposase lacking the first 55 amino acids posttranslationally forms nonproductive complexes with transposase, thus blocking its activity at IS50 inverted repeats (31). Superficially similar in setup, the shorter VirF₂₁ also lacks a large N-terminal portion of the longer VirF₃₀ protein, but here, the shorter form alone is sufficient to exert control at the level of *virF* transcription (Fig. 5A). Though the known C-terminal DNA-binding domain is present in both VirF variants, our data suggest different DNA recognition preferences. Further work will test whether N-terminal sequences affect binding properties of VirF₃₀ and whether protein folding differences in the shared domain can account for the observed specificity differences.

Since VirF₃₀ and VirF₂₁ originate from differential translation, we investigated the *virF* transcripts in more detail. Long (R1) and shorter (R2) *virF* mRNAs of lengths compatible with VirF₃₀ and VirF₂₁ were detected (Fig. 6A). Evaluation of deletions in the region upstream of the VirF₂₁ ORF, along with PE analyses, identified a new gene-internal *virF* promoter that drives the transcription of the *virF* R2 mRNA. *In vivo* and *in vitro* data support that the leaderless R2 is translated into VirF₂₁; plasmid vectors encoding R2 (start site, +309) support *in vivo* translation of VirF₂₁ (Fig. 7B). Moreover, leaderless translation of VirF₂₁ by R2 also occurs *in vitro* (Fig. 7C), and initiation complex formation occurs at the appropriate position (Fig. 7D).

In recent years, noncanonical translation initiation mechanisms have been reported, including so-called leaderless transcripts, i.e., those lacking a 5'-untranslated region (UTR) and an SD sequence (32–34). Most leaderless genes identified so far in *E. coli* reside in mobile DNA, including λ , P2, and Tn*1721*. The *virF* gene is also located within an IS-rich region on an extrachromosomal element, the large *Shigella*/EIEC invasive plasmid (9). Sequencing data for bacteria and archaea suggest that a leaderless model may not be uncommon (35, 36).

The mechanisms underlying synthesis and translation of llm-RNAs are not yet fully understood. Vesper et al. (37) showed that induction of the MazEF toxin-antitoxin (TA) system in E. coli produces a leaderless mRNA population and, simultaneously, specialized "stress" ribosomes with a preference to translate proteins from llmRNAs. The endoribonuclease MazF cleaves singlestranded mRNAs, sometimes at ACA sequences upstream of AUG start codons, generating llmRNA. MazF also cleaves 16S rRNA, removing the anti-SD sequence required for translation on canonical mRNAs. Thereby, a subpopulation of ribosomes is generated for selective translation on llmRNA (37). It is well established that Shigella bacteria sense and respond to environmental conditions within and outside the host, with corresponding reprogramming of transcription. Since VirF₂₁ modulates the intracellular level of VirF, this suggests that the transcription of the leaderless R2 mRNA could occur under conditions where the activation of the virulence program is undesirable. A possible coupling between stress conditions that might promote changes in R2 virF mRNA

transcription and/or preferential translation of leaderless R2 mRNA and effects on virulence gene regulation is an exciting possibility that we intend to pursue. In particular, the environmental cues that may regulate transcription of the shorter *virF* mRNA, and the translation of VirF₂₁ from the llmRNA under stress and infection-relevant conditions, will be addressed. In summary, this study has added new, entirely unexpected elements to the complex regulation of the *Shigella* virulence system and of its major regulator, the VirF protein.

MATERIALS AND METHODS

Oligodeoxyribonucleotides. Oligodeoxynucleotides used in this study (see Table S1 in the supplemental material) were purchased from Metabion.

Bacterial strains and general methods. Strains used in this study are listed in Table S2 in the supplemental material. Cloning was performed with strain DH10b. *E. coli* strain P90C λ B was obtained by transferring a P_{virB}-lacZ fusion from plasmid pRS415 via homologous recombination to the *lac* transducing phage λ RS45 and then integration (38) into the the λ *att* site of *E. coli* P90C λ A was previously described (see Table S2). Strains M90T-F3xFT and M90T Fd($\Delta virF$) were previously constructed (21).

Bacteria were grown aerobically in LB medium at 37°C. Antibiotics and chemicals were used at the following concentrations: ampicillin, 50 μ g/ml; chloramphenicol, 25 μ g/ml; kanamycin, 30 μ g/ml; streptomycin, 10 μ g/ml; tetracycline, 5 μ g/ml; 5-bromo-4-chloro-3-indolyl- β -Dgalactopyranoside, 20 mg/ml. β -Galactosidase assays were performed as described elsewhere (39). Reported values represent the means of at least three separate measurements. DNA isolation, PCR, restriction digests, cloning, and other DNA manipulation methods were performed as described previously (39). Plasmids are listed in Table S3 in the supplemental material. In addition, plasmid constructions are detailed in Text S1 in the supplemental material.

Analysis of virF mRNA. S. flexneri M90T Fd ($\Delta virF$) (Table S2) cells with or without pMYSH6504 were grown in LB broth at 37°C to an optical density at 600 nm of 0.4 to 0.5. Total RNA extraction and Northern blot assays with an α -³²P-labeled virF-specific probe were performed as described elsewhere (21). Loading controls entailed rRNA staining. Radioactivity was quantified using a Typhoon 9200 instrument (GE Healthcare).

qRT-PCR was performed using Power SYBR green PCR master mix on a 7300 real-time PCR system (Applied Biosystems) as described previously (19). The levels of *virB*, *icsA*, and *lacZ* transcripts were analyzed using the $2-\Delta\Delta CT$ (cycle threshold $[C_T]$) method (40), and results are reported that the fold increase relative to the reference. Primers for *mdh* (endogenous control) and for *virB*, *icsA*, and *lacZ* transcripts were designed by using Primer Express software v2.0 and validated. The following oligonucleotides were used (see Table S1 in the supplemental material): *mdhQF/mdhQR*, *virBQF/virBQR*, *icsAQL/icsAQR*, and *lacZQF/lacZQR*.

Primer extensions. Total RNA from exponentially growing plasmidcarrying *E. coli* strains was extracted (41). Total RNA (10 to 20 μ g) was hybridized with 5'-³²P-labeled ML-512 and ML-1314 primers. Reverse transcription experiments were done at 42°C using the reverse transcriptase ImProm-II (Promega). Reaction products were analyzed on an 8% polyacrylamide gel in parallel with sequencing reaction products obtained using the same primers.

DNase I footprinting. Supercoiled plasmid pMYSH6504 (42) (200 ng/sample) was preincubated for 20 min at room temperature with the indicated volumes of the translation mixture, which contained VirF₂₁ or control (no-template) PureSystem reagent in 30 μ l of binding buffer (40 mM Tris-HCl [pH 8.0], 50 mM KCl, 10 mM Mg-acetate, and 0.5 mM dithiothreitol). The DNA-protein complex was incubated with 1 U of DNase I for 40 s. After stopping the reaction, the DNA was precipitated and separately analyzed by primer extension on either DNA strand with 3 pmol of 5'-end-labeled primers ML-U30 or ML-U29 as described pre-

viously (14). The extension products and corresponding sequencing reactions were run on 7% sequencing gels and then fixed for 5 min (10% ethanol–6% acetic acid) and dried. Signals were detected using a phosphorimager screen.

Immunodetection of VirF protein. Western blot assays were carried out as described in reference 21. Incubation with primary antibodies (polyclonal halon anti-VirF, anti-FLAG [Sigma F1804]) was at 4°C in phosphate-buffered saline–Tween (PBS-T) containing 2% dried skim milk. Membranes were washed and incubated at room temperature for 1 h with a secondary anti-rabbit (1:10,000) or anti-mouse (1:5,000) horseradish peroxidase-conjugated antibody in PBS-T. After washing with PBS-T, membranes were developed for 5 min for enhanced chemiluminescence and visualized on a ChemiDoc XRS+ system.

RNA *in vitro* **transcription.** The *virF*-3XFT mRNAs R1 and R2 were transcribed for *in vitro* translation and toeprint assays. For *virF* mRNA R1-3XFT (start, +1), DNA templates contained a T7 promoter (PCR with primers ML-U1/ML-982). For *virF* mRNA R2-3XFT (start, +309), a fragment with a T7 promoter and a hammerhead ribozyme sequence in front of the *virF* sequence was produced by PCR (primers ML-U20/ML-982) on pAC-T7-HH-21-FT as the template (for the hammerhead sequence, Fig. S5 in the supplemental material). DNA templates were *in vitro* transcribed as described in reference 43. To obtain *virF* R2-3XFT, an additional ribozyme self-cleavage step was performed after *in vitro* transcription according methods described previously (44).

Toeprint assay. Toeprint assays were performed as in reference 45. Aliquots of 0.2 pmol of unlabeled *virF*-3xft mRNAs R1 and R2 were annealed with 0.5 pmol 5'-end-labeled ML-U25 or ML-U26 primer in water at 95°C for 1 min and chilled on ice for 2 min. After addition of renaturing buffer (20 mM Tris-HCl [pH 7.5], 20 mM MgCl₂, 100 mM NH₄Cl) and incubation for 10 min at 37°C, 2 pmol of 30S ribosomal subunits was added. After 15 min, 4 pmol of tRNA-fMet was added, and incubation continued for 20 min before cDNA synthesis with avian myeloblastosis virus reverse transcriptase (7.5 U; Invitrogen) and deoxynucleoside triphosphates (100 nM). Reactions were stopped by phenol-chloroform-isoamyl alcohol extraction followed by ethanol precipitation. The cDNAs and sequencing reactions were run on 8% denaturing polyacrylamide gels that were fixed for 5 min (10% ethanol–6% acetic acid) and dried for 1 h at 80°C. Signals were detected using a phoshorimager screen.

In vitro translation. To generate VirF₂₁ for DNase I footprinting, 500 ng of a PCR product containing a T7 promoter and the *virF₂₁* coding sequence was used as the template in the PureSystem Express (New England BioLabs [NEB]) transcription-translation system at 37°C for 4 h. VirF₂₁ translation was analyzed by immunoblotting using anti-VirF antibodies (see Fig. S4 in the supplemental material). For the *in vitro* translation of different *virF* mRNAs (Fig. 7C), each purified transcript was denatured for 2 min at 95°C, chilled for 1 min on ice, diluted in TMN (20 mM Tris-HCl [pH 7.5], 10 mM MgCl₂, 150 mM NaCl), and incubated for 15 min at room temperature. *In vitro* translation (mRNA at 50 nM) was performed with the PureSystem Express (NEB) translation system at 37°C. Translation products were analyzed by immunoblotting with anti-FLAG antibodies.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.01860-16/-/DCSupplemental.

Figure S1, PDF file, 0.1 MB. Figure S2, PDF file, 0.1 MB. Figure S3, PDF file, 0.1 MB. Figure S4, PDF file, 0.1 MB. Figure S5, PDF file, 0.1 MB. Table S1, PDF file, 0.04 MB. Table S2, PDF file, 0.1 MB. Table S3, PDF file, 0.05 MB. Text S1, PDF file, 0.02 MB.

ACKNOWLEDGMENTS

We thank Davide Roncarati for advice on DNaseI footprinting. We also thank Gioacchino Micheli and Mikael Sellin for critical reading of the manuscript. This work was supported by grants from Sapienza University, PRIN2012-WWJSX8K and PTR 24-2016 (B.C. and G.P.) and from the Swedish Research Council (E.G.H.W.).

FUNDING INFORMATION

This work was funded by grants from Sapienza University, from PRIN2012-WWJSX8K and from Institut Pasteur PTR-24-2016 to B.C. and G.P., and from the Swedish Research Council to E.G.H.W. This work, including the efforts of Maria Letizia Di Martino, was supported by grants from Sapienza University, from PRIN2012-WWJSX8K, from Istituto Pasteur Italia, and from the Swedish Research Council. This work, and the efforts of Cédric Romilly, was supported by the Swedish Research Council.

REFERENCES

- 1. The HC, Thanh DP, Holt KE, Thomson NR, Baker S. 2016. The genomic signatures of *Shigella* evolution, adaptation and geographical spread. Nat Rev Microbiol 14:235–250. http://dx.doi.org/10.1038/nrmicro.2016.10.
- Schroeder GN, Hilbi H. 2008. Molecular pathogenesis of *Shigella* spp.: controlling host cell signaling, invasion, and death by type III secretion. Clin Microbiol Rev 21:134–156. http://dx.doi.org/10.1128/CMR.00032-07.
- 3. Ray K, Marteyn B, Sansonetti PJ, Tang CM. 2009. Life on the inside: the intracellular lifestyle of cytosolic bacteria. Nat Rev Microbiol 7:333–340. http://dx.doi.org/10.1038/nrmicro2112.
- Prosseda G, Falconi M, Giangrossi M, Gualerzi CO, Micheli G, Colonna B. 2004. The virF promoter in Shigella: more than just a curved DNA stretch. Mol Microbiol 51:523–537. http://dx.doi.org/10.1046/j.1365 -2958.2003.03848.x.
- Falconi M, Colonna B, Prosseda G, Micheli G, Gualerzi CO. 1998. Thermoregulation of *Shigella* and *Escherichia coli* EIEC pathogenicity. A temperature-dependent structural transition of DNA modulates accessibility of *virF* promoter to transcriptional repressor H-NS. EMBO J 17: 7033–7043. http://dx.doi.org/10.1093/emboj/17.23.7033.
- Ogawa M, Handa Y, Ashida H, Suzuki M, Sasakawa C. 2008. The versatility of *Shigella* effectors. Nat Rev Microbiol 6:11–16. http:// dx.doi.org/10.1038/nrmicro1814.
- Di Martino ML, Falconi M, Micheli G, Colonna B, Prosseda G. 2016. The multifaceted activity of the VirF regulatory protein in the *Shigella* lifestyle. Front Mol Biosci 3:61. http://dx.doi.org/10.3389/fmolb .2016.00061.
- Penno C, Sansonetti P, Parsot C. 2005. Frameshifting by transcriptional slippage is involved in production of MxiE, the transcription activator regulated by the activity of the type III secretion apparatus in *Shigella flexneri*. Mol Microbiol 56:204–214. http://dx.doi.org/10.1111/j.1365 -2958.2004.04530.x.
- Buchrieser C, Glaser P, Rusniok C, Nedjari H, D'Hauteville H, Kunst F, Sansonetti P, Parsot C. 2000. The virulence plasmid pWR100 and the repertoire of proteins secreted by the type III secretion apparatus of *Shigella flexneri*. Mol Microbiol 38:760–771. http://dx.doi.org/10.1046/ j.1365-2958.2000.02179.x.
- Dorman CJ. 2004. H-NS: a universal regulator for a dynamic genome. Nat Rev Microbiol 2:391–400. http://dx.doi.org/10.1038/nrmicro883.
- Dorman CJ. 2007. H-NS, the genome sentinel. Nat Rev Microbiol 5:157–161. http://dx.doi.org/10.1038/nrmicro1598.
- Prosseda G, Mazzola A, Di Martino ML, Tielker D, Micheli G, Colonna B. 2010. A temperature-induced narrow DNA curvature range sustains the maximum activity of a bacterial promoter in vitro. Biochemistry 49: 2778–2785. http://dx.doi.org/10.1021/bi902003g.
- Falconi M, Prosseda G, Giangrossi M, Beghetto E, Colonna B. 2001. Involvement of FIS in the H-NS-mediated regulation of *virF* gene of *Shigella* and enteroinvasive *Escherichia coli*. Mol Microbiol 42:439–452. http://dx.doi.org/10.1046/j.1365-2958.2001.02646.x.
- 14. Tran CN, Giangrossi M, Prosseda G, Brandi A, Di Martino ML, Colonna B, Falconi M. 2011. A multifactor regulatory circuit involving H-NS, VirF and an antisense RNA modulates transcription of the virulence gene *icsA* of *Shigella flexneri*. Nucleic Acids Res 39:8122–8134. http://dx.doi.org/10.1093/nar/gkr521.

- 15. Tobe T, Yoshikawa M, Mizuno T, Sasakawa C. 1993. Transcriptional control of the invasion regulatory gene *virB* of *Shigella flexneri*: activation by *virF* and repression by H-NS. J Bacteriol 175:6142–6149.
- Kane KA, Dorman CJ. 2012. VirB-mediated positive feedback control of the virulence gene regulatory cascade of *Shigella flexneri*. J Bacteriol 194: 5264–5273. http://dx.doi.org/10.1128/JB.00800-12.
- Porter ME, Dorman CJ. 1997. Positive regulation of *Shigella flexneri* virulence genes by integration host factor. J Bacteriol 179:6537–6550.
- Hurt JK, Olgen S, Garcia GA. 2007. Site-specific modification of Shigella flexneri virF mRNA by tRNA-guanine transglycosylase in vitro. Nucleic Acids Res 35:4905–4913. http://dx.doi.org/10.1093/nar/gkm473.
- Giangrossi M, Prosseda G, Tran CN, Brandi A, Colonna B, Falconi M. 2010. A novel antisense RNA regulates at transcriptional level the virulence gene *icsA* of *Shigella flexneri*. Nucleic Acids Res 38:3362–3375. http:// dx.doi.org/10.1093/nar/gkq025.
- Barbagallo M, Di Martino ML, Marcocci L, Pietrangeli P, De Carolis E, Casalino M, Colonna B, Prosseda G. 2011. A new piece of the *Shigella* pathogenicity puzzle: spermidine accumulation by silencing of the *speG* gene. PLoS One 6:e27226. http://dx.doi.org/10.1371/journal .pone.0027226.
- Leuzzi A, Di Martino ML, Campilongo R, Falconi M, Barbagallo M, Marcocci L, Pietrangeli P, Casalino M, Grossi M, Micheli G, Colonna B, Prosseda G. 2015. Multifactor regulation of the MdtJI polyamine transporter in *Shigella*. PLoS One 10:e0136744. http://dx.doi.org/10.1371/ journal.pone.0136744.
- 22. Prosseda G, Di Martino ML, Campilongo R, Fioravanti R, Micheli G, Casalino M, Colonna B. 2012. Shedding of genes that interfere with the pathogenic lifestyle: the *Shigella* model. Res Microbiol 163:399–406. http://dx.doi.org/10.1016/j.resmic.2012.07.004.
- Emanuele AA, Garcia GA. 2015. Mechanism of action and initial, in vitro SAR of an inhibitor of the *Shigella flexneri* virulence regulator VirF. PLoS One 10:e0137410. http://dx.doi.org/10.1371/journal.pone.0137410.
- Porter ME, Dorman CJ. 2002. In vivo DNA-binding and oligomerization properties of the *Shigella flexneri* AraC-like transcriptional regulator VirF as identified by random and site-specific mutagenesis. J Bacteriol 184: 531–539. http://dx.doi.org/10.1128/JB.184.2.531-539.2002.
- Sakai T, Sasakawa C, Makino S, Yoshikawa M. 1986. DNA sequence and product analysis of the *virF* locus responsible for Congo red binding and cell invasion in *Shigella flexneri* 2a. Infect Immun 54:395–402.
- Kato J, Ito K, Nakamura A, Watanabe H. 1989. Cloning of regions required for contact hemolysis and entry into LLC-MK2 cells from *Shigella sonnei* form I plasmid: *virF* is a positive regulator gene for these phenotypes. Infect Immun 57:1391–1398.
- Porter ME, Mitchell P, Roe AJ, Free A, Smith DG, Gally DL. 2004. Direct and indirect transcriptional activation of virulence genes by an AraC-like protein, PerA from enteropathogenic *Escherichia coli*. Mol Microbiol 54:1117–1133. http://dx.doi.org/10.1111/j.1365-2958 .2004.04333.x.
- Fechter P, Chevalier C, Yusupova G, Yusupov M, Romby P, Marzi S. 2009. Ribosomal initiation complexes probed by toeprinting and effect of *trans*-acting translational regulators in bacteria. Methods Mol Biol 540: 247–263. http://dx.doi.org/10.1007/978-1-59745-558-9_18.
- Klucar L, Stano M, Hajduk M. 2010. phiSITE: database of gene regulation in bacteriophages. Nucleic Acids Res 38:D366–D370. http://dx.doi.org/ 10.1093/nar/gkp911.
- Ohashi H, Kanamori T, Shimizu Y, Ueda T. 2010. A highly controllable reconstituted cell-free system—a breakthrough in protein synthesis research. Curr Pharm Biotechnol 11:267–271. http://dx.doi.org/10.2174/ 138920110791111889.

- Mahnke Braam LA, Goryshin IY, Reznikoff WS. 1999. A mechanism for Tn5 inhibition. Carboxyl-terminal dimerization. J Biol Chem 274:86–92. http://dx.doi.org/10.1074/jbc.274.1.86.
- 32. Moll I, Grill S, Gualerzi CO, Bläsi U. 2002. Leaderless mRNAs in bacteria: surprises in ribosomal recruitment and translational control. Mol Microbiol 43:239–246. http://dx.doi.org/10.1046/j.1365-2958 .2002.02739.x.
- Malys N, McCarthy JE. 2011. Translation initiation: variations in the mechanism can be anticipated. Cell Mol Life Sci 68:991–1003. http:// dx.doi.org/10.1007/s00018-010-0588-z.
- Moll I, Engelberg-Kulka H. 2012. Selective translation during stress in Escherichia coli. Trends Biochem Sci 37:493–498. http://dx.doi.org/ 10.1016/j.tibs.2012.07.007.
- Kramer P, Gäbel K, Pfeiffer F, Soppa J. 2014. Haloferax volcanii, a prokaryotic species that does not use the Shine Dalgarno mechanism for translation initiation at 5'-UTRs. PLoS One 9:e94979. http://dx.doi.org/ 10.1371/journal.pone.0094979.
- 36. Shell SS, Wang J, Lapierre P, Mir M, Chase MR, Pyle MM, Gawande R, Ahmad R, Sarracino DA, Ioerger TR, Fortune SM, Derbyshire KM, Wade JT, Gray TA. 2015. Leaderless transcripts and small proteins are common features of the mycobacterial translational landscape. PLoS Genet 11:e1005641. http://dx.doi.org/10.1371/journal.pgen.1005641.
- 37. Vesper O, Amitai S, Belitsky M, Byrgazov K, Kaberdina AC, Engelberg-Kulka H, Moll I. 2011. Selective translation of leaderless mRNAs by specialized ribosomes generated by MazF in *Escherichia coli*. Cell 147: 147–157. http://dx.doi.org/10.1016/j.cell.2011.07.047.
- Simons RW, Houman F, Kleckner N. 1987. Improved single and multicopy lac-based cloning vectors for protein and operon fusions. Gene 53: 85–96. http://dx.doi.org/10.1016/0378-1119(87)90095-3.
- Prosseda G, Latella MC, Casalino M, Nicoletti M, Michienzi S, Colonna B. 2006. Plasticity of the P junc promoter of ISEc11, a new insertion sequence of the IS1111 family. J Bacteriol 188:4681–4689. http://dx.doi.org/ 10.1128/JB.00332-06.
- 40. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. Methods 25:402–408. http://dx.doi.org/10.1006/meth.2001.1262.
- 41. Campilongo R, Di Martino ML, Marcocci L, Pietrangeli P, Leuzzi A, Grossi M, Casalino M, Nicoletti M, Micheli G, Colonna B, Prosseda G. 2014. Molecular and functional profiling of the polyamine content in enteroinvasive *E. coli*: looking into the gap between commensal *E. coli* and harmful *Shigella*. PLoS One 9:e106589. http://dx.doi.org/10.1371/ journal.pone.0106589.
- 42. Sakai T, Sasakawa C, Makino S, Kamata K, Yoshikawa M. 1986. Molecular cloning of a genetic determinant for Congo red binding ability which is essential for the virulence of *Shigella flexneri*. Infect Immun 51: 476–482.
- 43. Romilly C, Lays C, Tomasini A, Caldelari I, Benito Y, Hammann P, Geissmann T, Boisset S, Romby P, Vandenesch F. 2014. A non-coding RNA Promotes bacterial persistence and decreases virulence by regulating a regulator in *Staphylococcus aureus*. PLoS Pathog 10:e1003979. http:// dx.doi.org/10.1371/journal.ppat.1003979.
- 44. Fechter P, Rudinger J, Giegé R, Théobald-Dietrich A. 1998. Ribozyme processed tRNA transcripts with unfriendly internal promoter for T7 RNA polymerase: production and activity. FEBS Lett 436:99–103. http:// dx.doi.org/10.1016/S0014-5793(98)01096-5.
- 45. Holmqvist E, Unoson C, Reimegård J, Wagner EG. 2012. A mixed double negative feedback loop between the sRNA MicF and the global regulator Lrp. Mol Microbiol 84:414–427. http://dx.doi.org/10.1111/ j.1365-2958.2012.07994.x.