

**Special Edition**  
"New Frontiers in Tinnitus, Hearing  
Loss and Hyperacusis"

## Editorial

### \*Corresponding author

**Massimo Ralli, MD, PhD**

Otolaryngologist  
Department of Oral and Maxillofacial  
Sciences  
Sapienza University of Rome  
Viale del Policlinico 155  
Rome 00186, Italy  
Tel. +39 333 8200853  
E-mail: [massimo.ralli@uniroma1.it](mailto:massimo.ralli@uniroma1.it)

### Special Edition 3

Article Ref. #: 1000OTLOJSE5e001

### Article History

Received: August 18<sup>th</sup>, 2016

Accepted: August 18<sup>th</sup>, 2016

Published: August 18<sup>th</sup>, 2016

### Citation

Ralli M. New frontiers in tinnitus, hearing loss and hyperacusis. *Otolaryngol Open J.* 2016; SE(5): Se1-Se2. doi: [10.17140/OTLOJ-SE-5-e001](https://doi.org/10.17140/OTLOJ-SE-5-e001)

### Copyright

©2016 Ralli M. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# New Frontiers in Tinnitus, Hearing Loss and Hyperacusis

**Massimo Ralli, MD, PhD\***

*Department of Oral and Maxillofacial Sciences, Sapienza University of Rome, Rome 00186, Italy*

Tinnitus, hearing loss and hyperacusis are widely spread conditions for which basic and clinical research will play a vital role in the future for prevention, diagnosis and treatment. Current scientific knowledge supports the presence of a strong pathophysiological and clinical link between the three, that may present singularly or in combination in the same subjects.<sup>1</sup>

Besides a proved correlation with hearing disorders, that often represent the trigger for its onset, tinnitus can also be linked to somatic disorders or psychological factors such as depression, anxiety, obsessive compulsive, mood, conversion, somatoform and other psychiatric disorders. Such association is worth further clinical investigation as psychiatric comorbidity has been found to be present in 14 to 80% of tinnitus patients and could surely represent an important element under the clinical management perspective of such patients.<sup>2</sup>

In 40% of patients with tinnitus, concomitant hyperacusis is present while 86% of patients with hyperacusis also report tinnitus. This testifies the strong connection between the two. It would be certainly interesting to further explore this to highlight the basis of such potential hypersensitivity that could be used for future therapeutic approaches.<sup>3</sup>

Hearing loss can follow noise exposure, aging, acquired or genetic disorders, the use of ototoxic drugs and other pathological conditions of the middle and the inner ear and the auditory pathways; among these, age-related and noise-induced hearing loss represent, in the opinion of the Author, very interesting conditions for future research.

Hearing loss following noise exposure is gaining increasing clinical attention among workers and adolescents; the World Health Organization (WHO) estimates that 1.1 billion young people worldwide are at risk of hearing loss due to unsafe listening practices.<sup>4</sup> Age-related hearing loss will probably be within the top leading causes of disease by 2030 and negatively affects mental health, interpersonal relations, and health-related quality of life (QoL).<sup>5</sup> On the contrary, compared to the past decades, hereditary and congenital hearing loss cases are now being diagnosed earlier thanks to the implementation of screening programs that allow early identification of these conditions; preventive medicine is also having a strong impact on prevalence of drug ototoxicity thanks to more clinical attention to the issue.

In the vast field of hearing loss, therapies are an interesting research field to explore: the progressively increasing knowledge of the pathophysiological mechanisms that underlie cell death and reparation, inner ear genetics, advances in stem cell research and the development of new technologies to deliver therapies to their specific target, such as viral vectors and nanoparticles, certainly represents a solid basis for research in this area.

Although considerable progresses have been made in the recent years, there are still wide-open pathways for research in tinnitus, hearing loss and hyperacusis. Among them, I'd like to put particular focus on basic and clinical research on the predisposing elements to acquired hearing loss in children and adults to favour understanding in prevention and treatment, on the exploration of the pathophysiological basis of tinnitus necessary for future possible treatments, and on the correlation between tinnitus and hyperacusis, an interesting direction that could rise questions and give answers along the way to identify target mechanisms able to

help in therapeutic strategies.

#### REFERENCES

1. Shore SE, Roberts LE, Langguth B. Maladaptive plasticity in tinnitus: Triggers, mechanisms and treatment. *Nat Rev Neurol*. 2016; 12(3): 150-160. doi: [10.1038/nrneurol.2016.12](https://doi.org/10.1038/nrneurol.2016.12)
2. Cianfrone G, Mazzei F, Salviati M, et al. Tinnitus holistic simplified classification (THoSC): A new assessment for subjective tinnitus, with diagnostic and therapeutic implications. *Ann Otol Rhinol Laryngol*. 2015; 124(7): 550-560. doi: [10.1177/0003489415570931](https://doi.org/10.1177/0003489415570931)
3. Schecklmann M, Landgrebe M, Langguth B, TRI Database Study Group. Phenotypic characteristics of hyperacusis in tinnitus. *PLoS One*. 2014; 9(1): e86944. doi: [10.1371/journal.pone.0086944](https://doi.org/10.1371/journal.pone.0086944)
4. Krug E, Cieza MA, Chadha S, et al. Hearing loss due to recreational exposure to loud sounds: A review. *World Health Organization*. 2015. Web site. [http://apps.who.int/iris/bitstream/10665/154589/1/9789241508513\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/154589/1/9789241508513_eng.pdf). Accessed August 17, 2016
5. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006; 3(11): e442. doi: [10.1371/journal.pmed.0030442](https://doi.org/10.1371/journal.pmed.0030442)