

Research Article

Operative Treatment of Chronic Achilles Tendinopathy in Sportive Patients

Matteo Guzzini, Lorenzo Proietti*, Riccardo Maria Lanzetti, Domenico Lupariello, Antonio Ponso, and Andrea Ferretti

Department of Orthopaedics and Traumatology, Sapienza University of Rome, Italy

***Corresponding author**

Lorenzo Proietti, Department of Orthopaedics and Traumatology, Sapienza University of Rome, Via Francesco Duodo, 10, Rome, Italy, Tel: 393934425123; Email: proiettilorenzo@hotmail.com

Submitted: 03 May 2017

Accepted: 12 May 2017

Published: 15 May 2017

ISSN: 2475-9112

Copyright

© 2017 Proietti et al.

OPEN ACCESS

Keywords

- Achilles tendon
- Tendinopathy
- Sportive
- Surgical treatment

Abstract

Introduction: The aim of this study is to present the long-term results of operative treatment of sportive patients with chronic Achilles tendinitis. Furthermore we evaluated the ability of the patients to return to their pre injury level of sports.

Materials and Methods: 90 sportive patients (94 tendons) treated surgically fulfilled the inclusion criteria. 81 patients (81 tendons) were available for follow-up. (Group A) consisted of 31 patients with peritendinitis. (Group B) included 9 patients with pure tendinosis. (Group C) consisted of 34 patients with insertional tendinopathy. (Group D) included 7 patients with insertional tendinopathy associated with peritendinitis, tendinosis

Results and Discussion: At final follow-up there were 61 excellent (75%); 10 good (12, 3%); 3 fair (3,7%) and 7 poor (8,6%) results. In (Group A), there were 24 excellent, 4 good, and 3 poor results. In (Group B), there were 8 excellent and 1 good results. In the (Group C), there were 24 excellent, 3 good, 3 fair and 4 poor results. In the (Group D), there were 5 excellent and 2 good results. Return to sport at the desired level of activity was observed in 66 cases (81,4%), 12 patients (14,8%) return to sport at lower level, and 3 patients (3,7%) was not able to return to sport after surgery at final follow up.

Conclusion: The techniques used lead to acceptable results in sportive patients. Furthermore, the ability of 96,2% of patients to return to sport with the 81,4% of them able to return to their pre-injury level of training and sports confirms that results.

INTRODUCTION

Disease of the Achilles tendon and its contiguous structures are the most common overuse syndromes, not only among athletes but also among people not involved in sports [1,2].

It is often described as an overuse injury seen in athletes and older individuals, with a higher male predilection. The incidence varies from 0.2% in the general population, up to 9% in recreational runners [3-5]. Chronic tendinopathies are most commonly thought to be a result of repetitive overuse injuries, which explains a tenfold increase in Achilles tendon injuries in runners compared to age-matched controls [6]. Achilles tendon pathology is classified in a group of distinct entities according to clinical and histological findings: peritendinitis, tendinosis, insertional tendinopathies, and mixed forms (tendinosis and peritendinitis associated with insertional tendinopathies) [7]. Peritendinitis involves inflammation only in the paratenon, a richly vascularized structure. Pure tendinosis is characterized by areas of degeneration of the tendon tissue, including foci of mucoid, fatty, cartilaginous, or calcified metaplasia, and is

frequently associated with partial or total ruptures of the Achilles tendon. In insertional tendinopathies, micro-tears of the bone tendon junction can occur, resulting in a chronic degenerative process. Moreover, the retro-calcaneal bursa, located between the posterior angle of the calcaneus and the Achilles tendon, may become inflamed and hypertrophic. The prominence of the calcaneus increases local pressure on the bursa, leading to inflammatory reaction and painful swelling on the ventral side of the Achilles tendon.

Because patients frequently experience pain on the medial side of the calcaneal tendon, and between 2 and 7cm above the calcaneal tuberosity, some authors suggest that the course of the plantaris tendon may have an impact on Achilles tendinopathy [8,9], five types of insertion were found for the plantaris tendon into the calcaneal tuberosity based on its shape, its relation to the calcaneal tendon and the exact point of insertion in the calcaneal tuberosity [10].

In spite of rest from exercise and various forms of conservative treatment, some patients with tendinopathy of the

Achilles tendon continue to have symptoms and, thus, operative treatment may be necessary.

The aim of this study is to presents the long-term results of operative treatment of sportive patients with chronic Achilles tendinitis. Furthermore we evaluated the ability of the patients to return to their preinjury level of training and sports.

MATERIALS AND METHODS

Between 1995 and 2010, 90 sportive patients (94 tendons) treated surgically at our institution fulfilled the inclusion criteria. 81 patients (81 tendons) were available for follow-up. Patients with total or gross partial rupture of the Achilles tendon were excluded from the study. Patients' age at surgery ranged from 17 to 54 (average 39) years. 59 were male and 22 female. All patients were involved in competitive or recreational sports prior to surgery. Patient distribution according to levels of sports was 47 competing athletes, 34 active recreational athletes. Duration of pre-operative symptoms ranged from 6 to 60 (average 28) months. All patients before surgery underwent to conservative treatment, like physiotherapy and shock wave, the surgical treatment was proposed after the failure of conservative treatment.

At an average follow-up of 13 (6-21) years, patients were questioned about their level of activity, symptoms compared with pre-operative status, and overall evaluation of the result. A physical examination was also performed. In accordance with Perugia's classification [7], patients were divided into four groups according to clinical as well as surgical findings. The (Group A) consisted of 31 patients with peritendinitis, affecting the tendon from 2 to 6 cm above the calcaneal insertion. The (Group B)

included 9 patients with pure tendinosis. The (Group C) consisted of 34 patients with insertional tendinopathy and symptoms localized to the retrocalcaneal bursa and posterior aspect of the calcaneus at the level of the Achilles insertion. The (group D) included 7 patients with insertional tendinopathy associated with peritendinitis, tendinosis, or both. A classification scale graded from 0 to 4 was used to evaluate subjective symptoms pre- and post-operatively. All patients were requested to estimate their subjective symptoms pre-operatively and the current status of their operated Achilles tendon Table 1. Post-operative results were evaluated according to the criteria used for evaluation of surgical treatment of patellar tendinopathy [11] Table 2.

SURGICAL TECHNIQUE

A longitudinal incision, medial to the Achilles tendon, was used in all cases. The extension of incision and the surgical technique used depended entirely on the observed lesion. In patients with peritendinitis, the tendon sheath was usually found to be hyperaemic, thickened and fibrous, as well as adherent to the tendon. This tissue was freed from the underlying tendon and excised. We were careful to leave the fatty tissue and mesotenon attached anterior to preserve the blood supply. In patients with tendinosis and intratendinous pathological processes, the tendon was carefully inspected and palpated for areas of thickening, defects or softening. A longitudinal division of the tendon was performed and the degenerated focus removed. In nine cases where an old partial tear was found, debridement was performed. The tendon was repaired by side-to-side suture. In cases with advanced degeneration, we reinforced the tendon with flaps of healthy tendon. In patients with insertional tendinopathy, the tendon insertion was carefully inspected. The deepest fibres of the tendon were released from the calcaneus and any calcification or metaplastic tissue removed Figure 1. The retrocalcaneal bursa was completely excised.

Usually it was enlarged and sometimes filled with fibrinous loose bodies. The postero-superior angle of the calcaneus was removed and the edges carefully smoothed. Finally it was ensured that there was no impingement of the calcaneus on the anterior side of the Achilles tendon. Drilling of the junction area and scarification of the tendon near the osteo-tendinous junction was done. In all groups a short leg plaster was used post-operatively for 2-4 weeks with weight bearing permitted after 2 weeks. In cases where excision or tendon repair had been used, immobilization was continued up to 6 weeks. After cast removal motion exercises were started, emphasizing passive dorsiflexion. A gradual program of swimming and stationary bicycling, along with isometric and isotonic strengthening of the calf muscles, was prescribed. Jogging was usually permitted at 8-12 weeks. Full return to a competitive level was allowed by 4 months.

RESULTS AND DISCUSSION

At final follow-up there were 61 excellent (75%); 10 good (12,3%); 3 fair (3,7%) and 7 poor (8,6%) results. When we classified the results according to diagnosis, it was found that in (Group A; 31patients), there were 24 excellent, 4 good, and 3 poor results. In (Group B;9 patients), there were 8 excellent and 1 good results. In the (Group C; 34 patients), there were 24 excellent, 3 good, 3 fair and 4 poor results. In the (Group D;

Table 1: Classification of Achilles tendon tendinopathies according to symptoms.

Stage 0	No Pain
Stage 1	Pain only during sport with no limitation of Sport performance
Stage 2	Pain only during sport, limiting sport participation, and sometimes at rest (particularly on rising in the morning)
Stage 3	Pain also during daily activities
Stage 4	Intolerable and constant pain

Table 2: Criteria used for evaluation of post-operative results.

Excellent	Stage 0 Pain free, able to return to previous level of sports or exertion Full ROM
Good	Stage 1 Occasional pain on exertion or after training, able to return to previous level of sports or exertion Full ROM
Fair	Improvement, but the final result as in Stage 2 or higher pain, but lesser than pre-operatively, participation in sport activities or performance decreased Mild Limitation of Rom
Poor	No improvement, as painful as before the operation, participation in sport activities ended and in exercise or work markedly reduced due to the operated tendon Limitation of ROM




Figure 1 Calcification and metaplastic tissue inside the tendon removed during surgical procedure.

7patients), there were 5 excellent and 2 good results. Return to sport at the desired level of activity was observed in 66 cases (81,4%), 12 patients (14,8%) return to sport at lower level, and 3 patients (3,7%) was not able to return to sport after surgery at final follow up.

Our study confirms excellent/good results in 71 sportive patients (87,3%) with the operative treatment; this data are in line with literature [12,13]. Success rates between 75% and 100% have been reported in many studies with open procedures [14-17]. Also about return to sport the present study confirm excellent results with an overall return to sport in 78 patients (96,2%), with 66 patients (81,4%) at the same or a higher level than before the procedure, also this results are in line with precedent studies, Calder et al showed in their study of 32 patients who underwent plantaris excision that 94% returned to sport by 11 weeks [18]. Alfredson [19] and Pearce et al. [20], have also shown good outcomes following this procedure.

In the treatment of tendinopathy of the Achilles tendon, Snook [21], Kvist [22] and, subsequently, Clancy [23] in their preliminary studies reported good results after simple surgical release by splitting and partial resection of the diseased peritendinous tissue. In these studies there was remarkable discrepancy between poor and good results. Differences in patient selection may explain deviating results. The authors report various surgical procedures and describe various lesions; however, their findings are inconsistent and difficult to interpret due to lack of an adequate description and a generally accepted terminology of the observed changes. Kvist [22] emphasized chronic para-tendonitis as a major cause of Achilles tendinopathy. The study was well documented with respect to changes in the paratenon, and results were excellent in 169 (84%) cases. The treated patients were a large number of top-

ranking athletes in international competitions. Surgery consisted of simple release of the tendon by splitting, and partial resection of the affected peritendinous tissue. Nelen et al., in 1989 [14], presented excellent/good results in 86% of the 93 cases of pure peritendinitis. Treatment consisted in release of the fascia of the lower legs as well as the peritenon. A resection of the affected tendon tissue was performed in 50 cases of pure tendinosis. In 26 cases side-to-side suture was performed, with excellent/good results in 73%. In 24 cases a turned-down tendon flap procedure was performed, with excellent/good results in 87%.

The classification proposed by Perugia et al. [4], and later reported by Puddu et al. [2], which we adopted in the present study, identifies three distinct forms of tendinopathies with different aetiology: peritendinitis (mainly inflammatory), tendinosis (mainly degenerative) and insertional tendinopathy (mainly mechanical). However, a relationship between the two forms affecting the tendon has been found. In previous biopsies of peritendineal tissues taken intra-operatively in Achilles peritendinitis patients, Perugia et al. [4], found marked degenerative and obstructive vascular changes. They suggested that if the tendon was allowed to suffer from hypoxia due to obstructive vascular changes, degenerative changes of the tendon tissue itself may arise. We could speculate that— as in two of the cases reported here—in several unsatisfactory results in Achilles peritendinitis, tendon degeneration with mild partial rupture of the tendon has sometimes been treated as chronic, pure peritendinitis. This suggests that simple release of peritenoneum should probably be performed before any intra-tendineal degeneration occurs. It must be emphasized that, in cases of pure peritendinitis, a surgical release of the paratenon is sufficient, but this is not true for peritendinitis associated with tendinosis. In peritendinitis associated with tendinosis, it is very important to promote tendon healing using multiple longitudinal incisions in order to encourage revascularization.

It is widely accepted that degeneration of the tendon can lead to partial or total rupture and, because small ruptures may be difficult to recognise both clinically and ultrasonographically, a careful inspection of the tendon belly is always recommended. Should a partial tear be identified, adequate repair is recommended. Denstad and Roaas [24] reported about 90% satisfactory results in cases of pure tendinosis with partial Achilles tendon ruptures treated with the resection of the affected tissue, side-to-side suture and a turned-down tendon flap. The results reported by Kvist [22], Nelen [14] and Denstad and Roaas [24] are more in agreement with those presented here. Regarding insertional tendinopathy reports on surgical treatment are difficult to interpret because of lack of adequate and generally accepted terminology. Most authors acknowledge the co-existence of various lesions. A biomechanically altered relationship between calcaneus and the Achilles tendon in insertional tendinopathy has been suggested to be the cause of the condition. Haglund [25] first gave the most accurate description through focusing on anatomical variants of the posterosuperior portion of the calcaneus and its relation to the Achilles bursa and tendon. Friction typically results in secondary inflammation of the bursa in the retrocalcaneal region, especially if there are alterations in the normal bony configuration of the calcaneus [26]. Moreover, muscular over-stress itself may, in

time, produce degenerative and metaplastic changes of the bone-tendon junction, typical for insertional tendinopathies, with microtears of the tendon junction and areas in which the tendon fibres seem to be torn and fragmented in the same way as in epicondylitis, jumper's knee, etc. Some authors have reported good results after a simple partial resection of the posterior upper corner of calcaneus [27,28]. In their opinion there is no difference in pain relief whether a large or small resection is used, but large resections can lead to more severe late complaints [26-28]. It is also recommended that all ridges of the calcaneus be carefully removed and rasped smooth so that no prominence is left posterior. On the basis of our experience, we believe that insertional tendinopathy is a complex condition consisting of various lesions mainly related to overuse and to impingement of the anterior portion of the tendon against the postero-superior corner of calcaneus. Therefore, the anatomical structures surrounding the affected osteo-tendinous junction are extremely important. The surgical treatment should include excision of the retrocalcaneal bursa, resection of the posterior superior corner of calcaneus, removal of degenerated tissues and calcifications, if any, drilling of the junction area and scarification of inferior part the tendon near the osteotendinous junction.

CONCLUSION

The present study shows that the techniques used lead to acceptable results in sportive patients, as also suggested by previous studies. Furthermore, the ability of 96,2% of patients to return to sport with the 81,4% of them able to return to their pre-injury level of training and sports confirms that results.

Our study, although a retrospective study, adhered to these basic guidelines set by such methodology, such as number of patients and, above all, by length of follow-up, which to the best of our knowledge is one of the longest in surgical treatment of Achilles tendinopathies.

REFERENCES

- Perugia L, Postacchini F, Ippolito E. The tendons. Editrices Kurtis. 1986.
- Puddu G, Ippolito E, Postacchini F. A classification of Achilles tendon disease. *Am J Sports Med.* 1976; 4: 145-150.
- Maffulli N, Kader D. Tendinopathy of tendo achillis. *J Bone Joint Surg Br.* 2002; 84: 1-8.
- de Jonge S, van den Berg C, de Vos RJ, van der Heide HJ, Weir A, Verhaar JA, et al. Incidence of midportion Achilles tendinopathy in the general population. *Br J Sports Med.* 2011; 45: 1026-1028.
- Lysholm J, Wiklander J. Injuries in runners. *Am J Sports Med.* 1987; 15: 168-171.
- Rompe JD, Furia JP, Maffulli N. Mid-portion Achilles tendinopathy--current options for treatment. *Disabil Rehabil.* 2008; 30: 1666-1676.
- Perugia L, Ippolito E, Postacchini F. A new approach to the pathology, clinical features and treatment of stress tendinopathy of the Achilles tendon. *Ital J Orthop Traumatol.* 1976; 2: 5-21.
- van Sterkenburg MN, Kerkhoffs GM, van Dijk CN. Good outcome after stripping the plantaris tendon in patients with chronic mid-portion Achilles tendinopathy. *Knee Surg Sports Traumatol Arthrosc.* 2011; 19: 1362-1366.
- van Sterkenburg MN, Kerkhoffs GM, Kleipool RP, Niek van Dijk C. The plantaris tendon and a potential role in mid-portion Achilles tendinopathy: an observational anatomical study. *J Anat.* 2011; 218: 336-341.
- Olewnik L, Wysiadecki G, Polgaj M, Topol M. Anatomic study suggests that the morphology of the plantaris tendon may be related to Achilles tendonitis. *Surg Radiol Anat.* 2017; 39: 69-75.
- Ferretti A, Conteduca F, Camerucci E, Morelli F. Patellar tendinosis: a follow-up study of surgical treatment. *J Bone Joint Surg Am.* 2002; 84: 2179-2185.
- Kvist H, Kvist M. The operative treatment of chronic calcaneal paratenonitis. *J Bone Joint Surg Br.* 1980; 62: 353-357.
- De Carbo WT, Bullock MJ. Midsubstance Tendinopathy, Surgical Management. *Clin Podiatr Med Surg.* 2017; 34: 175-193.
- Nelen G, Martens M, Burssens A. Surgical treatment of chronic Achilles tendinitis. *Am J Sports Med.* 1989; 17: 754-759.
- Schepesis AA, Leach RE. Surgical management of Achilles tendinitis. *Am J Sports Med.* 1987; 15: 308-315.
- Paavola M, Kannus P, Orava S, Pasanen M, Järvinen M. Surgical treatment for chronic Achilles tendinopathy: a prospective seven month follow up study. *Br J Sports Med.* 2002; 36: 178-182.
- Maffulli N, Testa V, Capasso G, Bifulco G, Binfield PM. Results of percutaneous longitudinal tenotomy for Achilles tendinopathy in middle- and long-distance runners. *Am J Sports Med.* 1997; 25: 835-840.
- Calder JD, Freeman R, Pollock N. Plantaris excision in the treatment of non-insertional Achilles tendinopathy in elite athletes. *Br J Sports Med.* 2015; 49: 1532-1534.
- Alfredson H. Midportion Achilles tendinosis and the plantaris tendon. *Br J Sports Med.* 2011; 45: 1023-1025.
- Pearce CJ, Carmichael J, Calder JD. Achilles tendinopathy and plantaris tendon release and division in the treatment of non-insertional Achilles tendinopathy. *Foot Ankle Surg.* 2012; 18: 124-127.
- Snook GA. Achilles tendon tenosynovitis in long-distance runners. *Med Sci Sports.* 1972; 4: 155-158.
- Kvist H, Kvist M. The operative treatment of chronic calcaneal paratenonitis. *J Bone Joint Surg Br.* 1980; 62: 353-357.
- Clancy WG, Neidhart D, Brand RL. Achilles tendonitis in runners: a report of five cases. *Am J Sports Med.* 1976; 4: 46-57.
- Denstad TF, Roaas A. Surgical treatment of partial Achilles tendon rupture. *Am J Sports Med.* 1979; 7: 15-17.
- Haglund P. To the clinic The Achillessehene *Z Orthop.* 1928; 49: 49.
- Heneghan MA, Pavlov H. The Haglund painful heel syndrome. Experimental investigation of cause and therapeutic implications. *Clin Orthop Relat Res.* 1984; 228-234.
- Huber HM. Prominence of the calcaneus: late results of bone resection. *J Bone Joint Surg Br.* 1992; 74: 315-316.
- Jones DC, James SL. Partial calcaneal osteotomy for retrocalcaneal bursitis. *Am J Sports Med.* 1984; 12: 72-73.

Cite this article

Guzzini M, Proietti L, Lanzetti RM, Lupariello D, Ponzo A, et al. (2017) Operative Treatment of Chronic Achilles Tendinopathy in Sportive Patients. *JSM Foot Ankle* 2(2): 1027.