

Autophagy and neurogenic muscle atrophy

Eur J Transl Myol 28 (3): 248-256, 2018

41. Bodine SC, Latres E, Baumhueter S, et al. Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science* 2001;294:1704-08.
42. Stitt TN, Drujan D, Clarke BA, et al. The igf-1/pi3k/akt pathway prevents expression of muscle atrophy-induced ubiquitin ligases by inhibiting foxo transcription factors. *Mol Cell* 2004;14:395-403.
43. Tang H, Inoki K, Lee M, et al. mTORC1 promotes denervation-induced muscle atrophy through a mechanism involving the activation of foxo and e3 ubiquitin ligases. *Sci Signal* 2014;7:ra18-ra18.
44. Zhang Y, Manning BD. mTORC1 signaling activates nrf1 to increase cellular proteasome levels. *Cell Cycle* 2015;14:2011-2017.
45. MacDonald EM, Andres-Mateos E, Mejias R, et al. Denervation atrophy is independent from akt and mtor activation and is not rescued by myostatin inhibition. *Dis Model Mech* 2014;7:471-81.
46. Malavaki CJ, Sakkas GK, Mitrou GI, et al. Skeletal muscle atrophy: disease-induced mechanisms may mask disuse atrophy. *J Muscle Res Cell Motil* 2015;36:405-21.
47. Huber K, Petzold J, Rehfeldt C, et al. Muscle energy metabolism: structural and functional features in different types of porcine striated muscles. *J Muscle Res Cell Motil* 2007;28:249-58.
48. Clavel S, Siffroi-Fernandez S, Coldefy AS, et al. Regulation of the intracellular localization of foxo3a by stress-activated protein kinase signaling pathways in skeletal muscle cells. *Mol Cell Biol* 2010;30:470-80.
49. Diebold I, Petry A, Burger M, et al. NOX4 mediates activation of foxo3a and matrix metalloproteinase-2 expression by urotensin-ii. *Mol Biol Cell* 2011;22:4424-34.
50. Patterson MF, Stephenson GMM, Stephenson DG. Denervation produces different single fiber phenotypes in fast- and slow-twitch hindlimb muscles of the rat. *Am J Physiol Physiol* 2006;291:C518-28.

Submission: July 11, 2018

Revision received: August 10, 2018

Acceptance: August 10, 2018

Non commercial use only