

**DEFINITION, EXISTENCE, STABILITY
AND UNIQUENESS OF THE SOLUTION
TO A SEMILINEAR ELLIPTIC PROBLEM
WITH A STRONG SINGULARITY AT $u = 0$**

DANIELA GIACHETTI, PEDRO J. MARTÍNEZ-APARICIO,
AND FRANÇOIS MURAT

Accepted for publication in Ann. Sc. Norm. Sup. Pisa
version prepared for typesetting
April 8, 2017

ABSTRACT. In this paper we consider a semilinear elliptic equation with a strong singularity at $u = 0$, namely

$$\begin{cases} u \geq 0 & \text{in } \Omega, \\ -\operatorname{div} A(x)Du = F(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

with $F(x, s)$ a Carathéodory function such that

$$0 \leq F(x, s) \leq \frac{h(x)}{\Gamma(s)} \quad \text{a.e. } x \in \Omega, \forall s > 0,$$

with h in some $L^r(\Omega)$ and Γ a $C^1([0, +\infty[)$ increasing function such that $\Gamma(0) = 0$ and $\Gamma'(s) > 0$ for every $s > 0$.

We introduce a notion of solution to this problem in the spirit of the solutions defined by transposition. This definition allows us to prove the existence and the stability of this solution, as well as its uniqueness when $F(x, s)$ is nonincreasing in s .

Contents

Abstract	1
1 Introduction	2
2 Assumptions	9
2.1 Notation	13
3 Definition of a solution to problem (2.1)	14
3.1 The space $\mathcal{V}(\Omega)$ of test functions	14
3.2 Definition of a solution to problem (2.1)	16

Key words and phrases. Semilinear equations at $u = 0$, singularity, existence, stability, uniqueness.

2010 Mathematics Subject Classification. 35J25, 35J67, 35J75.

2	D. GIACHETTI, P.J. MARTÍNEZ-APARICIO, AND F. MURAT	
4	Statements of the existence, stability and uniqueness results	20
5	A priori estimates	21
	5.1 A priori estimate of $G_k(u)$ in $H_0^1(\Omega)$	21
	5.2 A priori estimate of $\varphi DT_k(u)$ in $(L^2(\Omega))^N$ for $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$	
24		
	5.3 Control of the integral $\int_{\{u \leq \delta\}} F(x, u)v$	30
	5.4 A priori estimate of $\beta(u)$ in $H_0^1(\Omega)$	35
6	Proofs of the Stability Theorem 4.2 and of the Existence Theorem 4.1	
38		
	6.1 Proof of the Stability Theorem 4.2	38
	6.2 Proof of the Existence Theorem 4.1	45
7	Comparison Principle and proof of the Uniqueness Theorem 4.3	45
	Appendix A An useful lemma	49
	Acknowledgments	50
	References	50

1. INTRODUCTION

Position of the problem

In the present paper we deal with a semilinear problem with a strong singularity at $u = 0$, which consists in finding a function u which satisfies

$$(1.1) \quad \begin{cases} u \geq 0 & \text{in } \Omega, \\ -\operatorname{div} A(x)Du = F(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded open set of \mathbb{R}^N , $N \geq 1$, where A is a coercive matrix with coefficients in $L^\infty(\Omega)$, and where

$$F : (x, s) \in \Omega \times [0, +\infty[\rightarrow F(x, s) \in [0, +\infty]$$

is a Carathéodory function which satisfies

$$(1.2) \quad 0 \leq F(x, s) \leq \frac{h(x)}{\Gamma(s)} \text{ a.e. } x \in \Omega, \forall s > 0,$$

with

$$(1.3) \quad \begin{cases} h \geq 0, \\ h \in L^r(\Omega), r = \frac{2N}{N+2} \text{ if } N \geq 3, r > 1 \text{ if } N = 2, r = 1 \text{ if } N = 1, \\ \Gamma : s \in [0, +\infty[\longrightarrow \Gamma(s) \in [0, +\infty[\text{ is a } C^1([0, +\infty[) \text{ function} \\ \text{such that } \Gamma(0) = 0 \text{ and } \Gamma'(s) > 0 \forall s > 0. \end{cases}$$

The function F

A model for the function $F(x, s)$ (see a more general model in (2.2) and Remark 2.1 *viii*) below) is given by

$$(1.4) \quad \begin{cases} F(x, s) = f(x) \frac{(a + \sin(\frac{1}{s}))}{\exp(-\frac{1}{s})} + g(x) \frac{(b + \sin(\frac{1}{s}))}{s^\gamma} + l(x) \\ \text{a.e. } x \in \Omega, \forall s > 0, \end{cases}$$

where $\gamma > 0$, $a > 1$, $b > 1$ and where the functions f , g and l are nonnegative. In this model $\frac{(a + \sin(\frac{1}{s}))}{\exp(-\frac{1}{s})}$, as well as $\frac{(b + \sin(\frac{1}{s}))}{s^\gamma}$, are just examples of functions which can be replaced by any singularity $\frac{1}{\Gamma(s)}$ with Γ satisfying (1.3). Note that the behaviour of $F(x, s)$ for $s = 0$ can be very different according to the point x .

Note also that the function $F(x, s)$ is defined only for $s \geq 0$, and that in view of (1.2) and (1.3), $F(x, s)$ is finite for almost everywhere $x \in \Omega$ and for every $s > 0$, but that $F(x, s)$ can exhibit a singularity when $s > 0$ tends to 0 and when $h(x) > 0$. Let us emphasize the fact that the Carathéodory character of the function $F(x, s)$, which can take infinite values when $s = 0$, means in particular that for almost every $x \in \Omega$, the function $s \rightarrow F(x, s)$ is continuous not only for every $s > 0$ but also for $s = 0$.

Note finally that we do not require $F(x, s)$ to be nonincreasing in s , except when we deal with uniqueness and comparison results in the Uniqueness Theorem 4.3 and in Section 7 below.

The case of a mild singularity

In the present paper we consider the case of strong singularities, namely the case where $F(x, s)$ can have any (wild) behavior as s tends to zero, while in our previous paper [17] we restricted ourselves to the

case of mild singularities, namely the case where

$$(1.5) \quad \begin{cases} 0 \leq F(x, s) \leq h(x) \left(\frac{1}{s^\gamma} + 1 \right) & \text{a.e. } x \in \Omega, \forall s > 0 \\ \text{with } 0 < \gamma \leq 1. \end{cases}$$

When the function F satisfies (1.5), our definition of the solution to problem (1.1) is relatively classical, because it consists in looking for a function u such that

$$(1.6 \text{ mild}) \quad \begin{cases} u \in H_0^1(\Omega), \\ u \geq 0 \text{ a.e. in } \Omega, \\ \int_{\Omega} F(x, u)\varphi < +\infty \quad \forall \varphi \in H_0^1(\Omega), \varphi \geq 0, \\ \int_{\Omega} ADu D\varphi = \int_{\Omega} F(x, u)\varphi \quad \forall \varphi \in H_0^1(\Omega), \end{cases}$$

(see [17, Section 3]).

Definition of the solution for a strong singularity

In contrast, the definition of the solution to problem (1.1) that we use in the present paper is less classical. This definition is in our opinion one of the main originalities of the present paper.

We refer the reader to Section 3 below where this definition is given in details, but here we emphasize some of its main features.

When the function F satisfies (1.2) and (1.3), we will say (see Definition 3.6 below) that u is a solution to problem (1.1) if

$$(1.7 \text{ strong}) \quad \begin{cases} i) u \in L^2(\Omega) \cap H_{\text{loc}}^1(\Omega), \\ ii) u(x) \geq 0 \text{ a.e. } x \in \Omega, \\ iii) G_k(u) \in H_0^1(\Omega) \quad \forall k > 0, \\ iv) \varphi T_k(u) \in H_0^1(\Omega) \quad \forall k > 0, \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

where $G_k(s) = (s - k)^+$ and $T_k(s) = \inf(s, k) = s - G_k(s)$ for $s > 0$, and if

(1.8_{strong})

$$\left\{ \begin{array}{l} \forall v \in H_0^1(\Omega) \cap L^\infty(\Omega), v \geq 0, \\ \text{with } -\operatorname{div} {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i (-\operatorname{div} \hat{g}_i) + \hat{f} \text{ in } \mathcal{D}'(\Omega), \\ \text{where } I \text{ is finite, } \hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega), \hat{g}_i \in (L^2(\Omega))^N, \hat{f}_i \in L^1(\Omega), \\ \text{one has} \\ i) \int_{\Omega} F(x, u) v < +\infty, \\ ii) \int_{\Omega} {}^t A(x) Dv DG_k(u) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u)) + \int_{\Omega} \hat{f} T_k(u) = \\ = \langle -\operatorname{div} {}^t A(x) Dv, G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) Dv, T_k(u) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F(x, u) v \quad \forall k > 0, \end{array} \right.$$

where $\langle \langle \cdot, \cdot \rangle \rangle_{\Omega}$ is a notation introduced in (3.3) below.

Note that in this definition every term of (1.8_{strong} ii) makes sense in view of (1.7_{strong} iii) and (1.7_{strong} iv).

This definition (1.7_{strong}), (1.8_{strong}) strongly differs from the definition (1.6_{mild}). Indeed for example the assertion $u \in H_0^1(\Omega)$ of (1.6_{mild}) is replaced in (1.7_{strong}) by $u \in H_{\text{loc}}^1(\Omega)$, $G_k(u) \in H_0^1(\Omega)$ for every $k > 0$ and $\varphi T_k(u) \in H_0^1(\Omega)$ for every $k > 0$ and for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$. Here the assertion $G_k(u) \in H_0^1(\Omega)$ for every $k > 0$ in particular expresses the boundary condition $u = 0$ on $\partial\Omega$ (see Remark 3.7 ii) below.

Actually, the solution u does not in general belong to $H_0^1(\Omega)$, or in other terms u “does not belong to $H^1(\Omega)$ up to the boundary”, when the function F exhibits a strong singularity at $s = 0$. It is indeed proved by A.C. Lazer and P.J. McKenna in [22, Theorem 2] that when $f \in C^\alpha(\bar{\Omega})$, $\alpha > 0$, with $f(x) \geq f_0 > 0$ in Ω , in a domain Ω with $C^{2+\alpha}$ boundary, the solution u to the equation

$$\begin{cases} -\Delta u = \frac{f(x)}{u^\gamma} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

does not belong to $H^1(\Omega)$ when $\gamma > 3$, even if u belongs to $C^{2+\alpha}(\Omega) \cap C^0(\bar{\Omega})$ in this case.

The “space” defined by (1.7_{strong}) in which we look for a solution to (1.1) in the case of a strong singularity is therefore fairly different from

the space $H_0^1(\Omega)$ in which we look for a solution to (1.1) in the case of a mild singularity.

Another important difference between the two definitions appears as far as the partial differential equation in (1.1) is concerned. Indeed, if the formulation in the last line of (1.6_{mild}) is classical, with the use of test functions in $H_0^1(\Omega)$, the equation in (1.1) as formulated in (1.8_{strong}) involves test functions which belong to the vectorial space described in the three first lines of (1.8_{strong}), that we will denote by $\mathcal{V}(\Omega)$ in the rest of the present paper. Actually this space $\mathcal{V}(\Omega)$ consists in functions v such that $-\operatorname{div} {}^t A(x) Dv$ can be put in the usual duality between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$ with $G_k(u)$, and in a “formal duality” (through the notation $\langle\langle \cdot, \cdot \rangle\rangle_\Omega$ introduced in (3.3) below) with $T_k(u)$: indeed, when u satisfies (1.7_{strong}), one writes u as the sum $u = G_k(u) + T_k(u)$, where $G_k(u) \in H_0^1(\Omega)$ and where $\varphi T_k(u) \in H_0^1(\Omega)$ for every $k > 0$ and for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$; on the other hand $-\operatorname{div} {}^t A(x) Dv$, which of course belongs to $H^{-1}(\Omega)$ when $v \in H_0^1(\Omega)$, will be assumed to be the sum of a function $\hat{f} \in L^1(\Omega)$ and of a finite sum of functions $\hat{\varphi}_i(-\operatorname{div} \hat{g}_i)$, with $\hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega)$ and $-\operatorname{div} \hat{g}_i \in H^{-1}(\Omega)$, a fact which allows one to correctly define the “formal duality” $\langle\langle -\operatorname{div} {}^t A(x) Dv, T_k(u) \rangle\rangle_\Omega$ (see (3.3) in Definition 3.2 below).

This is a definition of the solution by transposition in the spirit of those introduced by J.-L. Lions and E. Magenes and by G. Stampacchia. Here again things are fairly different with respect to the case of a mild singularity.

Main results of the present paper

In the framework of the definition (1.7_{strong}), (1.8_{strong}) (i.e. of Definition 3.6 below) we are able to prove that there exists at least a solution to problem (1.1) (see the Existence Theorem 4.1 below). We also prove that this solution is stable with respect to variations of the right-hand side (see the Stability Theorem 4.2 below). Finally, if further to (1.2) and (1.3) we assume that the function $F(x, s)$ is nonincreasing with respect to s , then this solution is unique (see the Uniqueness Theorem 4.3 below).

In brief Definition 3.6 below provides a framework where problem (1.1) is well posed in the sense of Hadamard.

Moreover every solution u to problem (1.1) defined in this sense satisfies the following a priori estimates:

- an a priori estimate of $G_k(u)$ in $H_0^1(\Omega)$ for every $k > 0$ (see Proposition 5.1 below) which is formally obtained by using $G_k(u)$ as test function in (1.1); this implies an a priori estimate of u in $L^2(\Omega)$ (see Remark 5.2 below);
- an a priori estimate of $\varphi DT_k(u)$ in $(L^2(\Omega))^N$ for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$ (see Proposition 5.4 below) which is formally obtained by using $\varphi^2 T_k(u)$ and φ^2 as test functions in (1.1); this implies an a priori estimate of u in $H_{\text{loc}}^1(\Omega)$ (see Remark 5.7 below) and an a priori estimate of $\varphi T_k(u)$ in $H_0^1(\Omega)$ for every $k > 0$ and for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$ (see Remark 5.5 below);
- an a priori estimate of the integral $\int_{\{u \leq \delta\}} F(x, u)v$ for every $v \in \mathcal{V}(\Omega)$, $v \geq 0$, and every $\delta > 0$, which depends only on δ and on v (and also on u through the products $\hat{\varphi}_i Du$) by a constant which tends to zero when δ tends to zero (see Proposition 5.9 and Remark 5.10 below);
- an a priori estimate of $\beta(u)$ in $H_0^1(\Omega)$ (where the function β is defined from the function Γ by $\beta(s) = \int_0^s \sqrt{\Gamma'(t)} dt$) (see Proposition 5.13 below) which is formally obtained by using $\Gamma(u)$ as test function in (1.1).

The (mathematically correct) proofs of all the above a priori estimates are based on Definition (1.7_{strong}), (1.8_{strong}) and on the use of convenient test functions v in (1.8_{strong}).

Literature

There is a wide literature dealing with problem (1.1). We will not pretend to give here a complete list of references and we will concentrate on some papers which seem to us to be the most significant, also referring the interested reader to the references quoted there.

The problem (1.1) was initially proposed in 1960 in the pioneering work [16] of W. Fulks and J.S. Maybe as a model for several physical situations. The problem was then studied by many authors, among which we will quote the works of C.A. Stuart [25], M.G. Crandall, P.H. Rabinowitz and L. Tartar [11], J.I. Diaz, J.M. Morel and L. Oswald [14], M.M. Coclite and G. Palmieri [10], A.C. Lazer and P.J. McKenna [22], Y.S. Choi and P.J. McKenna [7], J. Davila and M. Montenegro [12], C.O. Alves, F.J.S.A. Correa and J.V.A. Goncalves [1], F. Cirstea, M. Ghergu and V. Radulescu [8], G.M. Coclite and M.M. Coclite [9], D. Arcoya and L. Moreno-Mérida [2], and F. Oliva and F. Petitta [23].

In most of the above quoted papers the authors look for a strong solution and use sub- and super-solutions. In particular in [22] A.C. Lazer

and P.J. McKenna work in $C^{2,\alpha}(\Omega)$ and $W^{2,q}(\Omega)$ and use methods of sub- and super-solutions, proving that when $F(x, s) = \frac{f(x)}{s^\gamma}$ with $\gamma > 0$, $f \in C^\alpha(\overline{\Omega})$ and $f(x) \geq f_0 > 0$ in a $C^{2,\alpha}$ domain Ω , then one has $c_1\phi_1(x) \leq u(x)^{\frac{\gamma+1}{2}} \leq c_2\phi_1(x)$ for two constants $0 < c_1 < c_2$, where ϕ_1 is the first (positive) eigenfunction of $-\operatorname{div} A(x)D$ in $H_0^1(\Omega)$; M.G. Crandall, P.H. Rabinowitz and L. Tartar [11] study the behaviours of $u(x)$ and $|Du(x)|$ at the boundary. Let us finally note that C. Stuart [24], as well as M.G. Crandall, P.H. Rabinowitz and L. Tartar [11], do not assume that $F(x, s)$ is nonincreasing in s .

More recently L. Boccardo and L. Orsina studied in [4] the problem in the framework of weak solutions in the sense of distributions. In that paper, the authors address the problem (1.1) with $F(x, s) = \frac{f(x)}{s^\gamma}$, where $\gamma > 0$ and where $f \geq 0$ belongs to $L^1(\Omega)$, or to other Lebesgue's spaces, or to the space of Radon's measures, and they prove existence and regularity as well as non existence results. In this work the strong maximum principle and the nonincreasing character of the function $F(x, s) = \frac{f(x)}{s^\gamma}$ with respect to s play prominent roles. The solution u to problem (1.1) is indeed required to satisfy $u(x) \geq c(\omega, u) > 0$ on every open set ω with $\overline{\omega} \subset \Omega$, and the framework in which the solution is searched for is the Sobolev's space $H_0^1(\Omega)$ or $H_{\text{loc}}^1(\Omega)$. Then L. Boccardo and J. Casado-Díaz proved in [3] the uniqueness of the solutions obtained by approximation and the stability of the solution with respect to the G -convergence of a sequence of matrices $A^\varepsilon(x)$ which are equicoercive and equibounded.

For a variational approach to the problem and extensions to the case of a nonlinear principal part, see A. Canino and M. Degiovanni [5], L.M. De Cave [13], and A. Canino, B. Sciunzi and A. Trombetta [6].

In the very recent preprint [23], F. Oliva and F. Petitta consider the case where $F(x, s) = f(x)g(s)$ with f a nonnegative measure and g a continuous function which is singular at $u = 0$; they prove in particular the existence of a solution $u \in L^1(\Omega) \cap W_{\text{loc}}^{1,1}(\Omega)$ which satisfies the equation in the sense of distributions, and, when g is nonincreasing, the uniqueness of such a solution by using convenient solutions to the adjoint problem.

Contributions of the present paper

In the present paper (as in [17], where mild solutions are considered), we obtain in particular an a priori estimate of the singular term in the region where the solution is close to zero. This estimate, which

is new, is an essential tool in our proof. Moreover, in the case of strong singularities, the main difficulty, with respect to the case of mild singularities, is that only local estimates in the energy space are available for the solutions (see [22]). In order to solve this difficulty, we prove that the solution satisfies $\varphi Du \in (L^2(\Omega))^N$ for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, which is a property of the solution which was not known before. We also introduce a convenient class of test functions, namely the class $\mathcal{V}(\Omega)$ defined in Subsection 3.1 below. The class $\mathcal{V}(\Omega)$ and the method of proof that we use seem to be rather flexible and can be adapted to other situations where other techniques would fail. This is the case for the existence of solutions when the equation involves a zeroth-order term which prevents the use of the strong maximum principle in the equation (see [20]), as well as for the homogenization of singular problems in perforated domains Ω^ε obtained from Ω by removing many small holes, when the Dirichlet boundary condition on $\partial\Omega^\varepsilon$ leads to the appearance of a “strange term” μu in Ω (see [19]).

One of the strong points of the present paper is Definition 3.6. This definition makes problem (1.1) well posed in the sense of Hadamard when the function $F(x, s)$ is nonincreasing in s , and allows us to perform in a mathematically correct way all the formal computations that we want to make on problem (1.1), even when the assumption that the function $F(x, s)$ is nonincreasing in s is not done. Two other strong points are the fact that we do not assume that the function $F(x, s)$ is nonincreasing with respect to s , except as far as uniqueness is concerned, and that we do not use the strong maximum principle.

A weak point of the present paper is however the fact that Definition 3.6 is a definition by transposition in the spirit of J.-L. Lions and E. Magenes and of G. Stampacchia, a feature which makes difficult (if not impossible, except maybe when $p = 2$) to extend it to the case of general nonlinear monotone operators.

2. ASSUMPTIONS

As said in the Introduction we study in this paper solutions to the following singular semilinear problem

$$(2.1) \quad \begin{cases} u \geq 0 & \text{in } \Omega, \\ -\operatorname{div} A(x)Du = F(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where a model for the function $F(x, s)$ is given by (1.4); another more general model is the following one

$$(2.2) \quad \begin{cases} F(x, s) = f(x) \frac{(a + \sin(S(s)))}{\exp(-S(s))} + g(x) \frac{(b + \sin(\frac{1}{s}))}{s^\gamma} + l(x) \\ \text{a.e. } x \in \Omega, \forall s > 0, \end{cases}$$

where $\gamma > 0$, $a > 1$, $b > 1$, where the function S satisfies

$$(2.3) \quad S \in C^1(]0, +\infty[), \quad S'(s) < 0 \quad \forall s > 0, \quad S(s) \rightarrow +\infty \text{ as } s \rightarrow 0,$$

and where the functions f , g and l are nonnegative and belong to $L^r(\Omega)$ with r defined in (2.6 *i*) (see Remark 2.1 *viii*) below).

In this section, we give the precise assumptions that we make on the data of problem (2.1).

We assume that Ω is an open bounded set of \mathbb{R}^N , $N \geq 1$ (no regularity is assumed on the boundary $\partial\Omega$ of Ω), that the matrix A is bounded and coercive, i.e. satisfies

$$(2.4) \quad A(x) \in (L^\infty(\Omega))^{N \times N}, \quad \exists \alpha > 0, \quad A(x) \geq \alpha I \quad \text{a.e. } x \in \Omega,$$

and that the function F satisfies

$$(2.5) \quad \begin{cases} F : (x, s) \in \Omega \times [0, +\infty[\rightarrow F(x, s) \in [0, +\infty] \text{ is a Carathéodory function,} \\ \text{i.e. } F \text{ satisfies} \\ \begin{cases} i) \forall s \in [0, +\infty[, x \in \Omega \rightarrow F(x, s) \in [0, +\infty] \text{ is measurable,} \\ ii) \text{ for a.e. } x \in \Omega, s \in [0, +\infty[\rightarrow F(x, s) \in [0, +\infty] \text{ is continuous,} \end{cases} \end{cases}$$

$$(2.6) \quad \begin{cases} i) \exists h, h(x) \geq 0 \text{ a.e. } x \in \Omega, h \in L^r(\Omega), \\ \text{with } r = \frac{2N}{N+2} \text{ if } N \geq 3, r > 1 \text{ if } N = 2, r = 1 \text{ if } N = 1, \\ ii) \exists \Gamma : s \in [0, +\infty[\rightarrow \Gamma(s) \in [0, +\infty[, \Gamma \in C^1([0, +\infty[), \\ \text{such that } \Gamma(0) = 0 \text{ and } \Gamma'(s) > 0 \quad \forall s > 0, \\ iii) 0 \leq F(x, s) \leq \frac{h(x)}{\Gamma(s)} \text{ a.e. } x \in \Omega, \forall s > 0. \end{cases}$$

Moreover, when we will prove comparison and uniqueness results (Proposition 7.1 and Theorem 4.3), we will assume that $F(x, s)$ is non-increasing in s , i.e. that

$$(2.7) \quad F(x, s) \leq F(x, t) \quad \text{a.e. } x \in \Omega, \quad \forall s, \forall t, \quad 0 \leq t \leq s.$$

Remark 2.1.

- *i*) If a function $\Gamma = \Gamma(s)$ satisfies (2.6 *ii*), then Γ is (strictly) increasing and satisfies $\Gamma(s) > 0$ for every $s > 0$; note that the function Γ can be either bounded or unbounded.

Observe also that if a function $F = F(x, s)$ satisfies (2.6) for $h(x)$ and $\Gamma(s)$, and if $\bar{\Gamma}(s)$ is a function which satisfies (2.6 *ii*) and $\bar{\Gamma}(s) \leq \Gamma(s)$, then $F(x, s)$ satisfies (2.6) for $h(x)$ and $\bar{\Gamma}(s)$.

- *ii*) The function $F = F(x, s)$ is a nonnegative Carathéodory function with values in $[0, +\infty]$ and not only in $[0, +\infty[$. But, in view of condition (2.6 *iii*), for almost every $x \in \Omega$, the function $F(x, s)$ can take the value $+\infty$ only when $s = 0$ (or, in other terms, $F(x, s)$ is finite for almost every $x \in \Omega$ when $s > 0$).
- *iii*) Note that (2.6 *ii*) and (2.6 *iii*) do not impose any restriction on the growth of $F(x, s)$ as s tends to zero. Moreover it can be proved (see [18, Section 3, Proposition 1]) that (2.6) is equivalent to the following assumption

$$(2.8) \quad \begin{cases} \forall k > 0, \exists h_k, h_k(x) \geq 0 \text{ a.e. } x \in \Omega, \\ h_k \in L^r(\Omega) \text{ with } r \text{ as in (2.6 } i) \text{ such that} \\ 0 \leq F(x, s) \leq h_k(x) \text{ a.e. } x \in \Omega, \forall s \geq k. \end{cases}$$

- *iv*) Note that no growth condition is imposed from below on $F(x, s)$ as s tends to zero. Indeed it can be proved (see [18, Section 3, Remark 2]) that for every given functions $G_1(s)$ and $G_2(s)$ which satisfy (2.6), with $h_1(x) = h_2(x) = 1$, $G_1(s) \leq G_2(s)$ and $\frac{G_2(s)}{G_1(s)}$ which tends to infinity as s tends to zero, there exist a function $F(s)$ and two sequences s_n^1 and s_n^2 which tend to zero such that $F(s_n^1) = G_1(s_n^1)$ and $F(s_n^2) = G_2(s_n^2)$.

Of course the growth of $F(x, s)$ as s tends to zero can (strongly) depend on the point $x \in \Omega$.

- *v*) Note that the growth condition (2.6 *iii*) is stated for every $s > 0$, while in (2.5) F is supposed to be a Carathéodory function defined for s in $[0, +\infty[$ and not only in $]0, +\infty[$. Indeed an indeterminacy $\frac{0}{0}$ appears in $\frac{h(x)}{\Gamma(s)}$ when $h(x) = 0$ and $s = 0$, while the growth and Carathéodory assumptions imply that

$$F(x, s) = 0 \quad \forall s \geq 0 \quad \text{a.e. on } \{x \in \Omega : h(x) = 0\}.$$

In contrast, when h is assumed to satisfy $h(x) > 0$ for almost every $x \in \Omega$, it is equivalent to have (2.6 *iii*) for every $s > 0$ or for every $s \geq 0$.

- *vi*) Let us observe that the functions $F(x, s)$ given in examples (1.4) and (2.2) satisfy assumption (2.6); indeed for these examples one has

$$0 \leq F(x, s) \leq \bar{h}(x) \left(\frac{1}{\bar{\Gamma}(s)} + 1 \right)$$

for some $\bar{h}(x)$ and $\bar{\Gamma}(s)$ which satisfy (2.6 *i*) and (2.6 *ii*); taking $\Gamma(s) = \frac{\bar{\Gamma}(s)}{1 + \bar{\Gamma}(s)}$ it is clear that $\Gamma(s)$ satisfies (2.6 *ii*) and that $F(x, s)$ satisfies (2.6).

- *vii*) The C^1 regularity of the function Γ which is assumed in (2.6 *ii*) is used to define the function β which appears in Proposition 5.13 (see (5.55)). The latest (regularity) result of $\beta(u)$ is in turn strongly used in the proofs of the Comparison Principle of Proposition 7.1 and of the Uniqueness Theorem 4.3.

This C^1 regularity could appear as a strong restriction on the function Γ , but it can actually be proved (see [18, Section 3]) that, given a function $\bar{\Gamma}$ such that for some $\delta > 0$ and some $M > 0$

$$\begin{cases} \bar{\Gamma} \in C^0([0, +\infty[), \bar{\Gamma}(0) = 0, \bar{\Gamma}(s) > 0 \quad \forall s > 0, \\ \bar{\Gamma}(s) \geq \delta \quad \forall s \geq M, \end{cases}$$

one can construct a function Γ which satisfies (2.6 *ii*) as well as $\Gamma(s) \leq \bar{\Gamma}(s)$ for every $s > 0$ and then use Remark 2.1 *i*) above.

- *vii*) As far as example (2.2) is concerned, note that when $a > 1$ and $b > 1$ and when the function S satisfies (2.3), the functions $\frac{(a + \sin(S(s)))}{\exp(-S(s))}$ and $\frac{(b + \sin(\frac{1}{s}))}{s^\gamma}$ are positive and continuous in $[0, +\infty[$ (this is no more the case when $a = 1$ or $b = 1$), and therefore satisfy (2.5). Note also that these functions are easily shown to be oscillatory (in the sense that they are not nondecreasing) when $a < \sqrt{2}$ and for every b . Note finally that every function Γ which satisfies (2.6 *ii*) can be written as $\Gamma(s) = \exp(-S(s))$ for a function S which satisfies (2.3). □

Remark 2.2. The function h which appears in hypothesis (2.6 *i*) is an element of $H^{-1}(\Omega)$. Indeed, when $N \geq 3$, the exponent $r = \frac{2N}{N+2}$ is nothing but the Hölder's conjugate $(2^*)'$ of the Sobolev's exponent 2^* , i.e.

$$(2.9) \quad \text{when } N \geq 3, \quad \frac{1}{r} = 1 - \frac{1}{2^*}, \quad \text{where } \frac{1}{2^*} = \frac{1}{2} - \frac{1}{N}.$$

Making an abuse of notation, we will set

$$(2.10) \quad \begin{cases} 2^* = \text{any } p \text{ with } 1 < p < +\infty \text{ when } N = 2, \\ 2^* = +\infty \text{ when } N = 1. \end{cases}$$

With this abuse of notation, h belongs to $L^r(\Omega) = L^{(2^*)'}(\Omega) \subset H^{-1}(\Omega)$ for all $N \geq 1$ since Ω is bounded.

This result is indeed a consequence of Sobolev's, Trudinger Moser's and Morrey's inequalities, which (with the above abuse of notation) assert that

$$(2.11) \quad \|v\|_{L^{2^*}(\Omega)} \leq C_S \|Dv\|_{(L^2(\Omega))^N} \quad \forall v \in H_0^1(\Omega) \text{ when } N \geq 1,$$

where C_S is a constant which depends only on N when $N \geq 3$, which depends on p and on Q when $N = 2$, and which depends on Q when $N = 1$, when Q is any bounded open set such that $\Omega \subset Q$. \square

Remark 2.3. Assumption (2.7), namely the fact that for almost every $x \in \Omega$ the function $s \rightarrow F(x, s)$ is nonincreasing in s , is not of the same nature as assumptions (2.5) and (2.6) on $F(x, s)$.

We will only use assumption (2.7) when proving comparison and uniqueness results, namely Proposition 7.1 and Theorem 4.3. In contrast, all the others results of the present paper, and in particular the existence and stability results stated in Theorems 4.1 and 4.2, as well as the a priori estimates of Section 5, do not use this assumption. \square

2.1. Notation

We denote by $\mathcal{D}(\Omega)$ the space of the $C^\infty(\Omega)$ functions whose support is compact and included in Ω , and by $\mathcal{D}'(\Omega)$ the space of distributions on Ω .

We denote by $\mathcal{M}_b^+(\Omega)$ the space of nonnegative bounded Radon measures on Ω .

Since Ω is bounded, $\|Dw\|_{L^2(\Omega)^N}$ is a norm equivalent to $\|w\|_{H^1(\Omega)}$ on $H_0^1(\Omega)$. We set

$$\|w\|_{H_0^1(\Omega)} = \|Dw\|_{(L^2(\Omega))^N} \quad \forall w \in H_0^1(\Omega).$$

For every $s \in \mathbb{R}$ and every $k > 0$ we define as usual

$$s^+ = \max\{s, 0\}, \quad s^- = \max\{0, -s\},$$

$$T_k(s) = \max\{-k, \min\{s, k\}\}, \quad G_k(s) = s - T_k(s).$$

For every measurable function $l : x \in \Omega \rightarrow l(x) \in [0, +\infty]$ we denote

$$\{l = 0\} = \{x \in \Omega : l(x) = 0\}, \quad \{l > 0\} = \{x \in \Omega : l(x) > 0\}.$$

Finally, in the present paper, we denote by φ and $\overline{\varphi}$ functions which belong to $H_0^1(\Omega) \cap L^\infty(\Omega)$, while we denote by ϕ and $\overline{\phi}$ functions which belong to $\mathcal{D}(\Omega)$.

3. DEFINITION OF A SOLUTION TO PROBLEM (2.1)

3.1. The space $\mathcal{V}(\Omega)$ of test functions

In order to introduce the notion of solution to problem (2.1) that we will use in the present paper, we define the following space $\mathcal{V}(\Omega)$ of test functions and a notation.

Definition 3.1. The space $\mathcal{V}(\Omega)$ is the space of the functions v which satisfy

$$(3.1) \quad v \in H_0^1(\Omega) \cap L^\infty(\Omega),$$

$$(3.2) \quad \begin{cases} \exists I \text{ finite, } \exists \hat{\varphi}_i, \exists \hat{g}_i, i \in I, \exists \hat{f}, \text{ with} \\ \hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega), \hat{g}_i \in (L^2(\Omega))^N, \hat{f} \in L^1(\Omega), \\ \text{such that } -\operatorname{div} {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i (-\operatorname{div} \hat{g}_i) + \hat{f} \text{ in } \mathcal{D}'(\Omega). \end{cases}$$

□

In the definition of $\mathcal{V}(\Omega)$ we use the notation $\hat{\varphi}_i$, \hat{g}_i , and \hat{f} to help the reader to identify the functions which enter in the definition of the functions of $\mathcal{V}(\Omega)$.

Note that $\mathcal{V}(\Omega)$ is a vector space.

Definition 3.2. When $v \in \mathcal{V}(\Omega)$ with

$$-\operatorname{div} {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i (-\operatorname{div} \hat{g}_i) + \hat{f} \quad \text{in } \mathcal{D}'(\Omega),$$

where I , $\hat{\varphi}_i$, \hat{g}_i and \hat{f} are as in (3.2), and when y satisfies

$$y \in H^1_{\text{loc}}(\Omega) \cap L^\infty(\Omega) \text{ with } \varphi y \in H_0^1(\Omega) \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega),$$

we use the following notation:

$$(3.3) \quad \langle \langle -\operatorname{div} {}^t A(x) Dv, y \rangle \rangle_\Omega = \sum_{i \in I} \int_\Omega \hat{g}_i D(\hat{\varphi}_i y) + \int_\Omega \hat{f} y.$$

□

Remark 3.3. In (3.2), the product $\hat{\varphi}_i (-\operatorname{div} \hat{g}_i)$ with $\hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega)$ and $\hat{g}_i \in (L^2(\Omega))^N$ is, as usual, the distribution on Ω defined, for every $\phi \in \mathcal{D}(\Omega)$, as

(3.4)

$$\langle \hat{\varphi}_i(-div \hat{g}_i), \phi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \langle -div \hat{g}_i, \hat{\varphi}_i \phi \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i \phi),$$

and the equality $-div {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i(-div \hat{g}_i) + \hat{f}$ holds in $\mathcal{D}'(\Omega)$.

In notation (3.3), the right-hand side is correctly defined since $\hat{\varphi}_i y \in H_0^1(\Omega)$ and since $y \in L^\infty(\Omega)$. In contrast the left-hand side $\langle \langle -div {}^t A Dv, y \rangle \rangle_{\Omega}$ is just a notation. \square

Remark 3.4. If $y \in H_0^1(\Omega) \cap L^\infty(\Omega)$, then $\varphi y \in H_0^1(\Omega)$ for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, so that for every $v \in \mathcal{V}(\Omega)$, $\langle \langle -div {}^t A Dv, y \rangle \rangle_{\Omega}$ is defined. Let us prove that actually one has

$$(3.5) \quad \begin{cases} \forall v \in \mathcal{V}(\Omega), \forall y \in H_0^1(\Omega) \cap L^\infty(\Omega), \\ \langle \langle -div {}^t A(x) Dv, y \rangle \rangle_{\Omega} = \langle -div {}^t A(x) Dv, y \rangle_{H^{-1}(\Omega), H_0^1(\Omega)}. \end{cases}$$

Indeed when $-div {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i(-div \hat{g}_i) + \hat{f}$, one has for every $\phi \in \mathcal{D}(\Omega)$ (see (3.3) and (3.4))

$$\begin{cases} \langle \langle -div {}^t A(x) Dv, \phi \rangle \rangle_{\Omega} = \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i \phi) + \int_{\Omega} \hat{f} \phi = \\ = \langle \sum_{i \in I} \hat{\varphi}_i(-div \hat{g}_i) + \hat{f}, \phi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \\ = \langle -div {}^t A(x) Dv, \phi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \int_{\Omega} {}^t A(x) Dv D\phi, \end{cases}$$

and therefore

$$(3.6) \quad \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i \phi) + \int_{\Omega} \hat{f} \phi = \int_{\Omega} {}^t A Dv D\phi, \quad \forall \phi \in \mathcal{D}(\Omega).$$

For every given function $y \in H_0^1(\Omega) \cap L^\infty(\Omega)$, taking a sequence $\phi_n \in \mathcal{D}(\Omega)$ such that

$$\phi_n \rightarrow y \quad \text{in } H_0^1(\Omega) \text{ strongly, } \phi_n \rightharpoonup y \text{ in } L^\infty(\Omega) \text{ weakly-star,}$$

and passing to the limit in (3.6) with $\phi = \phi_n$, we obtain (3.5). \square

Remark 3.5. Let us give some examples of functions which belong to $\mathcal{V}(\Omega)$.

- *i)* If $\varphi_1, \varphi_2 \in H_0^1(\Omega) \cap L^\infty(\Omega)$, then $\varphi_1 \varphi_2 \in \mathcal{V}(\Omega)$. Indeed $\varphi_1 \varphi_2 \in H_0^1(\Omega) \cap L^\infty(\Omega)$, and in the sense of distributions, one has

$$(3.7) \quad -div {}^t A(x) D(\varphi_1 \varphi_2) = \hat{\varphi}_1(-div \hat{g}_1) + \hat{\varphi}_2(-div \hat{g}_2) + \hat{f} \quad \text{in } \mathcal{D}'(\Omega),$$

with $\hat{\varphi}_1 = \varphi_1 \in H_0^1(\Omega) \cap L^\infty(\Omega)$, $\hat{\varphi}_2 = \varphi_2 \in H_0^1(\Omega) \cap L^\infty(\Omega)$,
 $\hat{g}_1 = {}^tA(x)D\varphi_2 \in L^2(\Omega)^N$, $\hat{g}_2 = {}^tA(x)D\varphi_1 \in (L^2(\Omega))^N$, and $\hat{f} =$
 $-{}^tA(x)D\varphi_2D\varphi_1 - {}^tA(x)D\varphi_1D\varphi_2 \in L^1(\Omega)$.

- *ii*) In particular, if $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, then $\varphi^2 \in \mathcal{V}(\Omega)$, with

$$(3.8) \quad -\operatorname{div} {}^tA(x)D\varphi^2 = \hat{\varphi}(-\operatorname{div} \hat{g}) + \hat{f} \text{ in } \mathcal{D}'(\Omega),$$

with $\hat{\varphi} = 2\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, $\hat{g} = {}^tA(x)D\varphi \in (L^2(\Omega))^N$ and
 $\hat{f} = -2{}^tA(x)D\varphi D\varphi \in L^1(\Omega)$.

- *iii*) If $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$ and has compact support, then $\varphi \in \mathcal{V}(\Omega)$.
Indeed

$$(3.9) \quad -\operatorname{div} {}^tA(x)D\varphi = \bar{\phi}(-\operatorname{div} {}^tA(x)D\varphi) \text{ in } \mathcal{D}'(\Omega),$$

for every $\bar{\phi} \in \mathcal{D}(\Omega)$, with $\bar{\phi} = 1$ on the support of φ .

- *iv*) In particular every $\phi \in \mathcal{D}(\Omega)$ belongs to $\mathcal{V}(\Omega)$.

□

3.2. Definition of a solution to problem (2.1)

We now give the definition of a solution to problem (2.1) that we will use in the present paper.

Definition 3.6. Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). We say that u is a solution to problem (2.1) if u satisfies

$$(3.10) \quad \begin{cases} i) u \in L^2(\Omega) \cap H_{\text{loc}}^1(\Omega), \\ ii) u(x) \geq 0 \text{ a.e. } x \in \Omega, \\ iii) G_k(u) \in H_0^1(\Omega) \quad \forall k > 0, \\ iv) \varphi T_k(u) \in H_0^1(\Omega) \quad \forall k > 0, \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

(3.11)

$$\left\{ \begin{array}{l} \forall v \in \mathcal{V}(\Omega), v \geq 0, \\ \text{with } -\operatorname{div} {}^t A(x) Dv = \sum_{i \in I} \hat{\varphi}_i (-\operatorname{div} \hat{g}_i) + \hat{f} \text{ in } \mathcal{D}'(\Omega), \\ \text{where } \hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega), \hat{g}_i \in (L^2(\Omega))^N, \hat{f} \in L^1(\Omega), \\ \text{one has} \\ i) \int_{\Omega} F(x, u) v < +\infty, \\ ii) \int_{\Omega} {}^t A(x) Dv DG_k(u) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u)) + \int_{\Omega} \hat{f} T_k(u) = \\ = \langle -\operatorname{div} {}^t A(x) Dv, G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) Dv, T_k(u) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F(x, u) v \quad \forall k > 0. \end{array} \right.$$

□

Remark 3.7.

- *i)* Definition 3.6 is a mathematically correct framework which gives a meaning to the solution to problem (2.1); in contrast (2.1) is only formal.

In this Definition 3.6, the requirement (3.10) is the "space" (which is not a vectorial space) to which the solution should belong, while the requirement (3.11), and specially (3.11 *ii*), expresses the partial differential equation in (2.1) in terms of (non standard) test functions in the spirit of the solutions defined by transposition by J.-L. Lions and E. Magenes and by G. Stampacchia.

- *ii)* Note that the statement (3.10 *iii*) formally contains the boundary condition " $u = 0$ on $\partial\Omega$ ". Indeed $G_k(u) \in H_0^1(\Omega)$ for every $k > 0$ formally implies that " $G_k(u) = 0$ on $\partial\Omega$ ", i.e. " $u \leq k$ on $\partial\Omega$ " for every $k > 0$, which implies " $u = 0$ on $\partial\Omega$ " since $u \geq 0$ in Ω .
- *iii)* In Section 5 below we will obtain a priori estimates for every solution u to problem (2.1) in the sense of Definition 3.6 in the various "spaces" which appear in (3.10).
- *iv)* Note finally that (very) formally, one has

$$\left\{ \begin{array}{l} \langle \langle -\operatorname{div} {}^t A(x) Dv, G_k(u) \rangle \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = \int_{\Omega} (-\operatorname{div} {}^t A(x) Dv) G_k(u) = \\ = \int_{\Omega} v (-\operatorname{div} A(x) DG_k(u)), \end{array} \right.$$

$$\left\{ \begin{aligned} & \langle \langle -\operatorname{div} {}^t A(x) Dv, T_k(u) \rangle \rangle_{\Omega} = \int_{\Omega} (-\operatorname{div} {}^t A(x) Dv) T_k(u) = \\ & = \int_{\Omega} v (-\operatorname{div} A(x) DT_k(u)), \end{aligned} \right.$$

so that (3.11 *ii*) formally means that

$$\left\langle \int_{\Omega} v (-\operatorname{div} {}^t A(x) Du) = \int_{\Omega} F(x, u)v \right\rangle \forall v \in \mathcal{V}(\Omega), v \geq 0.$$

Since every v can be written as $v = v^+ - v^-$ with $v^+ \geq 0$ and $v^- \geq 0$, one has formally (this is formal since we do not know whether v^+ and v^- belong to $\mathcal{V}(\Omega)$ when v belongs to $\mathcal{V}(\Omega)$)

$$\left\langle -\operatorname{div} A(x) Du = F(x, u) \right\rangle.$$

Observe that the above formal computation has no meaning in general, while (3.11 *ii*) has a perfectly correct mathematical sense when $v \in \mathcal{V}(\Omega)$ and when u satisfies (3.10). \square

The following Proposition 3.8 asserts that every solution to problem (2.1) in the sense of Definition 3.6 is a solution to (2.1) in the sense of distributions. Note that Proposition 3.8 does not say anything about the boundary conditions satisfied by u (see also Remark 3.7 *ii*).

Proposition 3.8. *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every solution to problem (2.1) in the sense of Definition 3.6 one has*

$$(3.12) \quad u \geq 0 \text{ a.e. in } \Omega, u \in H_{loc}^1(\Omega), F(x, u) \in L_{loc}^1(\Omega),$$

$$(3.13) \quad -\operatorname{div} A(x) Du = F(x, u) \text{ in } \mathcal{D}'(\Omega).$$

Proof. Since every $\phi \in \mathcal{D}(\Omega)$ belongs to $\mathcal{V}(\Omega)$ (see Remark 3.5 *iv*)), assumption (3.11 *i*) implies that

$$(3.14) \quad \int_{\Omega} F(x, u)\phi < +\infty, \quad \forall \phi \in \mathcal{D}(\Omega), \quad \phi \geq 0,$$

which together with $F(x, u) \geq 0$ implies that $F(x, u) \in L_{loc}^1(\Omega)$.

Therefore (3.12) holds true when u is a solution to problem (2.1) in the sense of Definition 3.6.

For what concerns (3.13), for every $\phi \in \mathcal{D}(\Omega)$ and $\bar{\phi} \in \mathcal{D}(\Omega)$ with $\bar{\phi} = 1$ on the support of ϕ , one has (see (3.9) with $\varphi = \phi$ as far as the

second term is concerned)

$$\begin{cases} \langle -\operatorname{div} {}^t A(x) D\phi, G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) D\phi, T_k(u) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} {}^t A(x) D\phi DG_k(u) + \int_{\Omega} {}^t A(x) D\phi D(\bar{\phi} T_k(u)) = \\ = \int_{\Omega} {}^t A(x) D\phi Du = \int_{\Omega} A(x) Du D\phi, \end{cases}$$

where we have used the fact that $u \in H_{\text{loc}}^1(\Omega)$ and that $\bar{\phi} = 1$ on the support of ϕ . This implies that when u is a solution to problem (2.1) in the sense of Definition 3.6, one has

$$(3.15) \quad \int_{\Omega} A(x) Du D\phi = \int_{\Omega} F(x, u) \phi \quad \forall \phi \in \mathcal{D}(\Omega), \quad \phi \geq 0,$$

namely

$$-\operatorname{div} A(x) Du \geq F(x, u) \text{ in } \mathcal{D}'(\Omega).$$

But (3.15) also implies that

$$\int_{\Omega} A(x) Du D\phi = \int_{\Omega} F(x, u) \phi, \quad \forall \phi \in \mathcal{D}(\Omega), \quad \phi \leq 0,$$

namely

$$-\operatorname{div} A(x) Du \leq F(x, u) \text{ in } \mathcal{D}'(\Omega).$$

This proves (3.13). \square

Remark 3.9. When u satisfies (3.10), then one has

$$(3.16) \quad \varphi Du \in (L^2(\Omega))^N \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega).$$

More precisely, when u satisfies (3.10 *i*) and (3.10 *iii*), assertion (3.10 *iv*) is equivalent to (3.16).

Indeed, if $y \in H_{\text{loc}}^1(\Omega)$ and $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, one has for every $k > 0$

$$\begin{cases} \varphi Dy = \varphi DT_k(y) + \varphi DG_k(y) \text{ in } (\mathcal{D}'(\Omega))^N, \\ D(\varphi T_k(y)) = \varphi DT_k(y) + T_k(y) D\varphi \text{ in } (\mathcal{D}'(\Omega))^N, \end{cases}$$

and therefore

$$\varphi Dy - D(\varphi T_k(y)) = \varphi DG_k(y) - T_k(y) D\varphi \text{ in } (\mathcal{D}'(\Omega))^N.$$

This in particular implies that

$$\varphi Dy - D(\varphi T_k(y)) \text{ belongs to } (L^2(\Omega))^N$$

when $y \in H_{\text{loc}}^1(\Omega)$, $G_k(y) \in H_0^1(\Omega)$ and $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$.

Therefore if $y \in H_{\text{loc}}^1(\Omega)$, $G_k(y) \in H_0^1(\Omega)$ and $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, one has

$$\varphi Dy \in (L^2(\Omega))^N \iff D(\varphi T_k(y)) \in (L^2(\Omega))^N,$$

which in view of Lemma A.1 below and of the inequality $-\varphi k \leq \varphi T_k(u) \leq +\varphi k$ implies, since $\varphi T_k(y) \in L^\infty(\Omega) \subset L^2(\Omega)$, that

$$\varphi D y \in (L^2(\Omega))^N \iff \varphi T_k(y) \in H_0^1(\Omega).$$

Taking $y = u$ gives the equivalence between (3.10 *iv*) and (3.16). \square

4. STATEMENTS OF THE EXISTENCE, STABILITY AND UNIQUENESS RESULTS

In this section we state results of existence, stability and uniqueness of the solution to problem (2.1) in the sense of Definition 3.6.

Theorem 4.1. (Existence). *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then there exists at least one solution u to problem (2.1) in the sense of Definition 3.6.*

The proof of Theorem 4.1 will be done in Subsection 6.2. It relies on a stability result (see Theorem 4.2 below) and on a priori estimates of $\|G_k(u)\|_{H_0^1(\Omega)}$ for every $k > 0$, of $\|\varphi DT_k(u)\|_{(L^2(\Omega))^N}$ for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$ and of $\int_{\{u \leq \delta\}} F(x, u)v$ for every $\delta > 0$ and every $v \in \mathcal{V}(\Omega)$, $v \geq 0$, which are satisfied by every solution u to problem (2.1) in the sense of Definition 3.6 (see Propositions 5.1, 5.4 and 5.9).

Theorem 4.2. (Stability). *Assume that the matrix A satisfies assumption (2.4). Let F_n be a sequence of functions and F_∞ be a function which all satisfy assumptions (2.5) and (2.6) for the same h and the same Γ . Assume moreover that*

$$(4.1) \quad \text{a.e. } x \in \Omega, F_n(x, s_n) \rightarrow F_\infty(x, s_\infty) \text{ if } s_n \rightarrow s_\infty, s_n \geq 0, s_\infty \geq 0.$$

Let u_n be any solution to problem (2.1)_n in the sense of Definition 3.6, where (2.1)_n is the problem (2.1) with $F(x, s)$ replaced by $F_n(x, s)$.

Then there exists a subsequence, still labelled by n , and a function u_∞ , which is a solution to problem (2.1)_{\infty} in the sense of Definition 3.6, such that

$$(4.2) \quad \begin{cases} u_n \rightarrow u_\infty \text{ in } L^2(\Omega) \text{ strongly, in } H_{loc}^1(\Omega) \text{ strongly and a.e. in } \Omega, \\ G_k(u_n) \rightarrow G_k(u_\infty) \text{ in } H_0^1(\Omega) \text{ strongly } \forall k > 0, \\ \varphi T_k(u_n) \rightarrow \varphi T_k(u_\infty) \text{ in } H_0^1(\Omega) \text{ strongly } \forall k > 0, \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

The proof of the Stability Theorem 4.2 will be done in Subsection 6.1.

Note that assumption (4.1) is some type of uniform convergence of the functions $F_n(x, s)$ to the function $F_\infty(x, s)$, which is nevertheless non standard since this function can take the value $+\infty$ for $s = 0$.

Finally, the following uniqueness result is an immediate consequence of the Comparison Principle stated and proved in Section 7 below. Note that both the Uniqueness Theorem 4.3 and the Comparison Principle of Proposition 7.1 are based on the nonincreasing character of the function $F(x, s)$ with respect to s .

Theorem 4.3. (Uniqueness). *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Assume moreover that the function $F(x, s)$ is nonincreasing with respect to s , i.e. satisfies assumption (2.7). Then the solution to problem (2.1) in the sense of Definition 3.6 is unique.*

Remark 4.4. When assumptions (2.4), (2.5), (2.6) as well as (2.7) hold true, Theorems 4.1, 4.2 and 4.3 together assert that problem (2.1) is well posed in the sense of Hadamard in the framework of Definition 3.6. \square

5. A PRIORI ESTIMATES

In this section we state and prove a priori estimates which are satisfied by every solution to problem (2.1) in the sense of Definition 3.6.

5.1. A priori estimate of $G_k(u)$ in $H_0^1(\Omega)$

Proposition 5.1. (A priori estimate of $G_k(u)$ in $H_0^1(\Omega)$). *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every u solution to problem (2.1) in the sense of Definition 3.6, one has*

$$(5.1) \quad \|G_k(u)\|_{H_0^1(\Omega)} = \|DG_k(u)\|_{(L^2(\Omega))^N} \leq \frac{C_S}{\alpha} \frac{\|h\|_{L^r(\Omega)}}{\Gamma(k)} \quad \forall k > 0,$$

where C_S is the (generalized) Sobolev's constant defined in (2.11).

Remark 5.2. Since Ω is bounded, using Poincaré's inequality

$$(5.2) \quad \|y\|_{L^2(\Omega)} \leq C_P(\Omega) \|Dy\|_{(L^2(\Omega))^N} \quad \forall y \in H_0^1(\Omega),$$

and writing $u = T_k(u) + G_k(u)$, one easily deduces from (5.1) that every solution u to problem (2.1) in the sense of Definition 3.6 satisfies the following a priori estimate in $L^2(\Omega)$

$$(5.3) \quad \|u\|_{L^2(\Omega)} \leq k|\Omega|^{\frac{1}{2}} + C_P(\Omega) \frac{C_S}{\alpha} \frac{\|h\|_{L^r(\Omega)}}{\Gamma(k)} \quad \forall k > 0,$$

which, taking $k = k_0$ for some k_0 fixed or minimizing the right-hand side in k , provides an a priori estimate of $\|u\|_{L^2(\Omega)}$ which does not depend on k ; unfortunately minimizing in k does not give an explicit constant for a general function Γ . \square

Remark 5.3. (Formal proof of Proposition 5.1). Taking formally $G_k(u)$ as test function in (2.1) one obtains

$$(5.4) \quad \int_{\Omega} A(x) DG_k(u) DG_k(u) = \int_{\Omega} F(x, u) G_k(u) \quad \forall k > 0.$$

Estimate (5.1) then follows from the growth condition (2.6 *iii*) and from the facts that $G_k(u) = 0$ on the set $\{u \leq k\}$, and that $\Gamma(s)$ is nondecreasing, so that

$$0 \leq F(x, u) G_k(u) \leq \frac{h(x)}{\Gamma(u)} G_k(u) \leq \frac{h(x)}{\Gamma(k)} G_k(u) \quad \text{a.e. } x \in \Omega,$$

and finally from the (generalized) Sobolev's inequality (2.11).

This formal computation will be made mathematically correct in the proof below. \square

Proof of Proposition 5.1.

Define for every k and n with $0 < k < n$ the function $S_{k,n}$ as

$$(5.5) \quad S_{k,n}(s) = \begin{cases} 0 & \text{if } 0 \leq s \leq k, \\ s - k & \text{if } k \leq s \leq n, \\ n - k & \text{if } n \leq s. \end{cases}$$

First step. In this step we will prove that

$$(5.6) \quad S_{k,n}(u) \in \mathcal{V}(\Omega).$$

Observe that for every $n > k$ we have

$$0 \leq S_{k,n}(u) \leq G_k(u) \quad \text{and} \quad |DS_{k,n}(u)| = \chi_{\{k \leq u \leq n\}} |Du| \leq |DG_k(u)|.$$

By (3.10 *iii*) and Lemma A.1 of Appendix A below this implies that

$$(5.7) \quad S_{k,n}(u) \in H_0^1(\Omega) \cap L^\infty(\Omega).$$

Let us prove that (5.6) holds true.

Let $\psi_k : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a C^1 nondecreasing function such that

$$\psi_k(s) = 0 \quad \text{for } 0 \leq s \leq \frac{k}{2} \quad \text{and} \quad \psi_k(s) = 1 \quad \text{for } s \geq k.$$

Since $u \in H_{\text{loc}}^1(\Omega)$ one has

$$D\psi_k(u) = \psi'_k(u) Du = \psi'_k(u) DG_{\frac{k}{2}}(u) \quad \text{in } \mathcal{D}'(\Omega),$$

from which using (3.10 *iii*) we deduce that $D\psi_k(u) \in (L^2(\Omega))^N$. Therefore $\psi_k(u) \in H^1(\Omega) \cap L^\infty(\Omega)$, and then, using again (3.10 *iii*), the inequality

$$0 \leq \psi_k(u) \leq \frac{4}{k} G_{\frac{k}{4}}(u),$$

(which results from $\frac{4}{k}G_{\frac{k}{4}}(s) \geq 1$ when $s \geq \frac{k}{2}$) and Lemma A.1 below, we obtain

$$(5.8) \quad \psi_k(u) \in H_0^1(\Omega) \cap L^\infty(\Omega).$$

On the other hand, in view of (5.7) one has $-\operatorname{div} {}^t A(x) DS_{k,n}(u) \in H^{-1}(\Omega)$, and one easily proves that

$$(5.9) \quad -\operatorname{div} {}^t A(x) DS_{k,n}(u) = \psi_k(u)(-\operatorname{div} {}^t A(x) DS_{k,n}(u)) \quad \text{in } \mathcal{D}'(\Omega),$$

which implies (5.6) with $\hat{\varphi} = \psi_k(u)$, $\hat{g} = {}^t A(x) DS_{k,n}(u)$ and $\hat{f} = 0$.

Second step. Since $S_{k,n}(u) \in \mathcal{V}(\Omega)$ by (5.6) and since $S_{k,n}(u) \geq 0$, we can use $S_{k,n}(u)$ as test function in (3.11 *ii*). We get, using (5.9),

$$\begin{cases} \int_{\Omega} {}^t A(x) DS_{k,n}(u) DG_k(u) + \int_{\Omega} {}^t A(x) DS_{k,n}(u) D(\psi_k(u) T_k(u)) = \\ = \langle -\operatorname{div} {}^t A(x) DS_{k,n}(u), G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \\ + \langle \langle -\operatorname{div} {}^t A(x) DS_{k,n}(u), T_k(u) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F(x, u) S_{k,n}(u), \end{cases}$$

where the second term of the left-hand side is zero since

$$D(\psi_k(u) T_k(u)) = \psi_k(u) DT_k(u) + T_k(u) D\psi_k(u) \quad \text{in } \mathcal{D}'(\Omega),$$

and since $DS_{k,n}(u) = 0$ on $\{u \leq k\}$, while $DT_k(u) = 0$ and $D\psi_k(u) = 0$ on $\{u \geq k\}$. This gives

$$(5.10) \quad \int_{\Omega} A(x) DG_k(u) DS_{k,n}(u) = \int_{\Omega} F(x, u) S_{k,n}(u).$$

Third step. Since $S_{k,n}(s) = 0$ for $s \leq k$, one has, using the growth condition (2.6 *iii*) and the (generalized) Sobolev's inequality (2.11)

$$(5.11) \quad \begin{cases} \int_{\Omega} F(x, u) S_{k,n}(u) \leq \int_{\Omega} \frac{h(x)}{\Gamma(u)} \chi_{\{u > k\}} S_{k,n}(u) \leq \int_{\Omega} \frac{h(x)}{\Gamma(k)} S_{k,n}(u) \leq \\ \leq \frac{\|h\|_{L^r(\Omega)}}{\Gamma(k)} \|S_{k,n}(u)\|_{L^{2^*}(\Omega)} \leq \frac{\|h\|_{L^r(\Omega)}}{\Gamma(k)} C_S \|DS_{k,n}(u)\|_{(L^2(\Omega))^N} \\ \forall k > 0, \forall n > k. \end{cases}$$

With the coercivity (2.4) of the matrix A , (5.10) and (5.11) imply that

$$(5.12) \quad \alpha \|DS_{k,n}(u)\|_{(L^2(\Omega))^N} \leq C_S \frac{\|h\|_{L^r(\Omega)}}{\Gamma(k)} \quad \forall k > 0, \forall n > k.$$

Therefore $S_{k,n}(u)$ is bounded in $H_0^1(\Omega)$ for $k > 0$ fixed independently of $n > k$, and the left-hand side (and therefore the right-hand side) of (5.10) is bounded independently of $n > k$. Since

(5.13)

$$S_{k,n}(u) \rightharpoonup G_k(u) \text{ in } H_0^1(\Omega) \text{ weakly and a.e. } x \in \Omega \text{ as } n \rightarrow +\infty,$$

applying Fatou's Lemma to the right-hand side of (5.10) one deduces that

$$(5.14) \quad \int_{\Omega} F(x, u) G_k(u) < +\infty \quad \forall k > 0.$$

Then using (5.13) in the left-hand side and Lebesgue's dominated convergence in the right-hand side of (5.10), one obtains (5.4).

Estimate (5.1) follows either from (5.12) and (5.13) or from (5.4). \square

5.2. A priori estimate of $\varphi DT_k(u)$ in $(L^2(\Omega))^N$ for $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$

Proposition 5.4. *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every u solution to problem (2.1) in the sense of Definition 3.6 one has*

(5.15)

$$\begin{cases} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N}^2 \leq \\ \leq \frac{32k^2}{\alpha^2} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 \|D\varphi\|_{(L^2(\Omega))^N}^2 + \frac{C_S^2 \|h\|_{L^r(\Omega)}^2}{\alpha^2 \Gamma(k)^2} \|\varphi\|_{L^\infty(\Omega)}^2 \\ \forall k > 0, \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

where C_S is the (generalized) Sobolev's constant defined in (2.11).

Remark 5.5. From the a priori estimate (5.15) and from $D(\varphi T_k(u)) = \varphi DT_k(u) + T_k(u) D\varphi$, one deduces that every solution u to problem (2.1) in the sense of Definition 3.6 satisfies the following a priori estimate of $\varphi T_k(u)$ in $H_0^1(\Omega)$

(5.16)

$$\begin{cases} \|\varphi T_k(u)\|_{H_0^1(\Omega)}^2 = \|D(\varphi T_k(u))\|_{(L^2(\Omega))^N}^2 \leq \\ \leq \left(\frac{64k^2}{\alpha^2} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 + 2k^2 \right) \|D\varphi\|_{(L^2(\Omega))^N}^2 + 2 \frac{C_S^2 \|h\|_{L^r(\Omega)}^2}{\alpha^2 \Gamma(k)^2} \|\varphi\|_{L^\infty(\Omega)}^2 \\ \forall k > 0, \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

\square

Remark 5.6. Using estimates (5.1) of $DG_k(u)$ and (5.15) of $\varphi DT_k(u)$ in $(L^2(\Omega))^N$, as well as $\varphi Du = \varphi DT_k(u) + \varphi DG_k(u)$, and $|DT_k(u)| |DG_k(u)| =$

0 almost everywhere in Ω , one deduces that every solution u to problem (2.1) in the sense of Definition 3.6 satisfies the following a priori estimate of φDu in $(L^2(\Omega))^N$

$$(5.17) \quad \begin{cases} \|\varphi Du\|_{(L^2(\Omega))^N}^2 = \|\varphi DT_k(u)\|_{(L^2(\Omega))^N}^2 + \|\varphi DG_k(u)\|_{(L^2(\Omega))^N}^2 \leq \\ \leq \frac{32k^2}{\alpha^2} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 \|D\varphi\|_{(L^2(\Omega))^N}^2 + 2 \frac{C_S^2}{\alpha^2} \frac{\|h\|_{L^r(\Omega)}^2}{\Gamma(k)^2} \|\varphi\|_{L^\infty(\Omega)}^2 \\ \forall k > 0, \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

which, taking $k = k_0$ for some k_0 fixed or minimizing the right-hand side in k , provides an a priori estimate of $\|\varphi Du\|_{(L^2(\Omega))^N}^2$ which does not depend on k ; unfortunately minimizing in k does not give an explicit constant for a general function Γ . \square

Remark 5.7. Since for every $\phi \in \mathcal{D}(\Omega)$ one has $D(\phi u) = \phi Du + (T_k(u) + G_k(u))D\phi$, which implies that $|D(\phi u)| \leq |\phi Du| + k|D\phi| + \|D\phi\|_{(L^\infty(\Omega))^N} |G_k(u)|$, and then using the inequality $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ and the a priori estimates (5.17) and (5.1) together with Poincaré's inequality (5.2), one deduces that every solution u to problem (2.1) in the sense of Definition 3.6 satisfies the following a priori estimate of u in $H_{\text{loc}}^1(\Omega)$

$$(5.18) \quad \begin{cases} \|\phi u\|_{H_0^1(\Omega)}^2 = \|D(\phi u)\|_{(L^2(\Omega))^N}^2 \leq \\ \leq 3 \left(\frac{32k^2}{\alpha^2} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 \|D\phi\|_{(L^2(\Omega))^N}^2 + 2 \frac{C_S^2}{\alpha^2} \frac{\|h\|_{L^r(\Omega)}^2}{\Gamma(k)^2} \|\phi\|_{L^\infty(\Omega)}^2 + \right. \\ \left. + k^2 \|D\phi\|_{L^2(\Omega)}^2 + C_P^2(\Omega) \frac{C_S^2}{\alpha^2} \frac{\|h\|_{L^r(\Omega)}^2}{\Gamma(k)^2} \|D\phi\|_{(L^\infty(\Omega))^N}^2 \right) \\ \forall k > 0, \quad \forall \phi \in \mathcal{D}(\Omega), \end{cases}$$

which, taking $k = k_0$ for some k_0 fixed or minimizing the right-hand side in k , provides an a priori estimate of $\|\phi u\|_{H_0^1(\Omega)}^2$ for every fixed $\phi \in \mathcal{D}(\Omega)$, i.e. an a priori estimate of $\|u\|_{H_{\text{loc}}^1(\Omega)}^2$, which does not depend on k . \square

Remark 5.8. (Formal proof of Proposition 5.4). The computation that we will perform in the present Remark is formal. We will make it mathematically correct in the proof below.

The idea of the proof of Proposition 5.4 is to formally use $\varphi^2 T_k(u)$ as test function in (2.1), where $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$. We formally get

$$(5.19) \quad \int_{\Omega} A(x) Du DT_k(u) \varphi^2 + 2 \int_{\Omega} A(x) Du D\varphi \varphi T_k(u) = \int_{\Omega} F(x, u) \varphi^2 T_k(u),$$

in the second term of which we write $Du = DT_k(u) + DG_k(u)$. Using the coercivity (2.4) of the matrix A we have

$$(5.20) \quad \begin{cases} \alpha \int_{\Omega} \varphi^2 |DT_k(u)|^2 \leq \\ \leq 2 \left| \int_{\Omega} A(x) DT_k(u) D\varphi \varphi T_k(u) \right| + 2 \left| \int_{\Omega} A(x) DG_k(u) D\varphi \varphi T_k(u) \right| + \\ + \int_{\Omega} F(x, u) \varphi^2 T_k(u) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

In this inequality we will use the estimate

$$(5.21) \quad \begin{cases} 2 \left| \int_{\Omega} A(x) DT_k(u) D\varphi \varphi T_k(u) \right| + 2 \left| \int_{\Omega} A(x) DG_k(u) D\varphi \varphi T_k(u) \right| \leq \\ \leq 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N}. \end{cases}$$

On the other hand, since $0 \leq T_k(u) \leq k$, and using formally φ^2 as test function in (2.1), we have

$$(5.22) \quad \begin{cases} 0 \leq \int_{\Omega} F(x, u) \varphi^2 T_k(u) \leq k \int_{\Omega} F(x, u) \varphi^2 = \\ = k \int_{\Omega} A(x) Du D\varphi^2 = 2k \int_{\Omega} A(x) Du D\varphi \varphi = \\ = 2k \int_{\Omega} A(x) DT_k(u) D\varphi \varphi + 2k \int_{\Omega} A(x) DG_k(u) D\varphi \varphi \leq \\ \leq 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N}. \end{cases}$$

Collecting together (5.20), (5.21) and (5.22) we obtain

$$(5.23) \quad \begin{cases} \alpha \int_{\Omega} \varphi^2 |DT_k(u)|^2 \leq \\ \leq 4k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 4k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N} \\ \forall k > 0, \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

Using Young's inequality in the first term of the right-hand side of (5.23) and the estimate (5.1) of $\|DG_k(u)\|_{(L^2(\Omega))^N}$ in the second term provides an estimate of $\|\varphi DT_k(u)\|_{(L^2(\Omega))^N}^2$. \square

Proof of Proposition 5.4.

First step. By (3.10 *iv*) we have $\varphi T_k(u) \in H_0^1(\Omega) \cap L^\infty(\Omega)$, which implies, using Remark 3.5 *i*), that

$$(5.24) \quad \varphi^2 T_k(u) = \varphi \varphi T_k(u) \in \mathcal{V}(\Omega),$$

with

$$\begin{cases} -\operatorname{div} {}^t A(x) D(\varphi^2 T_k(u)) = \\ = \varphi (-\operatorname{div} {}^t A(x) D(\varphi T_k(u))) - {}^t A(x) D(\varphi T_k(u)) D\varphi + \\ + (\varphi T_k(u)) (-\operatorname{div} {}^t A(x) D\varphi) - {}^t A(x) D\varphi D(\varphi T_k(u)) \quad \text{in } \mathcal{D}'(\Omega). \end{cases}$$

Since $\varphi^2 T_k(u) \in \mathcal{V}(\Omega)$ with $\varphi^2 T_k(u) \geq 0$ we can use $v = \varphi^2 T_k(u)$ as test function in (3.11 *ii*). Denoting by j the value of k which appears in (3.11 *ii*), we get

$$(5.25) \quad \begin{cases} \int_{\Omega} {}^t A(x) D(\varphi^2 T_k(u)) DG_j(u) + \\ + \int_{\Omega} {}^t A(x) D(\varphi T_k(u)) D(\varphi T_j(u)) - \int_{\Omega} {}^t A(x) D(\varphi T_k(u)) D\varphi T_j(u) + \\ + \int_{\Omega} {}^t A(x) D\varphi D(\varphi T_k(u)) T_j(u) - \int_{\Omega} {}^t A(x) D\varphi D(\varphi T_k(u)) T_j(u) = \\ = \langle -\operatorname{div} {}^t A(x) D(\varphi^2 T_k(u)), G_j(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \\ + \langle \langle -\operatorname{div} {}^t A(x) D(\varphi^2 T_k(u)), T_j(u) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F(x, u) \varphi^2 T_k(u) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \quad \forall j > 0, \end{cases}$$

where we observe that the fourth term of the left-hand side makes sense due to the fact that $\varphi T_k(u) \in H_0^1(\Omega) \cap L^\infty(\Omega)$ by (3.10 *iv*), which implies that $(\varphi T_k(u)) T_j(u) \in H_0^1(\Omega) \cap L^\infty(\Omega)$ again by (3.10 *iv*).

Since $u \in H_{\text{loc}}^1(\Omega)$, we can expand in $L_{\text{loc}}^1(\Omega)$ the integrands of each of the 5 terms of the left-hand side of (5.25). One obtains 13 terms

whose integrands belong to $L^1(\Omega)$ since $DG_j(u)$, $\varphi DT_k(u)$ and $\varphi DT_j(u)$ belong to $(L^2(\Omega))^N$. A simple but tedious computation leads from (5.25) to

$$(5.26) \quad \begin{cases} \int_{\Omega} A(x)DT_j(u)DT_k(u)\varphi^2 + \int_{\Omega} A(x)DG_j(u)DT_k(u)\varphi^2 + \\ + 2 \int_{\Omega} A(x)DT_j(u)D\varphi\varphi T_k(u) + 2 \int_{\Omega} A(x)DG_j(u)D\varphi\varphi T_k(u) = \\ = \int_{\Omega} F(x,u)\varphi^2 T_k(u) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \quad \forall j > 0. \end{cases}$$

Taking $j = k$ gives, since $|DG_k(u)||DT_k(u)| = 0$ almost everywhere in Ω ,

$$(5.27) \quad \begin{cases} \int_{\Omega} A(x)DT_k(u)DT_k(u)\varphi^2 + \\ + 2 \int_{\Omega} A(x)DT_k(u)D\varphi\varphi T_k(u) + 2 \int_{\Omega} A(x)DG_k(u)D\varphi\varphi T_k(u) = \\ = \int_{\Omega} F(x,u)\varphi^2 T_k(u) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

This result is nothing but (5.19), which had been formally obtained in Remark 5.8 by taking $\varphi^2 T_k(u)$ as test function in (2.1), but the proof that we just performed is mathematically correct. From (5.27) and the coercivity (2.4) of the matrix A we deduce that

$$(5.28) \quad \begin{cases} \alpha \int_{\Omega} \varphi^2 |DT_k(u)|^2 \leq \\ \leq 2 \left| \int_{\Omega} A(x)DT_k(u)D\varphi\varphi T_k(u) \right| + 2 \left| \int_{\Omega} A(x)DG_k(u)D\varphi\varphi T_k(u) \right| + \\ + \int_{\Omega} F(x,u)\varphi^2 T_k(u) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

which is nothing but (5.20) of the formal proof made in Remark 5.8.

Second step. As far as the first and the second terms of the right-hand side of (5.28) are concerned, we have, as in (5.21),

$$(5.29) \quad \begin{cases} 2 \left| \int_{\Omega} A(x)DT_k(u)D\varphi\varphi T_k(u) \right| + 2 \left| \int_{\Omega} A(x)DG_k(u)D\varphi\varphi T_k(u) \right| \leq \\ \leq 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N}. \end{cases}$$

On the other hand, since for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, φ^2 belongs to $\mathcal{V}(\Omega)$ (see Remark 3.5 *ii*) and since $\varphi^2 \geq 0$, using φ^2 as test function in (3.11 *ii*) gives

$$\begin{cases} \langle -\operatorname{div} {}^t A(x) D\varphi^2, G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) D\varphi^2, T_k(u) \rangle \rangle_\Omega = \\ = \int_\Omega F(x, u) \varphi^2, \end{cases}$$

which gives

$$(5.30) \quad \begin{cases} 2 \int_\Omega A(x) DG_k(u) D\varphi \varphi + 2 \int_\Omega A(x) DT_k(u) D\varphi \varphi = \\ = \int_\Omega F(x, u) \varphi^2 \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

Therefore we have, for the last term of the right-hand side of (5.28),

$$(5.31) \quad \begin{cases} 0 \leq \int_\Omega F(x, u) \varphi^2 T_k(u) \leq k \int_\Omega F(x, u) \varphi^2 = \\ = 2k \int_\Omega A(x) DT_k(u) D\varphi \varphi + 2k \int_\Omega A(x) DG_k(u) D\varphi \varphi \leq \\ \leq 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 2k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N}, \end{cases}$$

which is nothing but (5.22) of the formal proof made in Remark 5.8.

Third step. Collecting together (5.28), (5.29) and (5.31) we have proved that

$$(5.32) \quad \begin{cases} \alpha \int_\Omega \varphi^2 |DT_k(u)|^2 \leq \\ \leq 4k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N} + \\ + 4k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \|DG_k(u)\|_{(L^2(\Omega))^N} \\ \forall k > 0, \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega), \end{cases}$$

which is nothing but (5.23) of the formal proof made in Remark 5.8.

Using Young's inequality $XY \leq \frac{\alpha}{2} X^2 + \frac{1}{2\alpha} Y^2$ in the first term of the right-hand side of (5.32) and estimate (5.1) of $\|DG_k(u)\|_{(L^2(\Omega))^N}$ in

the second term gives

$$(5.33) \quad \begin{cases} \frac{\alpha}{2} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N}^2 \leq \\ \leq \frac{8k^2}{\alpha} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 \|D\varphi\|_{(L^2(\Omega))^N}^2 + \\ + 4k \|A\|_{(L^\infty(\Omega))^{N \times N}} \|D\varphi\|_{(L^2(\Omega))^N} \|\varphi\|_{L^\infty(\Omega)} \frac{C_S \|h\|_{L^r(\Omega)}}{\alpha \Gamma(k)}, \end{cases}$$

in which we use again Young's inequality $\frac{1}{\alpha}XY \leq \frac{1}{2\alpha}X^2 + \frac{1}{2\alpha}Y^2$ in the last term of the right-hand side. This gives

$$(5.34) \quad \begin{cases} \frac{\alpha}{2} \|\varphi DT_k(u)\|_{(L^2(\Omega))^N}^2 \leq \\ \leq \frac{16k^2}{\alpha} \|A\|_{(L^\infty(\Omega))^{N \times N}}^2 \|D\varphi\|_{(L^2(\Omega))^N}^2 + \frac{C_S^2 \|h\|_{L^r(\Omega)}^2}{2\alpha \Gamma(k)^2} \|\varphi\|_{L^\infty(\Omega)}^2, \end{cases}$$

i.e. (5.15).

This completes the proof of Proposition 5.4. \square

5.3. Control of the integral $\int_{\{u \leq \delta\}} F(x, u)v$

In this subsection we prove an a priori estimate (see (5.37) below) which is a key point in the proofs of our results.

For $\delta > 0$, we define the function $Z_\delta : s \in [0, +\infty[\rightarrow Z_\delta(s) \in [0, +\infty[$ by

$$(5.35) \quad Z_\delta(s) = \begin{cases} 1 & \text{if } 0 \leq s \leq \delta, \\ -\frac{s}{\delta} + 2 & \text{if } \delta \leq s \leq 2\delta, \\ 0 & \text{if } 2\delta \leq s. \end{cases}$$

Proposition 5.9. *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every u solution to problem (2.1) in the sense of Definition 3.6 and for every v such that*

$$(5.36) \quad \begin{cases} v \in \mathcal{V}(\Omega), v \geq 0, \\ \text{with } -\operatorname{div} {}^t A(x)Dv = \sum_{i \in I} \hat{\varphi}_i(-\operatorname{div} \hat{g}_i) + \hat{f} \text{ in } \mathcal{D}'(\Omega) \\ \text{where } \hat{\varphi}_i \in H_0^1(\Omega) \cap L^\infty(\Omega), \hat{g}_i \in L^2(\Omega)^N, \hat{f} \in L^1(\Omega), \end{cases}$$

one has

$$(5.37) \quad \begin{cases} \forall \delta > 0, \int_{\Omega} F(x, u) Z_{\delta}(u) v \leq \\ \leq \frac{3}{2} \left(\int_{\Omega} \left| \sum_{i \in I} \hat{g}_i D \hat{\varphi}_i + \hat{f} \right| \right) \delta + \sum_{i \in I} \int_{\Omega} Z_{\delta}(u) \hat{g}_i Du \hat{\varphi}_i. \end{cases}$$

Remark 5.10. Since $Z_{\delta}(s) \geq \chi_{\{u \geq \delta\}}(s)$ for every $s \geq 0$, estimate (5.37) provides an estimate of the integral $\int_{\{u \leq \delta\}} F(x, u) v$ as announced in the title of Subsection 5.3. \square

Remark 5.11. (Formal proof of Proposition 5.9). Estimate (5.37) can be formally obtained by using in (2.1) the test function $Z_{\delta}(u) v$, where Z_{δ} is defined by (5.35) and where v satisfies (5.36). One formally obtains

$$(5.38) \quad \int_{\Omega} F(x, u) Z_{\delta}(u) v = \int_{\Omega} A(x) Du Du Z'_{\delta}(u) v + \int_{\Omega} A(x) Du Dv Z_{\delta}(u),$$

and one observes that, denoting by Y_{δ} the primitive of the function Z_{δ} (see (5.40) below), one has,

$$\begin{cases} \int_{\Omega} A(x) Du Du Z'_{\delta}(u) v \leq 0, \\ \int_{\Omega} A(x) Du Dv Z_{\delta}(u) = \int_{\Omega} A(x) DY_{\delta}(u) Dv = \int_{\Omega} {}^t A(x) Dv DY_{\delta}(u) = \\ = \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i Y_{\delta}(u)) + \int_{\Omega} \hat{f} Y_{\delta}(u) = \\ = \int_{\Omega} \left(\sum_{i \in I} \hat{g}_i D \hat{\varphi}_i + \hat{f} \right) Y_{\delta}(u) + \sum_{i \in I} \int_{\Omega} Z_{\delta}(u) \hat{g}_i Du \hat{\varphi}_i, \end{cases}$$

where one finally uses $0 \leq Y_{\delta}(s) \leq \frac{3}{2} \delta$ to formally obtain estimate (5.37).

These formal computations will be made mathematically correct in the proof below. \square

As a consequence of Proposition 5.9 we have:

Proposition 5.12. *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every u solution to problem (2.1) in the sense of Definition 3.6 one has*

$$(5.39) \quad \int_{\{u=0\}} F(x, u) v = 0 \quad \forall v \in \mathcal{V}(\Omega), v \geq 0.$$

Proof of Proposition 5.12.

Let $\delta > 0$ tend to 0 in (5.37). In the left-hand side one has

$$\int_{\{u=0\}} F(x, u)v \leq \int_{\Omega} F(x, u)Z_{\delta}(u)v,$$

while in the right-hand side $Z_{\delta}(u)$ tends to $\chi_{\{u=0\}}$ almost everywhere in Ω ; but since $u \in H_{\text{loc}}^1(\Omega)$, one has $Du = 0$ almost everywhere on the set $\{u = 0\}$, and therefore, since $\hat{g}_i Du \hat{\varphi}_i \in L^1(\Omega)$, and since $0 \leq Z_{\delta}(u) \leq 1$, one has, by Lebesgue's dominated convergence Theorem,

$$\int_{\Omega} Z_{\delta}(u)\hat{g}_i Du \hat{\varphi}_i \rightarrow 0 \text{ as } \delta \rightarrow 0.$$

This proves (5.39). \square

Proof of Proposition 5.9.

In addition to the function $Z_{\delta}(s)$ defined by (5.35), we define for $\delta > 0$ the functions

$$(5.40) \quad Y_{\delta}(s) = \begin{cases} s & \text{if } 0 \leq s \leq \delta, \\ -\frac{s^2}{2\delta} + 2s - \frac{\delta}{2} & \text{if } \delta \leq s \leq 2\delta, \\ \frac{3}{2}\delta & \text{if } 2\delta \leq s, \end{cases}$$

$$(5.41) \quad R_{\delta}(s) = \begin{cases} 0 & \text{if } 0 \leq s \leq \delta, \\ \frac{s^2}{2\delta} - \frac{\delta}{2} & \text{if } \delta \leq s \leq 2\delta, \\ \frac{3}{2}\delta & \text{if } 2\delta \leq s. \end{cases}$$

Observe that Z_{δ} , Y_{δ} and R_{δ} are Lipschitz continuous and piecewise C^1 functions with

$$(5.42) \quad \begin{cases} Y'_{\delta}(s) = Z_{\delta}(s), \quad Z'_{\delta}(s) = -\frac{1}{\delta}\chi_{\{\delta < s < 2\delta\}}(s), \quad R'_{\delta}(s) = -sZ'_{\delta}(s), \\ Y_{\delta}(s) = sZ_{\delta}(s) + R_{\delta}(s), \quad 0 \leq Y_{\delta}(s) \leq \frac{3}{2}\delta, \quad \forall s \geq 0. \end{cases}$$

First step. In this first step we will prove that for every $\delta > 0$

$$(5.43) \quad \begin{cases} Y_{\delta}(u) \in H_{\text{loc}}^1(\Omega) \cap L^{\infty}(\Omega), \\ \varphi Y_{\delta}(u) \in H_0^1(\Omega) \quad \forall \varphi \in H_0^1(\Omega) \cap L^{\infty}(\Omega), \end{cases}$$

$$(5.44) \quad Z_{\delta}(u)v \in \mathcal{V}(\Omega) \quad \forall v \in \mathcal{V}(\Omega),$$

$$(5.45) \quad R_{\delta}(u) \in H_0^1(\Omega) \cap L^{\infty}(\Omega).$$

Since $u \in H_{\text{loc}}^1(\Omega)$ by (3.10 *i*), one has $Y_\delta(u) \in H_{\text{loc}}^1(\Omega) \cap L^\infty(\Omega)$ with

$$DY_\delta(u) = Y'_\delta(u)Du = Z_\delta(u)Du \text{ in } \mathcal{D}'(\Omega),$$

and for every $\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega)$, one has $\varphi Y_\delta(u) \in H_{\text{loc}}^1(\Omega) \cap L^\infty(\Omega)$ with

$$D(\varphi Y_\delta(u)) = \varphi Z_\delta(u)Du + Y_\delta(u)D\varphi \text{ in } \mathcal{D}'(\Omega),$$

which using (3.16) and the last assertion of (5.42) implies that $D(\varphi Y_\delta(u)) \in (L^2(\Omega))^N$, and therefore that $\varphi Y_\delta(u) \in H^1(\Omega) \cap L^\infty(\Omega)$. Since $0 \leq \varphi Y_\delta(u) \leq \frac{3}{2}\delta\varphi$, and since $\varphi \in H_0^1(\Omega)$, Lemma A.1 of Appendix A below completes the proof of (5.43).

On the other hand $Z_\delta(u) \in H_{\text{loc}}^1(\Omega)$ and

$$DZ_\delta(u) = -\frac{1}{\delta}\chi_{\{\delta < u < 2\delta\}}Du \text{ in } \mathcal{D}'(\Omega).$$

In view of (3.10 *iii*) with $k = \delta$, this implies that $DZ_\delta(u) \in (L^2(\Omega))^N$, and therefore

$$(5.46) \quad Z_\delta(u) \in H^1(\Omega) \cap L^\infty(\Omega),$$

which in turn implies that

$$(5.47) \quad Z_\delta(u)\varphi \in H_0^1(\Omega) \cap L^\infty(\Omega) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega).$$

Consider now $v \in \mathcal{V}(\Omega)$ which satisfies (5.36). Observe that $Z_\delta(u)v \in H_0^1(\Omega) \cap L^\infty(\Omega)$, and that

$$(5.48) \quad \begin{cases} -\text{div } {}^tA(x)D(Z_\delta(u)v) = \\ = Z_\delta(u)(-\text{div } {}^tA(x)Dv) - {}^tA(x)DvDZ_\delta(u) + \\ + v(-\text{div } {}^tA(x)DZ_\delta(u)) - {}^tA(x)DZ_\delta(u)Dv = \\ = \sum_{i \in I} Z_\delta(u)\hat{\varphi}_i(-\text{div } \hat{g}_i) + Z_\delta(u)\hat{f} - {}^tA(x)DvDZ_\delta(u) + \\ + v(-\text{div } {}^tA(x)DZ_\delta(u)) - {}^tA(x)DZ_\delta(u)Dv \text{ in } \mathcal{D}'(\Omega). \end{cases}$$

This proves that $Z_\delta(u)v \in \mathcal{V}(\Omega)$, namely (5.44).

Finally, since $u \in H_{\text{loc}}^1(\Omega)$ by (3.10 *i*), one has

$$R_\delta(u) \in H_{\text{loc}}^1(\Omega) \cap L^\infty(\Omega)$$

with

$$DR_\delta(u) = R'_\delta(u)Du = \frac{u}{\delta}\chi_{\{\delta < u < 2\delta\}}Du \text{ in } \mathcal{D}'(\Omega).$$

Therefore (3.10 *iii*) with $k = \delta$ implies that $DR_\delta(u) \in (L^2(\Omega))^N$, which proves that $R_\delta(u) \in H^1(\Omega) \cap L^\infty(\Omega)$. Then using the inequality $0 \leq R_\delta(u) \leq G_{\frac{\delta}{2}}(u)$, property (3.10 *iii*) with $k = \frac{\delta}{2}$ and Lemma A.1 of

Appendix A below completes the proof of (5.45).

Second step. In this step we fix $\delta > 0$ and k such that

$$(5.49) \quad k \geq 2\delta.$$

Since $Z_\delta(u)v \in \mathcal{V}(\Omega)$ for every $v \in \mathcal{V}(\Omega)$ (see (5.44)) with $Z_\delta(u)v \geq 0$ when $v \geq 0$, we can take $Z_\delta(u)v$ as test function in (3.11 *ii*), obtaining in view of (5.48)

$$(5.50) \quad \left\{ \begin{aligned} & \int_{\Omega} {}^tA(x)D(Z_\delta(u)v)DG_k(u) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(Z_\delta(u)\hat{\varphi}_i T_k(u)) + \\ & + \int_{\Omega} Z_\delta(u)\hat{f} T_k(u) - \int_{\Omega} {}^tA(x)DvDZ_\delta(u)T_k(u) + \\ & + \int_{\Omega} {}^tA(x)DZ_\delta(u)D(vT_k(u)) - \int_{\Omega} {}^tA(x)DZ_\delta(u)DvT_k(u) = \\ & = \langle -\operatorname{div} {}^tA(x)D(Z_\delta(u)v), G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \\ & + \langle \langle -\operatorname{div} {}^tA(x)D(Z_\delta(u)v), T_k(u) \rangle \rangle_{\Omega} = \\ & = \int_{\Omega} F(x, u)Z_\delta(u)v. \end{aligned} \right.$$

As far as the first term of the left-hand side of (5.50) is concerned, we have, since $k \geq 2\delta$,

$$(5.51) \quad \int_{\Omega} {}^tA(x)D(Z_\delta(u)v)DG_k(u) = 0;$$

indeed since $u \in H_{\text{loc}}^1(\Omega)$ by (3.10 *i*) one has

$$\left\{ \begin{aligned} & {}^tA(x)D(Z_\delta(u)v)DG_k(u) = \\ & = -{}^tA(x)DuDu \frac{1}{\delta} \chi_{\{\delta < u < 2\delta\}} \chi_{\{u > k\}} + {}^tA(x)DvDu Z_\delta(u) \chi_{\{u > k\}} \text{ in } \mathcal{D}'(\Omega), \end{aligned} \right.$$

where each term is zero almost everywhere when $k \geq 2\delta$.

As far as the fourth term of the left-hand side of (5.50) is concerned, we have, since $u \in H_{\text{loc}}^1(\Omega)$ and since $k \geq 2\delta$, using (5.42),

$$DZ_\delta(u)T_k(u) = DuZ'_\delta(u)T_k(u) = -DuR'_\delta(u) = -DR_\delta(u) \text{ in } \mathcal{D}'(\Omega),$$

which using the fact that $R_\delta(u) \in H_0^1(\Omega)$ (see (5.45)) and (5.36) implies that

$$(5.52) \quad \begin{cases} - \int_{\Omega} {}^t A(x) Dv DZ_\delta(u) T_k(u) = \int_{\Omega} {}^t A(x) Dv DR_\delta(u) = \\ = \langle -\operatorname{div} {}^t A(x) Dv, R_\delta(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = \\ = \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i R_\delta(u)) + \int_{\Omega} \hat{f} R_\delta(u). \end{cases}$$

As far as the fifth term of the left-hand side of (5.50) is concerned, we have, since $u \in H_{\text{loc}}^1(\Omega)$,

$$\begin{cases} \int {}^t A(x) DZ_\delta(u) D(vT_k(u)) = \\ = \int {}^t A(x) DZ_\delta(u) Dv T_k(u) + \int {}^t A(x) DZ_\delta(u) DT_k(u) v \text{ in } \mathcal{D}'(\Omega), \end{cases}$$

which, since $k \geq 2\delta$, implies that

$$(5.53) \quad \begin{cases} \int_{\Omega} {}^t A(x) DZ_\delta(u) D(vT_k(u)) = \\ = \int_{\Omega} {}^t A(x) DZ_\delta(u) Dv T_k(u) - \frac{1}{\delta} \int_{\{\delta < u < 2\delta\}} {}^t A(x) Du Du v. \end{cases}$$

Collecting together (5.50), (5.51), (5.52) and (5.53), we have proved that

$$(5.54) \quad \begin{cases} \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i (Z_\delta(u) T_k(u) + R_\delta(u))) + \int_{\Omega} \hat{f} (Z_\delta(u) T_k(u) + R_\delta(u)) = \\ = \int_{\Omega} F(x, u) Z_\delta(u) v + \frac{1}{\delta} \int_{\{\delta < u < 2\delta\}} {}^t A(x) Du Du v. \end{cases}$$

Since when $k \geq 2\delta$ one has (see (5.42))

$$\begin{cases} {}^t A(x) Du Du v \geq 0, \quad 0 \leq Y_\delta(u) \leq \frac{3}{2}\delta, \\ Z_\delta(u) T_k(u) + R_\delta(u) = Z_\delta(u) u + R_\delta(u) = Y_\delta(u), \\ \hat{g}_i D(\hat{\varphi}_i Y_\delta(u)) = \hat{g}_i D\hat{\varphi}_i Y_\delta(u) + \hat{g}_i Du Z_\delta(u) \hat{\varphi}_i, \end{cases}$$

estimate (5.37) follows from (5.54).

Proposition 5.9 is proved. \square

5.4. A priori estimate of $\beta(u)$ in $H_0^1(\Omega)$

This fourth a priori estimate is actually in some sense a regularity result, since it asserts that for every u solution to problem (2.1) in the sense of Definition 3.6, a certain function of u actually belongs to $H_0^1(\Omega)$. This property will be used in the proofs of the Comparison Principle 7.1 and of the Uniqueness Theorem 4.3.

Define the function $\beta : s \in [0, +\infty[\rightarrow \beta(s) \in [0, +\infty[$ by

$$(5.55) \quad \beta(s) = \int_0^s \sqrt{\Gamma'(t)} dt.$$

Proposition 5.13. *Assume that the matrix A and the function F satisfy (2.4), (2.5) and (2.6). Then for every u solution to problem (2.1) in the sense of Definition 3.6 one has*

$$(5.56) \quad \beta(u) \in H_0^1(\Omega),$$

with

$$(5.57) \quad \alpha \|D\beta(u)\|_{(L^2(\Omega))^N}^2 \leq \|h\|_{L^1(\Omega)}.$$

Remark 5.14. (Formal proof of Proposition 5.13). Estimate (5.57) can be obtained formally by taking $\Gamma(u)$ as test function in equation (2.1), using the coercivity (2.4) and the growth condition (2.6 *iii*), which implies that

$$0 \leq F(x, u)\Gamma(u) \leq \frac{h(x)}{\Gamma(u)}\Gamma(u) \leq h(x).$$

This formal computation will be made mathematically correct in the proof below. \square

Proof of Proposition 5.13.

As in (5.5), we define, for every δ and k with $0 < \delta < k$, the function $S_{\delta,k}$ as

$$(5.58) \quad S_{\delta,k}(s) = \begin{cases} 0 & \text{if } 0 \leq s \leq \delta, \\ s - \delta & \text{if } \delta \leq s \leq k, \\ k - \delta & \text{if } k \leq s. \end{cases}$$

As in the beginning of the proof of Proposition 5.1, one can prove that the function $\Gamma(S_{\delta,k}(u))$ belongs to $\mathcal{V}(\Omega)$ and that (compare with (5.9))

$$-div {}^t A(x) D\Gamma(S_{\delta,k}(u)) = \psi_\delta(u) (-div {}^t A(x) D\Gamma(S_{\delta,k}(u))) \quad \text{in } \mathcal{D}'(\Omega),$$

where $\psi_\delta(s)$ is a C^1 nondecreasing function such that

$$\psi_\delta(s) = 0 \text{ for } 0 \leq s \leq \frac{\delta}{2} \text{ and } \psi_\delta(s) = 1 \text{ for } s \geq \delta.$$

Since $\Gamma(S_{\delta,k}(u)) \in \mathcal{V}(\Omega)$ and since $\Gamma(S_{\delta,k}(u)) \geq 0$, we can use $\Gamma(S_{\delta,k}(u))$ as test function in (3.11 *ii*). We get

$$\left\{ \begin{aligned} & \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) DG_k(u) + \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) D(\psi_{\delta}(u) T_k(u)) = \\ & = \langle -\operatorname{div} {}^t A(x) D\Gamma(S_{\delta,k}(u)), G_k(u) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \\ & + \langle \langle -\operatorname{div} {}^t A(x) D\Gamma(S_{\delta,k}(u)), T_k(u) \rangle \rangle_{\Omega} = \\ & = \int_{\Omega} F(x, u) \Gamma(S_{\delta,k}(u)), \end{aligned} \right.$$

which can be written as

$$(5.59) \quad \left\{ \begin{aligned} & \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) DG_k(u) + \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) D\psi_{\delta}(u) T_k(u) + \\ & + \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) DT_k(u) \psi_{\delta}(u) = \int_{\Omega} F(x, u) \Gamma(S_{\delta,k}(u)). \end{aligned} \right.$$

Note that the two first integrals are zero, since $D\Gamma(S_{\delta,k}(u))$ is zero outside of the set $\{\delta < u < k\}$, while $DG_k(u)$ is zero outside of the set $\{u > k\}$ and $D\psi_{\delta}(u)$ is zero outside of the set $\{u < \delta\}$.

Note also that $\psi_{\delta}(u) = 1$ on the set $\{u \geq \delta\}$ while $DS_{\delta,k}(u) = 0$ outside of this set. Therefore the third term of (5.59) can be written as

$$(5.60) \quad \left\{ \begin{aligned} & \int_{\Omega} {}^t A(x) D\Gamma(S_{\delta,k}(u)) DT_k(u) \psi_{\delta}(u) = \\ & = \int_{\Omega} {}^t A(x) DS_{\delta,k}(u) DT_k(u) \Gamma'(S_{\delta,k}(u)) = \\ & = \int_{\Omega} {}^t A(x) DS_{\delta,k}(u) DS_{\delta,k}(u) \Gamma'(S_{\delta,k}(u)) \geq \\ & \geq \alpha \int_{\Omega} |DS_{\delta,k}(u)|^2 \Gamma'(S_{\delta,k}(u)) = \alpha \int_{\Omega} |D\beta(S_{\delta,k}(u))|^2. \end{aligned} \right.$$

For what concerns the right-hand side of (5.59), we have, using the growth condition (2.6 *iii*) and the fact that Γ is increasing,

$$(5.61) \quad F(x, u) \Gamma(S_{\delta,k}(u)) \leq \frac{h(x)}{\Gamma(u)} \Gamma(S_{\delta,k}(u)) \leq h(x).$$

By (5.59), (5.60) and (5.61) we get

$$(5.62) \quad \alpha \int_{\Omega} |D\beta(S_{\delta,k}(u))|^2 \leq \|h\|_{L^1(\Omega)} \quad \forall \delta, 0 < \delta < k.$$

Passing to the limit as δ tends to zero gives implies that

$$(5.63) \quad \alpha \|D\beta(T_k(u))\|_{(L^2(\Omega))^N}^2 \leq \|h\|_{L^1(\Omega)} \quad \forall k > 0.$$

On the other hand, the fact that for every δ with $0 < \delta < k$

$$(5.64) \quad \begin{cases} |D\beta(S_{\delta,k}(u))| = |\beta'(S_{\delta,k}(u))\chi_{\{\delta < u < k\}}Du| \leq \sup_{0 \leq s \leq k} \beta'(s) |DG_\delta(u)|, \\ 0 \leq \beta(S_{\delta,k}(u)) \leq \beta(G_\delta(u)), \end{cases}$$

condition (3.10 *iii*) and Lemma A.1 of Appendix A below imply that $\beta(S_{\delta,k}(u)) \in H_0^1(\Omega) \cap L^\infty(\Omega)$. The fact that $\beta(S_{\delta,k}(u))$ is bounded in $H_0^1(\Omega)$ by (5.62) and converges to $\beta(T_k(u))$ almost everywhere as δ tends to zero then implies that

$$(5.65) \quad \beta(T_k(u)) \in H_0^1(\Omega).$$

Since $\beta(T_k(u)) \in H_0^1(\Omega)$ is bounded in $H_0^1(\Omega)$ in view of (5.63) and converges to $\beta(u)$ almost everywhere in Ω (note that u is finite almost everywhere since $u \in L^2(\Omega)$), we have proved that $\beta(u)$ belongs to $H_0^1(\Omega)$ and satisfies (5.57).

Proposition 5.13 is proved. \square

6. PROOFS OF THE STABILITY THEOREM 4.2 AND OF THE EXISTENCE THEOREM 4.1

6.1. Proof of the Stability Theorem 4.2

First step. Since for every n the function $F_n(x, s)$ satisfies assumptions (2.5) and (2.6) for the same h and the same Γ , every solution u_n to problem (2.1) $_n$ in the sense of Definition 3.6 satisfies the a priori estimates (5.1), (5.3), (5.15), (5.16), (5.17), (5.18) and (5.37).

Therefore u_n is bounded in $L^2(\Omega)$ and in $H_{loc}^1(\Omega)$, and there exist a subsequence, still labelled by n , and a function u_∞ such that

$$(6.1) \quad \begin{cases} u_n \rightharpoonup u_\infty \text{ in } L^2(\Omega) \text{ weakly, in } H_{loc}^1(\Omega) \text{ weakly and a.e. in } \Omega, \\ G_k(u_n) \rightharpoonup G_k(u_\infty) \text{ in } H_0^1(\Omega) \text{ weakly } \forall k > 0, \\ \varphi T_k(u_n) \rightharpoonup \varphi T_k(u_\infty) \text{ in } H_0^1(\Omega) \text{ weakly } \forall k > 0, \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

Since $u_n \geq 0$, this function u_∞ satisfies $u_\infty \geq 0$.

Therefore u_∞ satisfies (3.10).

Fix now a function v

$$v \in \mathcal{V}(\Omega), v \geq 0, \text{ with } -\operatorname{div}^t A(x)Dv = \sum_{i \in I} \hat{\varphi}_i(-\operatorname{div} \hat{g}_i) + \hat{f} \text{ in } \mathcal{D}'(\Omega).$$

Using v as test function in (3.11 ii) $_n$, we obtain

$$(6.2) \quad \begin{cases} \int_{\Omega} {}^t A(x) Dv DG_k(u_n) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u_n)) + \int_{\Omega} \hat{f} T_k(u_n) = \\ = \langle -\operatorname{div} {}^t A(x) Dv, G_k(u_n) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) Dv, T_k(u_n) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F_n(x, u_n) v. \end{cases}$$

Since the left-hand side of (6.2) is bounded independently of n for every $k > 0$ fixed in view of the estimates (5.1) and (5.16), we have

$$\int_{\Omega} F_n(x, u_n) v \leq C(v) < +\infty \quad \forall n,$$

which using the almost everywhere convergence of u_n to u_{∞} , assumption (4.1) on the functions F_n and Fatou's Lemma gives

$$(6.3) \quad \int_{\Omega} F_{\infty}(x, u_{\infty}) v \leq C(v) < +\infty \quad \forall n,$$

namely (3.11 i) $_{\infty}$.

It remains to prove that (3.11 ii) $_{\infty}$ holds true and that the convergences in (6.1) are strong.

Second step. For $\delta > 0$ fixed, we write (6.2) as

$$(6.4) \quad \begin{cases} \int_{\Omega} {}^t A(x) Dv DG_k(u_n) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u_n)) + \int_{\Omega} \hat{f} T_k(u_n) = \\ = \langle -\operatorname{div} {}^t A(x) Dv, G_k(u_n) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \langle \langle -\operatorname{div} {}^t A(x) Dv, T_k(u_n) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} F_n(x, u_n) Z_{\delta}(u_n) v + \int_{\Omega} F_n(x, u_n) (1 - Z_{\delta}(u_n)) v. \end{cases}$$

It is easy to pass to the limit in the left-hand side of (6.4), obtaining

$$(6.5) \quad \begin{cases} \int_{\Omega} {}^t A(x) Dv DG_k(u_n) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u_n)) + \int_{\Omega} \hat{f} T_k(u_n) \rightarrow \\ \rightarrow \int_{\Omega} {}^t A(x) Dv DG_k(u_{\infty}) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u_{\infty})) + \int_{\Omega} \hat{f} T_k(u_{\infty}) \\ \text{as } n \rightarrow +\infty. \end{cases}$$

Concerning the first term of the right-hand side of (6.4) we use the a priori estimate (5.37), namely

$$\begin{cases} \forall \delta > 0, \int_{\{u_n \leq \delta\}} F_n(x, u_n) Z_\delta(u_n) v \leq \\ \leq \frac{3}{2} \left(\int_{\Omega} \left| \sum_{i \in I} \hat{g}_i D \hat{\varphi}_i + \hat{f} \right| \right) \delta + \sum_{i \in I} \int_{\Omega} Z_\delta(u_n) \hat{g}_i D u_n \hat{\varphi}_i, \end{cases}$$

in which we pass to the limit in n for $\delta > 0$ fixed. Since $Z_\delta(u_n) \hat{g}_i$ tends strongly to $Z_\delta(u_\infty) \hat{g}_i$ in $(L^2(\Omega))^N$ while $D u_n \hat{\varphi}_i$ tends weakly to $D u_\infty \hat{\varphi}_i$ in $(L^2(\Omega))^N$ (see (5.17)), we obtain

$$(6.6) \quad \begin{cases} \forall \delta > 0, \limsup_n \int_{\{u_n \leq \delta\}} F_n(x, u_n) Z_\delta(u_n) v \leq \\ \leq \frac{3}{2} \left(\int_{\Omega} \left| \sum_{i \in I} \hat{g}_i D \hat{\varphi}_i + \hat{f} \right| \right) \delta + \sum_{i \in I} \int_{\Omega} Z_\delta(u_\infty) \hat{g}_i D u_\infty \hat{\varphi}_i. \end{cases}$$

Since $Z_\delta(u_\infty)$ tends to $\chi_{\{u_\infty=0\}}$ almost everywhere in Ω as δ tends to zero, and since $u_\infty \in H_{\text{loc}}^1(\Omega)$ implies that $D u_\infty = 0$ almost everywhere on $\{x \in \Omega : u_\infty(x) = 0\}$, the right-hand side of (6.6) tends to 0 as δ tends to zero.

We have proved that the first term of the right-hand side of (6.4) satisfies

$$(6.7) \quad \limsup_n \int_{\{u_n \leq \delta\}} F_n(x, u_n) Z_\delta(u_n) v \rightarrow 0 \text{ as } \delta \rightarrow 0.$$

Third step. Let us now prove that

$$(6.8) \quad \int_{\{u_\infty=0\}} F_\infty(x, u_\infty) v = 0.$$

In view of assumption (4.1) on the convergence of the functions $F_n(x, s)$, of the continuity of the function Z_δ , and of the almost everywhere convergence of u_n to u_∞ , one has for every $\delta > 0$ fixed

$$(6.9) \quad F_n(x, u_n) Z_\delta(u_n) v \rightarrow F_\infty(x, u_\infty) Z_\delta(u_\infty) v \text{ a.e. } x \in \Omega \text{ as } n \rightarrow +\infty.$$

By Fatou's Lemma this implies that

$$(6.10) \quad \int_{\Omega} F_\infty(x, u_\infty) Z_\delta(u_\infty) v \leq \liminf_{n \rightarrow +\infty} \int_{\Omega} F_n(x, u_n) Z_\delta(u_n) v \quad \forall \delta > 0.$$

Since $Z_\delta(s)$ tends to $\chi_{\{s=0\}}$ for every $s \geq 0$ as δ tends to zero and since $F_\infty(x, u_\infty)v \in L^1(\Omega)$ by (6.3), the left-hand side of (6.10) tends to $\int_\Omega F_\infty(x, u_\infty)\chi_{\{u_\infty=0\}}v$ as δ tends to zero.

Combining the latest result with (6.10) and (6.7) proves (6.8).

Fourth step. We finally pass to the limit in n for $\delta > 0$ fixed in the second term of the right-hand side of (6.4), namely in the term $\int_\Omega F_n(x, u_n)(1 - Z_\delta(u_n))v$.

For that we observe that

$$0 \leq 1 - Z_\delta(s) \leq \chi_{\{s \geq \delta\}} \quad \forall s \geq 0 \quad \forall \delta > 0,$$

which combined with the growth condition (2.6 *iii*) and the fact that the function Γ is increasing gives

$$(6.11) \quad 0 \leq F_n(x, u_n)(1 - Z_\delta(u_n))v \leq \frac{h(x)}{\Gamma(u_n)}\chi_{\{u_n \geq \delta\}}v \leq \frac{h(x)}{\Gamma(\delta)}v.$$

Since the right-hand side of (6.11) belongs to $L^1(\Omega)$, and since, for every $\delta > 0$ fixed, one has, as in (6.9),

$$F_n(x, u_n)(1 - Z_\delta(u_n))v \rightarrow F_\infty(x, u_\infty)(1 - Z_\delta(u_\infty))v \text{ a.e. } x \in \Omega \text{ as } n \rightarrow +\infty,$$

Lebesgue's dominated convergence theorem implies that for every $\delta > 0$ fixed one has

$$(6.12) \quad \int_\Omega F_n(x, u_n)(1 - Z_\delta(u_n))v \rightarrow \int_\Omega F_\infty(x, u_\infty)(1 - Z_\delta(u_\infty))v \text{ as } n \rightarrow +\infty.$$

Since the right-hand side of (6.12) tends to $\int_\Omega F_\infty(x, u_\infty)(1 - \chi_{\{u_\infty=0\}})v$ as δ tends to zero, which is nothing but $\int_\Omega F_\infty(x, u_\infty)v$ in view of (6.8), we have proved that

$$(6.13) \quad \lim_n \int_\Omega F_n(x, u_n)(1 - Z_\delta(u_n))v \rightarrow \int_\Omega F_\infty(x, u_\infty)v \text{ as } \delta \rightarrow 0.$$

Fifth step. Collecting the results obtained in (6.5), (6.7) and (6.13), we pass to the limit in each term of (6.4), first in n for $\delta > 0$ fixed, and

then in δ . This proves that for every $v \in \mathcal{V}(\Omega)$, $v \geq 0$, one has

$$\begin{cases} \int_{\Omega} {}^t A(x) Dv DG_k(u_{\infty}) + \sum_{i \in I} \int_{\Omega} \hat{g}_i D(\hat{\varphi}_i T_k(u_{\infty})) + \int_{\Omega} \hat{f} T_k(u_{\infty}) = \\ = \int_{\Omega} F_{\infty}(x, u_{\infty}) v, \end{cases}$$

which is nothing but (3.11 *ii*) $_{\infty}$.

We have proved that u_{∞} is a solution to problem (2.1) $_{\infty}$ in the sense of Definition 3.6.

It only remains to prove that the convergences in (6.1) are strong (see (4.2)). To this aim it is sufficient to prove the two following strong convergences

$$(6.14) \quad G_k(u_n) \rightarrow G_k(u_{\infty}) \text{ in } H_0^1(\Omega) \text{ strongly } \forall k > 0,$$

$$(6.15) \quad \begin{cases} \varphi DT_k(u_n) \rightarrow \varphi DT_k(u_{\infty}) \text{ in } (L^2(\Omega))^N \text{ strongly } \forall k > 0, \\ \forall \varphi \in H_0^1(\Omega) \cap L^{\infty}(\Omega). \end{cases}$$

Indeed one has $u_n = T_k(u_n) + G_k(u_n)$, and the strong convergence of $T_k(u_n)$ in $L^2(\Omega)$ then follows from Lebesgue's dominated convergence Theorem and from the almost convergence of u_n (see (6.1)). These facts in turn imply the strong convergences of u_n in $L^2(\Omega)$, of $\varphi T_k(u_n)$ in $H_0^1(\Omega)$, and of u_n in $H_{\text{loc}}^1(\Omega)$.

Sixth step. As far as (6.14) is concerned, the strong convergence follows from the energy equality (5.4) $_n$, namely the fact that

$$(6.16) \quad \int_{\Omega} A(x) DG_k(u_n) DG_k(u_n) = \int_{\Omega} F_n(x, u_n) G_k(u_n) \quad \forall k > 0.$$

It is easy to pass to the limit in the right-hand side of (6.16). Indeed the inequality

$$0 \leq F_n(x, u_n) G_k(u_n) \leq \frac{h(x)}{\Gamma(u_n)} \chi_{\{u_n \geq k\}} G_k(u_n) \leq \frac{h(x)}{\Gamma(k)} G_k(u_n)$$

and the boundedness of $G_k(u_n)$ in $H_0^1(\Omega)$ (see (6.1)) imply that the sequence $F_n(x, u_n) G_k(u_n)$ is equintegrable in n ; since

$$F_n(x, u_n) G_k(u_n) \rightarrow F_{\infty}(x, u_{\infty}) G_k(u_{\infty}) \quad \text{a.e. } x \in \Omega,$$

using Vitali's Theorem one obtains that for every $k > 0$ fixed

$$(6.17) \quad \int_{\Omega} F_n(x, u_n) G_k(u_n) \rightarrow \int_{\Omega} F_{\infty}(x, u_{\infty}) G_k(u_{\infty}) \text{ as } n \rightarrow +\infty.$$

On the other hand, the energy equality $(5.4)_\infty$ asserts that

$$(6.18) \quad \int_{\Omega} A(x) DG_k(u_\infty) DG_k(u_\infty) = \int_{\Omega} F_\infty(x, u_\infty) G_k(u_\infty) \quad \forall k > 0.$$

Collecting together (6.16), (6.17) and (6.18), one has for every $k > 0$ fixed

$$\int_{\Omega} A(x) DG_k(u_n) DG_k(u_n) \rightarrow \int_{\Omega} A(x) DG_k(u_\infty) DG_k(u_\infty) \quad \text{as } n \rightarrow +\infty,$$

which, with the weak convergence in $(L^2(\Omega))^N$ of $DG_k(u_n)$ to $DG_k(u_\infty)$ (see (6.1)), implies the strong convergence (6.14).

Seventh step. As far as (6.15) is concerned, writing equation (5.27) for $F = F_n$ and $u = u_n$, one obtains

$$(6.19) \quad \begin{cases} \int_{\Omega} A(x) DT_k(u_n) DT_k(u_n) \varphi^2 = \\ = -2 \int_{\Omega} A(x) DT_k(u_n) D\varphi \varphi T_k(u_n) - 2 \int_{\Omega} A(x) DG_k(u_n) D\varphi \varphi T_k(u_n) + \\ + \int_{\Omega} F_n(x, u_n) \varphi^2 T_k(u_n) \quad \forall \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega). \end{cases}$$

Let us pass to the limit in the right-hand side of (6.19) as n tends to $+\infty$. In view of (6.1) one has

$$(6.20) \quad \begin{cases} -2 \int_{\Omega} A(x) DT_k(u_n) D\varphi \varphi T_k(u_n) - 2 \int_{\Omega} A(x) DG_k(u_n) D\varphi \varphi T_k(u_n) \rightarrow \\ \rightarrow -2 \int_{\Omega} A(x) DT_k(u_\infty) D\varphi \varphi T_k(u_\infty) - 2 \int_{\Omega} A(x) DG_k(u_\infty) D\varphi \varphi T_k(u_\infty) \\ \text{as } n \rightarrow +\infty. \end{cases}$$

As far as the last integral of the right-hand side of (6.19) is concerned, we write, for every fixed $\delta > 0$,

$$(6.21) \quad \begin{cases} \int_{\Omega} F_n(x, u_n) \varphi^2 T_k(u_n) = \\ = \int_{\Omega} F_n(x, u_n) Z_\delta(u_n) \varphi^2 T_k(u_n) + \int_{\Omega} F_n(x, u_n) (1 - Z_\delta(u_n)) \varphi^2 T_k(u_n). \end{cases}$$

For the first term of the right-hand side of (6.21), we use the a priori estimate (5.37) with $v = \varphi^2$; indeed $\varphi^2 \in \mathcal{V}(\Omega)$ (see (3.8)) with

$$-div {}^t A(x) D\varphi^2 = \hat{\varphi}(-div \hat{g}) + \hat{f} \text{ in } \mathcal{D}'(\Omega),$$

with $\hat{\varphi} = 2\varphi$, $\hat{g} = {}^tA(x)D\varphi$ and $\hat{f} = -2{}^tA(x)D\varphi D\varphi$; this yields, since $\hat{g}D\hat{\varphi} + \hat{f} = 0$,

$$(6.22) \quad \begin{cases} 0 \leq \int_{\Omega} F_n(x, u_n) Z_{\delta}(u_n) \varphi^2 T_k(u_n) \leq k \int_{\Omega} F_n(x, u_n) Z_{\delta}(u_n) \varphi^2 \leq \\ \leq 2k \int_{\Omega} Z_{\delta}(u_n) {}^tA(x) D\varphi D u_n \varphi = \\ = 2k \int_{\Omega} Z_{\delta}(u_n) {}^tA(x) D\varphi D T_k(u_n) \varphi + 2k \int_{\Omega} Z_{\delta}(u_n) {}^tA(x) D\varphi D G_k(u_n) \varphi. \end{cases}$$

We pass to the limit in the right-hand side of (6.22) first for $\delta > 0$ fixed as n tends to $+\infty$ thanks to (6.1), and then as δ tends to 0. Since $Z_{\delta}(u_{\infty})$ tends to $\chi_{\{u_{\infty}=0\}}$ in $L^{\infty}(\Omega)$ weak-star as δ tends to zero, and since $Du_{\infty} = 0$ a.e. on $\{u_{\infty} = 0\}$, we obtain that

$$(6.23) \quad \limsup_n \int_{\Omega} F_n(x, u_n) Z_{\delta}(u_n) \varphi^2 T_k(u_n) \rightarrow 0 \text{ as } \delta \rightarrow 0,$$

For the second term of the right-hand side of (6.21), we repeat the proof that we did above to prove (6.13), and we obtain that

$$(6.24) \quad \lim_n \int_{\Omega} F_n(x, u_n) (1 - Z_{\delta}(u_n)) \varphi^2 T_k(u_n) \rightarrow \int_{\Omega} F_{\infty}(x, u_{\infty}) \varphi^2 T_k(u_{\infty}) \text{ as } \delta \rightarrow 0.$$

On the other hand, writing equation (5.27) for $F = F_{\infty}$ and $u = u_{\infty}$, one has

$$(6.25) \quad \begin{cases} \int_{\Omega} A(x) D T_k(u_{\infty}) D T_k(u_{\infty}) \varphi^2 = \\ = -2 \int_{\Omega} A(x) D T_k(u_{\infty}) D \varphi \varphi T_k(u_{\infty}) - 2 \int_{\Omega} A(x) D G_k(u_{\infty}) D \varphi \varphi T_k(u_{\infty}) + \\ + \int_{\Omega} F_{\infty}(x, u_{\infty}) \varphi^2 T_k(u_{\infty}) \quad \forall \varphi \in H_0^1(\Omega) \cap L^{\infty}(\Omega). \end{cases}$$

Collecting together (6.19), (6.20), (6.21), (6.23), (6.24) and (6.25), one has for every $k > 0$ fixed

$$\begin{cases} \int_{\Omega} A(x) D T_k(u_n) D T_k(u_n) \varphi^2 \rightarrow \int_{\Omega} A(x) D T_k(u_{\infty}) D T_k(u_{\infty}) \varphi^2 \\ \text{as } n \rightarrow +\infty, \end{cases}$$

which, with the weak convergence in $(L^2(\Omega))^N$ of $\varphi D T_k(u_n)$ to $\varphi D T_k(u_{\infty})$ implies the strong convergence (6.15).

This completes the proof of the Stability Theorem 4.2. \square

6.2. Proof of the Existence Theorem 4.1

Consider the problem

$$(6.26) \quad \begin{cases} u_n \in H_0^1(\Omega), \\ \int_{\Omega} A(x)Du_nD\varphi = \int_{\Omega} T_n(F(x, u_n^+))\varphi \quad \forall \varphi \in H_0^1(\Omega), \end{cases}$$

where T_n is the truncation at height n .

Since

$$T_n(F(x, s^+)) : (x, s) \in \Omega \times]-\infty, +\infty[\rightarrow T_n(F(x, s^+)) \in [0, n]$$

is a Carathéodory function which is almost everywhere bounded by n , Schauder's fixed point theorem implies that problem (6.26) has at least a solution. Moreover, since $T_n(F(x, s^+)) \geq 0$, this solution is nonnegative by the weak maximum principle, so that $u_n \geq 0$, and $T_n(F(x, u_n^+)) = T_n(F(x, u_n))$.

Define now the function F_n by

$$F_n(x, s) = T_n(F(x, s)) \text{ a.e. } x \in \Omega, \forall s \in [0, +\infty[.$$

For every given n , the function F_n is bounded, and it is easy to see that every u_n solution to (6.26) is a solution of problem (2.1)_n in the sense of Definition 3.6, where (2.1)_n is the problem (2.1) where the function F has been replaced by F_n : indeed $u_n \geq 0$ belongs to $H_0^1(\Omega)$, and therefore satisfies (3.10) and (3.11 i); u_n also satisfies (3.11 ii) in view of

$$(6.27) \quad \begin{cases} \langle \langle -div {}^tA(x)Dv, T_k(u_n) \rangle \rangle_{\Omega} = \langle -div {}^tA(x)Dv, T_k(u_n) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} \\ \forall v \in \mathcal{V}(\Omega), \end{cases}$$

which results of (3.5) by taking $y = T_k(u_n) \in H_0^1(\Omega) \cap L^\infty(\Omega)$.

It is clear that the functions F_n satisfy (2.5) and (2.6) with the functions h and Γ which appear in the definition of the function F . Moreover it is not difficult to verify that the function F_n satisfy (4.1) with F_∞ given by $F_\infty(x, s) = F(x, s)$.

The Stability Theorem 4.2 then implies that there exists a subsequence and a function u_∞ which is a solution to problem (2.1) in the sense of Definition 3.6 such that the convergences (4.2) hold true.

This proves the Existence Theorem 4.1. □

7. COMPARISON PRINCIPLE
AND PROOF OF THE UNIQUENESS THEOREM 4.3

In this section we prove a comparison result which uses assumption (2.7), namely the fact that $F(x, s)$ is nonincreasing with respect to s .

Note that we never use this assumption in the rest of the present paper except as far as the Uniqueness Theorem 4.3 is concerned.

Proposition 7.1. (Comparison Principle). *Assume that the matrix A satisfies (2.4). Let $F_1(x, s)$ and $F_2(x, s)$ be two functions satisfying (2.5) and (2.6) (possibly for different functions h and Γ). Assume moreover that*

$$(7.1) \quad \text{either } F_1(x, s) \text{ or } F_2(x, s) \text{ is nonincreasing, i.e. satisfies (2.7),}$$

and that

$$(7.2) \quad F_1(x, s) \leq F_2(x, s) \text{ a.e. } x \in \Omega, \quad \forall s \geq 0.$$

Let u_1 and u_2 be any solutions in the sense of Definition 3.6 to problems (2.1)₁ and (2.1)₂, where (2.1)₁ and (2.1)₂ are (2.1) with $F(x, s)$ replaced respectively by $F_1(x, s)$ and $F_2(x, s)$. Then one has

$$(7.3) \quad u_1(x) \leq u_2(x) \text{ a.e. } x \in \Omega.$$

Remark 7.2. The proof of the Comparison Principle of Proposition 7.1 is based on the use of the test function $(B_1(T_k^+(u_1 - u_2)))^2$. A similar test function has been used by L. Boccardo and J. Casado-Díaz in [3] to prove the uniqueness of solution to problem (2.1) obtained by approximation.

Note that the use of this test function is allowed by the regularity property (5.56) proved in Proposition 5.13. \square

Proof of the Uniqueness Theorem 4.3.

Applying the Comparison Principle to the case where $F_1(x, s) = F_2(x, s) = F(x, s)$ with $F(x, s)$ satisfying (2.7) immediately proves the Uniqueness Theorem 4.3. \square

Proof of Proposition 7.1.

First step. Let $k > 0$ be fixed. Define φ by

$$(7.4) \quad \varphi = B_1(T_k^+(u_1 - u_2)),$$

where $B_1 : s \in [0, +\infty[\rightarrow B_1(s) \in [0, +\infty[$ is the function defined by

$$B_1(s) = \int_0^s \beta_1(t) dt \quad \forall s \geq 0,$$

where $\beta_1(s) = \int_0^s \sqrt{\Gamma_1'(t)} dt$ and Γ_1 is the function for which F_1 satisfies (2.6).

In this step we will prove that

$$(7.5) \quad \varphi \in H_0^1(\Omega) \cap L^\infty(\Omega).$$

Since u_1 and u_2 belong to $H_{\text{loc}}^1(\Omega)$, one has $T_k^+(u_1 - u_2) \in H_{\text{loc}}^1(\Omega) \cap L^\infty(\Omega)$; since $\beta_1 \in C^0([0, +\infty[)$, the function φ belongs to $H_{\text{loc}}^1(\Omega)$ and one has

$$(7.6) \quad D\varphi = \beta_1(T_k^+(u_1 - u_2))\chi_{\{0 < u_1 - u_2 < k\}}(Du_1 - Du_2) \text{ in } \mathcal{D}'(\Omega).$$

Since $0 \leq T_k^+(s_1 - s_2) \leq T_k(s_1)$ for $s_1 \geq 0$ and $s_2 \geq 0$ and since β_1 is nondecreasing, this implies that

$$|D\varphi| \leq \beta_1(T_k(u_1))(|Du_1| + |Du_2|).$$

But in view of (5.56), $\beta_1(T_k(u_1))$ belongs to $H_0^1(\Omega) \cap L^\infty(\Omega)$, and then the properties (3.16) for u_1 and u_2 imply that $D\varphi \in (L^2(\Omega))^N$, and therefore that $\varphi \in H^1(\Omega) \cap L^\infty(\Omega)$.

Since B_1 is nondecreasing one has $0 \leq \varphi \leq B_1(T_k(u_1))$. We now claim that

$$(7.7) \quad B_1(T_k(u_1)) \in H_0^1(\Omega),$$

which by Lemma A.1 of Appendix A below completes the proof of (7.5).

Let us now prove (7.7). As in (5.64), we observe that for every δ with $0 < \delta < k$ and for the function $S_{\delta,k}$ defined by (5.58) one has

$$\begin{cases} |DB_1(S_{\delta,k}(u_1))| = |\beta_1(S_{\delta,k}(u_1))\chi_{\{\delta < u_1 < k\}}Du_1| \leq \sup_{0 \leq s \leq k} \beta_1(s) |DG_\delta(u_1)|, \\ 0 \leq B_1(S_{\delta,k}(u_1)) \leq B_1(G_\delta(u_1)). \end{cases}$$

Then Lemma A.1 implies that $B_1(S_{\delta,k}(u_1)) \in H_0^1(\Omega) \cap L^\infty(\Omega)$. But $|DB_1(S_{\delta,k}(u_1))| \leq \beta_1(T_k(u_1))|Du_1|$, which belongs to $L^2(\Omega)$ because of (5.56) and (3.16). Therefore $\beta_1(S_{\delta,k}(u_1))$ converges to $\beta_1(T_k(u_1))$ in $H^1(\Omega)$ weakly as δ tends to zero.

This implies (7.7).

Second step. Since $\varphi^2 \in \mathcal{V}(\Omega)$ in view of (7.5) and of Remark 3.5 *ii*), we can take φ^2 as test function in (3.11 *ii*)₁ and (3.11 *ii*)₂. Taking the

difference of these two equations, we get

$$(7.8) \quad \left\{ \begin{array}{l} 2 \int_{\Omega} \varphi^t A(x) D\varphi D(G_k(u_1) - G_k(u_2)) + \\ + 2 \int_{\Omega} {}^t A(x) D\varphi D(\varphi(T_k(u_1) - T_k(u_2))) - \\ - 2 \int_{\Omega} {}^t A(x) D\varphi D\varphi(T_k(u_1) - T_k(u_2)) = \\ = \langle -\operatorname{div} {}^t A(x) D\varphi^2, G_k(u_1) - G_k(u_2) \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} + \\ + \langle \langle -\operatorname{div} {}^t A(x) D\varphi^2, T_k(u_1) - T_k(u_2) \rangle \rangle_{\Omega} = \\ = \int_{\Omega} (F_1(x, u_1) - F_2(x, u_2)) \varphi^2. \end{array} \right.$$

Expanding in $L_{\text{loc}}^1(\Omega)$ the integrands of the three first lines of (7.8), one realizes that their sum is nothing but $2\varphi^t A(x) D\varphi D(u_1 - u_2)$, which belongs to $L^1(\Omega)$ in view of (3.16). Therefore one has

$$(7.9) \quad 2 \int_{\Omega} \varphi^t A(x) D\varphi D(u_1 - u_2) = \int_{\Omega} (F_1(x, u_1) - F_2(x, u_2)) \varphi^2,$$

which is formally easily obtained by taking φ^2 as test function in (2.1)₁ and (2.1)₂ and making the difference.

Third step. Let us now prove that for φ given by (7.4) one has

$$(7.10) \quad (F_1(x, u_1) - F_2(x, u_2)) \varphi^2 \leq 0 \text{ a.e. in } x \in \Omega.$$

Since $F_1(x, u_1)$ and $F_2(x, u_2)$ belong to $L_{\text{loc}}^1(\Omega)$ (see (3.12)) and are therefore finite almost everywhere, one has

$$(F_1(x, u_1) - F_2(x, u_2)) \varphi^2 = 0 \text{ on the set where } \varphi^2 = 0,$$

since there are no indeterminacies of the types $(\infty - \infty)$ and $\infty \times 0$ in the latest formula.

On the set where $\varphi^2 > 0$, one has $u_1 > u_2$. If $F_1(x, s)$ is nonincreasing with respect to s , one has, using first this nonincreasing character and then assumption (7.2),

$$\left\{ \begin{array}{l} F_1(x, u_1) - F_2(x, u_2) \leq F_1(x, u_2) - F_2(x, u_2) \leq 0 \\ \text{on the set where } \varphi^2 > 0, \end{array} \right.$$

since no indeterminacy of the type $(\infty - \infty)$ appears.

This implies that

$$(F_1(x, u_1) - F_2(x, u_2)) \varphi^2 \leq 0 \text{ on the set where } \varphi^2 > 0.$$

The case where $F_2(x, s)$ is nonincreasing with respect to s is similar, using first assumption (7.2) and then this nonincreasing character.

We have proved (7.10).

Fourth step. Collecting together (7.9) and (7.10) gives

$$2 \int_{\Omega} B_1(T_k^+(u_1 - u_2)) \beta_1(T_k^+(u_1 - u_2))^t A(x) D T_k^+(u_1 - u_2) D(u_1 - u_2) \leq 0.$$

Defining the function M_1 by

$$M_1(s) = \int_0^s \sqrt{B_1(t) \beta_1(t)} dt \quad \forall s > 0,$$

this implies in view of the coercivity (2.4) of the matrix A that $D M_1(T_k^+(u_1 - u_2)) = 0$ on Ω . Therefore $M_1(T_k^+(u_1 - u_2))$ is a constant in each connected component ω of Ω , and since the function M_1 is strictly increasing, there exists a nonnegative constant C_ω such that $T_k^+(u_1 - u_2) = C_\omega$ in ω , and therefore $\beta_1(T_k^+(u_1 - u_2)) = \beta_1(C_\omega)$ in ω . Since $T_k^+(u_1 - u_2) \leq T_k(u_1)$ in Ω and since the function β_1 is nondecreasing, one has

$$0 \leq \beta_1(C_\omega) = \beta_1(T_k^+(u_1 - u_2)) \leq \beta_1(T_k(u_1)) \text{ in } \omega.$$

By the regularity property (5.56), the function $\beta_1(T_k(u_1))$ belongs to $H_0^1(\Omega)$ and therefore to $H_0^1(\omega)$ for every connected component ω of Ω . By Lemma A.1 this implies that $\beta_1(C_\omega) \in H_0^1(\Omega)$, and then that $C_\omega = 0$. This proves that $T_k^+(u_1 - u_2) = 0$ in Ω , which implies (7.3) since $k > 0$. \square

APPENDIX A. AN USEFUL LEMMA

In this Appendix we state and prove the following lemma which is used many times in the present paper.

Lemma A.1. *If $y \in H^1(\Omega)$ and if there exists \underline{y} and $\bar{y} \in H_0^1(\Omega)$ such that*

$$\underline{y} \leq y \leq \bar{y} \quad \text{a.e. in } \Omega,$$

then $y \in H_0^1(\Omega)$.

Remark A.2. Lemma A.1 is straightforward when $\partial\Omega$ is sufficiently smooth so that the traces of the functions of $H^1(\Omega)$ are defined. Note that we did not assume any smoothness of $\partial\Omega$. \square

Proof of Lemma A.1.

Since $0 \leq y - \underline{y} \leq \bar{y} - \underline{y}$, it is sufficient to consider the case where $\underline{y} = 0$, namely the case where

$$y \in H^1(\Omega) \text{ with } 0 \leq y \leq \bar{y}, \text{ where } \bar{y} \in H_0^1(\Omega).$$

Since $\bar{y} \in H_0^1(\Omega)$, there exists a sequence $\phi_n \in \mathcal{D}(\Omega)$, such that

$$\phi_n \rightarrow \bar{y} \text{ in } H_0^1(\Omega).$$

The function y_n defined by

$$y_n = \phi_n^+ - (\phi_n^+ - y)^+$$

belongs to $H^1(\Omega)$ and has compact support (included in the support of ϕ_n). Therefore

$$y_n \in H_0^1(\Omega),$$

and

$$y_n \rightarrow \bar{y}^+ - (\bar{y}^+ - y)^+ = y \quad \text{in } H^1(\Omega) \quad \text{as } n \rightarrow +\infty.$$

This implies that $y \in H_0^1(\Omega)$. \square

Acknowledgments. The authors would like to thank Gianni Dal Maso and Luc Tartar for their friendly help, and to thank Lucio Boccardo, Juan Casado-Díaz and Luigi Orsina for having introduced them to the topics of the present work. They also would like to thank their own institutions (Dipartimento di Scienze di Base e Applicate per l'Ingegneria, Facoltà di Ingegneria Civile e Industriale, Sapienza Università di Roma, Departamento de Matemática Aplicada y Estadística, Universidad Politécnica de Cartagena, and Laboratoire Jacques-Louis Lions, Université Pierre et Marie Curie Paris VI et CNRS) for providing the support of reciprocal visits which allowed them to perform the present work. The work of Pedro J. Martínez-Aparicio has been partially supported by the grant MTM2015-68210-P of the Spanish Ministerio de Economía y Competitividad (MINECO-FEDER), the FQM-116 grant of the Junta de Andalucía and the grant Programa de Apoyo a la Investigación de la Fundación Séneca-Agencia de Ciencia y Tecnología de la Región de Murcia 19461/PI/14.

REFERENCES

- [1] C.O. Alves, F.J.S.A. Correa and J.V.A. Goncalves, *Existence of solutions for some classes of singular Hamiltonian systems*. Adv. Nonlinear Stud. **5** (2005) 265–278.
- [2] D. Arcoya and L. Moreno-Mérida, *Multiplicity of solutions for a Dirichlet problem with a strongly singular nonlinearity*. Nonlinear Anal. **95** (2014), 281–291.
- [3] L. Boccardo and J. Casado-Díaz, *Some properties of solutions of some semilinear elliptic singular problems and applications to the G-convergence*. Asymptot. Anal. **86** (2014), 1–15.
- [4] L. Boccardo and L. Orsina, *Semilinear elliptic equations with singular nonlinearities*. Calc. Var. Partial Differential Equations **37** (2010), 363–380.
- [5] A. Canino and M. Degiovanni, *A variational approach to a class of singular semilinear elliptic equations*. J. Convex Anal. **11** (2004), 147–162.

- [6] A. Canino, B. Sciunzi and A. Trombetta, *Existence and uniqueness for p -Laplace equations involving singular nonlinearities*. NoDEA Nonlinear Differ. Equ. Appl. **23** (2016), in press, doi:10.1007/s00030-016-0361-6.
- [7] Y.S. Choi and P.J. McKenna, *A singular Gierer-Meinhardt system of elliptic equations*. Ann. Inst. Henri Poincaré Analyse non linéaire **4** (2000) 503–522.
- [8] F. Cirstea, M. Ghergu and V. Radulescu, *Combined effects of asymptotically linear and singular nonlinearities in bifurcation problem of Lane-Emden-Fowler type*. J. Math. Pures Appl. **84** (2005) 493–508.
- [9] G.M. Coclite and M.M. Coclite, *On a Dirichlet problem in bounded domains with singular nonlinearity*. Discrete Contin. Dyn. Syst. **33** (2013), 4923–4944.
- [10] M.M. Coclite and G. Palmieri, *On a singular nonlinear Dirichlet problem*. Comm. Partial Diff. Equations **14** (1989) 1315–1327.
- [11] M.G. Crandall, P.H. Rabinowitz and L. Tartar, *On a Dirichlet problem with a singular nonlinearity*. Comm. Partial Differential Equations **2** (1977), 193–222.
- [12] J. Davila and M. Montenegro, *Positive versus free boundary solutions to a singular elliptic equation*. J. Anal. Math. **90** (2003), 303–335.
- [13] L.M. De Cave, *Nonlinear elliptic equations with singular nonlinearities*. Asymptotic Analysis **84** (2013), 181–195.
- [14] J.I. Díaz, J.M. Morel and L. Oswald, *An elliptic equation with singular nonlinearity*. Comm. Partial Diff. Equations **12** (1987) 1333–1344.
- [15] P. Donato and D. Giachetti, *Existence and homogenization for a singular problem through rough surfaces*. SIAM J. Math. Anal. **48** (2016) 4047–4086.
- [16] W. Fulks and J.S. Maybee, *A singular nonlinear equation*. Osaka J. Math. **12** (1960), 1–19.
- [17] D. Giachetti, P.J. Martínez-Aparicio and F. Murat, *A semilinear elliptic equation with a mild singularity at $u = 0$: existence and homogenization*. J. Math. Pures Appl. **107** (2017), 41–77.
- [18] D. Giachetti, P.J. Martínez-Aparicio and F. Murat, *Advances in the study of singular semilinear elliptic problems*. In *Trends in differential equations and applications*, ed. by F. Ortégón Gallego, M.V. Redondo Neble and J.R. Rodríguez Galván. SEMA-SIMAI Springer Series **8**, Springer International Publishing Switzerland, (2016), 221–241.
- [19] D. Giachetti, P.J. Martínez-Aparicio and F. Murat, *Homogenization of a Dirichlet semilinear elliptic problem with a strong singularity at $u = 0$ in a domain with many small holes*. To appear.
- [20] D. Giachetti, P.J. Martínez-Aparicio and F. Murat, *Remarks on the definition of the solution to a semilinear elliptic problem with a strong singularity at $u = 0$* . To appear.
- [21] D. Giachetti, B. Vernescu and M.A. Vivaldi, *Asymptotic analysis of singular problems in perforated cylinders*. Differential Integral Equations **29** (2016), 531–562.
- [22] A.C. Lazer and P.J. McKenna, *On a singular nonlinear elliptic boundary-value problem*. Proc. Amer. Math. Soc. **111** (1991), 721–730.
- [23] F. Oliva and F. Petitta, *Finite and infinite energy solutions of singular elliptic problems: Existence and uniqueness*. To appear.

- [24] G. Stampacchia, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*. Ann. Inst. Fourier (Grenoble) **15** (1965), 189–258.
- [25] C.A. Stuart, *Existence and approximation of solutions of non-linear elliptic equations*. Math. Z. **147** (1976), 53–63.

DANIELA GIACHETTI

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L'INGEGNERIA
FACOLTÀ DI INGEGNERIA CIVILE E INDUSTRIALE
SAPIENZA UNIVERSITÀ DI ROMA
VIA SCARPA 16, 00161 ROMA, ITALY
E-mail address: daniela.giachetti@sbai.uniroma1.it

PEDRO J. MARTÍNEZ-APARICIO

DEPARTAMENTO DE MATEMÁTICA APLICADA Y ESTADÍSTICA
UNIVERSIDAD POLITÉCNICA DE CARTAGENA
PASEO ALFONSO XIII 52, 30202 CARTAGENA (MURCIA), SPAIN
E-mail address: pedroj.martinez@upct.es

FRANÇOIS MURAT

LABORATOIRE JACQUES-LOUIS LIONS ET CNRS
UNIVERSITÉ PIERRE ET MARIE CURIE
BOÎTE COURRIER 187, 75252 PARIS CEDEX 05, FRANCE
E-mail address: murat@ann.jussieu.fr