

1 Growth gap in hyperbolic groups and amenability

2 Rémi Coulon, Françoise Dal’Bo, Andrea Sambusetti

3 April 22, 2018

4 **Abstract**

5 We prove a general version of the amenability conjecture in the unified setting of a Gromov  
6 hyperbolic group  $G$  acting properly cocompactly either on its Cayley graph, or on a CAT(-1)-  
7 space. Namely, for any subgroup  $H$  of  $G$ , we show that  $H$  is co-amenable in  $G$  if and only if their  
8 exponential growth rates (with respect to the prescribed action) coincide. For this, we prove  
9 a quantified, representation-theoretical version of Stadlbauer’s amenability criterion for group  
10 extensions of a topologically transitive subshift of finite type, in terms of the spectral radii of  
11 the classical Ruelle transfer operator and its corresponding extension. As a consequence, we  
12 are able to show that, in our enlarged context, there is a gap between the exponential growth  
13 rate of a group with Kazhdan’s property (T) and the ones of its infinite index subgroups. This  
14 also generalizes a well-known theorem of Corlette for lattices of the quaternionic hyperbolic  
15 space or the Cayley hyperbolic plane.

16 **Contents**

17 **1 Introduction** **2**

18 **2 Preliminaries** **7**

19 2.1 Subshift of finite type . . . . . 7

20 2.2 Hyperbolic geometry . . . . . 9

21 **3 Horofunctions** **10**

22 **4 Dynamics in a hyperbolic group** **12**

23 4.1 Transitivity of Gromov’s geodesic flow . . . . . 12

24 4.2 Dynamics on the space of horofunctions . . . . . 14

25 4.3 Gromov’s coding . . . . . 16

26 **5 Potential and transfer operator** **20**

27 5.1 The data . . . . . 21

28 5.2 Transfer operator for the irreducible component . . . . . 21

29 5.3 Twisted transfer operator associated to a subgroup . . . . . 27

30 5.4 Proofs of the theorems . . . . . 30

1	<b>A An extension of Kesten’s criterion</b>	<b>31</b>
2	A.1 Function spaces . . . . .	31
3	A.2 Ruelle’s Perron Frobenius Theorem . . . . .	32
4	A.3 Twisted transfer operator . . . . .	33
5	A.4 Renormalization . . . . .	36
6	A.5 Invariant and almost invariant vectors . . . . .	36
7	A.6 Amenability and Kazhdan property (T) . . . . .	43
8	<b>B Roblin’s theorem</b>	<b>47</b>

## 9 1 Introduction

10 Amenability has a large number of equivalent formulations. In a seminal work dating back to 1959,  
 11 Kesten proved that a finitely generated group  $Q$  is amenable if and only if 1 is the spectral radius  
 12 of the Markov operator associated to a symmetric random walk on  $Q$  whose support generates  $Q$   
 13 [28]. Given a finite generating set  $S$  of  $Q$ , Grigorchuk [24] and Cohen [9] independently related the  
 14 spectral radius  $\rho$  for the random walk with uniform probability measure supported on  $S \cup S^{-1}$ , to  
 15 the exponential growth rates of the free group  $\mathbf{F}(S)$  and the kernel  $N$  of the canonical projection  
 16  $\mathbf{F}(S) \rightarrow Q$ . Recall that the *exponential growth rate* of a discrete group  $G$  of isometries of a proper  
 17 metric space  $X$ , denoted by  $\omega(G, X)$ , (or simply  $\omega_G$  if there is no ambiguity) is

$$18 \quad \omega(G, X) = \limsup_{r \rightarrow \infty} \frac{1}{r} \ln |\{g \in G \mid d(gx, x) \leq r\}|,$$

19 where  $x$  is any point of  $X$ . The Cohen-Grigorchuk formula states that

$$20 \quad \rho = \frac{\sqrt{e^{\omega_{\mathbf{F}(S)}}}}{1 + e^{\omega_{\mathbf{F}(S)}}} \left( \frac{\sqrt{e^{\omega_{\mathbf{F}(S)}}}}{e^{\omega_N}} + \frac{e^{\omega_N}}{\sqrt{e^{\omega_{\mathbf{F}(S)}}}} \right), \quad (1)$$

21 where  $\omega_N$  and  $\omega_{\mathbf{F}(S)}$  are the respective growth rates of  $N$  and  $\mathbf{F}(S)$  acting on the Cayley graph  
 22 of  $\mathbf{F}(S)$  with respect to  $S$ . This immediately yielded, by Kesten’s criterion, a characterization of  
 23 amenability for a group  $Q$  generated by a subset  $S$ , in terms of the growth of the relator subgroup:  
 24 the quotient  $Q = \mathbf{F}(S)/N$  is amenable if and only if  $\omega_N = \omega_{\mathbf{F}(S)} = \log(2|S| - 1)$ .

25 Almost at the same time, a geometric version of Kesten’s criterion, in terms of the bottom  
 26 of the  $L^2$ -spectrum of the Laplace operator, was discovered in Riemannian geometry by Brooks  
 27 [5, 6] (see also [7]): for any normal, Riemannian covering  $\hat{M} \rightarrow M$  of a Riemannian manifold  $M$   
 28 of *finite topological type*, if the automorphism group of  $\hat{M}$  is amenable, then  $\lambda_0(\hat{M}) = \lambda_0(M)$ ;  
 29 moreover, when  $\hat{M} = \tilde{M}$  is the universal covering of a convex-cocompact hyperbolic manifold  $M$   
 30 (or any Riemannian manifold  $M$  of finite topological type satisfying a “Cheeger-type” condition [6,  
 31 Theorem 2]), then the converse implication also holds. Realizing the base manifold as  $M = \tilde{M}/G$   
 32 and the cover as  $\hat{M} = \tilde{M}/N$  with  $N$  normal in  $G$ , this precisely says that the  $\lambda_0(\hat{M}) = \lambda_0(M)$   
 33 whenever the quotient group  $Q = G/N$  is amenable. Coupling this with Sullivan’s formula relating  
 34  $\lambda_0(M)$  to  $\omega_G$  for discrete subgroups of isometries of the hyperbolic space  $\mathbf{H}^n$ , it yields

$$35 \quad \lambda_0(M) = \begin{cases} \omega_G(n - 1 - \omega_G) & \text{if } \omega_G \geq \frac{n-1}{2} \\ \lambda_0(\mathbf{H}^n) & \text{if } \omega_G \leq \frac{n-1}{2} \end{cases}, \quad (2)$$

1 (where  $\omega_G$  is computed, this time, with respect to the action of  $G$  on the hyperbolic space).  
 2 Brooks then derived an analogue of Cohen-Grigorchuk statement for particular, discrete groups  
 3 of isometries of  $\mathbf{H}^n$ : for any normal Riemannian covering  $\tilde{M} = \tilde{M}/N$  of a convex-cocompact hy-  
 4 perbolic manifold  $M = \tilde{M}/G$  with  $\omega_G > (n-1)/2$ , the automorphism group  $Q = G/N$  of  $\tilde{M}$  is  
 5 amenable if and only if  $\omega_G = \omega_N$ . This holds, for instance, for uniform lattices. The result was also  
 6 extended to hyperbolic, non-uniform lattices in [35].

7 Beyond the evident formal analogies of these results – negatively curved cocompact groups look  
 8 like free groups at large scale, while the bottom of the Laplacian of  $M$  can be related to the spectral  
 9 radius of the Markov operator associated to particular random walks on  $G = \pi_1(M)$ , induced by  
 10 the heat kernel of  $\tilde{M}$  [5] – the two statements did not live on a common ground. This opened the  
 11 door to intensive research for a unifying, generalized setting, and a deeper understanding of the  
 12 relations between these results in dynamical terms.

13 The first statement in a general setting was given in 2005 by Roblin [34]. Given a discrete group  
 14  $G$  of isometries of a CAT(−1) space, he proved, using Patterson-Sullivan theory, that amenability  
 15 of the quotient  $G/N$  always implies the relation  $\omega_G = \omega_N$ . In this generality, it is worth to remark  
 16 that the reciprocal is not true. Indeed, there exist Kleinian, geometrically finite groups  $G$  – even  
 17 lattices in pinched, variable negative curvature [19, 20] – admitting a parabolic subgroup  $P$  with  
 18  $\omega_P = \omega_G$ . Such groups give easy counterexamples to the converse implication, by taking for  $N$  the  
 19 normal closure of  $P$  in  $G$ . Indeed in most of these cases  $G/N$  contains free subgroups and is not  
 20 amenable. The most accredited version of the *Amenability Problem* in the last decade can be stated  
 21 as follows. *Given a group  $G$  acting properly on a hyperbolic space  $X$  and a normal subgroup  $N$  of*  
 22  *$G$ , under which circumstances does the equality  $\omega_N = \omega_G$  imply that the quotient group  $Q = G/N$*   
 23 *is amenable?*

24 Clearly, for a group  $G$  acting on a general space  $X$ , an exact formula as (1) or (2) is hopeless.  
 25 Rather, one should expect that the equality  $\omega_N = \omega_G$  reflects the qualitative behavior of the  
 26 dynamics of  $G/N$  on the space  $X/N$ . Nevertheless an exact relation, in terms of the asymptotic  
 27 behavior of the spectral radii of random walks on  $G/N$  with probability measure supported by large  
 28 spheres, resists to this general setting, allowing to show the “easy part” of the implication above  
 29 even in the generality of cocompact group actions on Gromov-hyperbolic spaces (see for instance  
 30 Propositions B.5 and B.6 in Appendix B).

31 A substantial step forward in the solution of the Amenability Problem is due to Stadlbauer [36],  
 32 who generalized Kesten’s amenability criterion in terms of group extensions of topological Markov  
 33 chains. More precisely, he considered a topologically mixing subshift of finite type  $(\Sigma, \sigma)$  together  
 34 with a topologically transitive extension  $(\Sigma_\theta, \sigma_\theta)$  of this system by a locally constant evaluation  
 35 map  $\theta : \Sigma \rightarrow Q$  into a discrete group  $Q$ . He proved that  $Q$  is amenable if and only if the *Gurevič*  
 36 *pressures* of the two systems (with respect to a weakly symmetric potential with Hölder variations)  
 37 coincide<sup>1</sup>. As an application, Stadlbauer solved the Amenability Problem for the class of *essentially*  
 38 *free* groups  $G$  of isometries of  $\mathbf{H}^n$ , for the first time without assuming that  $\omega_G > (n-1)/2$ .  
 39 This result was recently generalized by Dougall and Sharp, using Stadlbauer’s criterion, to the  
 40 class of convex-cocompact groups of isometries of pinched, negatively curved Cartan-Hadamard  
 41 manifolds [22].

---

<sup>1</sup>Actually Stadlbauer’s criterion works in a slightly more general context which allow to consider symbolic dynamical systems over an infinite alphabet. Nevertheless in the context of hyperbolic groups a subshift of finite type is sufficient to conclude.

1 The first result of this paper solves the Amenability Problem in an enlarged context encompass-  
 2 ing two very different cases. The first one, of algebraic nature, concerns the growth of groups with  
 3 respect to the word metric. The second case, coming from the geometry, focuses on the action of a  
 4 group on a negatively curved Riemannian manifold or a  $\text{CAT}(-1)$  space. The aim is two-fold: to  
 5 give a self-contained proof of all these results in a unified setting, and to make clear the minimum  
 6 algebraic and geometric structure needed.

7 **Theorem 1.1** (see [Theorem 5.1](#)). *Let  $G$  be a group acting properly co-compactly by isometries on*  
 8 *a Gromov hyperbolic space  $X$ . We assume that one of the following holds. Either*

9 (i)  *$X$  is the Cayley graph of  $G$  with respect to a finite generating set, or*

10 (ii)  *$X$  is a  $\text{CAT}(-1)$  space.*

11 *Let  $H$  be a subgroup of  $G$ , and let  $\omega_G$  and  $\omega_H$  denote the exponential growth rates of  $G$  and  $H$*   
 12 *acting on  $X$ . The subgroup  $H$  is co-amenable in  $G$  if and only if  $\omega_H = \omega_G$ .*

13 Recall that the action of a group  $G$  on a space  $X$  is *amenable* if  $X$  admits a  $G$ -invariant mean, and  
 14 that  $H$  is called *co-amenable* in  $G$  if the action of  $G$  on the left coset space  $H \backslash G$  is amenable. When  
 15 the subgroup  $H$  is normal in  $G$  then  $H$  is co-amenable in  $G$  if and only if  $G/H$  is an amenable  
 16 group. Notice however that in the above theorem we do not assume that  $H$  is a normal subgroup.

17 Note that the  $\text{CAT}(-1)$  case in the above theorem extends the Riemannian convex-compact  
 18 situation studied by Dougall and Sharp [22]. Nevertheless, going from negatively curved manifolds  
 19 to  $\text{CAT}(-1)$  spaces is a substantial generalization. Indeed, Dougall and Sharp explicitly use the  
 20 Riemannian structure to encode the geodesic flow via Markov sections. To the best of our knowledge  
 21 there is no such coding for the geodesic flow on  $\text{CAT}(-1)$  spaces. We explain at the end of the  
 22 introduction our strategy to bypass this difficulty.

23 The easy part of [Theorem 1.2](#) is the “only if” implication. As we mentioned before, this direction  
 24 was proved for normal subgroups of discrete groups acting on  $\text{CAT}(-1)$  spaces by Roblin [34]. In  
 25 [35] Roblin and Tapie sketched how to extend the argument to the case of groups acting on a  
 26 Gromov hyperbolic space. Nevertheless, we decided to report in [Appendix B](#) a complete proof of  
 27 this fact via random walks, for general subgroups of Gromov hyperbolic groups, since this also gives  
 28 an exact formula which is similar to Sullivan’s one for the bottom of the Laplacian of hyperbolic  
 29 quotients (see [Theorem B.1](#)).

30 On the other hand, our proof of the converse implication is strongly inspired by Stadlbauer’s  
 31 work [36] and relies on a variation of his amenability criterion. However, our approach makes an  
 32 explicit use of representation theory and operator algebra, which was somehow hidden in [36]. We  
 33 hope that this point of view can enlighten the key conceptual arguments and clarify the exposition.  
 34 More precisely we take advantage of the Hulanicki–Reiter criterion for amenable actions: the action  
 35 of a discrete group  $G$  on a set  $Y$  is amenable if and only if the induced regular representation  
 36  $\lambda: G \rightarrow \mathcal{U}(\ell^2(Y))$  admits almost invariant vectors [3, Theorem G.3.2]. Assume now that  $(\Sigma_\theta, \sigma_\theta)$   
 37 is the extension of a subshift of finite type  $(\Sigma, \sigma)$  by a locally constant map  $\theta: \Sigma \rightarrow G$ . We associate  
 38 the classical Ruelle transfer operator  $\mathcal{L}$  to the original system  $(\Sigma, \sigma)$ . On the other hand, given an  
 39 action of  $G$  on a set  $Y$ , we endow the extended system  $(\Sigma_\theta, \sigma_\theta)$  with a *twisted transfer operator*  $\mathcal{L}_\lambda$   
 40 which is naturally related to the induced unitary representation  $\lambda: G \rightarrow \mathcal{U}(\ell^2(Y))$  (see [Section A.3](#)).  
 41 The twisted transfer operator acts on a subspace of the space of continuous functions  $C(\Sigma, \ell^2(Y))$   
 42 (the appropriate, Hölder regularity will be described in [Section A.1](#)). Using the uniform convexity of

1 Hilbert spaces, we relate the difference between the spectral radii  $\rho$  and  $\rho_\lambda$  of  $\mathcal{L}$  and  $\mathcal{L}_\lambda$  respectively,  
 2 to the existence of almost invariant vectors for the representation  $\lambda$ .

3 **Theorem 1.2** (see Theorems A.23 and A.25). *Let  $(\Sigma, \sigma)$  be a topologically transitive subshift of  
 4 finite type. Let  $F: \Sigma \rightarrow \mathbf{R}_+^*$  be a potential with  $\alpha$ -bounded Hölder variations (for some  $\alpha \in \mathbf{R}_+^*$ ),  
 5 and  $\mathcal{L}$  be the Ruelle transfer operator associated with  $F$ . Let  $G$  be a finitely generated group and  
 6  $\theta: \Sigma \rightarrow G$  a locally constant map. Assume that the extension  $(\Sigma_\theta, \sigma_\theta)$  of  $(\Sigma, \sigma)$  by  $\theta$  has the  
 7 visibility property. Then the following holds.*

8 (i) *For every finite subset  $S$  of  $G$  and every  $\varepsilon \in \mathbf{R}_+^*$ , there exists  $\eta \in \mathbf{R}_+^*$  with the following  
 9 property: if  $G$  acts on a set  $Y$ , and  $\mathcal{L}_\lambda$  is the corresponding twisted transfer operator, the  
 10 condition  $\rho_\lambda > (1 - \eta)\rho$  implies that the representation  $\lambda: G \rightarrow \mathcal{U}(\ell^2(Y))$  admits an  $(S, \varepsilon)$ -  
 11 invariant vector.*

12 (ii) *In particular, if  $\rho_\lambda = \rho$  then the action of  $G$  on  $Y$  is amenable.*

13 The second statement of this theorem easily follows from the first point, and is very similar  
 14 to the one obtained by Stadlbauer [36] when  $(\Sigma, \sigma)$  is a subshift of finite type. Let us highlight  
 15 a few important differences though. Unlike Stadlbauer's proof, our approach does not really use  
 16 the Gibbs measure but simply the Perron-Frobenius theorem. Therefore we do not ask the original  
 17 system  $(\Sigma, \sigma)$  to be *topologically mixing*, but just *topologically transitive* (this is much weaker, as in  
 18 many situations one can always reduce to an irreducible component of the system). Secondly, we  
 19 consider an extension  $(\Sigma_\theta, \sigma_\theta)$  of the initial system by the whole group  $G$ , and only assume that it  
 20 has the *visibility property* (which means that the extended flow visits almost the whole group  $G$ ),  
 21 whereas Stadlbauer extends the initial system by the quotient  $Q$ , and assumes that this extension  
 22 is *topologically transitive*. This is one of the key points which allows us to consider any subgroup  
 23 of a hyperbolic group and not only normal subgroups. Moreover, as we state our result in terms of  
 24 spectral radius instead of pressure, we do not need any kind of symmetry for the potential  $F$  (this  
 25 was already observed by Jaerisch [27]).

26 More importantly, our approach provides a *quantitative* version<sup>2</sup> of Stadlbauer's statement.  
 27 In this perspective, the first statement in the above theorem is close to some results of Dougall in  
 28 [21], which also includes more concrete representation theory (nevertheless she assumes mixing of  
 29 the initial system, and considers only normal subgroups to ensure the transitivity of the extended  
 30 system, as well as a condition called *linear visibility with reminders*, a bit stronger than ours, to  
 31 control the return times of the flow in a fixed cylinder). The quantitative version of the amenabil-  
 32 ity criterion (see Theorem A.28) makes apparent the following consequence for groups satisfying  
 33 Kazhdan's property (T).

34 **Theorem 1.3** (see Theorem 5.2). *Let  $G$  be a group with Kazhdan's property (T) acting properly  
 35 co-compactly by isometries on a hyperbolic space  $X$ . We assume that one of the following holds:*

36 (i) *either  $X$  is the Cayley graph of  $G$  with respect to a finite generating set,*

37 (ii) *or  $X$  is a CAT(-1) space.*

---

<sup>2</sup>In the proof of Theorem 1.2 (i), we choose to work with ultra-limit of Banach spaces: this has the advantage of simplifying the arguments involving almost invariant vectors. As a consequence, we do not provide a precise formula for  $\eta$  in terms of  $S$  and  $\varepsilon$ ; nevertheless, a careful reader could go through the arguments and make the relation between these quantities explicit.

1 *Then, there exists  $\eta > 0$  with the following property. Let  $H$  be a subgroup of  $G$ , and let  $\omega_G, \omega_H$*   
 2 *denote the exponential growth rates of  $G$  and  $H$  acting on  $X$ . If  $\omega_H > \omega_G - \eta$ , then  $H$  is a finite*  
 3 *index subgroup of  $G$ .*

4 We stress the fact that also in this statement  $H$  is not assumed to be normal. So, this gives the  
 5 following generalization of Corlette’s celebrated growth gap theorem [17] for subgroups of lattices  
 6 of rank one symmetric spaces of negative curvature possessing Kazhdan’s Property (T), i.e. the  
 7 quaternionic hyperbolic space  $\mathbf{H}_{\mathbb{H}}^n$  or the Cayley hyperbolic plane  $\mathbf{H}_{\mathbb{O}}^2$ . Corlette’s theorem was  
 8 generalized by Dougall [21] for convex-cocompact groups of isometries of pinched, negatively curved  
 9 Cartan-Hadamard manifolds. Our statement is an even further generalization which unifies the  
 10 combinatorial and the geometric point of views. Recall that, for a Gromov hyperbolic space, the  
 11 *visual dimension*  $\dim_{vis}(\partial X)$  is defined analogously to the Hausdorff dimension, but with respect  
 12 to the natural visual measures of  $\partial X$ , and it coincides with the exponential growth rate of any  
 13 cocompact group  $G$  of isometries of  $X$  – see for instance [32, 33]. We then have:

14 **Corollary 1.4.** *Let  $X$  be a CAT(−1) metric space admitting uniform lattices, whose isometry*  
 15 *group possesses Kazhdan’s property (T). Then, there exists  $\eta > 0$  such that for any subgroup  $H$  of*  
 16 *Isom( $X$ ), either  $H$  is a lattice, or the exponential growth rate of  $H$  is at most  $\dim_{vis}(\partial X) - \eta$ .*

17 An similar statement holds if  $X$  is the Cayley graph of a hyperbolic group. This result shall be  
 18 added to the list of geometric and dynamical rigidity consequences of property (T), such as Serre’s  
 19 fixed point-edge property for actions on trees [29, 38], or the local  $C^\infty$ -conjugacy rigidity of isometric  
 20 actions on compact Riemannian manifolds [23].

21 Let us now give an overview of the proof of Theorems 1.1 and 1.3. The main idea is to apply  
 22 our amenability criterion (Theorem 1.2) to the geodesic flow on  $X$ . However the criterion requires a  
 23 coding of this dynamical system, which may not exist for CAT(−1) spaces. To bypass this difficulty  
 24 we consider a geodesic flow not on the space  $X$  but rather on the Cayley graph  $\Gamma$  of the group  $G$   
 25 with respect to a finite generating set  $S$ . More precisely, if  $\mathcal{G}\Gamma$  stands for all bi-infinite geodesics  
 26  $\gamma: \mathbf{R} \rightarrow \Gamma$ , the flow  $\phi_t$  acts on  $\mathcal{G}\Gamma$  by shifting the time parameter by  $t$ . The issue is that this  
 27 dynamical system is rather pathological. Generally, any two points in the boundary at infinity  $\partial\Gamma$   
 28 of  $\Gamma$  are joined by infinitely many orbits of the flow.

29 As Gromov explained in [25] – see also Coornaert and Papadopoulos [15, 16] – one can restrict our  
 30 attention to an invariant subset of  $\mathcal{G}\Gamma$ : roughly speaking, all the bi-infinite geodesics whose labels  
 31 are minimal for the lexicographic order induced by some fixed, arbitrary order on the symmetric,  
 32 generating set  $S$  of  $G$ . Formally the system that we consider is the following. One introduces a  
 33 space  $\mathfrak{H}_0(\Gamma)$  of horofunctions on  $\Gamma$  (which generalize the Busemann functions) on which the group  
 34  $G$  acts. Any horofunction  $h \in \mathfrak{H}_0(\Gamma)$  naturally comes with a *preferred* gradient line starting at 1,  
 35 i.e. whose labelling is minimal for the fixed lexicographic order. Calling  $\theta(h)$  the first letter of our  
 36 preferred gradient line, the transformation  $T: \mathfrak{H}_0(\Gamma) \rightarrow \mathfrak{H}_0(\Gamma)$  is defined by sending  $h$  to  $\theta(h)^{-1}h$ .  
 37 Remarkably, the system  $(\mathfrak{H}_0(\Gamma), T)$  is conjugated to a subshift of finite type  $(\Sigma, \sigma)$ . Geometrically,  
 38 the suspension of  $(\Sigma, \sigma)$  should be thought of as analogue of the geodesic flow on the unit tangent  
 39 bundle of compact, negatively curved manifold  $M$ , while the suspension of the extension  $(\Sigma_\theta, \sigma_\theta)$   
 40 plays the role of the geodesic flow on the unit tangent bundle of its universal cover  $\tilde{M}$ . Nevertheless,  
 41 unlike in the Riemannian setting, this flow is neither mixing nor, a-priori, topologically transitive.  
 42 This reflects the fact that two points in the boundary at infinity of  $\Gamma$  can still be joined by finitely  
 43 many orbits of the flow.

1 Actually, the dynamical properties of  $(\mathfrak{H}_0(\Gamma), T)$  are very sensitive to the choice of the order on  $S$ .  
 2 For instance, if  $G$  is the direct product of the free group with a finite group, then  $\mathfrak{H}_0(\Gamma)$  naturally  
 3 splits into several disjoint “layers”  $\mathfrak{J}_0, \dots, \mathfrak{J}_n$ , where  $\mathfrak{J}_0$  is invariant under the action of  $T$ . Moreover,  
 4 depending on the choice of the order on  $S$ , the other layers  $\mathfrak{J}_i$  are either invariant under the action  
 5 of  $T$ , or mapped into  $\mathfrak{J}_0$ . For more details, we refer the reader to [Example 4.9](#).

6 To circumvent this difficulty, we are forced to restrict our study to an irreducible component  $\mathfrak{J}$   
 7 of the system  $(\mathfrak{H}_0(\Gamma), T)$ . The price to pay though, is that the extension of  $(\mathfrak{J}, T)$  by the map  
 8  $\theta: \mathfrak{H}_0(\Gamma) \rightarrow G$  may not “visit” the whole Cayley graph  $\Gamma$  of  $G$ . If it misses a large portion  
 9 of  $\Gamma$ , then this system will be useless for counting purposes. However, we show that there ex-  
 10 ists an irreducible component  $\mathfrak{J}$  whose extension has the visibility property (see [Definition 2.1](#)).  
 11 Our strategy to produce such an irreducible component is inspired by an idea of Constantine, La-  
 12 font and Thompson [10], and based on a construction of Gromov. Namely, in [25] Gromov build  
 13 from  $(\mathcal{G}\Gamma, \phi_t)$  a new flow  $(\widehat{\mathcal{G}}\Gamma, \psi_s)$  with enhanced properties – see also [30, 8]. The space  $\widehat{\mathcal{G}}\Gamma$   
 14 quasi-isometric to  $\Gamma$ , hence its boundary at infinity is homeomorphic to  $\partial\Gamma$ ; every two points in  $\partial\Gamma$   
 15 are joined by a *unique* orbit of the flow  $\psi_s$ ; there is a natural projection  $\mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$  which send every  
 16  $\phi_t$ -orbit homeomorphically onto a  $\psi_s$ -orbit. It turns out that the new flow  $(\widehat{\mathcal{G}}\Gamma, \psi_s)$  is topologically  
 17 transitive. In particular, it admits a dense orbit. The irreducible component  $\mathfrak{J}$  is, roughly speaking,  
 18 the closure of a lift of this dense orbit. The transitivity of  $(\widehat{\mathcal{G}}\Gamma, \psi_s)$  tells us that the the extension  
 19 of  $(\mathfrak{J}, T)$  passes uniformly near every point of  $\Gamma$ , hence ensuring the visibility property.

20 In order to apply our criterion ([Theorem 1.2](#)) to the system  $(\mathfrak{J}, T)$  we finally need to define a  
 21 potential  $F: \mathfrak{J} \rightarrow \mathbf{R}_+^*$  with bounded Hölder variations. Recall that the dynamical system  $(\mathfrak{J}, T)$  was  
 22 not build directly from the metric space  $X$  we are interested in: thus, the role of the potential is  
 23 to reflect the geometry of  $X$ . If  $X$  coincides with the Cayley graph of  $\Gamma$  (which corresponds to the  
 24 first case of [Theorem 1.1](#)), we simply take for  $F$  the constant map equal to 1. In this situation we  
 25 prove that the spectral radius of the corresponding Ruelle transfer operator  $\mathcal{L}$  is  $\rho = e^{\omega_G}$ , whereas  
 26 the one of the twisted transfer operator satisfies  $\rho_\lambda \geq e^{\omega_H}$ . Hence the conclusion of [Theorem 1.1](#)  
 27 directly follows from our amenability criterion. If  $X$  is a CAT(−1) space (which corresponds to  
 28 the second case of [Theorem 1.1](#)) we use a quasi-isometry between  $\Gamma$  and  $X$  to define a potential  $F$   
 29 describing the geometry of  $X$ . In this situation the CAT(−1) geometry is crucial to ensure that  $F$   
 30 has bounded Hölder variations. Once this is done, we provide as before estimates of the spectral  
 31 radii  $\rho$  and  $\rho_\lambda$  in terms of  $\omega_G$  and  $\omega_H$  and conclude by the amenability criterion.

32 **Acknowledgment.** The authors would like to thank the organizers of the *Ventotene International*  
 33 *Workshops* (2015) where this work was initiated. The first author acknowledges support from the  
 34 Agence Nationale de la Recherche under Grant *Dagger* ANR-16-CE40-0006-01. The first and second  
 35 authors are grateful to the *Centre Henri Lebesgue* ANR-11-LABX-0020-01 for creating an attractive  
 36 mathematical environment. Many thanks also go to the referees for their helpful comments and  
 37 corrections.

## 38 2 Preliminaries

### 39 2.1 Subshift of finite type

40 **Vocabulary.** Let  $\mathcal{A}$  be a finite set. We write  $\mathcal{A}^*$  for the set of all finite words over the alphabet  
 41  $\mathcal{A}$ . The length of a word  $w \in \mathcal{A}^*$  is denoted by  $|w|$ . Given  $n \in \mathbf{N}$ , the set of all words of length  $n$  is

denoted by  $\mathcal{A}^n$ . The set  $\mathcal{A}^{\mathbf{N}}$  is endowed with a distance  $d$  defined as follows: given  $x, y \in \Sigma$ , we let  $d(x, y) = e^{-n}$  where  $n$  is the length of the longest common prefix of  $x$  and  $y$ . Let  $\sigma: \mathcal{A}^{\mathbf{N}} \rightarrow \mathcal{A}^{\mathbf{N}}$  be the shift operator, i.e. the map sending the sequence  $(x_i)_{i \in \mathbf{N}}$  to  $(x_{i+1})_{i \in \mathbf{N}}$ . Let  $\Sigma$  be a *subshift* i.e. a closed  $\sigma$ -invariant subset  $\mathcal{A}^{\mathbf{N}}$ . A word  $w \in \mathcal{A}^*$  is *admissible* if it is the prefix of a sequence  $x = (x_i)$  in  $\Sigma$ . We denote by  $\mathcal{W}$  the set of all finite admissible words. For every  $n \in \mathbf{N}$ , we write  $\mathcal{W}^n = \mathcal{W} \cap \mathcal{A}^n$  for the set of admissible words of length  $n$ . Given  $w \in \mathcal{W}$ , the *cylinder of  $w$* , denoted by  $[w]$ , is the set of sequences  $x \in \Sigma$  such that the prefix of length  $|w|$  of  $x$  is exactly  $w$ . We refer to  $|w|$  as the *length* of the cylinder. From now on we assume that  $\Sigma$  is a *subshift of finite type*, i.e. there exists  $N \in \mathbf{N}$  with the following property: a sequence  $x \in \mathcal{A}^{\mathbf{N}}$  belongs to  $\Sigma$  if and only if every subword of length  $N$  of  $\Sigma$  belongs to  $\mathcal{W}^N$ . We say that  $(\Sigma, \sigma)$  is *irreducible* or *topologically transitive* if  $(\Sigma, \sigma)$  admits a dense orbit. As  $(\Sigma, \sigma)$  is a subshift of finite type it is equivalent to ask that one of the following holds.

- (i) For every  $x, y \in \Sigma$ , for every  $\varepsilon \in \mathbf{R}_+^*$ , there exists  $k \in \mathbf{N}$  and  $x' \in \Sigma$  such that  $d(x, x') \leq \varepsilon$  and  $\sigma^k x' = y$
- (ii) For every  $w, w' \in \mathcal{W}$ , there exists  $w_0 \in \mathcal{W}$  such that the concatenation  $ww_0w'$  is admissible

**Irreducible components.** We associate to  $(\Sigma, \sigma)$  an oriented graph  $\Gamma = (V, E)$  labeled by  $\mathcal{A}$  describing the dynamics of the shift. We still denote by  $N$  the integer given by the definition of a subshift of finite type. The vertex set  $V$  of  $\Gamma$  is simply  $\mathcal{W}^N$ . Given two words  $w_1, w_2 \in \mathcal{W}^N$  and a letter  $a \in \mathcal{A}$ , there is an edge from  $w_1$  to  $w_2$  labelled by  $a$  if  $w_1$  is the prefix of length  $N$  of  $aw_2$ . It follows from the definition of  $N$ , that the labelling of  $\Gamma$  induces a one-to-one correspondance between  $\Sigma$  and the set of infinite oriented paths in  $\Gamma$ . We now define an equivalence relation on  $\mathcal{W}^N$  seen as the vertex set of  $\Gamma$ . Two vertices  $w, w' \in \mathcal{W}^N$  are *communicating* and we write  $w \sim w'$  if there exists an oriented loop in  $\Gamma$  passing through  $w$  and  $w'$ . The corresponding equivalence classes  $V_1, \dots, V_m$  are called *communicating classes*.

Given  $i \in \llbracket 1, m \rrbracket$ , we write  $\Gamma_i$  for the full subgraph of  $\Gamma$  associated to  $V_i$ . The subspace  $\Sigma_i \subset \Sigma$  is defined as the set of all sequences of  $\mathcal{A}^{\mathbf{N}}$  labeling an infinite path in  $\Gamma_i$ . Observe that  $\Sigma_i$  is a closed  $\sigma$ -invariant subset of  $\Sigma$  and  $(\Sigma_i, \sigma)$  is a subshift of finite type. It follows from the construction that  $(\Sigma_i, \sigma)$  is irreducible.

We observe that for every sequence  $x \in \Sigma$ , there exists  $n_0 \in \mathbf{N}$ , and a unique  $i \in \llbracket 0, m \rrbracket$  such that for all integers  $n \geq n_0$ , the sequence  $\sigma^n x$  belongs to  $\Sigma_i$ . Indeed one can derive from the relation  $\sim$  a new graph  $\Gamma/\sim$  defined as follows. Its vertex set is the set of communicating classes  $V_1, \dots, V_m$ . There is an edge in  $\Gamma/\sim$  from  $V_i$  to  $V_j$ , if  $\Gamma$  contains an edge joining a vertex in  $V_i$  to a vertex in  $V_j$ . The graph  $\Gamma/\sim$  comes with a natural projection  $\Gamma \rightarrow \Gamma/\sim$ . It follows from the definition of  $\sim$  that  $\Gamma/\sim$  does not contain any oriented loop. Hence if  $c$  is an infinite path in  $\Gamma$ , its projection in  $\Gamma/\sim$  is ultimately constant. This means that there exists  $i \in \llbracket 1, m \rrbracket$  such that the path obtained from  $c$  by removing its first edges is contained in  $\Gamma_i$ . Hence if  $x$  is the sequence labelling  $c$ , there exists  $n_0 \in \mathbf{N}$ , such that for all integers  $n \geq n_0$ , the sequence  $\sigma^n x$  belongs to  $\Sigma_i$ . We say that  $\Sigma_i$  is the *asymptotic irreducible component of  $x$* .

**Group extension.** Let  $G$  be a finitely generated group. Let  $\theta: \Sigma \rightarrow G$  be a continuous map. We denote by  $\Sigma_\theta$  the direct product  $\Sigma_\theta = \Sigma \times G$  endowed with the product topology. We define a continuous map  $\sigma_\theta: \Sigma_\theta \rightarrow \Sigma_\theta$  by

$$\sigma_\theta(x, g) = (\sigma x, g\theta(x)), \quad \forall (x, g) \in \Sigma_\theta.$$



1 The dynamical system  $(\Sigma_\theta, \sigma_\theta)$  is called the *extension of  $(X, \sigma)$  by  $\theta$* . For every  $n \in \mathbf{N}$ , for every  
 2  $x \in \Sigma$  we define the cocycle  $\theta_n(x)$  by

$$3 \quad \theta_n(x) = \theta(x)\theta(\sigma x) \cdots \theta(\sigma^{n-1}x). \quad (3)$$

4 By convention  $\theta_0: \Sigma \rightarrow G$  is the constant map equal to 1 (the identity of  $G$ ). Hence we have

$$5 \quad \sigma_\theta^n(x, g) = (\sigma^n x, g\theta_n(x)).$$

6 **Definition 2.1.** We say that the extension  $(\Sigma_\theta, \sigma_\theta)$  has the *visibility property* if there exists a finite  
 7 subset  $U$  of  $G$  with the following property: for every  $g \in G$ , there is a point  $x \in \Sigma$ , an integer  
 8  $n \in \mathbf{N}$  and two elements  $u_1, u_2 \in U$  such that  $g = u_1\theta_n(x)u_2$ .

9 **Remark.** This definition is somewhat reminiscent of Dougall's *linear visibility with remainder*  
 10 property [21, Definition 3.1]. Nevertheless our notion is more flexible as we do not ask to control  
 11 the value of  $n$  in terms of the word length of  $g$ .

## 12 2.2 Hyperbolic geometry

13 **The four point inequality.** Let  $(X, d)$  be a geodesic metric space. The Gromov product of  
 14 three points  $x, y, z \in X$  is defined by

$$15 \quad \langle x, y \rangle_z = \frac{1}{2} \{d(x, z) + d(y, z) - d(x, y)\}.$$

16 We assume that the space  $X$  is  $\delta$ -hyperbolic, i.e. for every  $x, y, z, t \in X$ ,

$$17 \quad \langle x, z \rangle_t \geq \min \{ \langle x, y \rangle_t, \langle y, z \rangle_t \} - \delta, \quad (4)$$

18 **The boundary at infinity.** Let  $x_0$  be a base point of  $X$ . A sequence  $(y_n)$  of points of  $X$   
 19 *converges to infinity* if  $\langle y_n, y_m \rangle_{x_0}$  tends to infinity as  $n$  and  $m$  approach to infinity. The set  $\mathcal{S}$  of  
 20 such sequences is endowed with a binary relation defined as follows. Two sequences  $(y_n)$  and  $(z_n)$   
 21 are related if

$$22 \quad \lim_{n \rightarrow +\infty} \langle y_n, z_n \rangle_{x_0} = +\infty.$$

23 It follows from (4) that this relation is actually an equivalence relation. The *boundary at infinity*  
 24 of  $X$  denoted by  $\partial X$  is the quotient of  $\mathcal{S}$  by this relation. If the sequence  $(y_n)$  is an element in the  
 25 class of  $\xi \in \partial X$  we say that  $(y_n)$  *converges to  $\xi$*  and write

$$26 \quad \lim_{n \rightarrow +\infty} y_n = \xi.$$

27 Note that the definition of  $\partial X$  does not depend on the base point  $x_0$ . For more details about the  
 28 Gromov boundary, we refer the reader to [13, Chapitre 2]. The notation  $\partial^2 X$  stands for

$$29 \quad \partial^2 X = \partial X \times \partial X \setminus \text{diag}(\partial X),$$

30 where  $\text{diag}(\partial X) = \{(\xi, \xi) \mid \xi \in \partial X\}$  is the diagonal.

1 **Quasi-geodesics.** One major feature of hyperbolic spaces is the stability of quasi-geodesics also  
 2 known as Morse's Lemma.

3 **Definition 2.2.** Let  $\kappa \in \mathbf{R}_+^*$  and  $\ell \in \mathbf{R}_+$ . Let  $f: X_1 \rightarrow X_2$  be a map between two metric spaces.  
 4 We say that  $f$  is a  $(\kappa, \ell)$ -quasi-isometric embedding, if for every  $x, x' \in X_1$  we have

$$5 \quad \kappa^{-1}d(x, x') - \ell \leq d(f(x), f(x')) \leq \kappa d(x, x') + \ell.$$

6 A  $(\kappa, \ell)$ -quasi-geodesic of  $X$ , is a  $(\kappa, \ell)$ -quasi-isometric embedding of an interval of  $\mathbf{R}$  into  $X$ .

7 Given a  $(\kappa, \ell)$ -quasi-geodesic  $\gamma: \mathbf{R}_+ \rightarrow X$ , there exists a point  $\xi \in \partial X$  such that for every  
 8 sequence  $(t_n)$  diverging to infinity, we have

$$9 \quad \lim_{n \rightarrow +\infty} \gamma(t_n) = \xi,$$

10 see for instance [13, Chapitre 3, Théorème 2.2]. In this situation we consider  $\xi$  as the endpoint  
 11 at infinity of  $\gamma$  and write  $\gamma(\infty) = \xi$ . If  $\gamma: \mathbf{R} \rightarrow X$  is a bi-infinite  $(\kappa, \ell)$ -quasi-geodesic, we define  
 12  $\gamma(-\infty)$  in the same way.

13 **Proposition 2.3** ([13, Chapitre 3, Théorèmes 1.2 et 3.1]). *Let  $\kappa \in \mathbf{R}_+^*$  and  $\ell \in \mathbf{R}_+$ . There exists*  
 14  *$D = D(\kappa, \ell, \delta)$  in  $\mathbf{R}_+$  such that the Hausdorff distance between any two  $(\kappa, \ell)$ -quasi-geodesics with*  
 15 *the same endpoints (possibly in  $\partial X$ ) is at most  $D$ .*

16 **Group action.** Let  $x_0$  be a base point of  $X$ . Let  $G$  be a group acting properly by isometries on  
 17  $X$ . The *exponential growth rate of  $G$  acting on  $X$*  is the quantity

$$18 \quad \omega(G, X) = \limsup_{r \rightarrow \infty} \frac{1}{r} \ln |\{g \in G \mid d(gx_0, x_0) \leq r\}|. \quad (5)$$

19 Note that  $\omega(G, X)$  does not depend on  $x_0$ . It can also be interpreted as the critical exponent of the  
 20 Poincaré series

$$21 \quad \mathcal{P}_G(s) = \sum_{g \in G} e^{-sd(gx_0, x_0)}.$$

22 If there is no ambiguity, we simply write  $\omega_G$  instead of  $\omega(G, X)$ .

### 23 3 Horofunctions

24 In this section we recall the definition of horofunctions and gradient lines introduced by Gromov in  
 25 [25, Section 7.5]. We follow the exposition given by Coornaert and Papadopoulos in [14, 15]. Let  
 26  $(X, x_0)$  be a pointed geodesic  $\delta$ -hyperbolic space and  $G$  be a group acting by isometries on  $X$ .

27 **Horofunctions.** Let  $\varepsilon \in \mathbf{R}_+$ . A map  $f: X \rightarrow \mathbf{R}$  is  $\varepsilon$ -quasi-convex if for every geodesic  $\gamma: I \rightarrow X$ ,  
 28 for every  $a, b \in I$ , for every  $t \in [0, 1]$  we have

$$29 \quad f \circ \gamma(ta + (1-t)b) \leq tf \circ \gamma(a) + (1-t)f \circ \gamma(b) + \varepsilon.$$

30 The map  $f$  is *quasi-convex* if there exists  $\varepsilon \in \mathbf{R}_+$  such that  $f$  is  $\varepsilon$ -quasi-convex.

**Definition 3.1.** A *horofunction* is a quasi-convex map  $h: X \rightarrow \mathbf{R}$  satisfying the following distance-like property: for every  $x \in X$ , for every  $r \in \mathbf{R}$ , if  $r \leq h(x)$ , then

$$h(x) = r + d(x, h^{-1}(r)). \quad (6)$$

A *cocycle* is a map  $c: X \times X \rightarrow \mathbf{R}$  of the form  $c(x, y) = h(x) - h(y)$  where  $h$  is a horofunction. In this situation,  $h$  is called a *primitive* of  $c$ .

Let  $C(X)$  be the set of all continuous function  $f: X \rightarrow \mathbf{R}$  endowed with the topology of uniform convergence on compact subsets. We denote by  $C_*(X)$  the quotient of  $C(X)$  by the 1-dimensional closed subspace of constant functions. The space  $C_*(X)$  is endowed with the quotient topology. It can be identified with the set of all continuous functions  $f: X \rightarrow \mathbf{R}$  vanishing at  $x_0$ . This is the point of view that we adopt here. The action of  $G$  on  $X$  induces an action on  $C(X)$  and thus on  $C_*(X)$ . More precisely  $g \in G$ , and  $f \in C_*(X)$  the map  $g \cdot f$  is defined by

$$[g \cdot f](x) = f(g^{-1}x) - f(g^{-1}x_0), \quad \forall x \in X.$$

A horofunction is automatically 1-Lipschitz, hence continuous [15, Proposition 2.2]. Moreover, given a cocycle  $c$ , any two primitives  $h_1$  and  $h_2$  of  $c$  differ by a constant, i.e. there exists  $a \in \mathbf{R}$ , such that for every  $x \in X$ , we have  $h_2(x) - h_1(x) = a$ . Hence the set of all cocycles, or equivalently all horofunctions vanishing at  $x_0$ , embeds in  $C_*(X)$ . We denote it by  $\mathfrak{H}(X)$ . It is a compact subspace of  $C_*(X)$  [15, Proposition 3.9].

**Gradient lines.** The gradient lines are an important tool to track the behavior of horofunctions.

**Definition 3.2.** Let  $h \in \mathfrak{H}(X)$  be a horofunction. A *gradient line* for  $h$  or an  *$h$ -gradient line* is a path  $\gamma: I \rightarrow X$  parametrized by arc length such that for every  $t, t' \in I$  we have

$$h(\gamma(t)) - h(\gamma(t')) = t' - t.$$

If the interval  $I$  has the form  $I = \mathbf{R}_+$  we say that  $\gamma$  a  *$h$ -gradient ray*.

Let us recall a few properties of gradient lines. Let  $h \in \mathfrak{H}(X)$ . Every  $h$ -gradient line is a geodesic [15, Proposition 2.10]. If  $\gamma: I \rightarrow X$  be an  $h$ -gradient line, then for every  $g \in G$ , the path  $g\gamma$  is a gradient line for  $gh$  [15, Proposition 2.14]. For every  $x \in X$ , there exists an  $h$ -gradient ray starting at  $x$  [15, Proposition 2.13].

Let  $\gamma: \mathbf{R}_+ \rightarrow X$  be a geodesic ray. The *Busemann function* associated to  $\gamma$  is the map  $b_\gamma: X \rightarrow \mathbf{R}$  defined by

$$b_\gamma(x) = \lim_{t \rightarrow \infty} [d(x, \gamma(t)) - t], \quad \forall x \in X.$$

The map  $c_\gamma: X \times X \rightarrow \mathbf{R}$  defined by  $c_\gamma(x, y) = b_\gamma(x) - b_\gamma(y)$  is a cocycle [14, Chapter 3, Proposition 3.6]. Moreover  $\gamma$  is a gradient ray starting at  $\gamma(0)$  for this cocycle. However, one has to be careful that  $c_\gamma$  may admit other gradient lines starting at  $\gamma(0)$ . Similarly, given a horofunction  $h \in \mathfrak{H}(X)$ , the Busemann function associated to an  $h$ -gradient ray starting at  $x_0$  is not necessarily  $h$ .

1 **Comparison with the Gromov boundary.** Recall that  $\partial X$  is the Gromov boundary of  $X$ .

2 **Proposition 3.3** (Coornaert-Papadopoulos [15, Proposition 3.1]). *Let  $h \in \mathfrak{H}(X)$  be a horofunction.*  
 3 *Let  $\gamma_1: \mathbf{R}_+ \rightarrow X$  and  $\gamma_2: \mathbf{R}_+ \rightarrow X$  be two  $h$ -gradient rays. Then  $\gamma_1(\infty) = \gamma_2(\infty)$ .*

4 This gives rise to a map  $\pi: \mathfrak{H}(X) \rightarrow \partial X$  which associates to any  $h \in \mathfrak{H}(X)$  the endpoint at  
 5 infinity of any  $h$ -gradient ray.

6 **Proposition 3.4** (Coornaert-Papadopoulos [15, Proposition 3.3 and Corollary 3.8]). *The map*  
 7  *$\pi: \mathfrak{H}(X) \rightarrow \partial X$  is continuous,  $G$ -equivariant and onto. More precisely for every geodesic ray*  
 8  *$\gamma: \mathbf{R}_+ \rightarrow X$  starting at  $x_0$  the corresponding Busemann function  $b_\gamma$  is a preimage of  $\gamma(\infty)$  in*  
 9  *$\mathfrak{H}(X)$ . Two horofunctions  $h, h' \in \mathfrak{H}(X)$  have the same image in  $\partial X$  if and only if  $\|h - h'\|_\infty \leq 64\delta$ .*

## 10 4 Dynamics in a hyperbolic group

11 In this section we introduce a few dynamical systems to describe the “geodesic flow” of a hyperbolic  
 12 Cayley graph. Let  $G$  be a hyperbolic group and  $A$  a finite generating set of  $G$ . For simplicity we  
 13 assume that  $A$  is symmetric, i.e.  $A^{-1} = A$ . We denote by  $\Gamma(G, A)$  or simply  $\Gamma$  the Cayley graph of  
 14  $G$  with respect to  $A$ . We identify  $G$  with the vertex set of  $\Gamma$ . We consider 1 as a base point in  $\Gamma$ .  
 15 For every  $n \in \mathbf{N}$ , we write  $S(n)$  for the sphere of radius  $n$  in  $\Gamma$  centered at the identity, i.e. the set  
 16 of all elements  $g \in G$  such that  $d_\Gamma(1, g) = n$ . Similarly we write  $B(n)$  for the closed ball of radius  
 17  $n$ .

### 18 4.1 Transitivity of Gromov’s geodesic flow

19 We denote by  $\mathcal{G}\Gamma$  the set of all *parametrized* bi-infinite geodesic  $\gamma: \mathbf{R} \rightarrow \Gamma$  of  $\Gamma$ . This set is endowed  
 20 with a distance defined as follows: given two bi-infinite geodesics  $\gamma_1, \gamma_2: \mathbf{R} \rightarrow \Gamma$  we let

$$21 \quad d(\gamma_1, \gamma_2) = \int_{-\infty}^{\infty} e^{-|t|} d(\gamma_1(t), \gamma_2(t)) dt. \quad (7)$$

22 The action of  $G$  on  $\Gamma$  induces an action by isometries of  $G$  on  $\mathcal{G}\Gamma$ . One checks easily that the map  
 23  $\mathcal{G}\Gamma \rightarrow \Gamma$  sending  $\gamma$  to  $\gamma(0)$  is a  $G$ -equivariant quasi-isometry. The space  $\mathcal{G}\Gamma$  also comes with a flow  
 24  $\phi = (\phi_s)_{s \in \mathbf{R}}$  defined as follows: for every  $\gamma \in \mathcal{G}\Gamma$ , for every  $s \in \mathbf{R}$ , the geodesic  $\phi_s(\gamma): \mathbf{R} \rightarrow \Gamma$  is  
 25 given by

$$26 \quad \phi_s(\gamma)(t) = \gamma(s + t), \quad \forall t \in \mathbf{R}.$$

27 Starting from  $\mathcal{G}\Gamma$ , Gromov build a new hyperbolic space  $\widehat{\mathcal{G}}\Gamma$  that is quasi-isometric to  $\Gamma$ . In  
 28 particular it is hyperbolic and its boundary at infinity is homeomorphic to  $\partial\Gamma$ . Moreover  $\widehat{\mathcal{G}}\Gamma$  comes  
 29 with a flow so that any two distinct points of  $\partial\Gamma$  are joined by a *unique* orbit of the flow. The  
 30 construction is given in [25, Section 8.3], the details can be found in [8, 30]. We recall here the  
 31 main properties of this space.

32 (F1) The space  $\widehat{\mathcal{G}}\Gamma$  is geodesic and proper. It is endowed with a proper co-compact action by  
 33 isometries of  $G$  as well as a flow  $\psi = (\psi_s)_{s \in \mathbf{R}}$ . The flow and the action of  $G$  commute, i.e.  
 34 for every  $\widehat{\gamma} \in \widehat{\mathcal{G}}\Gamma$ , for every  $s \in \mathbf{R}$  and  $g \in G$  we have  $\psi_s(g\widehat{\gamma}) = g\psi_s(\widehat{\gamma})$ .

1 (F2) There exists a continuous  $G$ -equivariant quasi-isometric projection  $p: \mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$ . In particular  
 2  $p$  induces a homeomorphism  $p_\infty$  from  $\partial\Gamma$  onto the boundary at infinity of  $\widehat{\mathcal{G}}\Gamma$ . In addition,  
 3 for every geodesic  $\gamma \in \mathcal{G}\Gamma$ , the projection  $p$  maps the  $\phi$ -orbit of  $\gamma \in \mathcal{G}\Gamma$  homeomorphically  
 4 onto the  $\psi$ -orbit of  $\widehat{\gamma} = p(\gamma)$ .

5 (F3) For every point  $\widehat{\gamma} \in \widehat{\mathcal{G}}\Gamma$ , the map  $\mathbf{R} \rightarrow \widehat{\mathcal{G}}\Gamma$  sending  $s$  to  $\psi_s(\widehat{\gamma})$  is a quasi-isometric embedding  
 6 of  $\mathbf{R}$  into  $\widehat{\mathcal{G}}\Gamma$ . Hence for every point  $\widehat{\gamma} \in \widehat{\mathcal{G}}\Gamma$  one can associate two distinct points in the  
 7 boundary at infinity of  $\widehat{\mathcal{G}}\Gamma$  defined by

$$8 \quad \widehat{\gamma}(\infty) = \lim_{s \rightarrow \infty} \psi_s(\widehat{\gamma}) \quad \text{and} \quad \widehat{\gamma}(-\infty) = \lim_{s \rightarrow -\infty} \psi_s(\widehat{\gamma}).$$

9 By construction for every geodesic  $\gamma \in \mathcal{G}\Gamma$ , the homeomorphism  $p_\infty$  maps  $\gamma(\infty)$  and  $\gamma(-\infty)$   
 10 to  $\widehat{\gamma}(\infty)$  and  $\widehat{\gamma}(-\infty)$ , where  $\widehat{\gamma} = p(\gamma)$ .

11 (F4) The map  $\widehat{\mathcal{G}}\Gamma \rightarrow \partial^2\Gamma$  sending  $\widehat{\gamma}$  to  $(\widehat{\gamma}(-\infty), \widehat{\gamma}(\infty))$  induces a homeomorphism from  $\widehat{\mathcal{G}}\Gamma/\mathbf{R}$  onto  
 12  $\partial^2\Gamma$ . Actually  $\widehat{\mathcal{G}}\Gamma$  is homeomorphic to  $\partial^2\Gamma \times \mathbf{R}$ .

13 It is important to notice that in general  $p$  does not conjugate the flow, i.e.  $p \circ \phi_s \neq \psi_s \circ p$ .  
 14 Since the flow  $\psi$  and the action of  $G$  commute, the flow  $\psi$  induces a flow on  $\widehat{\mathcal{G}}\Gamma/G$  that we denote  
 15  $\overline{\psi} = (\overline{\psi}_s)_{s \in \mathbf{R}}$ .

16 **Proposition 4.1.** *The flow  $\overline{\psi}$  on  $\widehat{\mathcal{G}}\Gamma/G$  is topologically transitive, i.e. given any two non-empty  
 17 open subsets  $U$  and  $V$  of  $\widehat{\mathcal{G}}\Gamma/G$ , there exists  $s \in \mathbf{R}$  such that  $\overline{\psi}_s(U) \cap V$  is non-empty.*

18 The remainder of the section is dedicated to the proof of this proposition. We follow mostly the  
 19 strategy used in [2, Chapter III].

20 **Lemma 4.2.** *Let  $\gamma, \gamma' \in \mathcal{G}\Gamma$  such that  $\gamma(\infty) \neq \gamma'(-\infty)$ . There are sequences  $(\nu_n)$  of geodesics  
 21 in  $\mathcal{G}\Gamma$ ,  $(g_n)$  of elements in  $G$  and  $(t_n)$  of numbers in  $\mathbf{R}$  diverging to infinity with the following  
 22 properties.*

23 (i)  $(\nu_n)$  converges to a geodesic with the same endpoints as  $\gamma$ .

24 (ii)  $(g_n \phi_{t_n}(\nu_n))$  converges to a geodesic with the same endpoints as  $\gamma'$ .

25 *Proof.* For simplicity we write  $\xi_+$  and  $\xi_-$  for  $\gamma(\infty)$  and  $\gamma(-\infty)$ . Similarly we define  $\xi'_+$  and  $\xi'_-$ .  
 26 Since  $\xi_+ \neq \xi'_-$ , there exists a sequence  $(g_n)$  of elements of  $G$  such that for some (hence any)  $x \in \Gamma$   
 27 the sequence  $(g_n x)$  and  $(g_n^{-1} x)$  respectively converge to  $\xi'_-$  and  $\xi_+$  – see for instance [18, Chapter III,  
 28 Lemma 2.2.]. Nevertheless, the action of  $G$  on  $\Gamma \cup \partial\Gamma$  is a convergence action. Up to replacing  
 29  $(g_n)$  by a subsequence, we can assume that for every  $\xi \in \partial\Gamma \setminus \{\xi_+\}$ , the sequence  $(g_n \xi)$   
 30 converges to  $\xi'_-$  and for every  $\xi \in \partial\Gamma \setminus \{\xi'_-\}$ , the sequence  $(g_n^{-1} \xi)$  converges to  $\xi_+$  [37, Theorem 3A].  
 31 For every  $n \in \mathbf{N}$ , we denote by  $\nu_n: \mathbf{R} \rightarrow \Gamma$  a bi-infinite geodesic joining  $\xi_-$  to  $g_n^{-1} \xi'_+$ . We observe  
 32 that

33 ▶  $\nu_n(-\infty)$  and  $\nu_n(\infty)$  respectively converge to  $\xi_-$  and  $\xi_+$ , whereas

34 ▶  $g_n \nu_n(-\infty)$  and  $g_n \nu_n(\infty)$  respectively converge to  $\xi'_-$  and  $\xi'_+$ .

1 Geodesic triangles in  $\Gamma \cup \partial\Gamma$  are  $24\delta$ -thin [13, Chapitre 2 Proposition 2.2]. Up to passing again  
 2 to a subsequence we may assume that  $d(\gamma(0), \nu_n) \leq 24\delta$  and  $d(\gamma'(0), g_n\nu_n) \leq 24\delta$ . By shifting if  
 3 necessary the origin of  $\nu_n$  we can assume that for every  $n \in \mathbf{N}$ , there exists  $t_n \in \mathbf{R}$ , such that  
 4  $d(\gamma(0), \nu_n(0)) \leq 24\delta$  and  $d(\gamma'(0), g_n\nu_n(t_n)) \leq 24\delta$ . According to the Azelà-Ascoli theorem  $(\nu_n)$   
 5 converges to a geodesic  $\nu$ . It follows from our choice of  $(g_n)$  that  $\nu$  has the same endpoints as  $\gamma$ .  
 6 Similarly we obtain that  $(g_n\phi_{t_n}(\nu_n))$  converges to a geodesic with the same endpoints as  $\gamma'$ . Note  
 7 also that  $d(\nu_n(t_n), g_n^{-1}\gamma'(0))$  and  $d(\nu_n(0), \gamma(0))$  are uniformly bounded. As  $(g_n^{-1}\gamma'(0))$  converges to  
 8  $\xi_+$ , the sequence  $(t_n)$  has to diverge to  $\infty$ .  $\square$

9 *Proof of Proposition 4.1.* It suffices to show that for every non-empty open subset  $\widehat{U}$  and  $\widehat{V}$  of  $\widehat{\mathcal{G}}\Gamma$ ,  
 10 there exists  $s \in \mathbf{R}$ , and  $g \in G$  such that  $g\psi_s(\widehat{U}) \cap \widehat{V} \neq \emptyset$ . Let  $\widehat{U}$  and  $\widehat{V}$  be two non-empty open  
 11 subsets of  $\widehat{\mathcal{G}}\Gamma$ . We denote by  $\widehat{U}(\infty)$  the following subset of  $\partial\Gamma$

$$12 \quad \widehat{U}(\infty) = \left\{ \widehat{\gamma}(\infty) \mid \widehat{\gamma} \in \widehat{U} \right\}.$$

13 The set  $\widehat{V}(-\infty)$  is defined in a similar way. It follows from (F4) that  $\widehat{U}(\infty)$  and  $\widehat{V}(-\infty)$  are open  
 14 non-empty subsets of  $\partial\Gamma$ . In particular they are not reduced to a point. Hence there exists  $\widehat{\gamma} \in \widehat{U}$   
 15 and  $\widehat{\gamma}' \in \widehat{V}$  such that  $\widehat{\gamma}(\infty) \neq \widehat{\gamma}'(-\infty)$ . Let  $\gamma, \gamma' \in \mathcal{G}\Gamma$  be respective preimages of  $\widehat{\gamma}$  and  $\widehat{\gamma}'$ .  
 16 According to (F3),  $\gamma(\infty) \neq \gamma'(-\infty)$ . Applying Lemma 4.2, there are sequences  $(\nu_n)$  of geodesics  
 17 in  $\mathcal{G}\Gamma$ ,  $(g_n)$  of elements in  $G$  and  $(t_n)$  of numbers in  $\mathbf{R}$  diverging to infinity with the following  
 18 properties: the geodesic  $\nu_n$  converges to a geodesic  $\nu$  with the same endpoints as  $\gamma$ ; the geodesic  
 19  $\nu'_n = g_n\phi_{t_n}(\nu_n)$  converges to a geodesic  $\nu'$  with the same endpoints as  $\gamma'$ .

20 We now push these data in  $\widehat{\mathcal{G}}\Gamma$  using the projection  $p: \mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$ . For every  $n \in \mathbf{N}$ , we let  
 21  $\widehat{\nu}_n = p(\nu_n)$  and  $\widehat{\nu}'_n = p(\nu'_n)$ . We define  $\widehat{\nu}$  and  $\widehat{\nu}'$  in the same way. Since  $p$  maps homeomorphically  
 22  $\phi$ -orbits onto  $\psi$ -orbits, there exists a sequence  $(s_n)$  of numbers in  $\mathbf{R}$ , diverging to  $\infty$  such that for  
 23 every  $n \in \mathbf{N}$ , we have

$$24 \quad p(\phi_{t_n}(\nu_n)) = \psi_{s_n}(\widehat{\nu}_n).$$

25 By construction the endpoints of  $\widehat{\nu}$  are the same as those of  $\widehat{\gamma}$ . However there is a unique orbit of  
 26 the flow  $\psi$  joining two distinct point of  $\partial\Gamma$  – see (F4). Thus there exists  $s \in \mathbf{R}$  such that  $\widehat{\gamma} = \psi_s(\widehat{\nu})$ .  
 27 Recall that  $\nu_n$  converges to  $\nu$ . Since the projection  $p$  is continuous,  $\widehat{\nu}_n$  converges to  $\widehat{\nu}$ , hence  $\psi_s(\widehat{\nu}_n)$   
 28 converges to  $\widehat{\gamma}$ . Similarly we prove that there exists  $s' \in \mathbf{R}$  such that  $\psi_{s'}(\widehat{\nu}'_n) = g_n\psi_{s_n+s'}(\widehat{\nu}_n)$   
 29 converges to  $\widehat{\gamma}'$ . As  $\widehat{U}$  and  $\widehat{V}$  are open, there exists  $n_0 \in \mathbf{N}$  such that for every  $n \geq n_0$ , the point  
 30  $\psi_s(\widehat{\nu}_n)$  belongs to  $\widehat{U}$  while  $g_n\psi_{s_n+s'}(\widehat{\nu}_n)$  belongs to  $\widehat{V}$ . Hence

$$31 \quad g_n\psi_{s_n+s'-s}(\widehat{U}) \cap \widehat{V} \neq \emptyset, \quad \forall n \geq n_0,$$

32 which completes the proof of the proposition.  $\square$

33 **Corollary 4.3** (Compare Dal'bo [18, Chapter III, Theorem 4.2]). *There exists a point  $\widehat{\gamma} \in \widehat{\mathcal{G}}\Gamma$  such*  
 34 *that the image of  $\{\psi_s(\widehat{\gamma}) \mid s \in \mathbf{R}_+\}$  in  $\widehat{\mathcal{G}}\Gamma/G$  is dense.*

## 35 4.2 Dynamics on the space of horofunctions

36 **Definition 4.4.** A horofunction  $h \in \mathfrak{H}(\Gamma)$  is *integral* if its restriction to the vertex set  $G$  takes  
 37 integer values. We write  $\mathfrak{H}_0(\Gamma)$  (or simply  $\mathfrak{H}_0$ ) for the subset of  $\mathfrak{H}(\Gamma)$  consisting of all integral  
 38 horofunctions.

1 Note that  $\mathfrak{H}_0$  is closed,  $G$ -invariant subset of  $\mathfrak{H}(\Gamma)$ . Moreover, the Busemann function of any  
 2 geodesic ray starting at 1 is an integral horofunction. Consequently Proposition 3.4 leads to the  
 3 following statement

4 **Proposition 4.5.** *The projection  $\pi: \mathfrak{H}(\Gamma) \rightarrow \partial\Gamma$  maps  $\mathfrak{H}_0$  onto  $\partial\Gamma$ .*

5 We now define a map  $T: \mathfrak{H}_0 \rightarrow \mathfrak{H}_0$  which can be thought as an analogue of the first return map  
 6 for a section of the geodesic flow. To that end we endow the generating set  $A$  with an arbitrary  
 7 total order. The map  $\theta: \mathfrak{H}_0 \rightarrow G$  is the one sending  $h$  to the smallest element  $a \in A$  such that  
 8  $h(a) = h(1) - 1$ . The existence of such a generator  $a \in A$  is ensured by [15, Lemma 5.1].

9 **Definition 4.6.** The map  $T: \mathfrak{H}_0 \rightarrow \mathfrak{H}_0$  is the one sending  $h$  to  $\theta(h)^{-1}h$ .

10 This map is continuous [15, Proposition 5.6]. Before speaking of coding, let us recall a few  
 11 properties of the dynamical system  $(\mathfrak{H}_0, T)$ . Recall that the edges of  $\Gamma$  are labelled by elements  
 12 of  $A$ . Let  $h \in \mathfrak{H}_0$  be a horofunction and  $x$  a vertex of  $\Gamma$ . The lexicographic order on  $A^{\mathbf{N}}$  induces  
 13 a total order on the set of  $h$ -gradient rays starting at  $x$ . This set admits a smallest element [15,  
 14 Proposition 5.2] that we call the *minimal  $h$ -gradient ray starting at  $x$* . Assume that  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$  is  
 15 the minimal  $h$ -gradient ray starting at  $x$ . We have the following properties:

- 16 (i) for every  $n \in \mathbf{N}$ , the path  $\gamma_n: \mathbf{R}_+ \rightarrow \Gamma$  defined by  $\gamma_n(t) = \gamma(t+n)$  is the minimal  $h$ -gradient  
 17 ray starting at  $\gamma(n)$  [15, Proposition 5.3];
- 18 (ii) for every  $g \in G$ ,  $g\gamma$  is the minimal  $gh$ -gradient ray starting at  $gx$  [15, Proposition 5.4].

19 As usual we define for every  $n \in \mathbf{N}$ , the cocycle  $\theta_n: \mathfrak{H}_0 \rightarrow G$  by

$$20 \quad \theta_n(h) = \theta(h)\theta(Th) \cdots \theta(T^{n-1}h). \quad (8)$$

21 It follows from the previous two observations, that this cocycle has the following geometric inter-  
 22 pretation: if  $\gamma: \mathbf{R} \rightarrow \Gamma$  is the minimal  $h$ -gradient line starting at 1, then  $\gamma(n) = \theta_n(h)$ .

23 **Connection with Gromov's geodesic flow.** We relate here  $(\mathfrak{H}_0, T)$  to Gromov's geodesic flow  
 24 on  $\Gamma$ . Let  $h \in \mathfrak{H}_0$  be an integral horofunction. A bi-infinite  $h$ -gradient line  $\gamma: \mathbf{R} \rightarrow \Gamma$  is called  
 25 *primitive* if the following holds

- 26 (i)  $\gamma(0)$  is a vertex of  $\Gamma$ , and thus  $\gamma(\mathbf{Z})$  is contained in  $G$ .
- 27 (ii) For every integer  $m \in \mathbf{Z}$ , the path  $\gamma_m: \mathbf{R}_+ \rightarrow \Gamma$  sending  $t$  to  $\gamma(m+t)$  is the minimal  $h$ -gradient  
 28 ray starting at  $\gamma(m)$ .

29 Observe that in this case, for every  $k \in \mathbf{Z}$ , the path  $\phi_k(\gamma)$  is also a primitive  $h$ -gradient line. Similarly  
 30 for every  $g \in G$ , the path  $g\gamma$  is a primitive  $gh$ -gradient line [16, Lemma 2.5]. Given  $h \in \mathfrak{H}_0$  and  
 31  $\xi \in \partial\Gamma$ , there is always a primitive  $h$ -gradient line such that  $\gamma(-\infty) = \xi$  [16, Proposition 5.2].

32 Let  $\mathfrak{H}_{\mathbf{R}}$  be the subset of  $\mathfrak{H}_0 \times \mathcal{G}\Gamma$  that consists of all pairs  $(h, \gamma)$  where  $h \in \mathfrak{H}_0$  is an integral  
 33 horofunction and  $\gamma \in \mathcal{G}\Gamma$  a geodesic in the  $\phi$ -orbit of a primitive  $h$ -gradient line [16, Definition 4.1].  
 34 We endow  $\mathfrak{H}_{\mathbf{R}}$  with the topology induced by the product topology on  $\mathfrak{H}_0 \times \mathcal{G}\Gamma$ . The set  $\mathfrak{H}_{\mathbf{R}}$  comes  
 35 with a flow, that we still denote  $\phi = (\phi_t)_{t \in \mathbf{R}}$ , defined as follows: for every  $(h, \gamma) \in \mathfrak{H}_{\mathbf{R}}$ ,

$$36 \quad \phi_t(h, \gamma) = (h, \phi_t(\gamma)), \quad \forall t \in \mathbf{R}.$$

1 We call the dynamical system  $(\mathfrak{H}_{\mathbf{R}}, \phi)$  the *horoflow* of  $\Gamma$ . This system shall not be confused with the  
 2 horocyclic flow on a hyperbolic surface. The group  $G$  acts on  $\mathfrak{H}_{\mathbf{R}}$  as follows: for every  $(h, \gamma) \in \mathfrak{H}_{\mathbf{R}}$   
 3 we let

$$4 \quad g(h, \gamma) = (gh, g\gamma), \quad \forall g \in G.$$

5 Note that the horoflow and the action of  $G$  commutes. In order to compare  $(\mathfrak{H}_{\mathbf{R}}, \phi)$  with  $(\widehat{\mathcal{G}}\Gamma, \psi)$ ,  
 6 we define a  $G$ -equivariant continuous map

$$7 \quad q: \mathfrak{H}_{\mathbf{R}} \rightarrow \widehat{\mathcal{G}}\Gamma,$$

8 by composing the map  $\mathfrak{H}_{\mathbf{R}} \rightarrow \mathcal{G}\Gamma$  and the projection  $p: \mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$ . By construction, given any  
 9  $(h, \gamma) \in \mathfrak{H}_{\mathbf{R}}$ , the map  $q$  maps homeomorphically the  $\phi$ -orbit of  $(h, \gamma)$  onto the  $\psi$ -orbit of  $q(h, \gamma) =$   
 10  $p(\gamma)$ .

11 **Proposition 4.7.** *The map  $q: \mathfrak{H}_{\mathbf{R}} \rightarrow \widehat{\mathcal{G}}\Gamma$  is onto.*

12 *Proof.* Let  $\eta$  and  $\xi$  be two distinct points of  $\partial\Gamma$ . According to [Proposition 3.4](#) there exists an integral  
 13 horofunction  $h \in \mathfrak{H}_0$  such that  $\pi(h) = \xi$ . On the other hand, there exists a primitive  $h$ -gradient  
 14 line  $\gamma: \mathbf{R} \rightarrow \Gamma$  such that  $\gamma(-\infty) = \eta$  [[16](#), Proposition 5.2]. Being an  $h$ -gradient line,  $\gamma$  is such that  
 15  $\gamma(\infty) = \xi$ . Hence  $q$  maps the  $\phi$ -orbit of  $(h, \gamma)$  to the (unique)  $\psi$ -orbit in  $\widehat{\mathcal{G}}\Gamma$  joining  $\eta$  to  $\xi$ . This  
 16 works for any two distinct points  $\eta, \xi \in \partial\Gamma$ . Hence  $q$  is onto.  $\square$

17 **Discretization of the flow.** We denote by  $\mathfrak{H}_{\mathbf{Z}}$  the closed subset of  $\mathfrak{H}_{\mathbf{R}}$  containing all the pairs  
 18  $(h, \gamma)$  where  $\gamma$  is a primitive  $h$ -gradient line. Observe that  $\mathfrak{H}_{\mathbf{Z}}$  is  $G$ -invariant. Moreover the time  
 19 1 flow  $\phi_1$  on  $\mathfrak{H}_{\mathbf{R}}$  induces a homeomorphism of  $\mathfrak{H}_{\mathbf{Z}}$  onto itself [[16](#), Definition 2.6]. The system  
 20  $(\mathfrak{H}_{\mathbf{Z}}, \phi_1)$  is called the *discrete horoflow* of  $\Gamma$ . Let  $\bar{\mathfrak{H}}_{\mathbf{Z}}$  and  $\bar{\mathfrak{H}}_{\mathbf{R}}$  the quotients  $\mathfrak{H}_{\mathbf{Z}}/G$  and  $\mathfrak{H}_{\mathbf{R}}/G$   
 21 respectively. As the flow  $\phi$  and the action of  $G$  commute,  $\phi$  induces a flow  $\bar{\phi} = (\bar{\phi}_t)_{t \in \mathbf{R}}$  on the  
 22 space  $\bar{\mathfrak{H}}_{\mathbf{R}} = \mathfrak{H}_{\mathbf{R}}/G$ . Moreover  $\bar{\phi}_1$  induces a homeomorphism of  $\bar{\mathfrak{H}}_{\mathbf{Z}}$  onto itself  $\bar{\mathfrak{H}}_{\mathbf{Z}}$ . One can check  
 23 that  $(\bar{\mathfrak{H}}_{\mathbf{R}}, \bar{\phi})$  is the suspension of the system  $(\bar{\mathfrak{H}}_{\mathbf{Z}}, \bar{\phi}_1)$  [[16](#), Proposition 4.8]. Let  $r: \bar{\mathfrak{H}}_{\mathbf{Z}} \rightarrow \bar{\mathfrak{H}}_0$  be the  
 24 map sending  $(h, \gamma)$  to  $\gamma(0)^{-1}h$  (recall that  $\gamma$  is a primitive gradient line, hence  $\gamma(0)$  is a vertex of  $\Gamma$   
 25 which corresponds to a unique element of  $G$ ). We observe that  $r$  induces a map  $\bar{r}: \bar{\mathfrak{H}}_{\mathbf{Z}} \rightarrow \bar{\mathfrak{H}}_0$  such  
 26 that  $\bar{r} \circ \bar{\phi}_1 = T \circ \bar{r}$ . Actually  $(\bar{\mathfrak{H}}_{\mathbf{Z}}, \bar{\phi}_1)$  is conjugated to the canonical two sided shift induced by  
 27  $(\bar{\mathfrak{H}}_0, T)$  [[16](#), Propositions 2.5 and 2.22].

### 28 4.3 Gromov's coding

29 In [[25](#), Theorem 8.4.C] Gromov explains that  $(\bar{\mathfrak{H}}_0, T)$  is conjugated to a subshift of finite type. We  
 30 recall here Gromov's coding as it is detailed by Coornaert and Papadopoulos in [[15](#)].

31 **The alphabet.** Fix a real number  $R_0 \geq 100\delta + 1$  and an integer  $L_0 \geq 2R_0 + 32\delta + 1$ . Given a  
 32 subset  $S$  of  $\Gamma$  and a number  $r \in \mathbf{R}_+$ , we denote by  $\mathcal{N}_r(S)$  the  $r$ -neighborhood of  $S$ , i.e. the set

$$33 \quad \mathcal{N}_r(S) = \{x \in \Gamma \mid d(x, S) \leq r\}.$$

34 Let  $h \in \mathfrak{H}_0$  be a horofunction and  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$  the minimal  $h$ -gradient line starting at 1. The set  
 35  $V(h)$  is the  $R_0$ -neighborhood of  $\gamma$  restricted to  $[0, L_0]$ . In addition we define the map

$$36 \quad b(h): V(h) \rightarrow \mathbf{R},$$

37 to be the restriction of  $h$  to  $V(h)$  (recall that our horofunctions vanish at 1). The alphabet  $\mathcal{B}$  is the  
 38 set of functions  $b(h): V(h) \rightarrow \mathbf{R}$  where  $h$  runs over  $\mathfrak{H}_0$ . It is a finite set [[15](#), Proposition 6.2].



**The coding.** Let  $\sigma: \mathcal{B}^{\mathbf{N}} \rightarrow \mathcal{B}^{\mathbf{N}}$  the shift map, i.e. the map sending the sequence  $(b_n)_{n \in \mathbf{N}}$  to  $(b_{n+1})_{n \in \mathbf{N}}$ . We define a map  $j: \mathfrak{H}_0 \rightarrow \mathcal{B}^{\mathbf{N}}$  by sending a horofunction  $h \in \mathfrak{H}_0$  to the sequence  $(b_n)$  defined by

$$b_n = b(T^n h), \quad \forall n \in \mathbf{N}.$$

Let  $\Sigma$  be the image of  $j$ . One observes that  $j \circ T = \sigma \circ j$  [15, Lemma 6.3]. Moreover  $j: \mathfrak{H}_0 \rightarrow \mathcal{B}^{\mathbf{N}}$  induces a homeomorphism from  $\mathfrak{H}_0$  onto  $\Sigma$ , which is a subshift of finite type of  $\mathcal{B}^{\mathbf{N}}$  [15, Theorem 7.18].

**Remark 4.8.** From now on we implicitly identify  $\mathfrak{H}_0$  with its image  $\Sigma$ . In particular, we say that  $h_1, h_2 \in \mathfrak{H}_0$  belong to the same cylinder of length  $n$ , if  $j(h_1)$  and  $j(h_2)$  coincide on the first  $n$  letters. We endow  $\mathfrak{H}_0$  with the canonical distance on  $\mathcal{B}^{\mathbf{N}}$ : for every  $h_1, h_2 \in \mathfrak{H}_0$ , we let  $d(h_1, h_2) = e^{-n}$  where  $n$  is the largest integer such that  $h_1$  and  $h_2$  belong to the same cylinder of length  $n$ .

Let  $h \in \mathfrak{H}_0$ . By construction  $b(h)$  completely determines the restriction of  $h$  to the ball of radius  $R_0$  centered at 1. Hence  $\theta(h)$  only depends on  $b(h)$ , i.e. the first letter of  $j(h)$ . Consequently, if  $h_1, h_2 \in \mathfrak{H}_0$  belong to the same cylinder of length  $n$ , then  $\theta_n(h_1) = \theta_n(h_2)$ , or said differently the minimal  $h_1$ - and  $h_2$ -gradient line starting at 1 coincide on  $[0, n]$ .

**Choice of an irreducible component.** Unlike the geodesic flow on a negatively curved compact surface the dynamical system  $(\mathfrak{H}_0, T)$  is a priori not topologically mixing and even not topologically transitive. This can be a major issue to study its properties. The difficulty comes from the fact that two points in  $\Gamma \cup \partial\Gamma$  may be joined by multiple geodesics. This pathology can be illustrated by the following simple example.

**Example 4.9.** Let  $G = \mathbf{F}_2 \times B$  be the direct product of the free group generated by  $\{a_1, a_2\}$  and a non-trivial finite group  $B$ . We choose for the generating set  $A = \{a_1, a_1^{-1}, a_2, a_2^{-1}\} \cup B$  and write  $\Gamma$  for the corresponding Cayley graph. One can check easily that  $\mathfrak{H}_0(\Gamma)$ , contains one copy of  $\partial\mathbf{F}_2$  (the usual Gromov boundary of  $\mathbf{F}_2$ ) for each element  $b \in B$ . Said differently there is an embedding of  $\partial\mathbf{F}_2 \times B$  into  $\mathfrak{H}_0(\Gamma)$ . This subset is invariant under the action of  $G$ . More precisely, for every  $(h, b) \in \partial\mathbf{F}_2 \times B$ , for every  $g = (f, u)$  in  $G$ , we have

$$g \cdot (h, u) = (f \cdot h, ub).$$

Assume now that the order on  $A$  is such that the letters  $a_1, a_1^{-1}, a_2, a_2^{-1}$  are smaller than the one of  $B$ . Then for every  $b \in B$ , the ‘‘layer’’  $\partial\mathbf{F}_2 \times \{b\}$  is invariant under  $T$ . On the contrary if every letter of  $B$  is smaller than  $a_1, a_1^{-1}, a_2, a_2^{-1}$ , then  $T$  maps  $\partial\mathbf{F}_2 \times B$  onto  $\partial\mathbf{F}_2 \times \{1\}$ .

Nevertheless for our purpose, one does not need to work with the whole system  $(\mathfrak{H}_0, T)$ . It is sufficient to restrict our attention to an irreducible component of the system, as long as it visits almost all the group  $G$ . This is formalized by the visibility property (see Definition 2.1). The goal of this section is to prove that such an irreducible component exists (see Proposition 4.10). Our main tool is the space of the geodesic flow introduced by Gromov in [25, Section 8.3]. From now on,  $\Gamma$  is the Cayley graph of any hyperbolic group, as in the previous section.

We have seen that  $(\mathfrak{H}_0, T)$  is conjugated to a subshift of finite type. We write  $\mathfrak{I}_1, \dots, \mathfrak{I}_m$  for the irreducible components of  $(\mathfrak{H}_0, T)$  (see Section 2.1). Recall that for every  $h \in \mathfrak{H}_0$ , there exists  $i \in \llbracket 1, m \rrbracket$  and  $n_0 \in \mathbf{N}$  such that for every  $n \geq n_0$ , the horofunction  $T^n h$  belongs to  $\mathfrak{I}_i$  (which we call the asymptotic irreducible component of  $h$ ). We can now state the main result of this section:

**Proposition 4.10.** *There exists an irreducible component  $\mathfrak{I}_i$  of  $\mathfrak{H}_0$  such that the extension of  $(\mathfrak{I}_i, T)$  by  $\theta$  has the visibility property.*

*Proof.* According to [Corollary 4.3](#) there exists  $\hat{\gamma}$  in  $\widehat{\mathcal{G}}\Gamma$  such that the positive orbit defined by

$$\{\psi_s(\hat{\gamma}) \mid s \in \mathbf{R}_+\}$$

has a dense image in  $\widehat{\mathcal{G}}\Gamma/G$ . Recall that  $(\mathfrak{H}_{\mathbf{R}}, \phi)$  is the horoflow of  $\Gamma$  introduced in [Section 4.2](#). The map  $q: \mathfrak{H}_{\mathbf{R}} \rightarrow \widehat{\mathcal{G}}\Gamma$  being onto ([Proposition 4.7](#)), we can fix a pre-image  $(h, \gamma) \in \mathfrak{H}_{\mathbf{R}}$  of  $\hat{\gamma}$  by  $q$ . Without loss of generality we can assume that  $(h, \gamma)$  actually belongs to the space of the discrete horoflow  $\mathfrak{H}_{\mathbf{Z}}$ . We denote by  $h_0$  the image of  $(h, \gamma)$  by the map  $r: \mathfrak{H}_{\mathbf{Z}} \rightarrow \mathfrak{H}_0$ , i.e.  $h_0 = \gamma(0)^{-1}h$ . We choose for  $\mathfrak{I}_i$  the asymptotic irreducible component of  $h_0$ . In particular there exists  $K \in \mathbf{N}$ , such that for every integer  $k \geq K$ , we have  $T^k(h_0) \in \mathfrak{I}_i$ .

We now study the properties of the map  $\theta: \mathfrak{H}_0 \rightarrow G$  restricted to  $\mathfrak{I}_i$ . It is an exercise of hyperbolic geometry to prove that there exists  $R_0 \in \mathbf{R}_+$  with the following property: given any two points  $y, y' \in \Gamma$ , there exists a bi-infinite geodesic  $\nu \in \mathcal{G}\Gamma$  such that  $d(y, \nu) \leq R_0$  and  $d(y', \nu) \leq R_0$ . Recall that the map  $\mathcal{G}\Gamma \rightarrow \Gamma$  sending  $\nu$  to  $\nu(0)$  as well as the projection  $p: \mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$  are quasi-isometries. Hence there exists  $\kappa \geq 1$  and  $\varepsilon \geq 0$  such that for every  $\nu, \nu' \in \mathcal{G}\Gamma$  we have

$$d(\nu(0), \nu'(0)) \leq \kappa d(p(\nu), p(\nu')) + \varepsilon. \quad (9)$$

We define the finite set  $U$  by

$$U = \{u \in G \mid d(1, u) \leq \kappa + \varepsilon + R_0 + 1 + 50\delta\}.$$

We are going to prove that for every  $g \in G$ , there exists a horofunction  $h \in \mathfrak{I}_i$ , an integer  $n \in \mathbf{N}$  and two elements  $u_1, u_2 \in U$  such that  $g = u_1 \theta_n(h) u_2$ .

Let  $g \in G$ . There exists a geodesic  $\nu \in \mathcal{G}\Gamma$  such that  $d(1, \nu) \leq R_0$  and  $d(g, \nu) \leq R_0$ . Up to changing the parametrization of  $\nu$ , we can assume that  $d(1, \nu(0)) \leq R_0$  and  $d(g, \nu(m)) \leq R_0 + 1$  for some integer  $m \in \mathbf{N}$ . We denote by  $\hat{\nu}$  the image of  $\nu$  in  $\widehat{\mathcal{G}}\Gamma$ . According to our choice of  $\hat{\gamma}$ , there exist a sequence  $(g_n)$  of elements of  $G$  and a sequence  $(s_n)$  of times diverging to infinity, such that  $(g_n \psi_{s_n}(\hat{\gamma}))$  converges to  $\hat{\nu}$ .

The projection  $p: \mathcal{G}\Gamma \rightarrow \widehat{\mathcal{G}}\Gamma$  maps homeomorphically  $\phi$ -orbits onto  $\psi$ -orbits. Hence there exists a sequence  $(t_n)$  of times diverging to infinity such that for every  $n \in \mathbf{N}$ , the map  $p$  sends  $g_n \phi_{t_n}(\gamma)$  to  $g_n \psi_{s_n}(\hat{\gamma})$ . Combining (9) with the fact that  $(g_n \psi_{s_n}(\hat{\gamma}))$  converges to  $\hat{\nu}$  we get that for sufficiently large integer  $n$

$$d(g_n \gamma(t_n), \nu(0)) \leq \kappa + \varepsilon.$$

The convergence taking place in  $\widehat{\mathcal{G}}\Gamma$  also tells us that  $(g_n \gamma(\infty))$  and  $(g_n \gamma(-\infty))$  converge to  $\nu(\infty)$  and  $\nu(-\infty)$  respectively. Recall that the metric on  $\Gamma$  is  $8\delta$ -quasi-convex [[13](#), Chapitre 10, Corollaire 5.3] Combined with the previous inequality we get that for sufficiently large integer  $n$

$$d(g_n \gamma(t_n + m), \nu(m)) \leq \kappa + \varepsilon + 50\delta.$$

For every  $n \in \mathbf{N}$  we denote by  $k_n$  the integer the closest to  $t_n$  so that  $|t_n - k_n| \leq 1$ . Recall that  $\nu$  has been chosen to pass close by 1 and  $g$ . Combining all these facts together we finally get that there exists  $n_1 \in \mathbf{N}$  such that for all integer  $n \geq n_1$  we have

$$d(1, g_n \gamma(k_n)) \leq \kappa + \varepsilon + R_0 + 1 \quad \text{and} \quad d(g, g_n \gamma(k_n + m)) \leq \kappa + \varepsilon + R_0 + 1 + 50\delta.$$

1 Let  $n \geq n_1$ . Recall that  $\gamma$  is a *primitive*  $h$ -gradient line, while  $k_n$  and  $m$  are integers. Hence  $g_n\gamma(k_n)$   
 2 and  $g_n\gamma(k_n + m)$  are elements of  $G$ . We let  $u_n = g_n\gamma(k_n)$  and  $u'_n = \gamma(k_n + m)^{-1}g_n^{-1}g$ . The last  
 3 inequalities tell us that  $u_n$  and  $u'_n$  belong to  $U$ . In addition we claim that

$$4 \quad g = u_n \theta_m (T^{k_n}(h_0)) u'_n.$$

5 As we explained before, the map  $r: \mathfrak{H}_{\mathbf{Z}} \rightarrow \mathfrak{H}_0$  induces a map  $\bar{r}: \bar{\mathfrak{H}}_{\mathbf{Z}} \rightarrow \mathfrak{H}_0$  such that  $\bar{r} \circ \bar{\phi}_1 = T \circ \bar{r}$ .  
 6 It follows that the ray  $\rho_n: \mathbf{R}_+ \rightarrow \Gamma$  defined by  $\rho_n(t) = \gamma(k_n)^{-1}\gamma(k_n + t)$  is the (unique) minimal  
 7  $T^{k_n}(h_0)$ -gradient ray starting at 1. Consequently

$$8 \quad \theta_m (T^{k_n}(h_0)) = \rho_n(m) = \gamma(k_n)^{-1}\gamma(k_n + m) = u_n^{-1}g u_n'^{-1},$$

9 which completes the proof of our claim. This decomposition of  $g$  holds for any integer  $n \geq n_1$ .  
 10 However  $k_n$  is diverging to infinity. Hence if  $n$  is sufficiently large  $k_n \geq K$ , thus  $T^{k_n}(h_0)$  belongs  
 11 to the irreducible component  $\mathfrak{J}_i$ . Thus  $g$  can be decomposed as announced in the proposition.  $\square$

12 The next lemmas formalize the fact that the irreducible component “visits almost” all the group  
 13  $G$ . The statements are quite technical. Nevertheless the idea one has to keep in mind is the  
 14 following. Given a horofunction  $h \in \mathfrak{J}$ , one can follow its trajectory in  $G$  by looking at the  
 15 corresponding minimal  $h$ -gradient line  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$ . More precisely, given  $n \in \mathbf{N}$ , the point  $\gamma(n)$   
 16 can be thought at the position in  $G$  at time  $n$  of the orbit of  $h$ . The next lemmas say that at time  $n$   
 17 the positions reached by the flow almost embed in (respectively almost cover) the sphere  $S(n) \subset G$ .  
 18 These computations will be needed to estimate the spectral radii of various transfer operators (see  
 19 Lemmas 5.6, 5.7 and 5.9). Before stating the lemmas we introduce a few notations. Let  $n \in \mathbf{N}$ .  
 20 For every  $h_0 \in \mathfrak{J}$ , the set  $S_{\mathfrak{J}}(h_0, n)$  is

$$21 \quad S_{\mathfrak{J}}(h_0, n) = \{h \in \mathfrak{J} \mid T^n h = h_0\} = T^{-n}(h_0) \cap \mathfrak{J}.$$

22 In addition we define a map

$$23 \quad \begin{aligned} p_n: \mathfrak{H}_0 \times G \times G &\rightarrow G \\ (h, u_1, u_2) &\rightarrow u_1 \theta_n(h) u_2 \end{aligned}$$

24 **Lemma 4.11.** *Let  $h_0 \in \mathfrak{J}$ . For every  $n \in \mathbf{N}$ , the map  $\mathfrak{H}_0 \rightarrow G$  sending  $h$  to  $\theta_n(h)$  induces an*  
 25 *embedding from  $S_{\mathfrak{J}}(h_0, n)$  into  $S(n)$ .*

26 *Proof.* Let  $h \in \mathfrak{J}$  such that  $T^n h = h_0$ . Let  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$  be the minimal  $h$ -gradient line starting at  
 27 1. As we observed before  $\theta_n(h) = \gamma(n)$ . In particular  $\theta_n(h)$  belongs to  $S(n)$ . On the other hand,  
 28 it follows from the definition of the map  $T$  that  $h = \theta_n(h)T^n h = \theta_n(h)h_0$ . Hence  $h$  is completely  
 29 determined by  $\theta_n(h)$  which completes the proof of the lemma.  $\square$

30 **Lemma 4.12.** *There exist  $R, N \in \mathbf{N}$  with the following property. For every  $h_0 \in \mathfrak{J}$ , for every*  
 31  *$n \in \mathbf{N}$ , the sphere  $S(n) \subset G$  is contained in the image of the map*

$$32 \quad \bigsqcup_{k=n}^{n+N} \left( S_{\mathfrak{J}}(h_0, k) \times B(R) \times B(R) \right) \rightarrow G$$

33 *induced by the sequence  $(p_k)$ .*

*Proof.* We start by defining the constants  $R$  and  $N$ . Recall that the extension of  $(\mathcal{J}, T)$  by  $\theta: \mathcal{J} \rightarrow G$  has the visibility property, i.e. there is a finite set  $U$  such that for every  $g \in G$ , there exist a horofunction  $h \in \mathcal{J}$ , an integer  $n \in \mathbf{N}$ , and two elements  $u_1, u_2 \in U$  satisfying  $g = u_1 \theta_n(h) u_2$  ([Proposition 4.10](#)). We denote by  $L$  the maximal length (in  $\Gamma$ ) of an element of  $U$ . As the system  $(\mathcal{J}, T)$  is an irreducible subshift of finite type, there exists  $K$  with the following property: for every  $h_1, h_2 \in \mathcal{J}$ , for every  $n \in \mathbf{N}$ , there exists  $h'_1 \in \mathcal{J}$  and  $k \in \llbracket 0, K \rrbracket$  such that  $T^{n+k} h'_1 = h_2$  and  $h_1$  and  $h'_1$  belong to the same cylinder of length  $n$ . Finally we let

$$R = 5L + K \quad \text{and} \quad N = 2L + K.$$

We now fix a horofunction  $h_0 \in \mathcal{J}$  and an integer  $n \in \mathbf{N}$ . Let  $g \in S(n)$ . According to the visibility property there exist a horofunction  $h \in \mathcal{J}$ , an integer  $m \in \mathbf{N}$  and two elements  $u_1, u_2 \in U$  such that  $g = u_1 \theta_m(h) u_2$ . Recall that the length (in  $\Gamma$ ) of  $\theta_m(h)$  is  $m$ , hence  $|n - m| \leq 2L$ . As  $(\mathcal{J}, T)$  is irreducible, there exists  $h' \in \mathcal{J}$  and  $k \in \llbracket 0, K \rrbracket$  such that  $h$  and  $h'$  belongs to the same cylinder of length  $n + 2L$  and

$$T^{n+2L+k} h' = h_0.$$

Since  $m \leq n + 2L$ , we have  $\theta_m(h') = \theta_m(h)$ . It follows that

$$\theta_{n+2L+k}(h') = \theta_m(h') \theta_{n+2L-m+k}(T^m h') = \theta_m(h) \theta_{n+2L-m+k}(T^m h')$$

Consequently

$$g = u_1 \theta_{n+2L+k}(h') u'_2,$$

where

$$u'_2 = \left( \theta_{n+2L-m+k}(T^m h') \right)^{-1} u_2.$$

Observe that  $u_1$  and  $u_2$  belong to  $B(L) \subset B(R)$ . As we noticed  $n \leq m + 2L$ , thus  $n - m + 2L + k \leq 4L + K$ . Consequently  $u'_2$  belongs to  $B(R)$ . In other words  $p_{n+2L+k}$  maps the element  $(h', u_1, u'_2)$  of  $S_{\mathcal{J}}(h_0, n + 2L + k) \times B(R) \times B(R)$  to  $g$ .  $\square$

## 5 Potential and transfer operator

The goal of this section is to prove the following statements.

**Theorem 5.1.** *Let  $G$  be a group acting properly co-compactly by isometries on a hyperbolic space  $X$ . We assume that one of the following holds. Either*

- (i)  $X$  is the Cayley graph of  $G$  with respect to a finite generating set, or
- (ii)  $X$  is a CAT(-1) space.

Let  $H$  be a subgroup of  $G$ . We denote by  $\omega_G$  and  $\omega_H$  the exponential growth rates of  $G$  and  $H$  acting on  $X$ . The subgroup  $H$  is co-amenable in  $G$  if and only if  $\omega_H = \omega_G$ . In particular if  $H$  is a normal subgroup of  $G$ , the quotient  $G/H$  is amenable if and only if  $\omega_H = \omega_G$ .

**Theorem 5.2.** *Let  $G$  be a group with Kazhdan's property (T) acting properly co-compactly by isometries on a hyperbolic space  $X$ . We assume that one of the following holds. Either*

- (i)  $X$  is the Cayley graph of  $G$  with respect to a finite generating set, or

1 (ii)  $X$  is a CAT(-1) space.

2 There exists  $\varepsilon > 0$  with the following property. Let  $H$  be a subgroup of  $G$ . We denote by  $\omega_G$  and  
 3  $\omega_H$  the exponential growth rates of  $G$  and  $H$  acting on  $X$ . If  $\omega_H > \omega_G - \varepsilon$ , then  $H$  is a finite index  
 4 subgroup of  $G$ .

## 5 5.1 The data

6 Let  $G$  be a group acting properly co-compactly by isometries on a hyperbolic space  $X$ . As in the  
 7 statement of Theorems 5.1 and 5.2 we consider two cases.

8 **Case 1.** The space  $X$  is the Cayley graph of  $G$  with respect to a finite generating set  $A$ . In this  
 9 situation we denote by  $\Gamma$  a copy of  $X$ .

10 **Case 2.** The space  $X$  is CAT(-1). In this situation we fix an arbitrary finite generating set  $A$  of  
 11  $G$  and denote by  $\Gamma$  the Cayley graph of  $G$  with respect to  $A$ .

12 In both cases we may assume without loss of generality that  $A$  is symmetric. As we work with  
 13 two distinct metric spaces, namely the Cayley graph  $\Gamma$  and the space  $X$ , we use this section to  
 14 emphasize which objects are related to one or the other space.

15 **Data related to  $X$ .** The space  $X$  is the one that will carry the geometric information. We denote  
 16 by  $\delta$  its hyperbolicity constant. We fix a base point  $x_0 \in X$ . This allows us to identify  $C_*(X)$  with  
 17 the set of continuous maps vanishing at  $x_0$ . We denote by  $\pi_X: \mathfrak{H}(X) \rightarrow \partial X$  the projection studied  
 18 in Proposition 3.4. We denote by  $\omega_G$  the exponential growth rate of  $G$  acting on  $X$ .

19 **Data related to  $\Gamma$ .** The role of  $\Gamma$  is to provide a support for coding the geodesic flow. The space  
 20  $\mathfrak{H}_0 \subset \mathfrak{H}(\Gamma)$  refers to the integral horofunctions on the Cayley graph  $\Gamma$ . We denote by  $\pi_\Gamma: \mathfrak{H}_0 \rightarrow \partial\Gamma$   
 21 the projection coming from Proposition 4.5. The maps  $\theta: \mathfrak{H}_0 \rightarrow G$  and  $T: \mathfrak{H}_0 \rightarrow \mathfrak{H}_0$  are the ones  
 22 defined at the beginning of Section 4.2. For simplicity we denote by  $\mathfrak{J}$  the irreducible component  
 23 of  $(\mathfrak{H}_0, T)$  with the visibility property given by Proposition 4.10. Recall that for every  $n \in \mathbf{N}$ , the  
 24 sets  $S(n)$  and  $B(n)$  are respectively the sphere and the ball of radius  $n$ , measured in  $\Gamma$ .

25 **Comparing  $\Gamma$  and  $X$ .** Since  $G$  acts properly co-compactly on  $X$ , the orbit maps  $G \rightarrow X$  sending  
 26  $g$  to  $gx_0$  leads a  $(\kappa, \ell)$ -quasi-isometric embedding  $f: \Gamma \rightarrow X$ . This map induces a homeomorphism  
 27  $\partial\Gamma \rightarrow \partial X$  between the respective Gromov boundary of  $\Gamma$  and  $X$ . For simplicity we implicitly  
 28 identify  $\partial\Gamma$  and  $\partial X$ .

## 29 5.2 Transfer operator for the irreducible component

30 **Comparing horofunctions.** The first task is to define a potential  $F: \mathfrak{H}_0 \rightarrow \mathbf{R}_+^*$ . This potential  
 31 defined on the dynamical system  $(\mathfrak{H}_0, T)$  should reflect to geometry of  $X$ . In the first case – when  
 32  $X$  is actually the Cayley graph  $\Gamma$  of  $X$  – the geometry coincides with the dynamics, and we can  
 33 simply take for  $F$  the constant function equal to  $e^{-\omega_G}$ . In the second case – when  $X$  is an arbitrary  
 34 CAT(-1) space – the situation is more subtle. Indeed  $\mathfrak{H}(X)$  does not necessarily coincide with  $\mathfrak{H}(\Gamma)$   
 35 nor  $\mathfrak{H}_0$ . As the space  $X$  is CAT(-1), the set  $\mathfrak{H}(X)$  coincides with the usual visual boundary of  $X$ .  
 36 More precisely the map  $\pi_X: \mathfrak{H}(X) \rightarrow \partial X$  is a homeomorphism. On the other hand the projection  
 37  $\pi_\Gamma: \mathfrak{H}_0 \rightarrow \partial\Gamma$  is not always injective. Nevertheless we are going to build a map comparing  $\mathfrak{H}_0$  – the

horofunctions used for coding – to  $\mathfrak{H}(X)$  – the horofunctions capturing the geometry of  $X$ . This is the purpose of the next proposition. Actually we develop a framework that covers both cases simultaneously.

Recall that we identify  $(\mathfrak{H}_0, T)$  with an appropriate subshift of finite type of  $(\mathcal{B}^{\mathbf{N}}, \sigma)$ . This identification induces a distance on  $\mathfrak{H}_0$  (Remark 4.8) as defined at the beginning of Section 2.1. Namely the distance between two horofunctions  $h, h' \in \mathfrak{H}_0$  is  $d(h, h') = e^{-n}$ , where  $n$  is the largest integer such that the respective images of  $h$  and  $h'$  in  $\mathcal{B}^{\mathbf{N}}$  have the same first  $n$  letters. We denote by  $\mathcal{C}(\mathfrak{H}_0, \mathbf{C})$  for the space of continuous maps from  $\mathfrak{H}_0$  to  $\mathbf{C}$  while  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{C})$  stands for the space of functions with bounded  $\alpha$ -Hölder variations (see Section A.1).

**Proposition 5.3.** *There exists a  $G$ -equivariant comparison map  $\mathfrak{H}_0 \rightarrow \mathfrak{H}(X)$  which we denote  $h \mapsto h_X$  with the following properties.*

(i) *The following diagram commutes*

$$\begin{array}{ccc} \mathfrak{H}_0 & \longrightarrow & \mathfrak{H}(X) \\ \pi_\Gamma \downarrow & & \downarrow \pi_X \\ \partial\Gamma & \longrightarrow & \partial X \end{array}$$

(ii) *There exists  $\alpha \in \mathbf{R}_+^*$  such that the evaluation map defined by*

$$\begin{array}{ccc} \varphi: \mathfrak{H}_0 & \rightarrow & \mathbf{R} \\ h & \rightarrow & h_X \left( \theta(h)x_0 \right) \end{array}$$

*belongs to  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{R})$ .*

*Proof.* We distinguish two cases depending whether  $X$  is a Cayley graph or a CAT(−1) space.

**Case 1.** Assume first that  $X$  is the Cayley graph of  $G$  with respect to the generating set  $A$ . In this situation we defined  $\Gamma$  to be exactly  $X$ . In particular  $\mathfrak{H}_0$  can be seen as a subset of  $\mathfrak{H}(X)$ . We simply define the comparison map  $\mathfrak{H}_0 \rightarrow \mathfrak{H}(X)$  as the corresponding embedding. Point (i) becomes obvious. It follows from the very definition of  $\theta$  that  $\varphi$  is constant equal to  $-1$ . Thus it belongs to  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{R})$  for every  $\alpha \in \mathbf{R}_+^*$ .

**Case 2.** Assume now that  $X$  is a CAT(−1) space. In this situation  $\Gamma$  is the Cayley graph of  $G$  with respect to an arbitrary finite generating set. Given  $h \in \mathfrak{H}_0$  we define  $h_X$  to be the Busemann function at  $\xi = \pi_\Gamma(h)$  vanishing at  $x_0$ . By construction the diagram of Point (i) commutes.

Let  $D = D(\kappa, \ell, \delta)$  be the parameter given by the Morse lemma (Proposition 2.3). Let  $h, h' \in \mathfrak{H}_0$  such that  $d(h, h') < 1$ . We denote by  $n \in \mathbf{N}^*$  the largest integer such that  $h$  and  $h'$  belong to the same cylinder of length  $n$ , so that  $d(h, h') = e^{-n}$ . We write  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$  for the minimal  $h$ -gradient ray starting at 1 and let  $\xi = \gamma(\infty)$ . Let  $c: \mathbf{R}_+ \rightarrow X$  be the ray starting at  $x_0$  such that  $c(\infty) = \xi$ . Recall that the map  $f: \Gamma \rightarrow X$  induced by the orbit map is a  $(\kappa, \ell)$ -quasi-isometry. In particular  $f \circ \gamma$  is a  $(\kappa, \ell)$ -quasi-geodesic between  $x_0$  and  $\xi$ . It follows from the stability of quasi-geodesics (Proposition 2.3) that the Hausdorff distance between  $f \circ \gamma$  and  $c$  is bounded above by  $D$ . In a similar way we associate to  $h'$  a gradient ray  $\gamma': \mathbf{R}_+ \rightarrow \Gamma$  as well as a geodesic ray  $c': \mathbf{R}_+ \rightarrow X$ .

1 Since  $h$  and  $h'$  belong to the same cylinder of length  $n$ , the paths  $\gamma$  and  $\gamma'$  coincide on  $[0, n]$   
 2 (see [Remark 4.8](#)). In particular, for every  $k \in \llbracket 0, n \rrbracket$ , the point  $y_k = f \circ \gamma(k) = f \circ \gamma'(k)$  lies in the  
 3  $D$  neighborhood of both  $c$  and  $c'$ . As  $f$  is a  $(\kappa, \ell)$ -quasi-isometric embedding, we also observe that

$$4 \quad \kappa^{-1}k - \ell \leq d(x_0, y_k) \leq \kappa k + \ell.$$

5 It follows that there exists  $t \in \mathbf{R}_+$ , such that  $d(c(t), c'(t)) \leq 2D$  and  $t \geq \kappa^{-1}n - \ell - D$ . A standard  
 6 exercise of CAT(-1) geometry shows that

$$7 \quad d(c(t/2), c'(t/2)) \leq C_1 e^{-\frac{1}{2}t},$$

8 where  $C_1$  is a parameter that only depends on  $D$ . Recall that  $d(x_0, y_1) \leq \kappa + \ell$ . Another exercise  
 9 of CAT(-1) geometry shows that

$$10 \quad \left| h_X(y_1) - \left[ d(c(t/2), y_1) - t/2 \right] \right| \leq C_2 e^{-\frac{1}{2}t} \quad \text{and} \quad \left| h'_X(y_1) - \left[ d(c'(t/2), y_1) - t/2 \right] \right| \leq C_2 e^{-\frac{1}{2}t},$$

11 where  $C_2$  only depends on  $\kappa$  and  $\ell$ . Consequently

$$12 \quad |h_X(y_1) - h'_X(y_1)| \leq C_3 e^{-\frac{1}{2}t} \leq C_4 e^{-\frac{1}{2}\kappa^{-1}n} \leq C_4 d(h, h')^{\frac{1}{2}\kappa^{-1}}.$$

13 where  $C_3$  and  $C_4$  only depends on  $D$ ,  $\kappa$  and  $\ell$ . Nevertheless  $y_1$  is the point  $\theta(h)x_0 = \theta(h')x_0$ . Hence  
 14 the previous inequality exactly says that

$$15 \quad |\varphi(h) - \varphi(h')| \leq C_4 d(h, h')^{\frac{1}{2}\kappa^{-1}}.$$

16 This inequality holds for every  $h, h'$  such that  $d(h, h') < 1$ . Consequently  $\varphi$  belongs to the space  
 17  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{R})$  where  $\alpha = \kappa^{-1}/2$ , which proves Point (ii).  $\square$

18 **The potential  $F$ .** From now on, we fix the comparison map  $\mathfrak{H}_0 \rightarrow \mathfrak{H}(X)$ ,  $h \mapsto h_X$  given by  
 19 [Proposition 5.3](#). We keep the notations introduced in this statement. In particular the evaluation  
 20 map  $\varphi: \mathfrak{H}_0 \rightarrow \mathbf{R}$  sending  $h$  to  $h_X(\theta(h)x_0)$  belongs to  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{R})$ . The potential  $F: \mathfrak{H}_0 \rightarrow \mathbf{R}_+^*$  is  
 21 the map defined by

$$22 \quad F(h) = \exp\left(\omega_G \varphi(h)\right) = \exp\left(\omega_G h_X\left(\theta(h)x_0\right)\right), \quad \forall h \in \mathfrak{H}_0. \quad (10)$$

23 It directly follows from the previous proposition that  $\ln F$  belongs to  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{R})$

24 **Remark.** If  $X$  is simply the Cayley graph  $\Gamma$  of  $X$ , we previously observed that the evaluation  
 25 map  $\varphi$  is constant equal to  $-1$ . Hence the potential becomes  $F(h) = e^{-\omega_G}$ .

26 **Lemma 5.4.** *For every  $n \in \mathbf{N}$ , for every  $h \in \mathfrak{H}_0$ , we have*

$$27 \quad F_n(h) = \exp\left(\omega_G h_X\left(\theta_n(h)x_0\right)\right).$$

28 *Proof.* Let  $h \in \mathfrak{H}_0$ . It is sufficient to prove that for every  $n \in \mathbf{N}$ , we have

$$29 \quad \sum_{k=0}^{n-1} \varphi \circ T^k(h) = h_X\left(\theta_n(h)x_0\right).$$

1 The proof is by induction on  $n$ . By convention  $\theta_0(h) = 1$ . Since  $h_X$  vanishes at  $x_0$  the statement  
 2 obviously holds for  $n = 0$ . Assume now that the claim holds for some  $n \in \mathbf{N}$ . It follows from the  
 3 definition of  $T$  that  $T^n(h) = \theta_n(h)^{-1}h$ . As the comparison map  $\mathfrak{H}_0 \rightarrow \mathfrak{H}(X)$  is  $G$ -equivariant we  
 4 get

$$\begin{aligned} \varphi \circ T^n(h) &= [\theta_n(h)^{-1}h_X] \left( \theta(T^n(h))x_0 \right) = h_X \left( \theta_n(h)\theta(T^n(h))x_0 \right) - h_X \left( \theta_n(h)x_0 \right) \\ &= h_X \left( \theta_{n+1}(h)x_0 \right) - h_X \left( \theta_n(h)x_0 \right). \end{aligned}$$

5 The statement for  $n + 1$  now follows from the induction hypotheses.  $\square$

6 **Lemma 5.5.** *There exists a constant  $C \in \mathbf{R}_+^*$ , such that for every  $h \in \mathfrak{H}_0$ , for every  $n \in \mathbf{N}$ , we*  
 7 *have*

$$8 \quad \frac{1}{C} \leq \frac{F_n(h)}{\exp\left(-\omega_G d(\theta_n(h)x_0, x_0)\right)} \leq C.$$

9 *Proof.* According to Lemma 5.4, it suffices to show that there exists  $C' \in \mathbf{R}_+^*$  such that for every  
 10  $h \in \mathfrak{H}_0$ , for every  $n \in \mathbf{N}$  we have

$$11 \quad \left| h_X \left( \theta_n(h)x_0 \right) + d \left( \theta_n(h)x_0, x_0 \right) \right| \leq C'.$$

12 By the stability of quasi-geodesics, there exists  $D \in \mathbf{R}_+$  such that the Hausdorff distance between  
 13 two  $(\kappa, \ell)$ -quasi-geodesics of  $X$  joining the same endpoints (possibly in  $\partial X$ ) is at most  $D$ . Let  
 14  $h \in \mathfrak{H}_0$ . Let  $c: \mathbf{R}_+ \rightarrow X$  be geodesic ray between  $x_0$  and  $\xi = \pi_X(h_X)$ . We write  $b: X \rightarrow \mathbf{R}$  for the  
 15 corresponding Busemann function vanishing at  $x_0$ . Note that  $h_X$  and  $b$  are two horofunctions of  
 16  $X$  whose image by  $\pi_X: \mathfrak{H}(X) \rightarrow \partial X$  is  $\xi$ . It follows that  $\|h_X - b\|_\infty \leq 64\delta$  [15, Corollary 3.8]. Let  
 17  $\gamma: \mathbf{R}_+ \rightarrow \Gamma$  be the minimal  $h$ -gradient line starting at 1. Observe that  $f \circ \gamma$  is a  $(\kappa, \ell)$  quasi-geodesic  
 18 of  $X$ . Hence the Hausdorff distance between  $f \circ \gamma$  and  $c$  is at most  $D$ .

19 Let  $n \in \mathbf{N}$ . By construction the element  $\theta_n(h)$  lies on  $\gamma$ . Thus there exists  $t \in \mathbf{R}_+$ , such that  
 20  $d(\theta_n(h)x_0, c(t)) \leq D$ . It follows that

$$21 \quad |d(\theta_n(h)x_0, x_0) + b(c(t))| \leq D. \quad (11)$$

22 Recall that the Busemann function  $b$  is a 1-Lipschitz. Combined with the fact that  $\|h_X - b\|_\infty \leq 64\delta$   
 23 we get

$$24 \quad |h_X(\theta_n(h)x_0) - b(c(t))| \leq D + 64\delta.$$

25 Hence (11) becomes

$$26 \quad |h_X(\theta_n(h)x_0) + d(\theta_n(h)x_0, x_0)| \leq C',$$

27 where  $C' = 2D + 64\delta$ . Observes that  $C'$  neither depends on  $h$  of  $n$ , hence the proof is complete.  $\square$

28 **The transfer operator.** The transfer operator associated to the potential  $F$  is the operator  
 29  $\mathcal{L}: \mathcal{C}(\mathcal{J}, \mathbf{C}) \rightarrow \mathcal{C}(\mathcal{J}, \mathbf{C})$  defined by

$$30 \quad \mathcal{L}\Phi(h_0) = \sum_{Th=h_0} F(h)\Phi(h), \quad \forall \Phi \in \mathcal{C}(\mathcal{J}, \mathbf{C}), \quad \forall h_0 \in \mathcal{J}. \quad (12)$$



1 Note that the restriction map  $\mathcal{C}(\mathfrak{H}_0, \mathbf{C}) \rightarrow \mathcal{C}(\mathfrak{J}, \mathbf{C})$  induces a 1-Lipschitz map from  $H_\alpha^\infty(\mathfrak{H}_0, \mathbf{C}) \rightarrow$   
 2  $H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$ . Hence  $\ln F$  restricted to  $\mathfrak{J}$  belongs to  $H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$ . As we observed in the appendix  $\mathcal{L}$   
 3 induces a bounded operator of  $H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$ . Since the system  $(\mathfrak{J}, T)$  is irreducible, the spectral radii  
 4 of  $\mathcal{L}$  seen as an operator of  $\mathcal{C}(\mathfrak{J}, \mathbf{C})$  or  $H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$  are the same (Theorem A.6). We denote it by  $\rho$ .  
 5 According to (18) it can be computed as follows

$$6 \quad \rho = \limsup_{n \rightarrow \infty} \sqrt[n]{\|\mathcal{L}^n \mathbb{1}\|_\infty}. \quad (13)$$

7 **Computing  $\rho$ .** The goal of this section is to prove that  $\rho = 1$  (Proposition 5.8).

8 **Lemma 5.6.** *There exists  $A_1 \in \mathbf{R}_+^*$  such that for every  $n \in \mathbf{N}$ , we have*

$$9 \quad \|\mathcal{L}^n \mathbb{1}\|_\infty \leq A_1 \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)}.$$

10 *Proof.* We denote by  $C$  the constant given by Lemma 5.5. Let  $n \in \mathbf{N}$ . Let  $h_0 \in \mathfrak{J}$ . According to  
 11 Lemma 5.4 we have

$$12 \quad \mathcal{L}^n \mathbb{1}(h_0) = \sum_{T^n h = h_0} F_n(h) = \sum_{T^n h = h_0} \exp\left(\omega_G h_X\left(\theta_n(h)x_0\right)\right).$$

13 Applying Lemma 5.5 we get

$$14 \quad \mathcal{L}^n \mathbb{1}(h_0) \leq C \sum_{T^n h = h_0} \exp\left(-\omega_G d\left(\theta_n(h)x_0, x_0\right)\right). \quad (14)$$

15 By Lemma 4.11 the map  $\mathfrak{H}_0 \rightarrow G$  sending  $h$  to  $\theta_n(h)$  induces an embedding of  $\{h \in \mathfrak{J} \mid T^n h = h_0\}$   
 16 into  $S(n)$ . It follows that

$$17 \quad \mathcal{L}^n \mathbb{1}(h_0) \leq C \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)}.$$

18 This inequality holds for every  $h_0 \in \mathfrak{J}$ , thus

$$19 \quad \|\mathcal{L}^n \mathbb{1}\|_\infty \leq C \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)}. \quad \square$$

20 **Lemma 5.7.** *There exists  $A_2 \in \mathbf{R}_+^*$  such that for every  $n \in \mathbf{N}$ , we have*

$$21 \quad \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)} \leq A_2 \|\mathcal{L}^n \mathbb{1}\|_\infty.$$

22 *Proof.* We write  $C \in \mathbf{R}_+^*$ , and  $R, N \in \mathbf{N}$ , for the constants given by Lemma 5.5 and Lemma 4.12  
 23 respectively. Recall that the map  $f: \Gamma \rightarrow X$  induced by the orbit map is a  $(\kappa, \ell)$ -quasi-isometric  
 24 embedding. Let  $h_0 \in \mathfrak{J}$ . Let  $n \in \mathbf{N}$ . Let  $k \in \mathbf{N}$ . Applying Lemma 5.5 we observe that

$$\begin{aligned} \sum_{T^k h = h_0} \exp\left(-\omega_G d\left(\theta_k(h)x_0, x_0\right)\right) &\leq C \sum_{T^k h = h_0} \exp\left(\omega_G h_X\left(\theta_k(h)x_0\right)\right) \\ &\leq C \mathcal{L}^k \mathbb{1}(h_0). \end{aligned} \quad (15)$$

1 On the other hand if  $(u_1, u_2)$  belongs to  $B(R) \times B(R)$ , the triangle inequality tells us that

$$2 \quad |d(u_1\theta_k(h)u_2x_0, x_0) - d(\theta_k(h)x_0, x_0)| \leq d(u_1x_0, x_0) + d(u_2x_0, x_0).$$

3 However the map  $f: \Gamma \rightarrow X$  being a  $(\kappa, \ell)$ -quasi-isometric embedding, we get

$$4 \quad |d(u_1\theta_k(h)u_2x_0, x_0) - d(\theta_k(h)x_0, x_0)| \leq 2(\kappa R + \ell).$$

5 Summing (15) when  $(u_1, u_2)$  runs over  $B(R) \times B(R)$  and  $k$  over  $\llbracket n, n+N \rrbracket$  gives

$$\begin{aligned} & \sum_{k=n}^{n+N} \sum_{(u_1, u_2) \in B(R) \times B(R)} \sum_{T^k h = h_0} \exp\left(-\omega_G d(u_1\theta_k(h)u_2x_0, x_0)\right) \\ & \leq |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \sum_{k=n}^{n+N} \sum_{T^k h = h_0} \exp\left(-\omega_G d(\theta_k(h)x_0, x_0)\right) \\ & \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \sum_{k=n}^{n+N} \mathcal{L}^k \mathbb{1}(h_0). \end{aligned}$$

6 Lemma 4.12 provides a lower bound of the triple sum in the left hand side of the inequality, leading  
7 to

$$8 \quad \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)} \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \sum_{k=n}^{n+N} \mathcal{L}^k \mathbb{1}(h_0).$$

9 As the potential  $F$  and the function  $\mathbb{1}$  are positive we observe that for every  $k \geq n$ ,

$$10 \quad \mathcal{L}^k \mathbb{1}(h_0) \leq \|\mathcal{L}^k \mathbb{1}\|_\infty \leq \|\mathcal{L}^{k-n}\|_\infty \|\mathcal{L}^n \mathbb{1}\|_\infty.$$

11 Hence the previous inequality becomes

$$12 \quad \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)} \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \left( \sum_{k=0}^N \|\mathcal{L}^k\|_\infty \right) \|\mathcal{L}^n \mathbb{1}\|_\infty,$$

13 which is exactly the required inequality. □

14 **Proposition 5.8.** *The spectral radius of  $\mathcal{L}$  is  $\rho = 1$ .*

15 *Proof.* We form the series

$$16 \quad \Upsilon(s) = \sum_{n=0}^{\infty} e^{-sn} \|\mathcal{L}^n \mathbb{1}\|_\infty.$$

17 It follows from (13) that the critical exponent  $\Upsilon(s)$  is  $\ln \rho$ . Hence it suffices to prove this critical  
18 exponent is 0. Let  $s \in \mathbf{R}_+^*$ . According to Lemmas 5.6 and 5.7 there exists  $A_1, A_2$  such that for  
19 every  $n \in \mathbf{N}$

$$20 \quad A_1 \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)} \leq \|\mathcal{L}^n \mathbb{1}\|_\infty \leq A_2 \sum_{g \in S(n)} e^{-\omega_G d(gx_0, x_0)}.$$

1 Multiplying these inequalities by  $e^{-sn}$  and summing over  $n$  gives

$$2 \quad A_1 \sum_{n=0}^{\infty} \sum_{g \in S(n)} e^{-sn} e^{-\omega_G d(gx_0, x_0)} \leq \Upsilon(s) \leq A_2 \sum_{n=0}^{\infty} \sum_{g \in S(n)} e^{-sn} e^{-\omega_G d(gx_0, x_0)}.$$

3 The map  $f: \Gamma \rightarrow X$  being a  $(\kappa, \ell)$ -quasi-isometric embedding, for every  $n \in \mathbf{N}$ , for every  $g \in S(n)$   
4 we have

$$5 \quad \kappa^{-1} [d(gx_0, x_0) - \ell] \leq n \leq \kappa [d(gx_0, x_0) + \ell].$$

6 Consequently

$$7 \quad A_1 e^{-s\kappa\ell} \sum_{g \in G} e^{-(\omega_G + s\kappa)d(gx_0, x_0)} \leq \Upsilon(s) \leq A_2 e^{s\kappa^{-1}\ell} \sum_{g \in G} e^{-(\omega_G + s\kappa^{-1})d(gx_0, x_0)}.$$

8 This can be reformulated using the Poincaré series of  $G$  as

$$9 \quad A_1 e^{-s\kappa\ell} \mathcal{P}_G(\omega_G + s\kappa) \leq \Upsilon(s) \leq A_2 e^{s\kappa^{-1}\ell} \mathcal{P}_G(\omega_G + s\kappa^{-1}).$$

10 Recall that  $s \rightarrow \mathcal{P}_G(\omega_G + s)$  converges if  $s > 0$  and diverges if  $s < 0$ . It follows that the critical  
11 exponent of  $\Upsilon(s)$  equals 0.  $\square$

### 12 5.3 Twisted transfer operator associated to a subgroup

13 **Data associated to a subgroup.** Let  $H$  be a subgroup of  $G$ . We denote by  $Y$  the space of  
14 left cosets of  $H$  in  $G$ , i.e.  $Y = H \backslash G$ . We write  $y_0$  for the image of 1 in  $Y$ . In other words  $y_0$  is  
15 the coset  $H$ . We denote by  $\mathcal{H} = \ell^2(Y)$  the space of square summable functions from  $Y$  to  $\mathbf{C}$ . The  
16 group  $G$  acts on  $Y$  by right translations. It induces a unitary representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  defined

$$17 \quad [\lambda(g)\phi]y = \phi(y \cdot g), \quad \forall g \in G, \forall \phi \in \mathcal{H}.$$

18 We call  $\lambda$  the *regular representation of  $G$  relative to  $H$* . We denote by  $\omega_H$  the exponential  
19 growth rate of  $H$  acting on  $X$ .

20 **Twisted transfer operator.** We denote by  $\mathcal{C}(\mathfrak{J}, \mathcal{H})$  the set of continuous function from  $\mathfrak{J}$  to  
21  $\mathcal{H}$ . Similarly  $H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$  stands for the space of functions with bounded  $\alpha$ -Hölder variations (see  
22 [Section A.1](#)). As explained in this appendix, the representation  $\lambda$  leads to a twisted transfer operator  
23  $\mathcal{L}_\lambda: C(\mathfrak{J}, \mathcal{H}) \rightarrow C(\mathfrak{J}, \mathcal{H})$  defined by

$$24 \quad \mathcal{L}_\lambda \Phi(h_0) = \sum_{Th=h_0} F(h) \lambda(\theta(h))^{-1} \Phi(h), \quad \forall \Phi \in C(\mathfrak{J}, \mathcal{H}), \forall h_0 \in \mathfrak{J}.$$

25 This operator induces a bounded operator of  $H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$  ([Proposition A.8](#)). We write  $\rho_\lambda$  for the  
26 spectral radius of  $\mathcal{L}_\lambda$  seen as an operator of  $H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$ .

27 **Computing  $\rho_\lambda$ .** Our goal is to provide an estimate of  $\rho_\lambda$  in terms of  $\omega_G$  and  $\omega_H$ . Let us first  
28 remark that  $\rho_\lambda \leq \rho$  ([Corollary A.9](#)) that is in our setting  $\rho_\lambda \leq 1$  ([Proposition 5.8](#)). We now provide  
29 a lower bound for  $\rho_\lambda$ . The proof follows the same strategy as the one of [Proposition 5.8](#).

**Lemma 5.9.** *There exist  $B_2 \in \mathbf{R}_+^*$  and a function  $\Psi \in H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$  such that for every  $n \in \mathbf{N}$ , we have*

$$\sum_{g \in S(n) \cap H} e^{-\omega_G d(gx_0, x_0)} \leq B_2 \|\mathcal{L}_\lambda^n \Psi\|_\infty.$$

*Proof.* We write  $C \in \mathbf{R}_+^*$ , and  $R, N \in \mathbf{N}$ , for the constants given by [Lemma 5.5](#) and [Lemma 4.12](#) respectively. Recall that the map  $f: \Gamma \rightarrow X$  induced by the orbit map is a  $(\kappa, \ell)$ -quasi-isometric embedding. We denote by  $Z$  the following finite subset of  $Y$ .

$$Z = \{y_0 \cdot u \mid u \in B(R)\}.$$

The map  $\Psi: \mathfrak{J} \rightarrow \mathcal{H}$  is the constant function equal to the characteristic function  $\mathbb{1}_Z$  of  $Z$ . One observes easily that  $\Psi$  belongs to  $H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$ .

We now fix  $h_0 \in \mathfrak{J}$  and  $n \in \mathbf{N}$ . Let  $k \in \mathbf{N}$  and  $u_2 \in B(R)$ . Let us compute  $\mathcal{L}_\lambda^k \Psi(h_0)$  at the point  $y_0 \cdot u_2^{-1}$ . By definition we have

$$[\mathcal{L}_\lambda^k \Psi(h_0)](y_0 \cdot u_2^{-1}) = \sum_{T^k h = h_0} F_k(h) [\lambda(\theta_k(h)^{-1}) \Psi](y_0 \cdot u_2^{-1}) = \sum_{T^k h = h_0} F_k(h) \mathbb{1}_Z(y_0 \cdot u_2^{-1} \theta_k(h)^{-1}).$$

Recall that  $Z = y_0 \cdot B(R)$ . Hence the term

$$\mathbb{1}_Z(y_0 \cdot u_2^{-1} \theta_k(h)^{-1})$$

equals 1 if there exists  $u_1 \in B(R)$  such that  $u_1 \theta_k(h) u_2$  belongs to  $H$  and zero otherwise. Hence

$$\sum_{u_1 \in B(R)} \sum_{\substack{T^k h = h_0, \\ u_1 \theta_k(h) u_2 \in H}} F_k(h) \leq |B(R)| [\mathcal{L}_\lambda^k \Psi(h_0)](y_0 \cdot u_2^{-1}).$$

Recall that the potential  $F$  and the vector  $\mathbb{1}_Z$  are non-negative. It follows that

$$\sum_{u_1 \in B(R)} \sum_{\substack{T^k h = h_0, \\ u_1 \theta_k(h) u_2 \in H}} F_k(h) \leq |B(R)| \|\mathcal{L}_\lambda^k \Psi(h_0)\|_{\ell^2(Y)} \leq |B(R)| \|\mathcal{L}_\lambda^k \Psi\|_\infty.$$

Applying [Lemma 5.5](#) to the previous inequality, we observe (as in the proof of [Lemma 5.7](#)) that

$$\sum_{u_1 \in B(R)} \sum_{\substack{T^k h = h_0, \\ u_1 \theta_k(h) u_2 \in H}} \exp\left(-\omega_G d(\theta_k(h)x_0, x_0)\right) \leq C |B(R)| \|\mathcal{L}_\lambda^k \Psi\|_\infty. \quad (16)$$

On the other hand given  $(u_1, u_2) \in B(R) \times B(R)$ , the triangle inequality tells us that

$$|d(u_1 \theta_k(h) u_2 x_0, x_0) - d(\theta_k(h)x_0, x_0)| \leq d(u_1 x_0, x_0) + d(u_2 x_0, x_0).$$

The map  $f: \Gamma \rightarrow X$  being a  $(\kappa, \ell)$ -quasi-isometric embedding, we get

$$|d(u_1 \theta_k(h) u_2 x_0, x_0) - d(\theta_k(h)x_0, x_0)| \leq 2(\kappa R + \ell).$$

1 Summing (16) when  $u_2$  runs over  $B(R)$  and  $k$  over  $\llbracket n, n+N \rrbracket$  yields

$$\begin{aligned} & \sum_{k=n}^{n+N} \sum_{(u_1, u_2) \in B(R) \times B(R)} \sum_{\substack{T^k h = h_0, \\ u_1 \theta_k(h) u_2 \in H}} \exp\left(-\omega_G d(u_1 \theta_k(h) u_2 x_0, x_0)\right) \\ & \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \sum_{k=n}^{n+N} \|\mathcal{L}_\lambda^k \Psi\|_\infty. \end{aligned}$$

2 [Lemma 4.12](#) provides a lower bound of the triple sum in the left hand side of the inequality, leading  
3 to

$$4 \quad \sum_{g \in S(n) \cap H} e^{-\omega_G d(gx_0, x_0)} \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \sum_{k=n}^{n+N} \|\mathcal{L}_\lambda^k \Psi\|_\infty.$$

5 Observe that for every  $k \geq n$  we have,

$$6 \quad \|\mathcal{L}_\lambda^k \Psi\|_\infty \leq \|\mathcal{L}_\lambda^{k-n}\|_\infty \|\mathcal{L}_\lambda^n \Psi\|_\infty.$$

7 Hence the previous inequality becomes

$$8 \quad \sum_{g \in S(n) \cap H} e^{-\omega_G d(gx_0, x_0)} \leq C |B(R)|^2 e^{2\omega_G(\kappa R + \ell)} \left( \sum_{k=0}^N \|\mathcal{L}_\lambda^k\|_\infty \right) \|\mathcal{L}_\lambda^n \Psi\|_\infty,$$

9 which is exactly the required inequality.  $\square$

10 **Proposition 5.10.** *The spectral radius  $\rho_\lambda$  of  $\mathcal{L}_\lambda$  satisfies the following inequality*

$$11 \quad \rho_\lambda \geq \exp\left(\frac{\omega_H - \omega_G}{\kappa}\right).$$

12 *Proof.* According to [Lemma 5.9](#) there exist  $B_2 \in \mathbf{R}_+^*$  and a function  $\Psi \in H_\alpha^\infty(\mathcal{J}, \mathcal{H})$  such that for  
13 every  $n \in \mathbf{N}$  we have

$$14 \quad \sum_{g \in S(n) \cap H} e^{-\omega_G d(gx_0, x_0)} \leq B_2 \|\mathcal{L}_\lambda^n \Psi\|_\infty. \quad (17)$$

15 Recall that the canonical map  $H_\alpha^\infty(\mathcal{J}, \mathcal{H}) \rightarrow C(\mathcal{J}, \mathcal{H})$  is 1-Lipschitz (see [Section A.1](#)). Hence for  
16 every  $n \in \mathbf{N}$ , we have

$$17 \quad \|\mathcal{L}_\lambda^n \Psi\|_\infty \leq \|\mathcal{L}_\lambda^n \Psi\|_{\infty, \alpha} \leq \|\mathcal{L}_\lambda^n\|_{\infty, \alpha} \|\Psi\|_{\infty, \alpha}.$$

18 Hence

$$19 \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \ln \|\mathcal{L}_\lambda^n \Psi\|_\infty \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \ln \|\mathcal{L}_\lambda^n\|_{\infty, \alpha} \leq \ln \rho_\lambda.$$

20 The left hand side of the inequality can be interpreted as the critical exponent of the series

$$21 \quad \Upsilon_H(s) = \sum_{n=0}^{\infty} e^{-sn} \|\mathcal{L}_\lambda^n \Psi\|_\infty.$$

22 We use (17) exactly as we did in [Proposition 5.8](#) to prove that for every  $s \in \mathbf{R}$ ,

$$23 \quad \mathcal{P}_H(\omega_G + s\kappa) \leq B_2 e^{s\kappa\ell} \Upsilon_H(s),$$

24 where  $\mathcal{P}_H$  stands for the Poincaré series of  $H$ . Recall that  $s \rightarrow \mathcal{P}_H(s)$  diverges whenever  $s < \omega_H$ .  
25 Consequently the critical exponent of  $\Upsilon_H(s)$  is bounded below by  $(\omega_H - \omega_G)/\kappa$ , hence the result.  $\square$

1 **Corollary 5.11.** *If  $\omega_H = \omega_G$ , then  $\rho_\lambda = 1$ .*

2 *Proof.* It directly follows from the observation that  $\rho_\lambda \leq 1$ . □

3 **Remark.** The converse statement actually holds. It is a consequence of Theorems A.25 and  
 4 B.1. Indeed if  $\rho_\lambda = 1$ , then the group  $H$  is co-amenable in  $G$  (Theorem A.25), hence  $\omega_H = \omega_G$   
 5 (Theorem B.1). Nevertheless we are not aware of an upper bound of the spectral radius  $\rho_\lambda$  in the  
 6 spirit of Proposition 5.10 which would directly leads to the converse direction (and an alternative  
 7 proof of Roblin's theorem).

## 8 5.4 Proofs of the theorems

9 We are now in position to prove Theorems 5.1 and 5.2.

10 *Proof of Theorem 5.1.* The Cayley graph  $\Gamma$  of  $G$  is defined as in Section 5.1. This provides a subshift  
 11 of finite type  $(\mathfrak{H}_0, T)$  as detailed in Section 4.3 together with the labelling map  $\theta: \mathfrak{H}_0 \rightarrow G$  defined  
 12 in Section 4.2. We extract from this dynamical system an irreducible component  $\mathfrak{J}$ , such that the  
 13 extension of  $(\mathfrak{J}, T)$  by  $\theta$  has the visibility property (Proposition 4.10). Using the strategy developed  
 14 in Section 5.2 we define a potential  $F: \mathfrak{J} \rightarrow \mathbf{R}_+^*$  which belongs to  $H_\alpha^\infty(\mathfrak{J}, \mathbf{R})$  for some  $\alpha \in \mathbf{R}_+^*$ . We  
 15 denote by  $\mathcal{L}: H_\alpha^\infty(\mathfrak{J}, \mathbf{C}) \rightarrow H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$  the corresponding transfer operator. Its spectral radius is  
 16  $\rho = 1$  (Proposition 5.8).

17 Let  $H$  be a subgroup. We consider the set  $Y$  of left  $H$ -cosets of  $G$ . The group  $G$  acts on  $Y$  by  
 18 right translations. If the action is amenable, then it follows from Roblin's Theorem (Theorem B.1)  
 19 that  $\omega_H = \omega_G$ . Let us assume now that  $\omega_H = \omega_G$ . The action of  $G$  on  $Y$  induces a unitary  
 20 representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  where  $\mathcal{H}$  stands for  $\ell^2(Y)$ . This leads to a twisted transfer operator  
 21  $\mathcal{L}_\lambda: H_\alpha^\infty(\mathfrak{J}, \mathcal{H}) \rightarrow H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$ . Since  $\omega_H = \omega_G$ , Corollary 5.11 tells us that the spectral radius of  $\mathcal{L}_\lambda$   
 22 is  $\rho_\lambda = 1$ . In particular  $\rho_\lambda = \rho$ . It follows from the amenability criterion (Theorem A.25) that the  
 23 action of  $G$  on  $Y$  is amenable. □

24 *Proof of Theorem 5.2.* The dynamical system  $(\mathfrak{J}, T)$ , the extension map  $\theta: \mathfrak{J} \rightarrow G$ , the potential  
 25  $F \in H_\alpha^\infty(\mathfrak{J}, \mathbf{R})$  as well as the transfer operator  $\mathcal{L}: H_\alpha^\infty(\mathfrak{J}, \mathbf{C}) \rightarrow H_\alpha^\infty(\mathfrak{J}, \mathbf{C})$  are build as in the  
 26 previous proof. In particular the spectral radius of  $\mathcal{L}$  is  $\rho = 1$ . Let  $\eta \in \mathbf{R}_+^*$ , be the constant given  
 27 by Theorem A.28.

28 Let  $H$  be an infinite index subgroup of  $G$ . We consider the set  $Y$  of left  $H$ -cosets of  $G$ . The group  
 29  $G$  acts transitively on  $Y$  by right translations. It induces a unitary representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$   
 30 where  $\mathcal{H}$  stands for  $\ell^2(Y)$ . This leads to a twisted transfer operator  $\mathcal{L}_\lambda: H_\alpha^\infty(\mathfrak{J}, \mathcal{H}) \rightarrow H_\alpha^\infty(\mathfrak{J}, \mathcal{H})$ .  
 31 Since  $H$  has infinite index in  $G$ , Theorem A.28 tells us that the spectral radius  $\rho_\lambda$  of  $\mathcal{L}_\lambda$  is bounded  
 32 above by  $1 - \eta$ . However Proposition 5.10 provides a lower bound for  $\rho_\lambda$  in terms of  $\omega_G$  and  $\omega_H$ .  
 33 It yields  $\omega_H \leq \omega_G - \varepsilon$  where

$$34 \quad \varepsilon = \kappa |\ln(1 - \eta)|.$$

35 Note that  $\varepsilon$  does not depends on  $H$ , which completes the proof of the theorem. □

## 1 A An extension of Kesten's criterion

2 As explained in the introduction, this appendix is deeply inspired by the work of Stadlbauer. We  
 3 prove a variation of his amenability criterion. Our approach makes an explicit use of representation  
 4 theory and operator algebra, which were somehow hidden in [36]. In addition it provides precise  
 5 estimates that can be use to analyse groups with Kazhdan's property (T) (see [Theorem A.28](#)).  
 6 Similar results were also obtained by Dougall in [21].

7 In this section  $(\Sigma, \sigma)$  is a subshift of finite type of the alphabet  $\mathcal{A}$ . We use the same notations  
 8 as in [Section 2.1](#).

### 9 A.1 Function spaces

10 In order to study the dynamical system  $(\Sigma, \sigma)$  and its extensions, we define a family of function  
 11 spaces. To that end, we fix a Banach space  $(E, \|\cdot\|)$ . We endow the space  $\mathcal{C}(\Sigma, E)$  of continuous  
 12 function  $\Phi: \Sigma \rightarrow E$  with the norm  $\|\cdot\|_\infty$  defined by

$$13 \quad \|\Phi\|_\infty = \sup_{x \in \Sigma} \|\Phi(x)\|.$$

14 For our purpose, it will be rather convenient to work with smooth functions. Given  $\alpha > 0$  we  
 15 measure the  $\alpha$ -Hölder variations of  $\Phi$  by the quantity

$$16 \quad \Delta_\alpha(\Phi) = \sup_{x \neq y} \frac{\|\Phi(x) - \Phi(y)\|}{d(x, y)^\alpha}.$$

17 We define the norm  $\|\cdot\|_{\infty, \alpha}$  of  $\Phi$  by

$$18 \quad \|\Phi\|_{\infty, \alpha} = \|\Phi\|_\infty + \Delta_\alpha(\Phi).$$

19 **Definition A.1** (Functions with bounded variations). Let  $\alpha > 0$ . The space  $H_\alpha^\infty(\Sigma, E)$  is the set  
 20 of all maps  $\Phi: \Sigma \rightarrow E$  satisfying  $\|\Phi\|_{\infty, \alpha} < \infty$ .

21 It is a standard exercise to prove that  $H_\alpha^\infty(\Sigma, E)$  is a Banach space. Moreover the canonical  
 22 map  $H_\alpha^\infty(\Sigma, E) \rightarrow \mathcal{C}(\Sigma, E)$  is a 1-Lischitz embedding. Sometimes it is more convenient to focus on  
 23 the *local*  $\alpha$ -Hölder variations of a function. Given  $r \in \mathbf{R}_+^*$  and  $\Phi \in \mathcal{C}(\Sigma, E)$  we let

$$24 \quad \Delta_{\alpha, r}(\Phi) = \sup_{\substack{d(x, y) < r, \\ x \neq y}} \frac{\|\Phi(x) - \Phi(y)\|}{d(x, y)^\alpha}.$$

25 The next lemma explains how  $\Delta_{\alpha, r}(\Phi)$  depends on  $r$ .

26 **Lemma A.2.** *Let  $\alpha > 0$  and  $r \in \mathbf{R}_+^*$ . For every  $\Phi \in \mathcal{C}(\Sigma, E)$  we have*

$$27 \quad \Delta_\alpha(\Phi) - 2r^{-\alpha} \|\Phi\|_\infty \leq \Delta_{\alpha, r}(\Phi) \leq \Delta_\alpha(\Phi).$$

28 **Remark A.3.** In particular  $\Delta_\alpha(\Phi)$  is finite if and only if so is  $\Delta_{\alpha, r}(\Phi)$  for some (hence any)  
 29  $r \in \mathbf{R}_+^*$ .

1 *Proof.* Let  $\Phi \in \mathcal{C}(\Sigma, E)$ . The second inequality is obvious. Let us focus on the first one. Let  
 2  $x, y \in \Sigma$ . If  $d(x, y) < r$ , then by definition we have

$$3 \quad \|\Phi(x) - \Phi(y)\| \leq \Delta_{\alpha, r}(\Phi) d(x, y)^\alpha.$$

4 On the other hand if  $d(x, y) \geq r$ , the triangle inequality yields

$$5 \quad \|\Phi(x) - \Phi(y)\| \leq 2 \|\Phi\|_\infty \leq 2r^{-\alpha} \|\Phi\|_\infty d(x, y)^\alpha.$$

6 Consequently for every distinct  $x, y \in \Sigma$  we have

$$7 \quad \frac{\|\Phi(x) - \Phi(y)\|}{d(x, y)^\alpha} \leq \Delta_{\alpha, r}(\Phi) + 2r^{-\alpha} \|\Phi\|_\infty. \quad \square$$

8 The next lemmas are straightforward. Their proof is left to the reader.

9 **Lemma A.4.** *Let  $\Phi \in \mathcal{C}(\Sigma, \mathbf{R})$  such that  $\Phi(x) > 0$ , for every  $x \in \Sigma$ . The map  $\Phi$  belongs to  
 10  $H_\alpha^\infty(\Sigma, \mathbf{C})$  if and only if so does  $\ln \Phi$ .*

11 **Lemma A.5.** *Let  $f \in H^\infty(\Sigma, \mathbf{R})$  and  $\Phi \in H_\alpha^\infty(\Sigma, E)$ . The pointwise product function  $f\Phi$  belongs  
 12 to  $H_\alpha^\infty(\Sigma, E)$ . Moreover*

$$13 \quad \|f\Phi\|_{\infty, \alpha} \leq \|f\|_{\infty, \alpha} \|\Phi\|_{\infty, \alpha}.$$

## 14 A.2 Ruelle's Perron Frobenius Theorem

15 **Transfer operator.** We fix a potential  $F: \Sigma \rightarrow \mathbf{R}_+^*$  and assume that there exists  $\alpha > 0$  such  
 16 that  $\ln F \in H_\alpha^\infty(\Sigma, \mathbf{R})$ . For every  $n \in \mathbf{N}$ , for every  $x \in \Sigma$ , we let

$$17 \quad F_n(x) = F(x)F(\sigma x) \cdots F(\sigma^{n-1}x).$$

18 By convention  $F_0 = \mathbb{1}$ . To such a potential we associate a transfer operator  $\mathcal{L}: \mathcal{C}(\Sigma, \mathbf{C}) \rightarrow \mathcal{C}(\Sigma, \mathbf{C})$   
 19 defined by

$$20 \quad \mathcal{L}\Phi(x) = \sum_{\sigma y=x} F(y)\Phi(y) = \sum_{a \in \mathcal{A}} \mathbb{1}_{\sigma[a]}(x) F(ax)\Phi(ax).$$

21 One checks easily that the powers of  $\mathcal{L}$  are given by the following formula

$$22 \quad \mathcal{L}^n \Phi(x) = \sum_{\sigma^n y=x} F_n(y)\Phi(y) = \sum_{w \in \mathcal{W}^n} \mathbb{1}_{\sigma^n[w]}(x) F_n(wx)\Phi(wx).$$

23 It is a standard fact that  $\mathcal{L}$  defines a bounded operator of both  $\mathcal{C}(\Sigma, \mathbf{C})$  and  $H_\alpha^\infty(\Sigma, \mathbf{C})$  [26, Sec-  
 24 tion XII.2]. We write  $\rho_\infty$  and  $\rho$  for the spectral radius of  $\mathcal{L}$  seen as an operator of  $\mathcal{C}(\Sigma, \mathbf{C})$  and  
 25  $H_\alpha^\infty(\Sigma, \mathbf{C})$  respectively. One observes easily (see for instance [26, Section XII.2]) that

$$26 \quad \rho_\infty = \lim_{n \rightarrow \infty} \sqrt[n]{\|\mathcal{L}^n \mathbb{1}\|_\infty}. \quad (18)$$

27 Recall that by the Riesz representation theorem, the dual space of  $\mathcal{C}(\Sigma, \mathbf{R})$  can be identified with  
 28 the set of measures on  $\Sigma$ . We write  $\mathcal{L}^*$  for the dual operator of  $\mathcal{L}$ . From now on, unless mentioned  
 29 otherwise, we see  $\mathcal{L}$  as an operator on  $H_\alpha^\infty(\Sigma, \mathbf{C})$  rather than  $\mathcal{C}(\Sigma, \mathbf{C})$ . For a proof of the following  
 30 version of the Ruelle Perron-Frobenius Theorem we refer the reader to [26, Theorem XII.6] or [1,  
 31 Theorem 1.5].



**Theorem A.6** (Ruelle's Perron-Frobenius Theorem). *If  $(\Sigma, \sigma)$  is topologically transitive, then the following holds*

- (i)  $\rho = \rho_\infty$  is positive.
- (ii) There exists a probability measure  $\mu$  on  $\Sigma$  whose support is  $\Sigma$  such that  $\mathcal{L}^* \mu = \rho \mu$ .
- (iii)  $\rho$  is an eigenvalue of  $\mathcal{L}$ ; the corresponding eigenspace has dimension 1; it is spanned by a function  $h \in H_\alpha^\infty(\Sigma, \mathbf{C})$  such that  $h(x) > 0$ , for every  $x \in \Sigma$  and

$$\int_{\Sigma} h d\mu = 1.$$

- (iv)  $\mathcal{L}$  has only finitely many eigenvalue of modulus  $\rho$ ; the corresponding eigenspaces are finite dimensional; the rest of the spectrum of  $\mathcal{L}$  is included in a disc of radius strictly less than  $\rho$ .

### A.3 Twisted transfer operator

Let  $G$  be a finitely generated group. We fix a locally constant map  $\theta: \Sigma \rightarrow G$ . For every  $n \in \mathbf{N}$ , for every  $x \in \Sigma$  we write

$$\theta_n(x) = \theta(x)\theta(\sigma x) \cdots \theta(\sigma^{n-1}x).$$

By convention  $\theta_0$  is the constant map sending  $x$  to the identity  $1 \in G$ . We use this data to produce an extension  $(\Sigma_\theta, \sigma_\theta)$  of  $(\Sigma, \sigma)$  as follows. We let  $\Sigma_\theta = \Sigma \times G$  and define  $\sigma_\theta: \Sigma_\theta \rightarrow \Sigma_\theta$  by

$$\sigma_\theta(x, g) = (\sigma x, g\theta(x)).$$

Recall that the extension  $(\Sigma_\theta, \sigma_\theta)$  has the *visibility property* if there exists a finite subset  $U$  of  $G$  such that for every  $g \in G$ , there exists two elements  $u_1, u_2 \in U$ , a point  $x \in \Sigma$ , and an integer  $n \in \mathbf{N}$ , satisfying  $g = u_1 \theta_n(x) u_2$ .

We now fix a Banach space  $(E, \|\cdot\|)$ . We denote by  $\mathcal{B}(E)$  the space of bounded operators on  $E$  endowed with the operator norm, while  $\text{Isom}(E)$  stands for the set of *linear* isometries of  $E$ . Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a homomorphism. As the  $\theta: \Sigma \rightarrow G$  is locally constant the composition  $\lambda \circ \theta: \Sigma \rightarrow \mathcal{B}(E)$  has  $\alpha$ -Hölder bounded variations. For simplicity we make the following abuse of notations: given  $x \in \Sigma$  and  $n \in \mathbf{N}$ , we write  $\lambda(x)$  for  $\lambda \circ \theta(x)$  and  $\lambda_n(x)$  for  $\lambda \circ \theta_n(x)$ . In particular we have

$$\lambda_n(x) = \lambda(x)\lambda(\sigma x) \cdots \lambda(\sigma^{n-1}x).$$

The representation  $\lambda$  allows us to define a *twisted transfer operator*  $\mathcal{L}_\lambda: \mathcal{C}(\Sigma, E) \rightarrow \mathcal{C}(\Sigma, E)$  as follows

$$\mathcal{L}_\lambda \Phi(x) = \sum_{\sigma y=x} F(y)\lambda(y)^{-1}\Phi(y) = \sum_{a \in \mathcal{A}} \mathbb{1}_{\sigma[a]}(x)F(ax)\lambda(ax)^{-1}\Phi(ax).$$

A standard computation shows that the  $n$ -th power of  $\mathcal{L}_\lambda$  is given by

$$\mathcal{L}_\lambda^n \Phi(x) = \sum_{\sigma^n y=x} F_n(y)\lambda_n(y)^{-1}\Phi(y) = \sum_{w \in \mathcal{W}^n} \mathbb{1}_{\sigma^n[w]}(x)F_n(wx)\lambda_n(wx)^{-1}\Phi(wx).$$

If  $\lambda$  is the trivial representation of  $G$  in  $\mathbf{C}$ , we recover the usual Ruelle Perron-Frobenius operator defined in the previous section.

**Lemma A.7.** *The operator  $\mathcal{L}_\lambda: \mathcal{C}(\Sigma, E) \rightarrow \mathcal{C}(\Sigma, E)$  is bounded. More precisely for every  $n \in \mathbf{N}$ , we have*

$$\|\mathcal{L}_\lambda^n\|_\infty \leq \|\mathcal{L}^n \mathbf{1}\|_\infty.$$

*Proof.* Let  $n \in \mathbf{N}$ . Let  $\Phi \in \mathcal{C}(\Sigma, E)$ . It follows from the triangle inequality that for every  $x \in \Sigma$ ,

$$\|\mathcal{L}_\lambda^n \Phi(x)\| \leq \sum_{w \in \mathcal{W}^n} \mathbf{1}_{\sigma^n[w]}(x) F(w) \|\lambda_n(wx)^{-1} \Phi(wx)\| \leq \mathcal{L}^n \mathbf{1}(x) \|\Phi\|_\infty.$$

Hence  $\mathcal{L}_\lambda^n$  is a bounded operator and its operator norm is at most  $\|\mathcal{L}^n \mathbf{1}\|_\infty$ .  $\square$

**Proposition A.8.** *For every  $n \in \mathbf{N}$ , there exists  $C_n \in \mathbf{R}_+^*$  such that for every linear representation  $\lambda: G \rightarrow \text{Isom}(E)$  into a Banach space  $(E, \|\cdot\|)$ , for every  $\Phi \in H_\alpha^\infty(\Sigma, E)$  we have*

$$\Delta_\alpha(\mathcal{L}_\lambda^n \Phi) \leq e^{-n\alpha} \|\mathcal{L}^n \mathbf{1}\|_\infty \Delta_\alpha(\Phi) + C_n \|\Phi\|_\infty.$$

*Proof.* As  $(\Sigma, \sigma)$  is a subshift of finite type, there exists  $r > 0$  with the following property: for every  $x, y \in \Sigma$ , if  $d(x, y) < r$ , then for every  $n \in \mathbf{N}$ , for every  $w \in \mathcal{W}^n$ ,  $\mathbf{1}_{\sigma^n[w]}(x) = \mathbf{1}_{\sigma^n[w]}(y)$ . Said differently the words that can be added in front of  $x$  or  $y$  are the same. We will take advantage of this fact to estimate the local Hölder variations of the twisted transfer operator. Recall that  $\theta: \Sigma \rightarrow G$  is locally constant. Hence there exists  $m \in \mathbf{N}$  such that  $\theta$  is constant on every cylinder of length  $m$ .

Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a linear representation. Let  $n \in \mathbf{N}$ . Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$ . Let  $x, y \in \Sigma$  be two distinct points such that  $d(x, y) < r$ . We fix  $w \in \mathcal{W}^n$  such that  $\mathbf{1}_{\sigma^n[w]}(x) = \mathbf{1}_{\sigma^n[w]}(y)$  equals 1. A standard computation tells us that

$$\begin{aligned} F_n(wx) \lambda_n(wx)^{-1} \Phi(wx) - F_n(wy) \lambda_n(wy)^{-1} \Phi(wy) &= F_n(wx) \lambda_n(wx)^{-1} (\Phi(wx) - \Phi(wy)) \\ &\quad + F_n(wx) (\lambda_n(wx)^{-1} - \lambda_n(wy)^{-1}) \Phi(wy) \\ &\quad + (F_n(wx) - F_n(wy)) \lambda_n(wy)^{-1} \Phi(wy). \end{aligned}$$

Recall that the image of  $\lambda$  is contained in the isometry group of  $E$ . On the other hand we observe that  $d(wx, wy) = e^{-n} d(x, y)$ . Hence the triangle inequality yields

$$\begin{aligned} &\|F_n(wx) \lambda_n(wx)^{-1} \Phi(wx) - F_n(wy) \lambda_n(wy)^{-1} \Phi(wy)\| \\ &\leq \left( F_n(wx) \Delta_\alpha(\Phi) + F_n(wx) \Delta_\alpha(\lambda_n) \|\Phi\|_\infty + \Delta_\alpha(F_n) \|\Phi\|_\infty \right) e^{-n\alpha} d(x, y)^\alpha. \end{aligned}$$

This inequality holds for every  $w \in \mathcal{W}^n$  such that  $\mathbf{1}_{\sigma^n[w]}(x) = \mathbf{1}_{\sigma^n[w]}(y)$  – which equals  $\mathbf{1}_{\sigma^n[w]}(y)$  – does not vanish. We sum these inequalities to get

$$\frac{\|\mathcal{L}_\lambda^n \Phi(x) - \mathcal{L}_\lambda^n \Phi(y)\|}{d(x, y)^\alpha} \leq e^{-n\alpha} \|\mathcal{L}^n \mathbf{1}\|_\infty \Delta_\alpha(\Phi) + e^{-n\alpha} \left( |\mathcal{W}^n| \Delta_\alpha(F_n) + \|\mathcal{L}^n \mathbf{1}\|_\infty \Delta_\alpha(\lambda_n) \right) \|\Phi\|_\infty.$$

Recall that  $\theta$  is constant on any cylinder of length  $m$ . Hence  $\theta_n$  is constant on any cylinder of length  $n + m$ . It follows from [Lemma A.2](#) that  $\Delta_\alpha(\lambda_n) \leq 2e^{(n+m)\alpha}$ . Hence

$$\frac{\|\mathcal{L}_\lambda^n \Phi(x) - \mathcal{L}_\lambda^n \Phi(y)\|}{d(x, y)^\alpha} \leq e^{-n\alpha} \|\mathcal{L}^n \mathbf{1}\|_\infty \Delta_\alpha(\Phi) + \left( e^{-n\alpha} |\mathcal{W}^n| \Delta_\alpha(F_n) + 2e^{m\alpha} \|\mathcal{L}^n \mathbf{1}\|_\infty \right) \|\Phi\|_\infty.$$

1 This estimation holds for every distinct  $x, y \in \Sigma$  satisfying  $d(x, y) < r$ , hence the right hand side of  
 2 the last inequality is an upper bound of  $\Delta_{\alpha, r}(\mathcal{L}_\lambda^n \Phi)$ . Combined with [Lemma A.2](#) we get that

$$3 \quad \Delta_\alpha(\mathcal{L}_\lambda \Phi) \leq e^{-n\alpha} \|\mathcal{L}^n \mathbb{1}\|_\infty \Delta_\alpha(\Phi) + C_n \|\Phi\|_\infty,$$

4 where

$$5 \quad C_n = e^{-n\alpha} |\mathcal{W}^n| \Delta_\alpha(F_n) + 2e^{m\alpha} \|\mathcal{L}^n \mathbb{1}\|_\infty + 2r^{-\alpha}.$$

6 Observe that the parameter  $C_n$  neither depends on  $\lambda$  nor on  $\Phi$ , thus the proof of the proposition is  
 7 complete.  $\square$

8 Combined with [Lemma A.7](#), the previous proposition tells us that for every  $n \in \mathbf{N}$ , for every  
 9  $\Phi \in H_\alpha^\infty(\Sigma, E)$  we have

$$10 \quad \|\mathcal{L}_\lambda^n \Phi\|_{\infty, \alpha} \leq e^{-n\alpha} \|\mathcal{L}^n \mathbb{1}\|_\infty \|\Phi\|_{\infty, \alpha} + (C_n + \|\mathcal{L}^n \mathbb{1}\|_\infty) \|\Phi\|_\infty. \quad (19)$$

11 Hence we can view  $\mathcal{L}_\lambda$  as a bounded operator on  $H_\alpha^\infty(\Sigma, E)$ . This is the point of view that we will  
 12 adopt in the remainder of this appendix. In particular we denote by  $\|\mathcal{L}_\lambda\|_{\infty, \alpha}$  its corresponding  
 13 operator norm and  $\rho_\lambda$  its spectral radius. Recall that  $\rho_\infty$  stands for the spectral radius of  $\mathcal{L}$  seen  
 14 as an operator of  $\mathcal{C}(\Sigma, \mathbf{C})$ .

15 **Corollary A.9.** *The spectral radius  $\rho_\lambda$  of  $\mathcal{L}_\lambda$  is bounded above by  $\rho_\infty$ .*

16 *Proof.* Let  $\beta > \ln \rho_\infty$ . It follows from (18) that there exists  $n_0 \in \mathbf{N}$ , such that for every  $n \geq n_0$ .

$$17 \quad \|\mathcal{L}^n \mathbb{1}\|_\infty \leq e^{n\beta}.$$

18 We now fix  $n \geq n_0$ . According to [Proposition A.8](#) there exists  $C \in \mathbf{R}_+$  such that for every  
 19  $\Phi \in H_\alpha^\infty(\Sigma, E)$  we have

$$20 \quad \|\mathcal{L}_\lambda^n \Phi\|_{\infty, \alpha} \leq e^{n(\beta-\alpha)} \|\Phi\|_{\infty, \alpha} + C \|\Phi\|_\infty.$$

21 Let  $m \in \mathbf{N}$ . We are going to estimate  $\|\mathcal{L}_\lambda^{nm}\|_{\infty, \alpha}$ . To that end we choose  $\Phi \in H_\alpha^\infty(\Sigma, E)$  and  
 22  $k \in \llbracket 0, m-1 \rrbracket$ . Applying the previous inequality to  $\mathcal{L}_\lambda^{kn} \Phi$  we get

$$23 \quad \left\| \mathcal{L}_\lambda^{(k+1)n} \Phi \right\|_{\infty, \alpha} \leq e^{n(\beta-\alpha)} \|\mathcal{L}_\lambda^{kn} \Phi\|_{\infty, \alpha} + C \|\mathcal{L}_\lambda^{kn} \Phi\|_\infty.$$

24 which combined with [Lemma A.7](#) becomes

$$25 \quad \left\| \mathcal{L}_\lambda^{(k+1)n} \Phi \right\|_{\infty, \alpha} \leq e^{n(\beta-\alpha)} \|\mathcal{L}_\lambda^{kn} \Phi\|_{\infty, \alpha} + C e^{kn\beta} \|\Phi\|_\infty.$$

26 We multiply these inequalities by  $e^{-(k+1)n(\beta-\alpha)}$  and sum them when  $k$  runs over  $\llbracket 0, m-1 \rrbracket$ . It gives

$$27 \quad \|\mathcal{L}_\lambda^{nm} \Phi\|_{\infty, \alpha} \leq e^{nm(\beta-\alpha)} \|\Phi\|_{\infty, \alpha} + e^{nm\beta} \frac{C e^{-n(\beta-\alpha)}}{e^{n\alpha} - 1} \|\Phi\|_\infty.$$

28 Recall that  $\|\Phi\|_\infty \leq \|\Phi\|_{\infty, \alpha}$ . So we have proved that there exists  $D_n \in \mathbf{R}_+^*$  such that for every  
 29  $m \in \mathbf{N}$ , for every  $\Phi \in H^\infty(\Sigma, E)$

$$30 \quad \|\mathcal{L}_\lambda^{nm} \Phi\|_{\infty, \alpha} \leq e^{nm\beta} D_n \|\Phi\|_{\infty, \alpha}.$$

31 Hence for every  $m \in \mathbf{N}$ , we get

$$32 \quad \|\mathcal{L}_\lambda^{nm}\|_{\infty, \alpha} \leq e^{nm\beta} D_n.$$

33 Passing to the limit when  $m$  tends to infinity we obtain

$$34 \quad \ln \rho_\lambda = \lim_{m \rightarrow \infty} \frac{1}{nm} \ln \left( \|\mathcal{L}_\lambda^{nm}\|_{\infty, \alpha} \right) \leq \beta.$$

35 This inequality holds for every  $\beta \in \mathbf{R}$  such that  $\beta > \ln \rho_\infty$ , thus  $\rho_\lambda \leq \rho_\infty$ .  $\square$

## 1 A.4 Renormalization

2 In this section we study how the transfer operators are affected when replacing the potential  $F$  by  
 3 a rescaled and/or an homologous potential. This will allow us later to assume that  $\mathcal{L}$  has spectral  
 4 radius 1 and fixes the constant map  $\mathbb{1}$ , which will considerably lighten the notations.

5 **Rescaling a homologous potential.** Assume that the system  $(\Sigma, \sigma)$  is topologically transitive.  
 6 Let  $h \in H_\alpha^\infty(\Sigma, \mathbf{C})$  be the positive eigenvector given by [Theorem A.6](#). We define a new potential  
 7  $F': \Sigma \rightarrow \mathbf{R}_+^*$  by

$$8 \quad F'(x) = \frac{1}{\rho} \cdot \frac{h(x)}{h \circ \sigma(x)} F(x), \quad (20)$$

9 so that

$$10 \quad \ln F'(x) = -\ln \rho + \ln(h(x)) - \ln(h \circ \sigma(x)) + \ln F(x).$$

11 We know that  $\ln F$  belongs to  $H_\alpha^\infty(\Sigma, \mathbf{C})$ . The same holds for  $h$ . It follows that  $\ln F'$  belongs  
 12 to  $H_\alpha^\infty(\Sigma, \mathbf{C})$  ([Lemma A.4](#)). In particular,  $F'$  satisfies the same assumptions as  $F$ . We write  
 13  $\mathcal{L}': H_\alpha^\infty(\Sigma, \mathbf{C}) \rightarrow H_\alpha^\infty(\Sigma, \mathbf{C})$  and  $\mathcal{L}'_\lambda: H_\alpha^\infty(\Sigma, E) \rightarrow H_\alpha^\infty(\Sigma, E)$  for the corresponding usual and  
 14 twisted transfer operators.

15 **Comparing the operators.** Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$ . We observe that for every  $x \in \Sigma$

$$16 \quad \mathcal{L}'_\lambda \Phi(x) = \sum_{\sigma y=x} \frac{h(y)}{\rho h(x)} F(y) \lambda(y)^{-1} \Phi(y) = \frac{1}{\rho h(x)} \mathcal{L}_\lambda(h\Phi)(x).$$

17 In particular, since  $h$  is a positive eigenvector of  $\mathcal{L}$  for the eigenvalue  $\rho$ , we get that  $\mathcal{L}'\mathbb{1} = \mathbb{1}$ . It  
 18 follows from [\(18\)](#) combined with [Theorem A.6](#) that the spectral radius of  $\mathcal{L}'$  is 1. Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$ .  
 19 A proof by induction shows that for every  $n \in \mathbf{N}$ ,

$$20 \quad \rho^n h \mathcal{L}'_\lambda{}^n(\Phi) = \mathcal{L}_\lambda^n(h\Phi). \quad (21)$$

21 Since  $h$  is a positive continuous map on a compact set, there exists  $m, M \in \mathbf{R}_+^*$  such that for every  
 22  $x \in \Sigma$ , we have  $m \leq h(x) \leq M$ . Hence  $1/h$  also belongs to  $H_\alpha^\infty(\Sigma, \mathbf{C})$ . Combining [\(21\)](#) with  
 23 [Lemma A.5](#) we observe that there exists  $A_1, A_2 \in \mathbf{R}_+^*$  such that for every  $n \in \mathbf{N}$ ,

$$24 \quad A_1 \|\mathcal{L}'_\lambda{}^n\|_{\infty, \alpha} \leq \rho^n \|\mathcal{L}'_\lambda{}^n\|_{\infty, \alpha} \leq A_2 \|\mathcal{L}'_\lambda{}^n\|_{\infty, \alpha}.$$

25 Hence the spectral radius of  $\mathcal{L}'_\lambda$  is  $\rho_\lambda/\rho$ .

## 26 A.5 Invariant and almost invariant vectors

27 In this section we suppose that the system  $(\Sigma, \sigma)$  is topologically transitive. In addition we as-  
 28 sume that  $\mathcal{L}$  has spectral radius 1 and fixes  $\mathbb{1}$ . Under these assumptions, the operator norm of  
 29  $\mathcal{L}_\lambda: \mathcal{C}(\Sigma, E) \rightarrow \mathcal{C}(\Sigma, E)$  is at most 1 ([Lemma A.7](#)). Similarly [Proposition A.8](#) yields

30 **Proposition A.10.** *There exists  $C_1 \in \mathbf{R}_+^*$  such that for every linear representation  $\lambda: G \rightarrow$   
 31  $\text{Isom}(E)$  into a Banach space  $(E, \|\cdot\|)$ , for every  $\Phi \in H_\alpha^\infty(\Sigma, E)$  we have*

$$32 \quad \Delta_\alpha(\mathcal{L}_\lambda \Phi) \leq e^{-\alpha} \Delta_\alpha(\Phi) + C_1 \|\Phi\|_\infty, \quad (22)$$

1 We would like to understand the behavior  $\mathcal{L}_\lambda$  when 1 is a spectral value. To that end we study  
 2 invariant and almost invariant vectors of the twisted transfer operator. Given  $\varepsilon \in \mathbf{R}_+^*$  we say that  
 3  $\Phi \in H_\alpha^\infty(\Sigma, E)$  is an  $\varepsilon$ -invariant vector if

$$4 \quad \|\mathcal{L}_\lambda \Phi - \Phi\|_{\infty, \alpha} < \varepsilon \|\Phi\|_{\infty, \alpha}.$$

5 **Almost invariant vectors.** We start with some preliminary properties of almost invariants  
 6 vectors for the twisted transfer operator.

7 **Proposition A.11.** *There exists  $D_2, \eta \in \mathbf{R}_+^*$  with the following property. For every linear repre-*  
 8 *sentation  $\lambda: G \rightarrow \text{Isom}(E)$  into a Banach space  $(E, \|\cdot\|)$ , if  $\Phi \in H_\alpha^\infty(\Sigma, E)$  is an  $\eta$ -invariant vector*  
 9 *of  $\mathcal{L}_\lambda$ , then  $\Delta_\alpha(\Phi) \leq D_2 \|\Phi\|_\infty$ .*

10 *Proof.* Let  $C_1$  be the parameter given by [Proposition A.10](#). We fix  $\eta \in \mathbf{R}_+^*$  such that  $\eta < 1 - e^{-\alpha}$ .  
 11 Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a linear representation into a Banach space  $(E, \|\cdot\|)$ . Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$   
 12 be an  $\eta$ -invariant vector of  $\mathcal{L}_\lambda$ . The triangle inequality combined with [\(22\)](#) yields

$$13 \quad \Delta_\alpha(\Phi) \leq \Delta_\alpha(\mathcal{L}_\lambda \Phi) + \Delta_\alpha(\mathcal{L}_\lambda \Phi - \Phi) \leq e^{-\alpha} \Delta_\alpha(\Phi) + C_1 \|\Phi\|_\infty + \eta \|\Phi\|_{\infty, \alpha}.$$

14 Hence

$$15 \quad \Delta_\alpha(\Phi) \leq \frac{C_1 + \eta}{1 - (e^{-\alpha} + \eta)} \|\Phi\|_\infty. \quad \square$$

16 **Invariants vectors.** We now detail the behavior of invariant vectors for the twisted transfer  
 17 operator.

18 **Lemma A.12.** *Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$ . If  $\mathcal{L}_\lambda \Phi = \Phi$ , then the map  $\Sigma \rightarrow \mathbf{R}$  sending  $x$  to  $\|\Phi(x)\|$  is*  
 19 *constant.*

20 *Proof.* We denote by  $\Psi: \Sigma \rightarrow \mathbf{R}$  the map defined by  $\Psi(x) = \|\Phi(x)\|$ . It is an element of  $H_\alpha^\infty(\Sigma, \mathbf{R})$ .  
 21 Let  $x_0 \in \Sigma$  such that

$$22 \quad \Psi(x_0) = \sup_{x \in \Sigma} \Psi(x).$$

23 Such a point exists as  $\Psi$  is a continuous function on a compact set. Let  $x \in \Sigma$ . Let  $\varepsilon \in \mathbf{R}_+^*$ .  
 24 Since the system  $(\Sigma, \sigma)$  is topologically transitive, there exists  $n \in \mathbf{N}$  and  $w_0 \in \mathcal{W}^n$  such that  $w_0 x_0$   
 25 belongs to  $\Sigma$  and  $d(x, w_0 x_0) \leq \varepsilon$ . According to the triangle inequality we have

$$26 \quad \Psi(x_0) \leq \mathcal{L}^n \Psi(x_0) \leq \sum_{w \in \mathcal{W}^n} \mathbb{1}_{\sigma^n[w]}(x_0) F_n(w x_0) \Psi(w x_0).$$

27 Recall that  $\mathcal{L}^n \mathbb{1} = \mathbb{1}$ . Hence the right hand side is a convex combination of terms of the form  
 28  $\Psi(w x_0)$ , all of them being bounded above by  $\Psi(x_0)$ . Consequently  $\Psi(w_0 x_0) = \Psi(x_0)$ . Since  $\Psi$  has  
 29 bounded variations we get

$$30 \quad \|\Psi(x) - \Psi(x_0)\| = \|\Psi(x) - \Psi(w_0 x_0)\| \leq \Delta_\alpha(\Psi) d(x, w_0 x_0)^\alpha \leq \Delta_\alpha(\Psi) \varepsilon^\alpha.$$

31 This inequality holds for every sufficiently small positive  $\varepsilon$  hence  $\Psi(x) = \Psi(x_0)$ . This proves that  
 32  $\Psi$  is a constant function equal to  $\Psi(x_0)$ .  $\square$

1 **Lemma A.13.** *Assume that  $E$  is strictly convex. Let  $\Phi \in H_\alpha^\infty(\Sigma, E)$ . If  $\mathcal{L}_\lambda \Phi = \Phi$ , then for every*  
 2  *$x \in \Sigma$ , for every  $n \in \mathbf{N}$ , we have*

$$3 \quad \lambda_n(x)\Phi(\sigma^n x) = \Phi(x).$$

4 *Proof.* It follows from [Lemma A.12](#) that for every  $x \in \Sigma$ , we have  $\|\Phi(x)\| = \|\Phi\|_\infty$ . Let  $x \in \Sigma$  and  
 5  $n \in \mathbf{N}$ . We let  $z = \sigma^n x$ . Observe that

$$6 \quad \Phi(z) = \mathcal{L}_\lambda^n \Phi(z) = \sum_{\sigma^n y = z} F_n(y) \lambda_n(y)^{-1} \Phi(y).$$

7 Recall that  $\mathcal{L}^n \mathbb{1} = \mathbb{1}$ . Hence the right hand side is a convex combination of vectors of the form  
 8  $\lambda_n(y)^{-1} \Phi(y)$ . Since the image of  $\lambda$  is contained in the isometry group of  $E$ , their norm is the same as  
 9 the one of  $\Phi(z)$ , namely  $\|\Phi\|_\infty$ . The space  $E$  being strictly convex, we get that  $\Phi(z) = \lambda_n(y)^{-1} \Phi(y)$ ,  
 10 for every  $y \in \sigma^{-n}(\{z\})$ . This holds in particular for  $y = x$ , hence the result.  $\square$

11 **From the transfer operator to the representation.** The goal is now to prove that if  $\mathcal{L}$  and  $\mathcal{L}_\lambda$   
 12 have the same spectral radius, then  $\lambda$  admits almost invariant vectors. We first cover the case when  
 13 1 is an *eigenvalue* of  $\mathcal{L}_\lambda$  ([Proposition A.14](#)). In this situation we combine a convexity argument  
 14 taking place in  $E$  with the visibility property to prove the existence of a non-zero vector  $\phi_0 \in E$   
 15 that is fixed by a finite index subgroup  $G_0$  of  $G$ . The second step deals with the general situation,  
 16 i.e. when  $\mathcal{L}_\lambda$  admits almost invariant vectors ([Proposition A.18](#)). The proof of this proposition is  
 17 by contradiction. Negating the statement provides a family of counterexamples. Then, using an  
 18 ultra-limit argument, we are able to produce a new twisted transfer operator for which 1 is an  
 19 eigenvalue, therefore reducing the general case to the previous one.

20 **Proposition A.14.** *We assume that the extension of  $(\Sigma, \sigma)$  by  $\theta$  has the visibility property. There*  
 21 *exists a finite index subgroup  $G_0$  of  $G$  such that for every representation  $\lambda: G \rightarrow \text{Isom}(E)$  into a*  
 22 *uniformly convex Banach space  $(E, \|\cdot\|)$  the following holds. If 1 is an eigenvalue of the twisted*  
 23 *transfer operator  $\mathcal{L}_\lambda$ , then the representation  $\lambda$  restricted to  $G_0$  has a non-zero invariant vector*  
 24  *$\phi_0 \in E$ .*

25 *Proof.* We start the proof by introducing a few auxiliary objects that will lead to the definition  
 26 of  $G_0$ . The most important point is that these objects do not involve the representation of  $G$ .  
 27 Let  $D_2 \in \mathbf{R}$  be the constant given by [Proposition A.11](#). We fix an integer  $m \in \mathbf{N}$  such that  
 28  $e^{-m\alpha} D_2 < 1/2$ . Up to increasing the value of  $m$  we can assume that  $\theta: \Sigma \rightarrow G$  is constant when  
 29 restricted to any cylinder of length  $m$ . Since  $(\Sigma, \sigma)$  is an irreducible subshift of finite type, there  
 30 exists a finite subset  $\mathcal{W}_0 \subset \mathcal{W}$  with the following property: for every  $w, w' \in \mathcal{W}$ , there exists  
 31  $w_0 \in \mathcal{W}_0$  such that  $ww_0w'$  is admissible. We denote by  $N$  the length of the longest word in  $\mathcal{W}_0$ .  
 32 As  $\theta$  is locally constant, the set

$$33 \quad U = \{\theta_k(x) \mid x \in \Sigma, k \in \llbracket 0, m + N \rrbracket\}$$

34 is finite. It follows from the visibility property, there exists a finite subset  $U'$  of  $G$  with the following  
 35 property: for every  $g \in G$ , there exists two elements  $u'_1, u'_2 \in U'$ , a point  $x \in \Sigma$ , and an integer  
 36  $n \in \mathbf{N}$ , satisfying  $g = u'_1 \theta_n(x) u'_2$ . Finally, we let

$$37 \quad K = |U|^2 |U'|^2.$$

38 Since  $G$  is finitely generated, it has only finitely many finite index subgroups whose index does not  
 39 exceed  $K$ . We define  $G_0$  as the intersection of these subgroups. It is a finite index subgroup of  $G$ .

1 Let us now fix a representation  $\lambda: G \rightarrow \text{Isom}(E)$  into a uniformly convex Banach space  $(E, \|\cdot\|)$   
 2 such that 1 is an eigenvalue of the twisted transfer operator  $\mathcal{L}_\lambda$ . We choose a non-zero eigenvector  
 3  $\Phi \in H_\alpha^\infty(\Sigma, E)$  of  $\mathcal{L}_\lambda$ , i.e.  $\mathcal{L}_\lambda\Phi = \Phi$ . It follows that the map  $x \rightarrow \|\Phi(x)\|$  is constant ([Lemma A.12](#))  
 4 and  $\Delta_\alpha(\Phi) \leq D_2\|\Phi\|_\infty$  ([Proposition A.11](#)).

5 We write  $v_1, \dots, v_\ell$  for the collection of admissible words of length  $m$ . Let  $i \in \llbracket 1, \ell \rrbracket$ . We denote  
 6 by  $C_i$  the closure of the convex hull of  $\Phi([v_i])$ . Since  $E$  is uniformly convex, the zero vector  $0 \in E$   
 7 admits a unique projection on  $C_i$  that we denote by  $\phi_i$ . It follows from our choice of  $m$ , that for  
 8 every  $x, x' \in [v_i]$ , we have

$$9 \quad \|\Phi(x) - \Phi(x')\| \leq \Delta_\alpha(\Phi)e^{-m\alpha} < \frac{1}{2} \|\Phi(x)\|.$$

10 Consequently  $C_i$  does not contain the zero vector and therefore  $\phi_i$  is non-zero. We define  $H_i$  as  
 11 the pre-image by  $\lambda: G \rightarrow \text{Isom}(E)$  of the stabilizer of  $\phi_i$ . We are going to show that  $H_i$  is a finite  
 12 index subgroup of  $G$ . To that end we start by proving the following lemma.

13 **Lemma A.15.** *For every  $x \in \Sigma$ , for every  $n \in \mathbf{N}$ , there exists two elements  $u_1, u_2 \in U$  such that*  
 14  *$u_1\theta_n(x)u_2 \in H_i$ .*

15 *Proof.* Let  $x \in \Sigma$  and  $n \in \mathbf{N}$ . For simplicity we let  $g = \theta_n(x)$ . Let  $w$  be the prefix of length  
 16  $n + m$  of  $x$ . Since  $\theta: \Sigma \rightarrow G$  is constant when restricted on any cylinder of length  $m$ , we note that  
 17  $\theta_n(x') = \theta_n(x)$  for every  $x' \in [w]$ . By the very definition of  $\mathcal{W}_0$ , there exists  $w_1, w_2 \in \mathcal{W}_0$  such  
 18 that  $v_i w_1 w w_2 v_i$  is admissible. We denote by  $p$  and  $q$  the length of  $v_i w_1$  and  $w w_2$  respectively and  
 19 observe that  $p \leq m + N$  and  $q \leq n + m + N$ .

20 We now claim that there exist  $u_1, u_2 \in U$  such that for every  $y_1 \in [v_i]$ , there exists  $y_2 \in [v_i]$   
 21 satisfying

$$22 \quad \lambda(u_1 g u_2) \Phi(y_1) = \Phi(y_2).$$

23 Choose  $y_1 \in [v_i]$ . It follows from our choice of  $w_1$  and  $w_2$  that  $y_2 = v_i w_1 w w_2 y_1$  and  $z = w w_2 y_1$  are  
 24 two well defined points of  $\Sigma$ . Moreover  $z = \sigma^p y_2$  and  $y_1 = \sigma^q z$ . According to [Lemma A.13](#) we have

$$25 \quad \lambda_{p+q}(y_2) \Phi(y_1) = \lambda_{p+q}(y_2) \Phi(\sigma^{p+q} y_2) = \Phi(y_2).$$

26 Note that  $y_2$  belongs to  $[v_i]$ . Hence it suffices to show that  $\theta_{p+q}(y_2)$  can be written  $u_1 g u_2$  where  $u_1$   
 27 and  $u_2$  do not depend on  $y_1$ . The cocycle relation that  $\theta_{p+q}$  satisfies yields

$$28 \quad \theta_{p+q}(y_2) = \theta_p(y_2) \theta_n(\sigma^p y_2) \theta_{q-n}(\sigma^{p+n} y_2) = \theta_p(y_2) \theta_n(z) \theta_{q-n}(\sigma^n z).$$

29 By construction  $z$  belongs to  $[w]$ , hence  $\theta_n(z) = \theta_n(x) = g$ . Observe that  $p$  and  $q - n$  are bounded  
 30 above by  $m + N$ . Hence  $u_1 = \theta_p(y_2)$  and  $u_2 = \theta_{q-n}(\sigma^n z)$  belong to  $U$ . The proof of the claim  
 31 will be complete if we can prove that  $u_1$  and  $u_2$  do not depend on  $y_1$ . Consider  $y'_1$  another element  
 32 of  $[v_i]$  and let as previously  $y'_2 = v_i w_1 w w_2 y'_1$  and  $z = w w_2 y'_1$ . We observe that  $z$  and  $z'$  belong to  
 33 the same cylinder, namely  $[w w_2 v_i]$  whose length is bounded below by  $q + m$ . Hence  $\sigma^n z$  and  $\sigma^n z'$   
 34 coincide on the first  $q - n + m$  letters. Since  $\theta: \Sigma \rightarrow G$  is constant on any cylinder of length  $m$ , we  
 35 get  $\theta_{q-n}(\sigma^n z) = \theta_{q-n}(\sigma^n z')$ . The same argument shows that  $\theta_p(y_2) = \theta_p(y'_2)$ , which completes the  
 36 proof of our claim.

37 It follows from the claim that  $\lambda(u_1 g u_2)$  maps  $C_i$  into itself as well. In particular,  $\lambda(u_1 g u_2) \phi_i$   
 38 belongs to  $C_i$ . However  $\lambda(u_1 g u_2)$  being a linear isometry we have  $\|\lambda(u_1 g u_2) \phi_i\| = \|\phi_i\|$ . Recall

1 that we defined  $\phi_i$  as the projection of 0 onto  $C_i$ . By unicity of the projection ( $E$  is uniformly  
 2 convex) we get  $\lambda(u_1 g u_2) \phi_i = \phi_i$ . In other words  $u_1 g u_2 = u_1 \theta_n(x) u_2$  belongs to  $H_i$  which completes  
 3 the proof of the lemma.  $\square$

4 As the extension of  $(\Sigma, \sigma)$  by  $\theta$  satisfies the visibility property, the previous lemma has the  
 5 following consequence:

$$6 \quad G = \bigcup_{u_1, u_2 \in U, u'_1, u'_2 \in U'} u'_1 u_1^{-1} H_i u_2^{-1} u'_2 = \bigcup_{u_1, u_2 \in U, u'_1, u'_2 \in U'} \left[ (u'_1 u_1^{-1}) H_i (u'_1 u_1^{-1})^{-1} \right] (u'_1 u_1^{-1} u_2^{-1} u'_2).$$

7 In other words  $G$  can be covered by finitely many cosets of conjugates of  $H_i$ . Moreover the number  
 8 of these cosets is bounded above by

$$9 \quad K = |U|^2 |U'|^2.$$

10 According to [31, Lemma 4.1],  $H_i$  is a finite index subgroup of  $G$  and  $[G : H_i] \leq K$ . It follows from  
 11 its definition that  $G_0$  is a subgroup of  $H_i$ , thus the representation  $\lambda$  restricted to  $G_0$  has a non-zero  
 12 invariant vector, namely  $\phi_i$ .  $\square$

13 **Definition A.16.** We say that a collection  $\mathcal{E}$  of Banach spaces is *uniformly convex* if for every  
 14  $\varepsilon > 0$  there exists  $\eta > 0$  such that for every space  $(E, \|\cdot\|)$  of  $\mathcal{E}$ , for every unit vectors  $\phi, \phi' \in E$ , if  
 15  $\|\phi - \phi'\| \geq \varepsilon$ , then  $\|\phi + \phi'\| \leq 2(1 - \eta)$ .

16 **Remark.** This definition not only asks that each space  $E \in \mathcal{E}$  is uniformly convex, but also that  
 17 the parameters quantifying their rotundity work for all spaces simultaneously.

18 **Definition A.17.** Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a representation of  $G$  into a Banach space. Let  $S$  be a  
 19 finite subset of  $G$  and  $\varepsilon \in \mathbf{R}_+^*$ . A vector  $\phi \in E$  is  $(S, \varepsilon)$ -invariant (with respect to  $\lambda$ ) if

$$20 \quad \sup_{s \in S} \|\lambda(s)\phi - \phi\| < \varepsilon \|\phi\|.$$

21 The representation  $\lambda: G \rightarrow \text{Isom}(E)$  *almost has invariant vectors* if for every finite subset  $S$  of  $G$ ,  
 22 for every  $\varepsilon \in \mathbf{R}_+^*$  there exists an  $(S, \varepsilon)$ -invariant vector.

23 **Proposition A.18.** *We assume that the extension of  $(\Sigma, \sigma)$  by  $\theta$  has the visibility property. There*  
 24 *exists a finite index subgroup  $G_0$  of  $G$  with the following property. Let  $\mathcal{E}$  be a uniformly convex*  
 25 *collection of Banach spaces. For every finite subset  $S_0$  of  $G_0$ , for every  $\varepsilon \in \mathbf{R}_+^*$ , there exists*  
 26  *$\eta \in \mathbf{R}_+^*$ , such that the following holds. Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a representation of  $G$  into a*  
 27 *Banach space  $(E, \|\cdot\|)$  of  $\mathcal{E}$ . If the twisted transfer operator  $\mathcal{L}_\lambda$  has an  $\eta$ -invariant vector, then the*  
 28 *representation  $\lambda$  admits an  $(S_0, \varepsilon)$ -invariant vector.*

29 Before giving the proof, we recall some useful material regarding ultra-limit of Banach spaces.

30 **Ultra-limit of Banach spaces.** Let  $\omega: \mathcal{P}(\mathbf{N}) \rightarrow \{0, 1\}$  be a *non-principal ultra-filter*, i.e. a  
 31 finitely additive map which vanishes on any finite subset of  $\mathbf{N}$  and such that  $\omega(\mathbf{N}) = 1$ . We say  
 32 that a property  $\mathcal{P}_n$  is true  $\omega$ -almost surely ( $\omega$ -as) if  $\omega(\{n \in \mathbf{N} \mid \mathcal{P}_n \text{ holds}\}) = 1$ . A real sequence  
 33  $(u_n)$  is  $\omega$ -essentially bounded ( $\omega$ -eb) if there exists  $M \in \mathbf{R}$  such that  $|u_n| \leq M$   $\omega$ -as. We say that  
 34 the  $\omega$ -limit of  $(u_n)$  is  $\ell \in \mathbf{R}$  and write  $\lim_\omega u_n = \ell$ , if for every  $\varepsilon > 0$ , we have  $d(u_n, \ell) < \varepsilon$   $\omega$ -as.  
 35 Any real sequence which is  $\omega$ -eb admits an  $\omega$ -limit [4].



1 Let  $(E_n, \|\cdot\|)$  be a sequence of Banach spaces. We define the ultra-product  $\Pi_\omega E_n$  as the set

$$2 \quad \Pi_\omega E_n = \{(\phi_n) \in \Pi_{\mathbf{N}} E_n \mid \|\phi_n\| \omega\text{-eb}\}.$$

3 We endow  $\Pi_\omega E_n$  with the following equivalence relation:  $(\phi_n) \equiv (\phi'_n)$  if  $\lim_\omega \|\phi_n - \phi'_n\| = 0$ .

4 **Definition A.19.** The  $\omega$ -limit of the sequence  $(E_n, \|\cdot\|)$  that we denote  $\lim_\omega E_n$  is the quotient  
5 of  $\Pi_\omega E_n$  by the equivalence relation  $\equiv$ .

6 If  $(\phi_n)$  is an element of  $\Pi_\omega E_n$  we write  $\lim_\omega \phi_n$  for its image in  $\lim_\omega E_n$ . The set  $\lim_\omega E_n$   
7 naturally comes with a structure of vector space characterized as follows.

8 (i)  $(\lim_\omega \phi_n) + (\lim_\omega \phi'_n) = \lim_\omega (\phi_n + \phi'_n)$ , for every  $(\phi_n), (\phi'_n) \in \Pi_\omega E_n$ ;

9 (ii)  $c(\lim_\omega \phi_n) = \lim_\omega (c\phi_n)$ , for every  $(\phi_n) \in \Pi_\omega E_n$  and  $c \in \mathbf{C}$ .

10 The space  $\lim_\omega E_n$  carries also a norm characterized by

$$11 \quad \left\| \lim_\omega \phi_n \right\| = \lim_\omega \|\phi_n\|, \quad \forall (\phi_n) \in \Pi_\omega E_n.$$

12 One can check that  $(\lim_\omega E_n, \|\cdot\|)$  is a Banach space. The next lemma is a straightforward exercise.

13 **Lemma A.20.** *If  $(E_n, \|\cdot\|)$  is a uniformly convex collection of Banach spaces, then  $(\lim_\omega E_n, \|\cdot\|)$   
14 is uniformly convex.*

15 *Proof of Proposition A.18.* We denote by  $G_0$  the finite index subgroup of  $G$  given by [Proposi-](#)  
16 [tion A.14](#). We denote by  $D_2$  and  $\eta$  the parameters given by [Proposition A.11](#). Let  $\mathcal{E}$  be a uniformly  
17 convex collection of Banach spaces.

18 **A family of counterexamples.** Let  $S_0$  be a finite subset of  $G_0$  and  $\varepsilon \in \mathbf{R}_+^*$ . Let  $(\eta_n)$  be a  
19 sequence of positive real numbers converging to zero. Assume that the proposition is false. It  
20 means that for every  $n \in \mathbf{N}$ , there exists a representation  $\lambda_n: G \rightarrow \text{Isom}(E_n)$  where  $(E_n, \|\cdot\|)$   
21 belongs to  $\mathcal{E}$ , with the following properties:

22 (i) the twisted transfer operator  $\mathcal{L}_{\lambda_n}$  has an  $\eta_n$ -invariant vector  $\Phi_n \in H_\alpha^\infty(\Sigma, E_n)$ ;

23 (ii) the representation  $\lambda_n$  does not admit any  $(S_0, \varepsilon)$ -invariant vector.

24 In the remainder of the proof, we write for simplicity  $\mathcal{L}_n$  instead of  $\mathcal{L}_{\lambda_n}$ .

25 **Almost invariant vectors for the twisted operator.** Without loss of generality we can as-  
26 sume that  $\|\Phi_n\|_\infty = 1$ , for every  $n \in \mathbf{N}$ . Recall that  $D_2, \eta \in \mathbf{R}_+^*$  are the parameter given by  
27 [Proposition A.11](#). Since  $(\eta_n)$  converges to 0, up to throwing away the first terms of the sequence,  
28 we can assume that  $\eta_n \leq \eta$  for every  $n \in \mathbf{N}$ . It follows now from [Proposition A.11](#) that for every  
29  $n \in \mathbf{N}$ , we have

$$30 \quad \Delta_\alpha(\Phi_n) \leq D_2 \|\Phi_n\|_\infty \leq D_2. \quad (23)$$

31 We now fix  $k \in \mathbf{N}$  such that  $e^{-k\alpha} D_2 < 1/2$ . We write  $v_1, \dots, v_\ell$  for the collection of all admissible  
32 words of length  $k$ . Up to passing to a subsequence we may assume that there exists  $i \in \llbracket 1, \ell \rrbracket$  such  
33 that the map  $x \rightarrow \|\Phi_n(x)\|$  achieved its maximum in  $[v_i]$ . By reordering the elements  $v_1, \dots, v_\ell$  we  
34 can actually assume that  $i = 1$ . In particular it follows from (23) and our choice of  $k$ , that for every  
35  $n \in \mathbf{N}$ , for every  $x \in [v_1]$ , we have  $\|\Phi_n(x)\| \geq 1/2$ .

1 **The Banach space  $E_\infty$ .** We now fix a non-principal ultra-filter  $\omega$ . The limit space  $E_\infty = \lim_\omega E_n$   
 2 is a uniformly convex Banach space ([Lemma A.20](#)). The next step is to define a representation  
 3  $\lambda_\infty : G \rightarrow \text{Isom}(E_\infty)$ . Given  $g \in G$  and a vector  $\phi = \lim_\omega \phi_n$  of  $E_\infty$ , we let

$$4 \quad \lambda_\infty(g)\phi = \lim_\omega [\lambda_n(g)\phi_n].$$

5 One easily checks that  $\lambda_\infty(g)$  is a well-defined linear isometry of  $E_\infty$ . Moreover that map  $\lambda_\infty : G \rightarrow$   
 6  $\text{Isom}(E_\infty)$  obtained in this way is a homomorphism. In particular, one can consider the twisted  
 7 transfer operator

$$8 \quad \mathcal{L}_{\lambda_\infty} : H_\alpha^\infty(\Sigma, E_\infty) \rightarrow H_\alpha^\infty(\Sigma, E_\infty).$$

9 that for simplicity we denote  $\mathcal{L}_\infty$ .

10 **The eigenvector  $\Phi_\infty$ .** We now use the sequence  $(\Phi_n)$  to produce an eigenvector of  $\mathcal{L}_\infty$ . Recall  
 11 that for every  $n \in \mathbf{N}$ , for every  $x \in \Sigma$ , we have  $\|\Phi_n(x)\| \leq 1$ . Hence we can define a map  
 12  $\Phi_\infty : \Sigma \rightarrow E_\infty$  as follows

$$13 \quad \Phi_\infty(x) = \lim_\omega \Phi_n(x), \quad \forall x \in \Sigma.$$

14 Note that  $\Phi_\infty$  is bounded. More precisely,  $\|\Phi_\infty\|_\infty \leq 1$ . Recall that for every  $\Delta_\alpha(\Phi_n) \leq D_2$  for  
 15 every  $n \in \mathbf{N}$ . It directly follows that  $\Delta_\alpha(\Phi_\infty) \leq D_2$ . Hence  $\Phi_\infty$  belongs to  $H_\alpha^\infty(\Sigma, E_\infty)$ . We also  
 16 observe that  $\Phi_\infty$  is non-trivial. Indeed by construction  $\|\Phi_n(x)\| \geq 1/2$ , for every  $n \in \mathbf{N}$ , for every  
 17  $x \in [v_1]$ . It follows that  $\Phi_\infty$  restricted to  $[v_1]$  does not vanish. Finally, we claim that  $\mathcal{L}_{\lambda_\infty} \Phi_\infty = \Phi_\infty$ .  
 18 Let  $x \in \Sigma$ . Since  $\mathcal{L}\mathbf{1} = \mathbf{1}$  we can write

$$19 \quad \mathcal{L}_n \Phi_n(x) - \Phi_n(x) = \sum_{\sigma y=x} F(y) \left[ \lambda_n(y)^{-1} \Phi_n(y) - \Phi_n(x) \right], \quad \forall n \in \mathbf{N}$$

20 and

$$21 \quad \mathcal{L}_\infty \Phi_\infty(x) - \Phi_\infty(x) = \sum_{\sigma y=x} F(y) \left[ \lambda_\infty(y)^{-1} \Phi_\infty(y) - \Phi_\infty(x) \right].$$

22 It follows from the definition of  $\lambda_\infty$  and  $\Phi_\infty$  that

$$23 \quad \mathcal{L}_\infty \Phi_\infty(x) - \Phi_\infty(x) = \lim_\omega \left[ \mathcal{L}_n \Phi_n(x) - \Phi_n(x) \right] = 0,$$

24 which completes the proof of our claim.

25 **Almost invariant vector for  $\lambda_n$ .** The previous discussion shows that 1 is an eigenvalue of  
 26  $\mathcal{L}_\infty : H_\alpha^\infty(\Sigma, E_\infty) \rightarrow H_\alpha^\infty(\Sigma, E_\infty)$ . It follows from [Proposition A.14](#) that the limit representation  
 27  $\lambda_\infty : G \rightarrow \text{Isom}(E_\infty)$  restricted to the finite index subgroup  $G_0$  admits a non-zero invariant  $\phi_\infty$ .  
 28 Such a vector can be written  $\phi_\infty = \lim_\omega \phi_n$ , where  $(\phi_n) \in \Pi_\omega E_n$ . Since  $\phi_\infty$  is non zero, we can  
 29 assume without loss of generality that  $\|\phi_n\| = 1$ , for every  $n \in \mathbf{N}$ . Since  $S_0$  is contained in  $G_0$ , for  
 30 every  $g_0 \in S_0$ , we have

$$31 \quad \lim_\omega [\lambda_n(g_0)\phi_n] = \lambda_\infty(g_0)\phi_\infty = \phi_\infty = \lim_\omega \phi_n.$$

32 The set  $S_0$  being finite, the vector  $\phi_n$  is an  $(S_0, \varepsilon)$ -invariant vector (with respect to  $\lambda_n$ )  $\omega$ -as. This  
 33 contradicts our initial assumption and completes the proof of the proposition.  $\square$

**Corollary A.21.** *We assume that the extension of  $(\Sigma, \sigma)$  by  $\theta$  has the visibility property. There exists a finite index subgroup  $G_0$  of  $G$  with the following property. Let  $\mathcal{E}$  be a uniformly convex collection of Banach spaces. For every finite subset  $S_0$  of  $G_0$ , for every  $\varepsilon \in \mathbf{R}_+^*$  there exists  $\eta \in \mathbf{R}_+^*$  such that the following holds. Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a representation of  $G$  into a Banach space  $(E, \|\cdot\|)$  of  $\mathcal{E}$ . If the spectral radius  $\rho_\lambda$  of the twisted transfer operator  $\mathcal{L}_\lambda$  satisfies  $\rho_\lambda > 1 - \eta$ , then there exists  $\beta \in \mathbf{R}$  such that the representation  $e^{i\beta}\lambda$  admits an  $(S_0, \varepsilon)$ -invariant vector.*

*Proof.* Let  $\lambda: G \rightarrow \text{Isom}(E)$  be a representation of  $G$  into a Banach space  $(E, \|\cdot\|)$ . Let  $\eta > 0$ . We claim that if  $\rho_\lambda > 1 - \eta$ , then there exists  $\beta \in \mathbf{R}$ , such that the operator  $e^{i\beta}\mathcal{L}_\lambda$  admits an  $\eta$ -invariant vector. Since  $\rho_\lambda$  is the spectral radius of  $\mathcal{L}_\lambda$ , there exists  $\beta \in \mathbf{R}$ , such that  $\rho_\lambda e^{-i\beta}$  is a point in the boundary of  $\text{Spec}(\mathcal{L}_\lambda)$ . According to [11, Proposition 6.7], there exists  $\Phi \in H_\alpha^\infty(\Sigma, E)$  such that

$$\|\mathcal{L}_\lambda \Phi - \rho_\lambda e^{-i\beta} \Phi\|_{\infty, \alpha} < (\rho_\lambda - 1 + \eta) \|\Phi\|_{\infty, \alpha}. \quad (24)$$

Combined with the triangle inequality, it yields

$$\begin{aligned} \|e^{i\beta}\mathcal{L}_\lambda \Phi - \Phi\|_{\infty, \alpha} &\leq \|\mathcal{L}_\lambda \Phi - \rho_\lambda e^{-i\beta} \Phi\|_{\infty, \alpha} + \|\rho_\lambda \Phi - \Phi\|_{\infty, \alpha} \\ &< (\rho_\lambda - 1 + \eta) \|\Phi\|_{\infty, \alpha} + (1 - \rho_\lambda) \|\Phi\|_{\infty, \alpha}. \end{aligned}$$

Hence  $\|e^{i\beta}\mathcal{L}_\lambda \Phi - \Phi\|_{\infty, \alpha} < \eta \|\Phi\|_{\infty, \alpha}$ , which completes the proof of our claim. Observe that the operator  $e^{i\beta}\mathcal{L}_\lambda$  can be seen as the twisted transfer operator  $\mathcal{L}_{\lambda'}$  associated to the representation  $\lambda': G \rightarrow \text{Isom}(E)$  defined by  $\lambda'(g) = e^{i\beta}\lambda(g)$ . The corollary is now a direct consequence of Proposition A.18  $\square$

## A.6 Amenability and Kazhdan property (T)

In this section we focus on representations induced by a group actions. Let  $Y$  be a set. Let  $\mathcal{H} = \ell^2(Y)$  be the set of functions  $\phi: Y \rightarrow \mathbf{C}$  which are square summable. It carries a natural structure of Hilbert space. A vector  $\phi \in \mathcal{H}$  is *non-negative*, if  $\phi(y) \in \mathbf{R}_+$  for every  $y \in Y$ . Given any vector  $\phi \in \mathcal{H}$ , we defined its *modulus* to be the vector  $|\phi| \in \mathcal{H}$  defined by  $|\phi|(y) = |\phi(y)|$  for every  $y \in Y$ . Observe that  $\| |\phi| \| = \|\phi\|$ .

Let  $G$  be a group acting on  $Y$ . The action of  $G$  induces a unitary representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  defined as follows: for every  $g \in G$ , for every  $\phi \in \mathcal{H}$ ,

$$[\lambda(g)\phi](y) = \phi(g^{-1}y), \quad \forall y \in Y.$$

Observe that for every  $\phi \in \mathcal{H}$ , for every  $g \in G$  we have  $|\lambda(g)\phi| = \lambda(g)|\phi|$ .

**Lemma A.22.** *Let  $Y$  be a metric space endowed with an action of  $G$ . Let  $\mathcal{H} = \ell^2(Y)$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the unitary representation induced by the action of  $G$ . Let  $\beta \in [0, 2\pi)$ . Let  $S$  be a finite subset of  $G$  and  $\varepsilon \in \mathbf{R}_+^*$ . If  $\phi \in \mathcal{H}$  is  $(S, \varepsilon)$ -invariant with respect to  $e^{i\beta}\lambda$ , then  $|\phi|$  is non-negative and  $(S, \varepsilon)$ -invariant with respect to  $\lambda$ .*

*Proof.* Given any two vectors  $\phi_1, \phi_2 \in \mathcal{H}$ , one checks easily that their modulus satisfies

$$\| |\phi_1| - |\phi_2| \| \leq \|\phi_1 - \phi_2\|.$$

1 Let  $\phi \in \mathcal{H}$ , be an  $(S, \varepsilon)$ -invariant with respect to  $e^{i\beta} \lambda$ . Combining our various observations on  
 2 modulus vector, we get that for every  $g \in S$ ,

$$3 \quad \|\lambda(g) |\phi| - |\phi|\| = \| |e^{i\beta} \lambda(g) \phi| - |\phi| \| \leq \|e^{i\beta} \lambda(g) \phi - \phi\| \leq \varepsilon \|\phi\| = \varepsilon \| |\phi| \|.$$

4 Hence  $|\phi|$  is a non-negative  $(S, \varepsilon)$ -invariant with respect to  $\lambda$  □

5 The main result of this section is the following statement.

6 **Theorem A.23.** *Let  $(\Sigma, \sigma)$  be an irreducible subshift of finite type. Let  $F: \Sigma \rightarrow \mathbf{R}_+^*$  be a potential  
 7 with  $\alpha$ -bounded Hölder variations for some  $\alpha \in \mathbf{R}_+^*$ . Let  $\mathcal{L}$  be the corresponding transfer operator  
 8 and  $\rho$  its spectral radius. Let  $G$  be a finitely generated group and  $\theta: \Sigma \rightarrow G$  be a locally constant  
 9 map. We assume that the corresponding extension  $(\Sigma_\theta, \sigma_\theta)$  has the visibility property. For every  
 10 finite subset  $S$  of  $G$  and every  $\varepsilon \in \mathbf{R}_+^*$  there exists  $\eta \in \mathbf{R}_+^*$  with the following property.*

11 *Let  $Y$  be a set endowed with an action of  $G$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the induced unitary rep-  
 12 resentation, where  $\mathcal{H} = \ell^2(Y)$ . Let  $\rho_\lambda$  be the spectral radius of the twisted transfer operator  
 13  $\mathcal{L}_\lambda: H_\alpha^\infty(\Sigma, \mathcal{H}) \rightarrow H_\alpha^\infty(\Sigma, \mathcal{H})$ . If  $\rho_\lambda > (1 - \eta)\rho$ , then the representation  $\lambda$  admits an  $(S, \varepsilon)$  in-  
 14 variant vector.*

15 *Proof of Theorem A.23.* The strategy of the proof is the following. First we renormalize the po-  
 16 tential so that we can assume that the spectral radius of  $\mathcal{L}$  is  $\rho = 1$  and  $\mathbb{1}$  is an invariant vector of  
 17  $\mathcal{L}$ . Applying Corollary A.21 we get a finite index subgroup  $G_0$  of  $G$  such that the representation  $\lambda$   
 18 when restricted to  $G_0$  admits a certain almost invariant vector  $\phi$ . We finally take advantage of the  
 19 structure of  $\ell^2(Y)$  to average the orbit of  $\phi$ , and thus get an almost invariant vector with respect  
 20 to  $\lambda$ .

21 **Renormalization of the potential.** We start with a reduction argument: we claim that without  
 22 loss of generality we can assume that  $\mathcal{L}$  has spectral radius 1 and fixes  $\mathbb{1}$ . Assume indeed that  
 23 the result has been proved in this context and let us explain how to deduce the general case. Let  
 24  $h \in H_\alpha^\infty(\Sigma, \mathbf{C})$  be the positive eigenvector of  $\mathcal{L}$  given by the Ruelle Perron-Frobenius Theorem  
 25 (Theorem A.6). Following the strategy of Section A.4 we define a new potential  $F': \Sigma \rightarrow \mathbf{R}_+^*$  by

$$26 \quad F'(x) = \frac{1}{\rho} \cdot \frac{h(x)}{h \circ \sigma(x)} F(x).$$

27 We write  $\mathcal{L}'$  for the corresponding transfer operator. As we observed  $\mathcal{L}'$  has spectral radius 1 and  
 28 fixes  $\mathbb{1}$ . It follows from our assumption that we can apply Theorem A.23 to this operator.

29 Let  $S$  be a finite subset of  $G$  and  $\varepsilon \in \mathbf{R}_+^*$ . Let  $\eta \in \mathbf{R}_+^*$  be the parameter given by Theorem A.23  
 30 (with the additional assumption that the transfer operator has spectral radius 1 and  $\mathbb{1}$  as an  
 31 eigenvector) applied to the potential  $F'$ . Suppose now that  $Y$  is a space endowed with an action of  
 32  $G$  and denote by  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  the induced unitary representation, where  $\mathcal{H} = \ell^2(Y)$ . Assume that  
 33 the spectral radius of  $\mathcal{L}_\lambda$  satisfies  $\rho_\lambda > (1 - \eta)\rho$ . It follows from the discussion of Section A.4 that  
 34 the spectral radius  $\rho'_\lambda$  of  $\mathcal{L}'_\lambda$  satisfies  $\rho'_\lambda = \rho_\lambda / \rho$ . In particular  $\rho'_\lambda > 1 - \eta$ . Theorem A.23 applied  
 35 to the potential  $F'$  tells us that  $\lambda$  admits an  $(S, \varepsilon)$  invariant vector, which completes the proof of  
 36 our claim.

1 **Finite index subgroup with almost invariant vectors.** From now on we assume that  $\mathcal{L}$  has  
 2 spectral radius 1 and fixes  $\mathbb{1}$ . Let  $S$  be a finite subset of  $G$  and  $\varepsilon \in \mathbf{R}_+^*$ . We denote by  $G_0$  the finite  
 3 index subgroup of  $G$  given by [Corollary A.21](#). We denote by  $u_1, \dots, u_m$  a set of representatives of  
 4  $G/G_0$ . For every  $g \in S$ , there exists a permutation  $\sigma_g: \llbracket 1, m \rrbracket \rightarrow \llbracket 1, m \rrbracket$  such that for all  $i \in \llbracket 1, m \rrbracket$ ,  
 5 we have

$$6 \quad u_{\sigma_g(i)}^{-1} g u_i \in G_0.$$

7 We now define a finite subset  $S_0$  of  $G_0$  as

$$8 \quad S_0 = \left\{ u_{\sigma_g(i)}^{-1} g u_i \mid g \in S, i \in \llbracket 1, m \rrbracket \right\}.$$

9 Note that the set of all Hilbert spaces is a uniformly convex collection of Banach spaces. According  
 10 to [Corollary A.21](#), there exists  $\eta \in \mathbf{R}_+^*$  with the following property. Let  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be a unitary  
 11 representation of  $G$ . If the spectral radius of the twisted transfer operator  $\mathcal{L}_\lambda$  is larger than  $1 - \eta$ ,  
 12 then there exists  $\beta \in \mathbf{R}$  such that the representation  $e^{i\beta} \lambda$  admits an  $(S_0, \varepsilon/\sqrt{m})$ -invariant vector.

13 **Representation induced by an action.** Let  $Y$  be a set endowed with an action of  $G$  and  
 14  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the induced unitary representation, where  $\mathcal{H} = \ell^2(Y)$ . Assume that the spectral  
 15 radius  $\rho_\lambda$  of the twisted transfer operator  $\mathcal{L}_\lambda$  is at least  $1 - \eta$ . According to the very definition  
 16 of  $\eta$ , there exists  $\beta \in \mathbf{R}$ , such that the representation  $e^{i\beta} \lambda$  admits an  $(S_0, \varepsilon/\sqrt{m})$ -invariant vector  
 17  $\phi$ . It follows from [Lemma A.22](#) that  $|\phi|$  is an  $(S_0, \varepsilon/\sqrt{m})$ -invariant vector with respect to the  
 18 representation  $\lambda$ . We now let

$$19 \quad \bar{\phi} = \frac{1}{m} \sum_{i=1}^m \lambda_n(u_i) |\phi|.$$

20 Let  $g \in S$ . The computation yields

$$21 \quad m\lambda(g)\bar{\phi} = \sum_{i=1}^m \lambda(gu_i) |\phi| = \sum_{i=1}^m \lambda(u_{\sigma_g(i)}) \lambda(u_{\sigma_g(i)}^{-1} g u_i) |\phi|.$$

22 On the other hand, reindexing the sum defining  $\bar{\phi}$  gives

$$23 \quad m\bar{\phi} = \sum_{i=1}^m \lambda(u_{\sigma_g(i)}) |\phi|$$

24 Recall that for every  $i \in \llbracket 1, m \rrbracket$ , the element  $u_{\sigma_g(i)}^{-1} g u_i$  belongs to  $S_0$ . Thus the triangle inequality  
 25 yields

$$26 \quad \|\lambda(g)\bar{\phi} - \bar{\phi}\| \leq \frac{1}{m} \sum_{i=1}^m \left\| \lambda(u_{\sigma_g(i)}^{-1} g u_i) |\phi| - |\phi| \right\| < \frac{\varepsilon}{\sqrt{m}}.$$

27 This inequality holds for every  $g \in G$ . Observe that  $\bar{\phi}$  is obtained by averaging non-negative vectors  
 28 of  $\mathcal{H}$  all of them having norm 1. It follows that the norm of  $\bar{\phi}$  is bounded below by  $1/\sqrt{m}$ . Hence  
 29  $\bar{\phi}$  is an  $(S, \varepsilon)$ -invariant vector with respect to  $\lambda$ .  $\square$

1 **Amenability.** There are numerous equivalent definition of amenability. The one that is the most  
2 adapted four our purpose can be formulated in terms of regular representation.

3 **Definition A.24.** The action of  $G$  on  $Y$  is amenable if and only if the representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$   
4 admits almost invariant vectors. The group  $G$  is *amenable* if its action on itself is amenable.

5 **Theorem A.25** (Amenability criterion). *Let  $(\Sigma, \sigma)$  be an irreducible subshift of finite type. Let*  
6  *$F: \Sigma \rightarrow \mathbf{R}_+^*$  be a potential with  $\alpha$ -bounded Hölder variations for some  $\alpha \in \mathbf{R}_+^*$ . Let  $\mathcal{L}$  be the*  
7 *corresponding transfer operator and  $\rho$  its spectral radius. Let  $G$  be a finitely generated group and*  
8  *$\theta: \Sigma \rightarrow G$  be a locally constant map. We assume that the corresponding extension  $(\Sigma_\theta, \sigma_\theta)$  has the*  
9 *visibility property. Let  $Y$  be a set endowed with an action of  $G$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the induced*  
10 *unitary representation, where  $\mathcal{H} = \ell^2(Y)$ . Let  $\rho_\lambda$  be the spectral radius of the twisted transfer*  
11 *operator  $\mathcal{L}_\lambda: H_\alpha^\infty(\Sigma, \mathcal{H}) \rightarrow H_\alpha^\infty(\Sigma, \mathcal{H})$  defined by*

$$12 \quad \mathcal{L}_\lambda \Phi(x) = \sum_{\sigma y=x} F(y) \lambda(y)^{-1} \Phi(y).$$

13 *The following statements are equivalent.*

14 (i) *The action of  $G$  on  $Y$  is amenable.*

15 (ii)  *$\rho$  belongs to  $\text{Spec}(\mathcal{L}_\lambda)$ .*

16 (iii)  *$\rho_\lambda = \rho$ .*

17 *Proof.* Reasoning as in the beginning of the proof of [Theorem A.23](#) we observe that without loss  
18 of generality we can assume that  $\mathcal{L}$  has spectral radius 1 and fixes  $\mathbf{1}$ . We start with (ii) $\Rightarrow$ (iii)  
19 Recall that  $\rho_\lambda \leq 1$  ([Corollary A.9](#)). Hence if 1 belongs to  $\text{Spec}(\mathcal{L}_\lambda)$ , then  $\rho_\lambda = 1$ . We now focus  
20 on (iii) $\Rightarrow$ (i). Assume that  $\rho_\lambda = 1$ . It follows from [Theorem A.23](#) that  $\lambda$  almost admits invariants  
21 vectors. Hence the action of  $G$  on  $Y$  is amenable. We are left to prove (i) $\Rightarrow$ (ii). Assume that the  
22 action of  $G$  on  $Y$  is amenable. According to [Corollary A.9](#) it is sufficient to prove that  $\rho_\lambda \geq 1$ . Let  
23  $n \in \mathbf{N}$ . Let  $\varepsilon > 0$ . Since  $\theta: \Sigma \rightarrow G$  is locally constant, the set

$$24 \quad S = \{\theta_n(x) \mid x \in \Sigma\}$$

25 is finite. Since the action of  $G$  is amenable, there exists an  $(S, \varepsilon)$ -invariant vector  $\phi \in \mathcal{H} \setminus \{0\}$ .  
26 Without loss of generality we can assume that  $\|\phi\| = 1$ . We define a map  $\Phi: \Sigma \rightarrow \mathcal{H}$  by letting  
27  $\Phi(x) = \phi$ , for every  $x \in \Sigma$ . Obviously  $\|\Phi\|_\infty = 1$  and  $\Delta_\alpha(\Phi) = 0$ , hence  $\Phi$  belongs to  $H_\alpha^\infty(\Sigma, \mathcal{H})$ .  
28 Using the fact that  $\mathcal{L}\mathbf{1} = \mathbf{1}$ , we can write for every  $x \in \Sigma$ ,

$$29 \quad \|\mathcal{L}_\lambda^n \Phi(x) - \Phi(x)\| \leq \sum_{\sigma^n y=x} F_n(y) \|\lambda_n(y)^{-1} \Phi(y) - \Phi(x)\| = \sum_{\sigma^n y=x} F_n(y) \|\lambda_n(y)\phi - \phi\| < \varepsilon.$$

30 This proves that  $\|\mathcal{L}_\lambda^n \Phi - \Phi\|_\infty < \varepsilon$ . In particular, we get

$$31 \quad \|\mathcal{L}_\lambda^n \Phi\|_{\infty, \alpha} \geq \|\mathcal{L}_\lambda^n \Phi\|_\infty > \|\Phi\|_\infty - \varepsilon \geq 1 - \varepsilon.$$

32 Recall that  $\|\Phi\|_{\infty, \alpha} = 1$ . Thus we have proved that the norm of  $\mathcal{L}_\lambda^n$  – see as an operator of  $H_\alpha^\infty(\Sigma, \mathcal{H})$   
33 – is larger than  $1 - \varepsilon$ . This holds for every  $\varepsilon > 0$ . Hence for every  $n \in \mathbf{N}$ , we have

$$34 \quad \|\mathcal{L}_\lambda^n\|_{\infty, \alpha} \geq 1.$$

35 Consequently

$$36 \quad \rho_\lambda = \lim_{n \rightarrow \infty} \sqrt[n]{\|\mathcal{L}_\lambda^n\|_{\infty, \alpha}} \geq 1. \quad \square$$

1 **Kazhdan property (T).** Let us recall first the definition of property (T).

2 **Definition A.26.** A discrete group  $G$  has *Kazhdan property (T)*, if there exists a finite subset  $S$   
 3 of  $G$  and  $\varepsilon \in \mathbf{R}_+^*$  with the following property. Any unitary representation  $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$  into a  
 4 Hilbert space which admits  $(S, \varepsilon)$ -invariant vectors has a non-zero invariant vector. Such a pair  
 5  $(S, \varepsilon)$  is called a Kazhdan pair.

6 Let us also recall the following useful statement.

7 **Lemma A.27.** *Assume that the action of  $G$  on  $Y$  is transitive. The set  $Y$  is finite if and only if*  
 8 *the representation  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  admits a non-zero invariant vector.*

9 **Theorem A.28.** *Let  $(\Sigma, \sigma)$  be an irreducible subshift of finite type. Let  $F: \Sigma \rightarrow \mathbf{R}_+^*$  be a potential*  
 10 *with  $\alpha$ -bounded Hölder variations for some  $\alpha \in \mathbf{R}_+^*$ . Let  $\mathcal{L}$  be the corresponding transfer operator*  
 11 *and  $\rho$  its spectral radius. Let  $G$  be a finitely generated group with Kazhdan property (T) and*  
 12  *$\theta: \Sigma \rightarrow G$  be a locally constant map. We assume that the corresponding extension  $(\Sigma_\theta, \sigma_\theta)$  has the*  
 13 *visibility property. There exists  $\eta > 0$  with the following property.*

14 *Let  $Y$  be an infinite set endowed with an transitive action of  $G$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the induced*  
 15 *unitary representation, where  $\mathcal{H} = \ell^2(Y)$ . Let  $\rho_\lambda$  be the spectral radius of the twisted transfer*  
 16 *operator  $\mathcal{L}_\lambda: H_\alpha^\infty(\Sigma, \mathcal{H}) \rightarrow H_\alpha^\infty(\Sigma, \mathcal{H})$  defined by*

$$17 \quad \mathcal{L}_\lambda \Phi(x) = \sum_{\sigma y=x} F(y) \lambda(y)^{-1} \Phi(y).$$

18 *Then  $\rho_\lambda \leq (1 - \eta)\rho$ .*

19 *Proof.* Let  $(S, \varepsilon)$  be a Kazhdan pair of  $G$ . Let  $\eta > 0$  be the constant given by [Theorem A.23](#).  
 20 Let  $Y$  be a set endowed with an transitive action of  $G$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$  be the induced unitary  
 21 representation, where  $\mathcal{H} = \ell^2(Y)$ . Let  $\mathcal{L}_\lambda$  be the corresponding twisted transfer operator. Assume  
 22 that  $\rho_\lambda > (1 - \eta)\rho$ . It follows from [Theorem A.23](#) that  $\lambda$  has an  $(S, \varepsilon)$ -invariant vector. Since  $(S, \varepsilon)$   
 23 is a Kazhdan pair, it follows that  $\lambda$  has a non-zero invariant vector. However the action of  $G$  on  $Y$   
 24 is transitive. Hence  $Y$  is finite ([Lemma A.27](#)).  $\square$

## 25 B Roblin's theorem

26 In this appendix, we provide a proof of Roblin's Theorem. Note that the statement below does not  
 27 require  $H$  to be a normal subgroup. The proof is probably well-known from the specialists in the  
 28 field, however we did not find it in the literature. It relies on a rather simple counting argument in  
 29 a hyperbolic space.

30 **Theorem B.1** (compare with Roblin [[34](#), Théorème 2.2.2]). *Let  $G$  be a group acting properly*  
 31 *co-compactly on a hyperbolic space  $X$ . Let  $H$  be a subgroup of  $G$ . We denote by  $\omega_G$  and  $\omega_H$  the*  
 32 *exponential growth rates of  $G$  and  $H$  acting on  $X$ . If  $H$  is co-amenable in  $G$ , then  $\omega_H = \omega_G$ .*

33 Let  $G$  be a group acting properly co-compactly on a hyperbolic space. Let  $\omega_G$  be the exponential  
 34 growth rate of  $G$  acting on  $X$ . We fix a base point  $o \in X$ . Given  $r \in \mathbf{R}_+$  we define the *ball* of  
 35 radius  $r$  to be

$$36 \quad B(r) = \{g \in G \mid d(go, o) \leq r\}.$$

1 Coornaert [12] proved that there exists  $C_1 \in \mathbf{R}_+^*$  such that for every  $r \in \mathbf{R}_+$ ,

$$2 \quad e^{\omega_G r} \leq |B(r)| \leq C_1 e^{\omega_G r}. \quad (25)$$

3 Let  $\delta \in \mathbf{R}_+$  be the hyperbolicity constant of  $X$ . Up to increasing the value of  $\delta$  we can always  
4 assume that the following holds:

5 (i) The diameter of  $X/G$  is at most  $\delta$ . In particular, for every  $x, y \in X$ , there exists  $g \in G$  such  
6 that  $d(x, gy) \leq \delta$ .

7 (ii)  $1 - C_1 e^{-\omega_G \delta} > 0$ .

8 For every  $r \in \mathbf{R}_+$ , we denote by  $S(r) = B(r) \setminus B(r - \delta)$  the *sphere* of radius  $r$ .

9 **Lemma B.2.** *There exists  $C_2 \in \mathbf{R}_+^*$  with the following property. Given  $\ell, r \in \mathbf{R}_+$  and  $x \in X$ , we  
10 denote by  $U$  the set of elements  $g \in B(\ell)$  such that  $\langle go, x \rangle_o \geq r$ . The cardinality of  $U$  is bounded  
11 above by*

$$12 \quad |U| \leq C_2 e^{\omega_G(\ell-r)}.$$

13 *Proof.* We fix a geodesic  $[o, x]$  from  $o$  to  $x$  and write  $y$  for the point of  $[o, x]$  at distance  $r$  from  $o$ .  
14 According to our choice of  $\delta$  that there exists  $h \in G$  such that  $d(y, ho) \leq \delta$ . Note that  $d(ho, o) \geq r - \delta$ .  
15 Let  $g \in U$ . It follows from the four point inequality (4) that

$$16 \quad \langle go, o \rangle_{ho} \leq \langle go, o \rangle_y + \delta \leq 2\delta$$

17 Consequently

$$18 \quad d(h^{-1}go, o) = d(go, ho) = d(go, o) - d(ho, o) + 2 \langle go, o \rangle_{ho} \leq \ell - r + 5\delta.$$

19 Thus  $h^{-1}U$  is contained in  $B(\ell - r + 5\delta)$  and the result follows from (25).  $\square$

20 Let  $\ell \in \mathbf{R}_+$ . We denote by  $\mu_\ell$  the probability measure on  $G$  which is uniformly distributed on  
21  $S(\ell)$ . It follows from (25) that for every  $g \in S(\ell)$  we have

$$22 \quad \frac{1}{C_1} e^{-\omega_G \ell} \leq \mu_\ell(g) \leq \frac{1}{C_3} e^{-\omega_G \ell}, \quad (26)$$

23 where  $C_3 = 1 - C_1 e^{-\omega_G \delta}$ . Our first task is to provide an estimate for the  $n$ -th convolution product  
24 of  $\mu_\ell$ . Later we will let  $\ell$  tends to infinity. Thus we will be particularly careful to control these  
25 estimates in terms of  $\ell$ . More precisely we are going to prove the following statement.

26 **Proposition B.3.** *There exists  $D \in \mathbf{R}_+^*$  such that for every  $\ell \in \mathbf{R}_+$ , for every  $n \in \mathbf{N}$ , for every  
27  $g \in G$ , we have*

$$28 \quad \mu_\ell^{*n}(g) \leq D^n \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( -\omega_G \frac{n\ell + d(go, o)}{2} \right).$$

29 In order to prove this proposition, we introduce the following sets that will allow us to track the  
30 orbits of the random walk. For every  $\ell \in \mathbf{R}_+$ , for every  $n \in \mathbf{N}$ , for every  $g \in G$ , we let

$$31 \quad \mathcal{O}_\ell(g, n) = \{(u_1, \dots, u_n) \in S(\ell)^n \mid u_1 \cdots u_n = g\}.$$

32 Note that if  $d(go, o) > n\ell$ , then  $\mathcal{O}_\ell(g, n)$  is empty. We adopt the convention that a product of  
33 elements of  $G$  indexed by the empty set is trivial. It follows that  $\mathcal{O}_\ell(g, 0)$  is empty if  $g$  is non trivial  
34 and reduced to a single element (the empty tuple) if  $g = 1$ .



**Lemma B.4.** *There exists  $D_0 \in \mathbf{R}_+^*$  such that for every  $\ell \in \mathbf{R}_+$ , for every  $n \in \mathbf{N}$ , for every  $g \in G$ , we have*

$$|\mathcal{O}_\ell(g, n)| \leq D_0^n \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( \omega_G \frac{n\ell - d(go, o)}{2} \right).$$

*Proof.* Let  $C_2$  be the constant given by [Lemma B.2](#). We let

$$D_0 = C_2 e^{2\omega_G \delta}.$$

Let  $\ell \in \mathbf{R}_+$ . We are going to prove the result by induction on  $n$ . If  $n = 0$ , it follows from our convention that for every  $g \in G$ , the set  $\mathcal{O}_\ell(g, 0)$  contains at most 1 element, hence the result.

Assume now that the statement holds for some  $n \in \mathbf{N}$ . Let  $g \in G$ . For every element  $u = (u_1, \dots, u_{n+1})$  of  $\mathcal{O}_\ell(g, n+1)$  we let  $g_u = u_1 \cdots u_n = g u_{n+1}^{-1}$  (according to our convention  $g_u$  is trivial if  $n = 0$ ). For every  $k \in \mathbf{N}$  such that  $k\delta \leq \ell$ , we denote by  $P_k$  the set of elements  $u \in \mathcal{O}_\ell(g, n+1)$  such that

$$k\delta \leq \langle g_u o, o \rangle_{go} < (k+1)\delta.$$

Note that if  $u = (u_1, \dots, u_{n+1})$  is an element of  $\mathcal{O}_\ell(g, n+1)$ , then  $\langle g_u o, o \rangle_{go} \leq d(u_{n+1} o, o) \leq \ell$ . Hence the collection  $(P_k)$  forms a partition of  $\mathcal{O}_\ell(g, n+1)$ . We are now going to estimate the cardinality of each of these sets.

Let  $k \in \mathbf{N}$  such that  $k\delta \leq \ell$ . We write  $U_k$  for the image of  $P_k$  by the projection  $P_k \rightarrow S(\ell)$  sending  $(u_1, \dots, u_{n+1})$  to  $u_{n+1}$ . It follows from [Lemma B.2](#) that

$$|U_k| \leq C_2 e^{\omega_G(\ell - k\delta)} \leq D_0 e^{-\omega_G(k+2)\delta} e^{\omega_G \ell}.$$

Let  $u_{n+1}$  be an element of  $U_k$  and  $u = (u_1, \dots, u_{n+1})$  a pre-image of  $u_{n+1}$  in  $P_k$ . By definition  $(u_1, \dots, u_n)$  is an element of  $\mathcal{O}_\ell(g u_{n+1}^{-1}, n)$ , whose cardinality can be bounded from above using the induction hypotheses. It follows that

$$\begin{aligned} |P_k| &\leq \sum_{u_{n+1} \in U_k} |\mathcal{O}_\ell(g u_{n+1}^{-1}, n)| \\ &\leq \sum_{u_{n+1} \in U_k} D_0^n \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( \omega_G \frac{n\ell - d(g u_{n+1}^{-1} o, o)}{2} \right). \end{aligned} \tag{27}$$

Observe that for any  $u_{n+1} \in U_k$  we have  $\langle g u_{n+1}^{-1} o, o \rangle_{go} < (k+1)\delta$ . Hence

$$d(g u_{n+1}^{-1} o, o) \geq d(go, o) + d(u_{n+1} o, o) - 2 \langle g u_{n+1}^{-1} o, o \rangle_{go} \geq d(go, o) + \ell - 2(k+2)\delta.$$

Consequently (27) becomes

$$\begin{aligned} |P_k| &\leq \sum_{u_{n+1} \in U_k} e^{\omega_G(k+2)\delta} D_0^n \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( \omega_G \frac{(n-1)\ell - d(go, o)}{2} \right) \\ &\leq |U_k| e^{\omega_G(k+2)\delta} D_0^n \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( \omega_G \frac{(n-1)\ell - d(go, o)}{2} \right). \end{aligned}$$

We now use the above estimate of  $|U_k|$  to get

$$|P_k| \leq D_0^{n+1} \left( \frac{\ell}{\delta} + 1 \right)^n \exp \left( \omega_G \frac{(n+1)\ell - d(go, o)}{2} \right).$$

1 Note that this estimate does not depend on  $k$ . Moreover there are at most  $\ell/\delta + 1$  integer  $k \in \mathbf{N}$   
 2 such that  $k\delta \leq \ell$ . Since  $(P_k)$  forms a partition of  $\mathcal{O}_\ell(g, n+1)$  we obtain

$$3 \quad |\mathcal{O}_\ell(g, n+1)| \leq \sum_{k\delta \leq \ell} |P_k| \leq D_0^{n+1} \left(\frac{\ell}{\delta} + 1\right)^{n+1} \exp\left(\omega_G \frac{(n+1)\ell - d(go, o)}{2}\right).$$

4 Hence the statement holds for  $n+1$ , which completes the proof of the proposition.  $\square$

5 *Proof of Proposition B.3.* We denote by  $C_3$  and  $D_0$  the constants given by (26) and Lemma B.4  
 6 respectively and let  $D = D_0/C_3$ . Let  $\ell \in \mathbf{R}_+$ . Let  $n \in \mathbf{N}$  and  $g \in G$ . It follows from the definition  
 7 of the convolution that

$$8 \quad \mu_\ell^{*n}(g) = \sum_{(u_1, \dots, u_n) \in \mathcal{O}_\ell(g, n)} \mu_\ell(u_1) \cdots \mu_\ell(u_n).$$

9 Combining (26) and Lemma B.4, the previous equality becomes

$$\begin{aligned} \mu_\ell^{*n}(g) &\leq \left(\frac{D_0}{C_3}\right)^n \left(\frac{\ell}{\delta} + 1\right)^n \exp\left(-\omega_G \frac{n\ell + d(go, o)}{2}\right) \\ &\leq D^n \left(\frac{\ell}{\delta} + 1\right)^n \exp\left(-\omega_G \frac{n\ell + d(go, o)}{2}\right). \end{aligned} \quad \square$$

10 We now fix a subgroup  $H$  of  $G$  and write  $\omega_H$  for the exponential growth rate of  $H$  acting on  
 11  $X$ . We denote by  $Y$  the set of left  $H$ -cosets in  $G$ . The group  $G$  acts on  $Y$  by right translations.  
 12 We write  $\mathcal{H} = \ell^2(Y)$  for the set of square summable functions from  $Y$  to  $\mathbf{C}$  and  $\lambda: G \rightarrow \mathcal{U}(\mathcal{H})$   
 13 for the regular representation of  $G$  relative to  $H$ . Given  $\ell \in \mathbf{R}_+$ , we consider the random walk  
 14 on  $Y$  associated to the probability measure  $\mu_\ell$ . Said differently for every  $y \in Y$  and  $g \in G$  the  
 15 probability of going from  $y$  to  $y \cdot g$  is  $\mu_\ell(g)$ . Let  $y_0$  be the point of  $Y$  corresponding to  $H$ . Note that  
 16 its stabilizer is exactly  $H$ . Hence the probability  $p_\ell(n)$  that after  $n$ -step, the random walk starting  
 17 to  $y_0$  goes back to  $y_0$  is exactly

$$18 \quad p_\ell(n) = \mu_\ell^{*n}(H).$$

19 We associate to this random walk a Markov operator  $M_\ell$  on  $\mathcal{H}$ .

$$20 \quad M_\ell \phi = \sum_{g \in G} \mu_\ell(g) \lambda(g) \phi, \quad \forall \phi \in \mathcal{H}.$$

21 Since  $\mu_\ell$  is symmetric,  $M_\ell$  is a self-adjoint operator. It follows that its spectral radius  $\rho_\ell$  can be  
 22 computed as follows – see for instance [39, Lemma 10.1].

$$23 \quad \rho_\ell = \limsup_{n \rightarrow \infty} \sqrt[n]{p_\ell(n)} = \limsup_{n \rightarrow \infty} \sqrt[n]{\mu_\ell^{*n}(H)}.$$

24 The next proposition relates the spectral radius  $\rho_\ell$  to the critical exponents  $\omega_H$  and  $\omega_G$ .

25 **Proposition B.5.** *The growth rates of  $H$  and  $G$  acting on  $X$  satisfy the following inequality*

$$26 \quad \ln \rho_\infty \leq \max \left\{ -\frac{1}{2} \omega_G, \omega_H - \omega_G \right\},$$

27 where  $\rho_\infty = \limsup_{\ell \rightarrow \infty} \sqrt[\ell]{\rho_\ell}$ .

1 *Proof.* We fix  $\varepsilon \in \mathbf{R}_+^*$  such that if  $\omega_H < \omega_G/2$ , then  $\omega_H + \varepsilon < \omega_G/2$ . It follows from the definition  
2 of exponential growth rate that there exists  $A \in \mathbf{R}_+^*$  such that for every  $r \in \mathbf{R}_+$ ,

$$3 \quad |H \cap S(r)| \leq |H \cap B(r)| \leq Ae^{(\omega_H + \varepsilon)r}. \quad (28)$$

4 We write  $D$  for the constant given by [Proposition B.3](#). Let  $\ell \in \mathbf{R}_+$  and  $n \in \mathbf{N}$ . Our first task is to  
5 bound  $\mu_\ell^{*n}(H)$  from above. To that end we partition  $H$  according to the length of its elements.

$$6 \quad \mu_\ell^{*n}(H) = \sum_{k \in \mathbf{N}} \sum_{h \in H \cap S(k\delta)} \mu_\ell^{*n}(h).$$

7 Note that if  $n\ell > k\delta$ , then the probabilities  $\mu_\ell^{*n}(h)$  vanish. Using [Proposition B.3](#), we get

$$8 \quad \mu_\ell^{*n}(H) \leq \sum_{k\delta \leq n\ell} |H \cap S(k\delta)| D^n \left(\frac{\ell}{\delta} + 1\right)^n \exp\left(-\omega_G \frac{n\ell + (k-1)\delta}{2}\right).$$

9 Combined with (28) it yields

$$10 \quad \mu_\ell^{*n}(H) \leq Ae^{\frac{1}{2}\omega_G\delta} D^n \left(\frac{\ell}{\delta} + 1\right)^n e^{-\frac{1}{2}\omega_G n\ell} \sum_{k\delta \leq n\ell} e^{(\omega_H + \varepsilon - \frac{1}{2}\omega_G)k\delta}. \quad (29)$$

11 We now distinguish two cases. Assume first that  $\omega_H < \omega_G$ . It follows from our choice of  $\varepsilon$  that  
12  $\omega_H + \varepsilon - \omega_G/2 < 0$ . Hence a (rather brutal !) majoration in (29) gives

$$13 \quad \mu_\ell^{*n}(H) \leq Ae^{\frac{1}{2}\omega_G\delta} D^n \left(\frac{\ell}{\delta} + 1\right)^n \left(\frac{n\ell}{\delta} + 1\right) e^{-\frac{1}{2}\omega_G n\ell}.$$

14 Consequently

$$15 \quad \ln \rho_\ell \leq \ln D + \ln \left(\frac{\ell}{\delta} + 1\right) - \frac{1}{2}\omega_G \ell.$$

16 This inequality holds for every  $\ell \in \mathbf{R}_+$ . Consequently

$$17 \quad \ln \rho_\infty = \limsup_{\ell \rightarrow \infty} \frac{1}{\ell} \ln \rho_\ell \leq -\frac{1}{2}\omega_G,$$

18 which completes the first case. Assume now that  $\omega_H \geq \omega_G/2$ . Computing the sum in (29) we get

$$19 \quad \mu_\ell^{*n}(H) \leq Ae^{\frac{1}{2}\omega_G\delta} D^n \left(\frac{\ell}{\delta} + 1\right)^n e^{-\frac{1}{2}\omega_G n\ell} \frac{e^{(\omega_H + \varepsilon - \frac{1}{2}\omega_G)(n\ell + \delta)} - 1}{e^{(\omega_H + \varepsilon - \frac{1}{2}\omega_G)\delta} - 1}.$$

20 Consequently

$$21 \quad \ln \rho_\ell \leq \ln D + \ln \left(\frac{\ell}{\delta} + 1\right) + (\omega_H + \varepsilon - \omega_G)\ell.$$

22 Since this inequality holds for every  $\ell \in \mathbf{R}_+$ , we get  $\ln \rho_\infty \leq \omega_H + \varepsilon - \omega_G$ . This last inequality  
23 holds for every  $\varepsilon > 0$ , hence the result.  $\square$

24 *Proof of Theorem B.1.* Assume now that  $H$  is co-amenable in  $G$ . According to Kesten's criterion,  
25 the spectral radius of any of the Markov operator  $M_\ell$  is 1. It follows from [Proposition B.5](#) that  
26  $\omega_H \geq \omega_G$ . The other inequality is obvious.  $\square$

**Remark.** The exact same strategy can be used to provide a lower bound for  $\mu_\ell^{*n}$  of the same kind than the one given in [Proposition B.3](#). This leads to the more general version of [Proposition B.5](#)

**Proposition B.6.** *The limit  $\rho_\infty = \lim_{\ell \rightarrow \infty} \sqrt[\ell]{\rho_\ell}$  exists. Moreover*

$$\ln \rho_\infty = \max \left\{ -\frac{1}{2} \omega_G, \omega_H - \omega_G \right\}.$$

## References

- [1] V. Baladi. *Positive transfer operators and decay of correlations*, volume 16 of *Advanced Series in Nonlinear Dynamics*. World Scientific Publishing Co., Inc., River Edge, NJ, 2000.
- [2] W. Ballmann. *Lectures on spaces of nonpositive curvature*, volume 25 of *DMV Seminar*. Birkhäuser Verlag, Basel, Basel, 1995.
- [3] M. E. B. Bekka, P. de la Harpe, and A. Valette. *Kazhdan's property (T)*, volume 11 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2008.
- [4] N. Bourbaki. *Éléments de mathématique. Topologie générale. Chapitres 1 à 4*. Hermann, Paris, 1971.
- [5] R. Brooks. The fundamental group and the spectrum of the Laplacian. *Commentarii Mathematici Helvetici*, 56(4):581–598, 1981.
- [6] R. Brooks. The bottom of the spectrum of a Riemannian covering. *Journal für die Reine und Angewandte Mathematik. [Crelle's Journal]*, 357(357):101–114, 1985.
- [7] M. Burger. Spectre du laplacien, graphes et topologie de Fell. *Commentarii Mathematici Helvetici*, 63(2):226–252, 1988.
- [8] C. Champetier. Petite simplification dans les groupes hyperboliques. *Toulouse. Faculté des Sciences. Annales. Mathématiques. Série 6*, 3(2):161–221, 1994.
- [9] J. M. Cohen. Cogrowth and amenability of discrete groups. *Journal of Functional Analysis*, 48(3):301–309, 1982.
- [10] D. Constantine, J.-F. Lafont, and D. Thompson. The weak specification property for geodesic flows on CAT(-1) spaces. *arXiv.org*, June 2016.
- [11] J. B. Conway. *A course in functional analysis*, volume 96 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, New York, NY, 1985.
- [12] M. Coornaert. Mesures de Patterson-Sullivan sur le bord d'un espace hyperbolique au sens de Gromov. *Pacific Journal of Mathematics*, 159(2):241–270, 1993.
- [13] M. Coornaert, T. Delzant, and A. Papadopoulos. *Géométrie et théorie des groupes*, volume 1441 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1990.
- [14] M. Coornaert and A. Papadopoulos. *Symbolic dynamics and hyperbolic groups*, volume 1539 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1993.

- 
- 1 [15] M. Coornaert and A. Papadopoulos. Horofunctions and symbolic dynamics on Gromov hyper-  
2 bolic groups. *Glasgow Mathematical Journal*, 43(3):425–456, 2001.
- 3 [16] M. Coornaert and A. Papadopoulos. Symbolic coding for the geodesic flow associated to a  
4 word hyperbolic group. *Manuscripta Mathematica*, 109(4):465–492, 2002.
- 5 [17] K. Corlette. Hausdorff dimensions of limit sets. I. *Inventiones Mathematicae*, 102(3):521–541,  
6 1990.
- 7 [18] F. Dal’Bo. *Geodesic and horocyclic trajectories*. Universitext. Springer-Verlag London, Ltd.,  
8 London; EDP Sciences, Les Ulis, London, 2011.
- 9 [19] F. Dal’Bo, J.-P. Otal, and M. Peigné. Séries de Poincaré des groupes géométriquement finis.  
10 *Israel Journal of Mathematics*, 118(1):109–124, 2000.
- 11 [20] F. Dal’Bo, M. Peigné, J.-C. Picaud, and A. Sambusetti. Convergence and counting in infinite  
12 measure. *Université de Grenoble. Annales de l’Institut Fourier*, 67(2):483–520, 2017.
- 13 [21] R. Dougall. Critical exponents of normal subgroups, the spectrum of group extended transfer  
14 operators, and Kazhdan distance. *arXiv.org*, Feb. 2017.
- 15 [22] R. Dougall and R. Sharp. Amenability, Critical Exponents of Subgroups and Growth of Closed  
16 Geodesics. *arXiv.org*, Nov. 2014.
- 17 [23] D. Fisher and G. Margulis. Almost isometric actions, property (T), and local rigidity. *Inven-*  
18 *tiones Mathematicae*, 162(1):19–80, 2005.
- 19 [24] R. I. Grigorchuk. Symmetrical random walks on discrete groups. In *Multicomponent random*  
20 *systems*, pages 285–325. Dekker, New York, 1980.
- 21 [25] M. Gromov. Hyperbolic groups. In *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New  
22 York, 1987.
- 23 [26] H. Hennion and L. Hervé. *Limit theorems for Markov chains and stochastic properties of dy-*  
24 *namical systems by quasi-compactness*, volume 1766 of *Lecture Notes in Mathematics*. Springer-  
25 Verlag, Berlin, Berlin, Heidelberg, 2001.
- 26 [27] J. Jaerisch. Group-extended Markov systems, amenability, and the Perron-Frobenius operator.  
27 *Proceedings of the American Mathematical Society*, 143(1):289–300, 2015.
- 28 [28] H. Kesten. Full Banach mean values on countable groups. *Mathematica Scandinavica*, 7:146–  
29 156, 1959.
- 30 [29] G. A. Margulis. On the decomposition of discrete subgroups into amalgams. *Selecta Mathe-*  
31 *matica Sovietica*, 1(2):197–213, 1981.
- 32 [30] F. Mathéus. Flot géodésique et groupes hyperboliques d’après M. Gromov. In *Séminaire*  
33 *de Théorie Spectrale et Géométrie, No. 9, Année 1990–1991*, pages 67–87. Univ. Grenoble I,  
34 Saint-Martin-d’Hères, 1991.
- 35 [31] B. H. Neumann. Groups covered by finitely many cosets. *Publicationes Mathematicae Debrecen*,  
36 3:227–242 (1955), 1954.

- 1 [32] F. Paulin. Un groupe hyperbolique est déterminé par son bord. *Journal of the London Math-*  
2 *ematical Society. Second Series*, 54(1):50–74, 1996.
- 3 [33] F. Paulin. On the critical exponent of a discrete group of hyperbolic isometries. *Differential*  
4 *Geometry and its Applications*, 7(3):231–236, 1997.
- 5 [34] T. Roblin. Un théorème de Fatou pour les densités conformes avec applications aux revêtements  
6 galoisiens en courbure négative. *Israel Journal of Mathematics*, 147(1):333–357, 2005.
- 7 [35] T. Roblin and S. Tapie. Exposants critiques et moyennabilité. In *Géométrie ergodique*, pages  
8 61–92. Enseignement Math., Geneva, Hoboken, NJ, USA, 2013.
- 9 [36] M. Stadlbauer. An extension of Kesten’s criterion for amenability to topological Markov chains.  
10 *Advances in Mathematics*, 235:450–468, 2013.
- 11 [37] P. Tukia. Convergence groups and Gromov’s metric hyperbolic spaces. *New Zealand Journal*  
12 *of Mathematics*, 23(2):157–187, 1994.
- 13 [38] Y. Watatani. Property T of Kazhdan implies property FA of Serre. *Mathematica Japonica*,  
14 27(1):97–103, 1982.
- 15 [39] W. Woess. *Random walks on infinite graphs and groups*, volume 138 of *Cambridge Tracts in*  
16 *Mathematics*. Cambridge University Press, Cambridge, 2000.

17 *Rémi Coulon*

18 Univ Rennes, CNRS  
19 IRMAR - UMR 6625  
20 F-35000 Rennes, France  
21 `remi.coulon@univ-rennes1.fr`  
22 `http://rcoulon.perso.math.cnrs.fr`

23 *Françoise Dal’Bo*

24 Univ Rennes, CNRS  
25 IRMAR - UMR 6625  
26 F-35000 Rennes, France  
27 `francoise.dalbo@univ-rennes1.fr`  
28 `https://perso.univ-rennes1.fr/francoise.dalbo`

29 *Andrea Sambusetti*

30 Dipartimento di Matematica  
31 Sapienza Università di Roma  
32 P.le Aldo Moro 5, 00185 Roma, Italy  
33 `sambuset@mat.uniroma1.it`  
34 `http://www1.mat.uniroma1.it/people/sambusetti/andreas_webpage/home.html`