



RESEARCH PAPER

DETERMINATION OF ORDER IN LINEAR FRACTIONAL
DIFFERENTIAL EQUATIONS

Mirko D'Ovidio ¹, Paola Loreti ²,
Alireza Momenzadeh ³, Sima Sarv Ahrabi ⁴

Abstract

The order of fractional differential equations (FDEs) has been proved to be of great importance in an accurate simulation of the system under study. In this paper, the orders of some classes of linear FDEs are determined by using the asymptotic behaviour of their solutions. Specifically, it is demonstrated that the decay rate of the solutions is influenced by the order of fractional derivatives. Numerical investigations are conducted into the proven formulas.

MSC 2010: Primary 34A08; Secondary 26A33, 33E12, 34E05

Key Words and Phrases: fractional differential equations, Mittag-Leffler type functions, asymptotic analysis, order estimation

1. Introduction

The practical significance of fractional calculus has been recently discerned as a vastly superior method of describing the long-memory processes and developed remarkably over the last few years [3, 8, 16, 21, 24, 5]. In particular, fractional differential equations (FDEs) have been proven extremely important for more accurately modelling of many physical phenomena [6, 23, 2, 20, 15, 26].

Inverse problems to FDEs occur in many branches of science. Such problems have been investigated in, for instance, fractional diffusion equation [7, 13, 19, 25, 28] and inverse boundary value problem for semi-linear

fractional telegraph equation [18]. In [13], it has been specifically shown that determination of β , the order of fractional differential operator, is definitely crucial to the appropriate simulation of the anomalous diffusion in order to specify that the transport phenomenon exhibits sub-diffusion or super-diffusion (respectively for $\beta < 1$ and $\beta > 1$). The authors of [13] have presented a theorem, the idea behind which is the determination of the order of a fractional diffusion equation. It has been the motivation behind this paper for proving formulas indicating the relationship between the fractional order and the asymptotic behaviour of the solutions to several different classes of linear FDEs.

The paper is organized as follows. Section 2 is allocated for recalling some bases of fractional calculus. In Section 3, a few special cases of linear FDEs are considered for which the fractional orders are determined. In Section 4, two examples are given to illustrate the correctness of the obtained results.

2. Preliminaries

For the sake of convenience, some basic definitions in fractional calculus and some useful properties are reviewed.

2.1. Fractional integral and derivatives. The definitions of the Riemann-Liouville integral and derivative, and also of the Caputo fractional derivative are represented in summary in some basic books as [16], [22], etc.

DEFINITION 2.1. Let $t_0 \in \mathbb{R}$ and $f : (t_0, +\infty) \rightarrow \mathbb{R}$ be continuous and integrable in every finite interval (t_0, t) . The Riemann-Liouville fractional integral of order $\beta \in \mathbb{C}$ ($\Re(\beta) > 0$) of the function f is defined by

$$\left(I_{t_0}^\beta f\right)(t) = \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t - \tau)^{\beta-1} f(\tau) d\tau, \quad t > t_0. \quad (2.1)$$

DEFINITION 2.2. Let t_0 be a real number, $\beta \in \mathbb{C}$ ($\Re(\beta) > 0$), $n = [\Re(\beta)] + 1$, and let the function $f : (t_0, +\infty) \rightarrow \mathbb{R}$ be continuous and integrable in every finite interval (t_0, t) . The Riemann-Liouville fractional derivative of order β of the function f is defined by

$$\left(D_{t_0}^\beta f\right)(t) = \frac{1}{\Gamma(n - \beta)} \left(\frac{d}{dt}\right)^n \int_{t_0}^t (t - \tau)^{n-\beta-1} f(\tau) d\tau, \quad t > t_0. \quad (2.2)$$

The Caputo differential operator plays a major role in physical phenomena due to the fact that initial conditions for the FDEs with Caputo derivative are the same as those for integer-order differential equations.

DEFINITION 2.3. Let $t_0 \in \mathbb{R}$, $\beta \in \mathbb{C}$ ($\Re(\beta) > 0$), $n - 1 < \Re(\beta) < n$ ($n \in \mathbb{N}$), and let the function f be continuous and have n continuous derivatives in the interval $(t_0, +\infty)$. The Caputo fractional derivative of order β of the function f is defined by

$$\left(D_t^\beta f\right)(t) = \frac{1}{\Gamma(n - \beta)} \int_{t_0}^t (t - \tau)^{n-\beta-1} f^{(n)}(\tau) d\tau, \quad t > t_0. \quad (2.3)$$

Next, we mention also the sequential fractional derivative (see [16, 22]), defined by

$$\mathcal{D}^{n\beta} u(t) = \underbrace{D^\beta D^\beta \dots D^\beta}_n u(t), \quad (2.4)$$

where D^β could be Riemann-Liouville, Caputo, or any other type of fractional derivative not discussed here.

2.2. Mittag-Leffler function and its derivatives. The two-parameter Mittag-Leffler (M-L) function appeared first in an article by Wiman [27] and was studied by Agarwal and Humbert [1, 14]. It is defined by

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(k\alpha + \beta)}, \quad \alpha, \beta, z \in \mathbb{C}, \quad \Re(\alpha) > 0, \quad (2.5)$$

and $E_{\alpha,1}(z)$ is simply denoted by $E_\alpha(z)$, the original one-parameter function of M-L. In the case α and β are real and positive numbers, the series converges for all values of the argument z , therefore, the M-L function $E_{\alpha,\beta}(z)$ is an entire function of the order α^{-1} . Detailed information on the Mittag-Leffler type functions, some extensions and their properties can be found in, for instance, [16], [12].

The Mittag-Leffler function satisfies the following differentiation formula [22]:

$$\frac{d^n}{dz^n} \left(z^{\beta-1} E_{\alpha,\beta}(\lambda z^\alpha) \right) = z^{\beta-n-1} E_{\alpha,\beta-n}(\lambda z^\alpha), \quad n \in \mathbb{N}, \lambda \in \mathbb{R}, \quad (2.6)$$

and the following practical formulas can be directly derived from (2.6):

$$\frac{d}{dz} E_\alpha(\lambda z^\alpha) = \frac{1}{z} E_{\alpha,0}(\lambda z^\alpha) = \lambda z^{\alpha-1} E_{\alpha,\alpha}(\lambda z^\alpha), \quad (2.7)$$

$$\frac{d}{dz} \left(z^{\beta-1} E_{\alpha,\beta}(\lambda z^\alpha) \right) = z^{\beta-2} E_{\alpha,\beta-1}(\lambda z^\alpha), \quad (2.8)$$

$$\frac{d}{dz} (z E_{\alpha,2}(\lambda z^\alpha)) = E_\alpha(\lambda z^\alpha). \quad (2.9)$$

2.3. Asymptotic expansion of Mittag-Leffler function. The asymptotic behaviour of the Mittag-Leffler function $E_{\alpha,\beta}(z)$ ($|z| \rightarrow \infty$) is complicated for $\alpha > 0$ and diverges greatly for $0 < \alpha < 2$ and $\alpha \geq 2$. This topic has been discussed in [22] with scrupulous attention to detail. Here, the asymptotic behaviour of the M-L function is only briefly stated for the case $0 < \alpha < 2$.

Let $0 < \alpha < 2$, $\beta, z \in \mathbb{C}$ and μ be an arbitrary real number such that $\frac{\pi}{2}\alpha < \mu < \min(\pi, \pi\alpha)$, then:

$$E_{\alpha,\beta}(z) = - \sum_{k=1}^n \frac{1}{\Gamma(\beta - k\alpha) z^k} + O(|z|^{-n-1}), \tag{2.10}$$

$|z| \rightarrow \infty, \quad \mu \leq |\arg(z)| \leq \pi.$

Consequently, the following expansions can be obtained from (2.10):

$$E_{\alpha}(\lambda z^{\alpha}) = - \frac{z^{-\alpha}}{\lambda \Gamma(1 - \alpha)} + O\left(\frac{1}{|\lambda|^2 z^{2\alpha}}\right), \quad z \rightarrow \infty, \tag{2.11}$$

$$E_{\alpha,\alpha}(\lambda z^{\alpha}) = \frac{\alpha z^{-2\alpha}}{\lambda^2 \Gamma(1 - \alpha)} + O\left(\frac{1}{|\lambda|^3 z^{3\alpha}}\right), \quad z \rightarrow \infty. \tag{2.12}$$

where z is a positive real number, $0 < \alpha < 2$, and $\lambda < 0$.

3. Main results

In this section the fractional orders of several classes of linear FDEs are determined by using the asymptotic behaviour of Mittag-Leffler functions.

THEOREM 3.1. *Let $0 < \beta \leq 1$ and $t_0 > 0$. Consider the sequential linear differential equation in the sense of Riemann-Liouville derivative:*

$$\mathcal{D}_{t_0}^{2\beta} u + a_1 \mathcal{D}_{t_0}^{\beta} u + a_0 u = 0, \tag{3.1}$$

with the initial condition $u(t_0) = u_0$ and $\mathcal{D}_{t_0}^{\beta} u(t_0) = u_1$, and let a_0 and a_1 are reals such that r_1 and r_2 , the roots of the characteristic equation $r^2 + a_1 r + a_0 = 0$, are distinct and negative real numbers. Then, the following formula holds

$$\beta = -1 - \lim_{t \rightarrow \infty} \frac{tu'}{u}. \tag{3.2}$$

P r o o f. The solution of (3.1) is as follows ([16]):

$$u(t) = c_1 t^{\beta-1} E_{\beta,\beta}(r_1 t^{\beta}) + c_2 t^{\beta-1} E_{\beta,\beta}(r_2 t^{\beta}), \tag{3.3}$$

where c_1 and c_2 depend on the initial conditions. The first derivative of $u(t)$ can be calculated by using (2.8) as below:

$$u'(t) = c_1 t^{\beta-2} E_{\beta,\beta-1}(r_1 t^{\beta}) + c_2 t^{\beta-2} E_{\beta,\beta-1}(r_2 t^{\beta}). \tag{3.4}$$

The asymptotic expansions of $u(t)$ and $u'(t)$ can be obtained by using (2.10),

$$u(t) = \frac{-t^{-\beta-1}}{\Gamma(-\beta)} \left(\frac{c_1}{r_1^2} + \frac{c_2}{r_2^2} \right) + c_1 t^{\beta-1} O(|r_1|^{-3} t^{-3\beta}) + c_2 t^{\beta-1} O(|r_2|^{-3} t^{-3\beta}), \tag{3.5}$$

$$u'(t) = \frac{(\beta+1)t^{-\beta-2}}{\Gamma(-\beta)} \left(\frac{c_1}{r_1^2} + \frac{c_2}{r_2^2} \right) + c_1 t^{\beta-2} O(|r_1|^{-3} t^{-3\beta}) + c_2 t^{\beta-2} O(|r_2|^{-3} t^{-3\beta}). \tag{3.6}$$

Thus,

$$-\lim_{t \rightarrow \infty} \frac{tu'}{u} = -\lim_{t \rightarrow \infty} \frac{\frac{(\beta+1)t^{-\beta-1}}{\Gamma(-\beta)} \left(\frac{c_1}{r_1^2} + \frac{c_2}{r_2^2} \right)}{\frac{-t^{-\beta-1}}{\Gamma(-\beta)} \left(\frac{c_1}{r_1^2} + \frac{c_2}{r_2^2} \right)} = \beta + 1. \tag{3.7}$$

□

THEOREM 3.2. *Let $0 < \beta < \frac{1}{2}$ and $\gamma, \mu \in \mathbb{R}$ such that $0 < \gamma < \mu^2$, and let $D_t^\beta u$ indicate the Caputo differentiation operator. For the initial value problem*

$$D_t^{2\beta} u(t) + 2\mu D_t^\beta u(t) + \gamma u(t) = 0, \tag{3.8}$$

with the initial condition $u(0) = 1$, and also for the sequential linear differential equation of fractional order

$$\mathcal{D}_t^{2\beta} u + 2\mu \mathcal{D}_t^\beta u + \gamma u = 0, \tag{3.9}$$

with the initial condition $\mathcal{D}_t^\beta u(0) = 0$ and $u(0) = 1$, the following formula holds:

$$\beta = -\lim_{t \rightarrow \infty} \frac{tu'}{u}. \tag{3.10}$$

REMARK 3.1. If $\mathcal{D}_t^\beta u(0) = 0$, then $c_1 r_1 + c_2 r_2 = 0$ and $\mathcal{D}_t^{2\beta} u = D_t^{2\beta} u$. The case of $\mathcal{D}_t^\beta u(0) \neq 0$ leads to $c_1 r_1 + c_2 r_2 \neq 0$. Thus, the coefficients c_1 and c_2 are not the same as those represented in the proof of Theorem 3.2 and therefore must be calculated.

P r o o f. The solutions of (3.8) and (3.9) are represented by [9]

$$u(t) = c_1 E_\beta(r_1 t^\beta) + c_2 E_\beta(r_2 t^\beta), \tag{3.11}$$

where the coefficients c_1 and c_2 are respectively equal to $\frac{1}{2} \left(1 + \frac{\mu}{\sqrt{\mu^2 - \gamma}} \right)$ and $\frac{1}{2} \left(1 - \frac{\mu}{\sqrt{\mu^2 - \gamma}} \right)$, and the parameters r_1 and r_2 are equal to $-\mu +$

$\sqrt{\mu^2 - \gamma} < 0$ and $-\mu - \sqrt{\mu^2 - \gamma} < 0$ respectively. Equation (3.11) can be rewritten in the following form by using (2.11)

$$u(t) = -\frac{t^{-\beta}}{\Gamma(1-\beta)} \left(\frac{c_1}{r_1} + \frac{c_2}{r_2} \right) + c_1 O\left(r_1^{-2} t^{-2\beta}\right) + c_2 O\left(r_2^{-2} t^{-2\beta}\right). \tag{3.12}$$

The first derivative of (3.11) can be obtained by referring to (2.7)

$$u'(t) = c_1 r_1 t^{\beta-1} E_{\beta,\beta}\left(r_1 t^\beta\right) + c_2 r_2 t^{\beta-1} E_{\beta,\beta}\left(r_2 t^\beta\right). \tag{3.13}$$

Equation (2.12) is used for applying the asymptotic behaviour of Mittag-Leffler function to (3.13):

$$u'(t) = \frac{\beta t^{-\beta-1}}{\Gamma(1-\beta)} \left(\frac{c_1}{r_1} + \frac{c_2}{r_2} \right) + t^{\beta-1} \left(c_1 r_1 O\left(r_1^{-3} t^{-3\beta}\right) + c_2 r_2 O\left(r_2^{-3} t^{-3\beta}\right) \right). \tag{3.14}$$

Therefore,

$$-\lim_{t \rightarrow \infty} \frac{tu'}{u} = -\lim_{t \rightarrow \infty} t \frac{\frac{\beta t^{-\beta-1}}{\Gamma(1-\beta)} \left(\frac{c_1}{r_1} + \frac{c_2}{r_2} \right)}{-\frac{t^{-\beta}}{\Gamma(1-\beta)} \left(\frac{c_1}{r_1} + \frac{c_2}{r_2} \right)} = \beta. \tag{3.15}$$

□

THEOREM 3.3. *Let $1 < \beta < 2$, and r be a negative real number. For the fractional differential equation with Caputo derivative*

$$D_t^\beta u - ru = 0, \tag{3.16}$$

with the initial condition $u(0) = 1$ and $u'(0) = 1$, the following relationship is held true:

$$\beta = 1 - \lim_{t \rightarrow \infty} \frac{tu'}{u}. \tag{3.17}$$

P r o o f. The solution to (3.16) is

$$u(t) = E_\beta\left(rt^\beta\right) + tE_{\beta,2}\left(rt^\beta\right). \tag{3.18}$$

The first derivative of $u(t)$ could be calculated by referring to (2.7) and (2.9)

$$u'(t) = rt^{\beta-1} E_{\beta,\beta}\left(rt^\beta\right) + E_\beta\left(rt^\beta\right). \tag{3.19}$$

The asymptotic expansions of $u(t)$ and $u'(t)$ are respectively

$$u(t) = -\frac{t^{-\beta}}{r\Gamma(1-\beta)} \left(1 + \frac{t}{1-\beta} \right) + O\left(|r|^{-2} t^{-2\beta}\right) + tO\left(|r|^{-2} t^{-2\beta}\right), \tag{3.20}$$

and

$$u'(t) = \frac{t^{-\beta}}{r\Gamma(1-\beta)} \left(\frac{\beta}{t} - 1 \right) + rt^{\beta-1}O\left(|r|^{-3}t^{-3\beta}\right) + O\left(|r|^{-2}t^{-2\beta}\right), \quad (3.21)$$

therefore,

$$\lim_{t \rightarrow \infty} \frac{tu'}{u} = \lim_{t \rightarrow \infty} \frac{\frac{t^{-\beta}}{r\Gamma(1-\beta)}(\beta - t)}{-\frac{t^{-\beta}}{r\Gamma(1-\beta)}\left(1 + \frac{t}{1-\beta}\right)} = 1 - \beta. \quad (3.22)$$

□

THEOREM 3.4. *Let $0 < \beta < 1$, and r be a negative real number. For (3.16) with the initial condition $u(0) = 1$, the following is held true*

$$\beta = - \lim_{t \rightarrow \infty} \frac{tu'}{u}. \quad (3.23)$$

P r o o f. The solution to (3.16) is as follows:

$$u(t) = E_{\beta}\left(rt^{\beta}\right), \quad (3.24)$$

where $0 < \beta < 1$, [17]. The first derivative of $u(t)$ could be calculated by referring to (2.7):

$$u'(t) = rt^{\beta-1}E_{\beta,\beta}\left(rt^{\beta}\right). \quad (3.25)$$

The asymptotic expansions of $u(t)$ and $u'(t)$ are respectively

$$u(t) = -\frac{t^{-\beta}}{r\Gamma(1-\beta)} + O\left(|r|^{-2}t^{-2\beta}\right), \quad (3.26)$$

and

$$u'(t) = \frac{\beta t^{-\beta-1}}{r\Gamma(1-\beta)} + rt^{\beta-1}O\left(|r|^{-3}t^{-3\beta}\right), \quad (3.27)$$

then

$$\lim_{t \rightarrow \infty} \frac{tu'}{u} = \lim_{t \rightarrow \infty} \frac{\frac{\beta t^{-\beta}}{r\Gamma(1-\beta)}}{-\frac{t^{-\beta}}{r\Gamma(1-\beta)}} = -\beta. \quad (3.28)$$

□

4. Numerical investigation

EXAMPLE 4.1. Consider the initial value problem

$$D_t^{2\beta}u + 2D_t^{\beta}u + 0.7u = 0, \quad t \geq 0, \quad 0 < \beta < \frac{1}{2}, \quad (4.1)$$

with the initial condition $u(0) = 1$. The solution to (4.1) is the following:

$$u(t) = c_1E_{\beta}\left(r_1t^{\beta}\right) + c_2E_{\beta}\left(r_2t^{\beta}\right), \quad (4.2)$$

where $r_1 = -0.4523$, $r_2 = -1.5477$, $c_1 = 1.4129$, $c_2 = -0.4129$. The graph of $-\frac{tu'}{u}$, which has been evaluated for several different values of β , is shown

in Fig. 1. It is obvious that $-\frac{tu'}{u}$ tends asymptotically to β , as t goes to infinity. The term $-\frac{tu'}{u}$ is numerically evaluated by means of the MATLAB code “ml.m” [10] which is based on the numerical inversion of the Laplace transform of Mittag-Leffler function [11]. Numerical results are in good agreement with the formula introduced in Theorem 3.2 and the rate of the convergence of $-\frac{tu'}{u}$ is greatly influenced by the value of β .




FIGURE 1. Graph of $-\frac{tu'}{u}$ for $\beta = 0.35$, $\beta = 0.40$, and $\beta = 0.45$.

EXAMPLE 4.2. Consider the fractional differential equation

$$\begin{cases} D_t^\beta u + 2u = 0, \\ u(0) = 1, \\ u'(0) = 1, \end{cases} \tag{4.3}$$

where $1 < \beta < 2$ and $D_t^\beta u$ is in the sense of Caputo derivative. The solution is in the form of

$$u(t) = E_\beta(-2t^\beta) + tE_{\beta,2}(-2t^\beta). \tag{4.4}$$

According to Theorem 3.3, the term $1 - \frac{tu'}{u}$ tends to β as t goes to the infinity. The numerical evaluation of $1 - \frac{tu'}{u}$ has been conducted by using (4.4) and its derivative. The results are shown in Fig. 2. As it can be observed, $1 - \frac{tu'}{u}$ converges to β with a rate affected by the value of β , i.e. the convergence will be faster if the fractional order β tends to 2.




FIGURE 2. Graph of $1 - \frac{tu'}{u}$ for $\beta = 1.2$, $\beta = 1.5$, and $\beta = 1.7$.

5. Conclusion

Inverse problems occur in many branches of science and have been also examined in the models described by FDEs. For instance, determination of the order of fractional systems has been indicated to be of such crucial importance that it could influence how anomalous diffusion equations must be appropriately simulated. Thus, in this paper, the solutions to several classes of linear FDEs represented, for which the order determination has been demonstrated by using asymptotic behaviour of Mittag-Leffler type functions. The formulas in Theorems 3.1-3.4 have been numerically examined for different values of β . The results show the accuracy of the obtained formulas.

Acknowledgements

The authors are grateful to Professor Masahiro Yamamoto for pointing out the papers [13], [25], [7].

References

- [1] R.P. Agarwal, A propos d'une note de M. Pierre Humbert. *CR Acad. Sci. Paris* **236**, No 21 (1953), 2031–2032.
- [2] O.P. Agrawal, O. Defterli, D. Baleanu, Fractional optimal control problems with several state and control variables. *J. Vib. Control.* **16**, No 13 (2010), 1967–1976.
- [3] D. Baleanu, Z.B. Güvenç, J.T. Machado, *New Trends in Nanotechnology and Fractional Calculus Applications*. Springer, New York (2010).
- [4] M. Caputo, Linear models of dissipation whose Q is almost frequency independent—II. *Geophys. J. Int.* **13**, No 5 (1967), 529–539.
- [5] M. Caputo, M. Fabrizio, A new definition of fractional derivative without singular kernel. *Progr. Fract. Differ. Appl.* **1**, No 2 (2015), 1–13.
- [6] A. Carpinteri, F. Mainardi, *Fractals and Fractional Calculus in Continuum Mechanics*. Springer, New York (1997).
- [7] J. Cheng, J. Nakagawa, M. Yamamoto, T. Yamazaki, Uniqueness in an inverse problem for a one-dimensional fractional diffusion equation. *Inverse Probl.* **25**, No 11 (2009), # 115002.
- [8] K. Diethelm, *The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operators of Caputo Type*. Springer, Heidelberg (2010).
- [9] M. D'Ovidio, E. Orsingher, B. Toaldo, Time-changed processes governed by space-time fractional telegraph equations. *Stoch. Anal. Appl.* **32**, No 6 (2014), 1009–1045.
- [10] R. Garrappa, The Mittag-Leffler function, *MATLAB Central File Exchange* (2014).
- [11] R. Garrappa, Numerical evaluation of two and three parameter Mittag-Leffler functions. *SIAM J. Numer. Anal.* **53**, No 3 (2015), 1350–1369.
- [12] R. Gorenflo, A.A. Kilbas, F. Mainardi, and S.V. Rogosin, *Mittag-Leffler Functions, Related Topics and Applications*. Springer Monographs in Mathematics, Springer-Verlag, Berlin-Heidelberg (2014).
- [13] Y. Hatano, J. Nakagawa, S. Wang, M. Yamamoto, Determination of order in fractional diffusion equation. *J. Math-for-Ind.* **5A** (2013), 51–57.
- [14] P. Humbert, Quelques résultats relatifs à la fonction de Mittag-Leffler. *Comptes rendus hebdomadaires des seances de l'academie des sciences* **236**, No 15 (1953), 1467–1468.

- [15] R. Kamocki, Pontryagin maximum principle for fractional ordinary optimal control problems. *Math. Methods Appl. Sci.* **37**, No 11 (2014), 1668–1686.
- [16] A. Kilbas, H. Srivastava, J. Trujillo, *Theory and Applications of Fractional Differential Equations*. Elsevier, Amsterdam (2006).
- [17] S.D. Lin, C.H. Lu, Laplace transform for solving some families of fractional differential equations and its applications. *Adv. Difference Equ.* **2013**, No 1 (2013), 137.
- [18] H. Lopushanska, V. Rapita, Inverse coefficient problem for semi-linear fractional telegraph equation. *Electr. J. Differ. Equ.* **2015**, No 153 (2015), 1–13.
- [19] A.O. Lopushansky, H.P. Lopushanska, One inverse problem for the diffusion-wave equation in bounded domain. *Ukr. Math. J.* **66**, No 5 (2014), 743–757.
- [20] R.L. Magin, Fractional calculus models of complex dynamics in biological tissues. *Comput. Math. Appl.* **59**, No 5 (2010), 1586–1593.
- [21] F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*. Imperial College Press, London (2010).
- [22] I. Podlubny, *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*. Academic Press, San Diego (1998).
- [23] Y.A. Rossikhin, M.V. Shitikova, Application of fractional calculus for dynamic problems of solid mechanics: novel trends and recent results. *Appl. Mech. Rev.* **63**, No 1 (2010), # 010801.
- [24] J. Sabatier, O.P. Agrawal, J.A. Tenreiro Machado, *Advances in Fractional Calculus*. Springer, Dordrecht (2007).
- [25] K. Sakamoto, M. Yamamoto, Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.* **382**, No 1 (2011), 426–447.
- [26] D. Valério, J.T. Machado, V. Kiryakova, Some pioneers of the applications of fractional calculus. *Fract. Calc. Appl. Anal.* **17**, No 2 (2014), 552–578; DOI: 10.2478/s13540-014-0185-1; <https://www.degruyter.com/view/j/fca.2014.17.issue-2/issue-files/fca.2014.17.issue-2.xml>.
- [27] A. Wiman, Über den fundamentalsatz in der theorie der funktionen $E_a(x)$. *Acta Math.* **29**, No 1 (1905), 191–201.
- [28] Y. Zhang, X. Xu, Inverse source problem for a fractional diffusion equation. *Inverse Probl.* **27**, No 3 (2011), Art. # 035010.

^{1,2,4} *Department of Basic and Applied Sciences for Engineering (SBAI)*
Sapienza University of Rome
Via Antonio Scarpa, 16
00161 Rome, ITALY

¹ *e-mail: mirko.dovidio@sbai.uniroma1.it*

Received: July 16, 2017

² *e-mail: paola.loreti@sbai.uniroma1.it*

⁴ *e-mail: sima.sarvahrabi@sbai.uniroma1.it*

³ *Department of Information Engineering,*
Electronics and Telecommunications (DIET)
Sapienza University of Rome
Via Eudossiana, 18
00184 Rome, ITALY
e-mail: momenzadeh@diet.uniroma1.it

Please cite to this paper as published in:

Fract. Calc. Appl. Anal., Vol. **21**, No 4 (2018), pp. 937–948,

DOI: 10.1515/fca-2018-0051 , at <https://www.degruyter.com/view/j/fca>