

Figure 1. Cardiac magnetic resonance (MR) showing moderate symmetric left ventricular hypertrophy (LVH), which mostly involves the anterior section of interventricular septum (IVS) in its basal and medial part (septal thickness of anterior-basal part is 19 mm; inferior-basal part is 14 mm; anterior-medial part is 16 mm; inferior-basal part is 14 mm), in basal portion of anterior wall (thickness 18 mm) and in middle portion of lateral wall (thickness 16 mm). The remaining part of myocardium has a thickness between 10 and 15 mm. This hypertrophy is responsible for a mild tightening/obstruction of the left ventricular outflow tract and a modest acceleration of flow, without an atypical systolic anterior motion of the mitral valve (SAM). Mild aortic and mitral valve regurgitation can be seen. LVH = left ventricular hypertrophy, IVS = interventricular septum, MR = magnetic resonance.

regular. Her respiratory rate was 20 breaths/min and her lungs were clear at the auscultation. The electrocardiogram (ECG) showed sinus rhythm; there were signs of left ventricular hypertrophy (LVH) with negative T waves in lateral leads, V5 and V6. The plasma proBNP concentration was 87 pg/mL, which was in normal range (normal values < 100 pg/ml). Other data, including electrolytes, troponin, creatinine kinase, creatinine were within the normal range. The chest X-ray film no revealed enlargement of the cardiac silhouette. A 2D transthoracic echocardiogram showed a dilatation of the left atrium (24 cm²) with severe LVH (end-diastolic intraventricular septum thickness—17 mm) and preserved left ventricular ejection fraction (LVEF—60%). Suspecting the presence of a cardiomyopathy a cardiac magnetic resonance (MR) imaging with gadolinium was performed and showed symmetric LVH with a prevalent involvement of the anterior portion of interventricular septum. A minimum left ventricular outflow obstruction was reported without an atypical systolic anterior motion of the

mitral valve. A supposed diagnosis of HCM was made (Fig. 1). Heart muscle impairment was investigated by cardiac catheterization and angiography. Coronary angiography revealed no coronary artery stenosis. Catheterization was followed by a left ventricle endomyocardial biopsy extracting 4 fragments, which were processed for histology and electron microscopy. At the left ventricle endomyocardial biopsy myocytes demonstrated features of hypercontraction of myofilaments with formation of contraction bands and occasional cellular necrosis. Wide intracellular and perivascular fibrosis was seen. A significant number of arterioles showed thickened tunica with resulting vessel lumen reduction. Further, the endocardium was thickened with prominence of smooth muscle cells (Fig. 2). An electron microscopy showed at higher magnification the catecholamine-induced diffuse contraction band necrosis. This data was suggestive of catecholamine-induced LVH (Fig. 2).

A hormonal screening test for secondary arterial hypertension revealed an abnormal increase of 24-hours urinary metanephrines




Figure 2. Microscopic findings of biopsy specimen from the endomyocardium of left ventricle showing markedly hypertrophied fibers, contraction bands and replacement fibrosis.

Table 1**Endocrinological data of the patient at diagnosis and after laparoscopy surgical treatment.**

	At diagnosis	After surgery	Normal values
PLASMA			
Plasma renin activity	2.0	2.7	0.2–2.7 ng/mL/h
Plasma aldosterone concentration	141	207	30–160 pg/mL
Plasma cortisol	9	11.1	4.5–21 µg/dL
Plasma cortisol post-overnight DXM-test	12	–	< 50 pg/mL
URINE			
Free cortisol	126.3	78.3	58–403 µg/24h
Urine aldosterone concentration	19.0	30.7	2.8–34 µg/24
Metanephrines	396; 421*	143	0–350 µg/24h
Vanillyl mandelic acid	6.7; 5.1*	3.2	1–10 µg/24h

DXM = dexamethasone: 1 mg at 11:00 pm

* Second sample collection

(396 and 421 µg/24h; normal values 2–350 µg/24h) suggesting the presence of a PHEO (Table 1). A computed tomography (CT) scan showed an enlargement of the left adrenal gland with globe shape. The strongest uptake of ^{123}I -Metaiodobenzylguanidine (^{123}I -MIBG) by the left adrenal lesion was observed in ^{123}I -MIBG scintigraphy (Fig. 3).

Pre-operative management was begun with fluids and alpha-adrenergic receptor blockers, resulting in BP control. The patient underwent left laparoscopic adrenalectomy with no significant hemodynamic alterations during the procedure. The adrenal gland measured 3.5 x 2.5 x 2.0 cm and it had an altered profile due to the presence of a yellow nodule surrounded by a subtle fibrous capsule. Nodule was made of 2 cell populations arranged in nests of trabeculae: 1 minor cell line had eosinophilic granular cytoplasm, the other had clear vacuolated cytoplasm. The remaining left adrenal gland showed areas of cortical hyperplasia (Fig. 4). On the basis of these findings a histological diagnosis of cortical adenoma with adrenocortical hyperplasia was primarily made. Secondly, immunohistochemical staining revealed diffused positivity for synaptophysin and chromogranin. At electron

microscopy the tumor cells contained lipid vacuoles and smooth endoplasmic reticulum, consisted with adrenocortical differentiation, along multiple intracytoplasmic adrenaline-type, dense core neurosecretory granules (Fig. 5).

Considering these results a diagnosis of adrenocortical adenoma with mixed nodules enriched in adrenaline-type secretory granules was made.

The patient was discharged medicated with irbesartan 150 mg/day, amlodipine 5 mg/day, and hydrochlorothiazide 12.5 mg/day.

At 6 months follow up, the patient's 24-hours urinary metanephrines levels were normalized (143 µg/24h) and the transthoracic echocardiography after surgery showed a reduction of end-diastolic intraventricular septum thickness (14 mm) (Table 2) associated to normalization of BP values (130/80 mm Hg).

3. Discussion

This is a unique case, to our knowledge, of a catecholamine-induced HCM induced by a cortical adrenal adenoma with




Figure 3. Scintigraphy with ^{123}I -metaiodobenzylguanidine (^{123}I -MIBG) showed raised activity within the left adrenal gland, concordant with the mass.




Figure 4. Immunohistochemical staining for synaptophysin in cortical adrenal gland specimen.

adrenaline-neurosecretory granules inside. The appearance of these metabolites, stored into nodules enriched in multiple secretory granules, explains the strongest uptake of the left adrenal lesion in ¹²³I-MIBG scintigraphy. Previously, false-positive results of MIBG imaging were reported in adrenal adenomas showing augmented tracer uptake.^[6,7,8] We support that our case cannot be listed in those previously described because we demonstrated, by using electron microscopy, the existence of a complex network of neurosecretory granules, that are directly responsible of this clinical scenario.

Left ventricular endomyocardial biopsy has been crucial, in this case, to testify catecholamine-induced LVH. Use of endomyocardial biopsy in HCM is still controversial and is recommended only in infiltrative or storage diseases.^[9] In patients with true HCM, left ventricular endomyocardial biopsy can be specifically useful to analyze the underlying causes of cardiac deterioration,^[10] to demonstrate histological changes during the course of the disease, and finally to provide myocardial tissue for research purposes.^[11]

Catecholamine-induced cardiomyopathy is defined by a reversible left ventricular dysfunction without evidence of obstructive coronary artery disease. Acute and chronic myocardial damage are the results of an exogenous or endogenous catecholamine excess.

Overactivation of the sympatho-adrenergic system is an essential process to support cardiovascular system during stressful conditions in critical illnesses.^[12] Although, its prolonged and excessive stimulation is detrimental to cardiovascular system. Abnormal amounts of catecholamines, via the beta-1 adrenoceptor transduction pathway, induce directly intracellular




Figure 5. Electron microscopy showed neurosecretory granules in cortico-medulla nodules.

calcium overload in cardiomyocytes.^[13] Furthermore, these metabolites can also indirectly damage the heart, in particular via their oxidation during stressful situations, which generate oxygen radicals that provoke coronary artery spasm, arrhythmias and cardiac dysfunction.^[14] On the other hand, long-term exposure to catecholamines down-regulate the expression of beta-adrenergic receptors, inducing the suboptimal function of myofibers and decreasing the number of contracting units. As the consequent result, PHEO-associated cardiomyopathy is histologically characterized by contraction band necrosis, neutrophil infiltration, and fibrosis.^[15]

Similar mechanisms of myocardial toxicity, which originate from a hyper catecholaminergic state, have been described in case of cocaine abuse, including marked oxidative stress and mitochondrial dysfunction. Instead, chronic use of cocaine may result in various degrees of systolic and diastolic dysfunction, cardiac hypertrophy and dilatation.^[16] Histological features, such as in those cases of endogenous excess of

Table 2

Comparison of 2D Echocardiographic features before and 6 months after laparoscopy surgical treatment.

	At diagnosis	After surgery
Left Ventricle (LV) posterior wall thickness in diastole (mm)	13.0	12.2
Interventricular septum (IVS) thickness in diastole, (mm)	17.0	15.0
Systolic anterior motion of the mitral valve (SAM)	Absent	Absent
Mitral regurgitation	Mild	Absent
Mean LV outflow gradient (mm Hg)	35	22
Ejection fraction (%)	60	60

IVS = Interventricular septum, LV = Left Ventricle, SAM = Systolic anterior motion.

catecholamines, include loss of myofibrils, multiple foci of band contraction necrosis and fibrosis.^[17]

Treatment of PHEO-associated cardiomyopathy is curative surgical resection of catecholamine-producing tumor. After the appropriate excision of PHEO, cardiomyopathy may definitely reverse.^[18]

In summary, we have described a rare case of catecholamine-induced cardiomyopathy due to an adrenal adenoma mixed with nodules enriched in epinephrine-type secreting granules. The patient has provided informed consent for publication of this case report.

Author contributions

Data curation: Federica Olmati, Gaia Oliviero, Maria Bonvicini, Antonio Ciardi, Gino Iannucci.

Investigation: Vincenza Saracino, Martina Mezzadri.

Methodology: Valeria Bisogni, Antonio Concistré.

Supervision: Giorgio de Toma, Claudio Letizia.

Validation: Giorgio de Toma, Andrea Frustaci, Claudio Letizia.

Writing – original draft: Luigi Petramala, Andrea Frustaci,

Claudio Letizia, Federica Olmati.

Writing – review & editing: Luigi Petramala, Federica Olmati.

References

- [1] Plouin PF, Gimenez-Roqueplo AP. Pheochromocytomas and secreting paragangliomas. *Orphanet J Rare Dis* 2006;1:49.
- [2] Sever PS, Roberts JC, Snell ME. Pheochromocytoma. *Clin Endocrinol Metab* 1980;9:543–68.
- [3] Bravo EL, Gifford RW Jr. Current concepts. Pheochromocytoma: diagnosis, localization and management. *N Engl J Med* 1984;311:1298–303.
- [4] Stolk RF, Bakx C, Mulder J, et al. Is the excess cardiovascular morbidity in pheochromocytoma related to blood pressure or to catecholamines? *J Clin Endocrinol Metab* 2013;98:1100–6.
- [5] Batisse-Lignier M, Pereira B, Motreff P, et al. Acute and chronic pheochromocytoma-induced cardiomyopathies: different prognoses?: a systematic analytical review. *Medicine [Baltimore]* 2015;94:e2198.
- [6] Alsabeh R, Mazoujian G, Goates J, et al. Adrenal cortical tumors clinically mimicking pheochromocytoma. *Am J Clin Pathol* 1995;104:382–90.
- [7] Sone H, Okuda Y, Nakamura Y, et al. Radioiodinated metaiodobenzylguanidine scintigraphy for pheochromocytoma. A false-positive case of adrenocortical adenoma and literature review. *Horm Res* 1996;46:138–42.
- [8] Letizia C, De Toma G, Massa R, et al. False-positive diagnosis of adrenal pheochromocytoma on iodine-123-MIBG scan. *J Endocrinol Invest* 1998;21:779–83.
- [9] Isabelle M, Vergeade A, Moritz F, et al. NADPH oxidase inhibition prevents cocaine-induced up-regulation of xanthine oxidoreductase and cardiac dysfunction. *J Mol Cell Cardiol* 2007;42:326–32.
- [10] Frustaci A, Verardo R, Caldarulo M, et al. Myocarditis in hypertrophic cardiomyopathy patients presenting acute clinical deterioration. *Eur Heart J* 2007;28:733–40.
- [11] Cooper LT, Baughman KL, Feldman AM, et al. The role of endomyocardial biopsy in the management of cardiovascular disease: a scientific statement from the American Heart Association, the American College of Cardiology, and the European Society of Cardiology. *Circulation* 2007;116:2216–33.
- [12] Kassim TA, Clarke DD, Mai VQ, et al. Catecholamine-induced cardiomyopathy. *Endocr Pract* 2008;14:1137–49.
- [13] Adameova A, Abdellatif Y, Dhalla NS. Role of the excessive amounts of circulating catecholamines and glucocorticoids in stress-induced heart disease. *Can J Physiol Pharmacol* 2009;87:493–514.
- [14] Schwartz BG, Rezkalla S, Kloner RA. Cardiovascular effects of cocaine. *Circulation* 2010;122:2558–69.
- [15] Chimenti C, Frustaci A. Contribution and risks of left ventricular endomyocardial biopsy in patients with cardiomyopathies: a retrospective study over a 28-year period. *Circulation* 2013;128:1531–41.
- [16] Liaudet L, Calderari B, Pacher P. Pathophysiological mechanisms of catecholamine and cocaine-mediated cardiotoxicity. *Heart Fail Rev* 2014;19:815–24.
- [17] de Miguel V, Arias A, Paissan A, et al. Catecholamine-induced myocarditis in pheochromocytoma. *Circulation* 2014;129:1348–9.
- [18] Zieleń P, Klisiewicz A, Januszewicz A, et al. Pheochromocytoma-related 'classic' takotsubo cardiomyopathy. *J Hum Hypertens* 2010;24:363–6.