

Axion phenomenology and θ -dependence from $N_f = 2 + 1$ lattice QCD

Claudio Bonati,^{a,b} Massimo D'Elia,^{a,b} Marco Mariti,^{a,b} Guido Martinelli,^{c,d}
 Michele Mesiti,^{a,b} Francesco Negro,^b Francesco Sanfilippo^e and Giovanni Villadoro^f

^a*Dipartimento di Fisica dell'Università di Pisa,
 Largo Pontecorvo 3, I-56127 Pisa, Italy*

^b*INFN sezione di Pisa,
 Largo Pontecorvo 3, I-56127 Pisa, Italy*

^c*Dipartimento di Fisica dell'Università di Roma "La Sapienza",
 Piazzale Aldo Moro 5, I-00185 Roma, Italy*

^d*INFN Sezione di Roma La Sapienza,
 Piazzale Aldo Moro 5, I-00185 Roma, Italy*

^e*School of Physics and Astronomy, University of Southampton,
 SO17 1BJ Southampton, U.K.*

^f*Abdus Salam International Centre for Theoretical Physics,
 Strada Costiera 11, 34151, Trieste, Italy*

E-mail: claudio.bonati@df.unipi.it, massimo.delia@unipi.it,
mariti@df.unipi.it, guido.martinelli@roma1.infn.it,
mesiti@pi.infn.it, fnegro@pi.infn.it, f.sanfilippo@soton.ac.uk,
giovanni.villadoro@ictp.it

ABSTRACT: We investigate the topological properties of $N_f = 2 + 1$ QCD with physical quark masses, both at zero and finite temperature. We adopt stout improved staggered fermions and explore a range of lattice spacings $a \sim 0.05 - 0.12$ fm. At zero temperature we estimate both finite size and finite cut-off effects, comparing our continuum extrapolated results for the topological susceptibility χ with predictions from chiral perturbation theory. At finite temperature, we explore a region going from T_c up to around $4T_c$, where we provide continuum extrapolated results for the topological susceptibility and for the fourth moment of the topological charge distribution. While the latter converges to the dilute instanton gas prediction the former differs strongly both in the size and in the temperature dependence. This results in a shift of the axion dark matter window of almost one order of magnitude with respect to the instanton computation.

KEYWORDS: CP violation, Lattice QCD

ARXIV EPRINT: [1512.06746](https://arxiv.org/abs/1512.06746)

Contents

1	Introduction	1
1.1	General framework	2
1.2	Summary of main results and paper organization	4
2	Numerical setup	5
2.1	Discretization adopted	5
2.2	Determination of the topological content	5
3	Numerical results	7
3.1	Zero temperature	8
3.2	Finite temperature	11
4	Implications for axion phenomenology	18
5	Conclusions	20

1 Introduction

Axions are among the most interesting candidates for physics beyond the Standard Model. Their existence has been advocated long ago [1–4] as a solution to the so-called strong-CP problem through the Peccei-Quinn (PQ) mechanism. It was soon realized that they could also explain the observed dark matter abundance of the visible Universe [5–7]. However, a reliable computation of the axion relic density requires a quantitative estimate of the parameters entering the effective potential of the axion field, in particular its mass and self-couplings as a function of the temperature T of the thermal bath.

The purpose of this study is to obtain predictions from the numerical simulations of Quantum Chromodynamics (QCD) on a lattice. Our results, which are summarized at the end of this section, suggest a possible shift of the axion dark matter window by almost one order of magnitude with respect to instanton computations. This shift is a consequence of the much slower decrease of the axion mass with the temperature in comparison to the dilute instanton gas prediction. Our present simulations are however limited to a range of temperatures not exceeding 600 MeV: the main obstruction is represented by the freezing of the topological modes on fine lattices, which afflicts present lattice QCD algorithms. For a more complete understanding of axion dynamics at finite T , in the future a major effort must be undertaken to reach higher temperatures.

1.1 General framework

Given the strong bounds on its couplings, the axion field can be safely treated as a non-dynamical external field. Its potential is completely determined by the dependence of the QCD partition function on the θ -angle, which enters the pure gauge part of the QCD Euclidean Lagrangian as

$$\mathcal{L}_\theta = \frac{1}{4} F_{\mu\nu}^a(x) F_{\mu\nu}^a(x) - i\theta q(x), \tag{1.1}$$

where

$$q(x) = \frac{g^2}{64\pi^2} \epsilon_{\mu\nu\rho\sigma} F_{\mu\nu}^a(x) F_{\rho\sigma}^a(x) \tag{1.2}$$

is the topological charge density. The θ -dependent part of the free energy density can be parametrized as follows

$$\mathcal{F}(\theta, T) \equiv F(\theta, T) - F(0, T) = \frac{1}{2} \chi(T) \theta^2 s(\theta, T) \tag{1.3}$$

where $\chi(T)$ is the topological susceptibility at $\theta = 0$,

$$\chi = \int d^4x \langle q(x) q(0) \rangle_{\theta=0} = \frac{\langle Q^2 \rangle_{\theta=0}}{\mathcal{V}} \tag{1.4}$$

($Q = \int d^4x q(x)$ is the global topological charge and $\mathcal{V} = V/T$), while $s(\theta, T)$ is a dimensionless even function of θ such that $s(0, T) = 1$. The quadratic term in θ , $\chi(T)$, is proportional to the axion mass, while non-linear corrections in θ^2 , contained in $s(\theta, T)$, provide information about axion interactions. In particular, assuming analyticity around $\theta = 0$, $s(\theta, T)$ can be expanded as follows [8]

$$s(\theta, T) = 1 + b_2(T) \theta^2 + b_4(T) \theta^4 + \dots, \tag{1.5}$$

where the coefficients b_n are proportional to the cumulants of the topological charge distribution. For instance b_2 , which is related to quartic interactions terms in the axion potential, can be expressed as

$$b_2 = - \frac{\langle Q^4 \rangle_{\theta=0} - 3 \langle Q^2 \rangle_{\theta=0}^2}{12 \langle Q^2 \rangle_{\theta=0}}. \tag{1.6}$$

The function $\mathcal{F}(\theta, T)$, related to the topological properties of QCD, is of non-perturbative nature and hence not easy to predict reliably with analytic methods. This is possible only in some specific regimes: chiral perturbation theory (ChPT) represents a valid approach only in the low temperature phase; at high- T , instead, a possible analytic approach is the Dilute Instanton Gas Approximation (DIGA). DIGA predictions can in fact be classified in two groups: those that make only use of the DIGA hypothesis itself (i.e. that just relies on the existence of weakly interacting objects of topological charge one), and those that exploit also perturbation theory, the latter being expected to hold only at asymptotically high values of T . Using only the dilute gas approximation one can show that the θ -dependence of the free energy is of the form (see e.g. [9, 10])

$$F(\theta, T) - F(0, T) \simeq \chi(T) (1 - \cos \theta), \tag{1.7}$$

and thus $b_2^{\text{DIGA}} = -1/12$, $b_4^{\text{DIGA}} = 1/360$ and so on. Using also perturbation theory it is possible to obtain an explicit form for the dependence of the topological susceptibility on the temperature. To leading order, for $N_f^{(l)}$ light quark flavors of mass m_l , one obtains (see e.g. [9, 10])

$$\chi(T) \sim T^{4-\beta_0} \left(\frac{m_l}{T}\right)^{N_f^{(l)}}, \quad (1.8)$$

where $\beta_0 = 11N_c/3 - 2N_f/3$. Only part of the NLO corrections to this expression are known (see [11] or [12] for a summary, [13] for the $N_f = 0$ case).

As an alternative, a fully non-perturbative approach, which is based completely on the first principles of the theory, is represented by lattice QCD simulations. In fact, extensive studies have been carried out regarding the θ -dependence of pure gauge theories. It was shown in ref. [14], and later confirmed in refs. [13, 15, 16], that the form of the free energy in eq. (1.7) describes with high precision the physics of the system for $T \gtrsim 1.15 T_c$, while for $T \lesssim T_c$ everything is basically independent of the temperature, thus strengthening the conclusion $\chi(T < T_c) \approx \chi(T = 0)$ obtained in previous studies [17–21]. In refs. [13, 22] it was also shown that the temperature dependence of the topological susceptibility is correctly reproduced by eq. (1.8) for temperatures just above T_c , even if the overall normalization is about a factor ten larger than the perturbative prediction.

A realistic study of θ -dependence aimed at being relevant to axion phenomenology requires the numerical simulation of lattice QCD including dynamical quarks with physical masses. Apart from the usual computational burden involved in the numerical simulation of light quarks, that represents a challenge from at least two different but interrelated points of view.

Because of the strict connection, in the presence of light fermions, between the topological content of gauge configurations and the spectrum of the fermion matrix (in particular regarding the presence of zero modes), a reliable study of topological quantities requires a discretization of the theory in which the chiral properties of fermions fields are correctly implemented. For standard discretizations, such properties are recovered only for small enough lattice spacings, so that a careful investigation of the continuum limit becomes essential. Indeed only recently it was possible to measure the dependence of the topological susceptibility on the quark masses to a sufficient accuracy to be compared with the prediction of chiral perturbation theory [23–27].

On the other hand, it is well known that, as the continuum limit is approached, it becomes increasingly difficult to correctly sample the topological charge distribution, because of the large energy barriers developing between configurations belonging to different homotopy classes, i.e. to different topological sectors, which can be hardly crossed by standard algorithms [28–32]. That causes a loss of ergodicity which, in principle, can spoil any effort to approach the continuum limit itself.

Combining these two problems together, the fact that a proper window exists, in which the continuum limit can be taken and still topological modes can be correctly sampled by current state-of-the-art algorithms, is highly non-trivial. In a finite temperature study, since the equilibrium temperature is related to the inverse temporal extent of the lattice by $T = 1/(N_t a)$, the fact that one cannot explore arbitrarily small values of the lattice

spacing a , because of the above mentioned sampling problem, limits the range of explorable temperatures from above.

In the present study we show that, making use of current state-of-the-art algorithms, one can obtain continuum extrapolated results for the θ -dependence of QCD at the physical point, in a range of temperatures going up to about $4 T_c$, where $T_c \sim 155$ MeV is the pseudo-critical temperature at which chiral symmetry restoration takes place. Then we discuss the consequences of such results to axion phenomenology in a cosmological context.

Our investigation is based on the numerical simulation of $N_f = 2 + 1$ QCD, adopting a stout staggered fermions discretization with physical values of the quark masses and the tree level improved Symanzik gauge action. First, we consider simulations at zero temperature and various different values of the lattice spacing, in a range $\sim 0.05 - 0.12$ fm and staying on a line of constant physics, in order to identify a proper scaling window where the continuum limit can be taken without incurring in severe problems with the freezing of topological modes. Results are then successfully compared to the predictions of chiral perturbation theory. We also show that, for lattice spacings smaller than those explored by us, the freezing problem becomes severe, making the standard Rational Hybrid Monte-Carlo (RHMC) algorithm useless.

For a restricted set of lattice spacings, belonging to the scaling window mentioned above, we perform finite temperature simulations, obtaining continuum extrapolated results for χ and b_2 in a range of T going up to around 600 MeV. These results are then taken as an input to fix the parameters of the axion potential in the same range of temperatures and perform a phenomenological analysis.

1.2 Summary of main results and paper organization

Our main results are the following. Up to T_c the topological susceptibility is almost constant, and compatible with the prediction from ChPT. Above T_c the value of b_2 rapidly converges to what predicted by DIGA computations; on the contrary the dependence of the topological susceptibility on T shows significant deviations from DIGA. This has a significant impact on axion phenomenology, in particular it results in a shift of the axion dark matter window by almost one order of magnitude with respect to the instanton computation. Since the T -dependence is much milder than expected from instanton calculations, it becomes crucial, for future studies, to investigate the system for higher values of T , something which also claims for the inclusion of dynamical charm quarks and for improved algorithms, capable to defeat or at least alleviate the problem of the freezing of topological modes.

The paper is organized as follows. In section 2, we discuss the setup of our numerical simulations, in particular the lattice discretization adopted and the technique used to extract the topological content of gauge configurations. In section 3 we present our numerical analysis and continuum extrapolated result for the θ -dependence of the free energy density, both at zero and finite temperature. Section 4 is dedicated to the analysis of the consequences of our results in the context of axion cosmology. Finally, in section 5, we draw our conclusions and discuss future perspectives.

β	a [fm]	am_s
3.750	0.1249	5.03×10^{-2}
3.850	0.0989	3.94×10^{-2}
3.938	0.0824	3.30×10^{-2}
4.020	0.0707	2.81×10^{-2}
4.140	0.0572	2.24×10^{-2}

Table 1. Bare parameters used in this work, from [36, 37] or spline interpolation of data thereof. The systematic uncertainty on the lattice spacing determination is 2–3% and the light quark mass is fixed by using $m_s/m_l = 28.15$.

2 Numerical setup

2.1 Discretization adopted

The action of $N_f = 2 + 1$ QCD is discretized by using the tree level improved Symanzik gauge action [33, 34] and the stout improved staggered Dirac operator. Explicitly, the Euclidean partition function is written as

$$\mathcal{Z} = \int \mathcal{D}U e^{-S_{\text{YM}}} \prod_{f=u,d,s} \det \left(M_{\text{st}}^f[U] \right)^{1/4}, \quad (2.1)$$

$$S_{\text{YM}} = -\frac{\beta}{3} \sum_{i,\mu \neq \nu} \left(\frac{5}{6} W_{i;\mu\nu}^{1 \times 1} - \frac{1}{12} W_{i;\mu\nu}^{1 \times 2} \right), \quad (2.2)$$

$$(M_{\text{st}}^f)_{i,j} = am_f \delta_{i,j} + \sum_{\nu=1}^4 \frac{\eta_{i;\nu}}{2} \left[U_{i;\nu}^{(2)} \delta_{i,j-\hat{\nu}} - U_{i-\hat{\nu};\nu}^{(2)\dagger} \delta_{i,j+\hat{\nu}} \right]. \quad (2.3)$$

The symbol $W_{i;\mu\nu}^{n \times m}$ denotes the trace of the $n \times m$ Wilson loop built using the gauge links departing from the site in position i along the positive μ, ν directions. The gauge matrices $U_{i;\mu}^{(2)}$, used in the definition of the staggered Dirac operator, are related to the gauge links $U_{i;\mu}$ (used in S_{YM}) by two levels of stout-smearing [35] with isotropic smearing parameter $\rho = 0.15$.

The bare parameters β , m_s and $m_l \equiv m_u = m_d$ were chosen in such a way to have physical pion mass $m_\pi \approx 135$ MeV and physical m_s/m_l ratio. The values of the bare parameters used in this study to move along this line of constant physics are reported in table 1. Most of them were determined in [36, 37], the remaining have been extrapolated by using a cubic spline interpolation. The lattice spacings reported in table 1 have a 2–3% of systematic uncertainty, as discussed in [36, 37] (see also [38]).

2.2 Determination of the topological content

In order to expose the topological content of the gauge configurations, we adopt a gluonic definition of the topological charge density, measured after a proper smoothing procedure, which has been shown to provide results equivalent to definitions based on fermionic operators [39–43]. The basic underlying idea is that, close enough to the continuum limit,

the topological content of gauge configurations becomes effectively stable under a local minimization of the gauge action, while ultraviolet fluctuations are smoothed off.

A number of smoothing algorithms has been devised along the time. A well known procedure is cooling [44–48]: an elementary cooling step consists in the substitution of each link variable $U_{i; \mu}$ with the $SU(3)$ matrix that minimizes the Wilson action, in order to reduce the UV noise. While for the case of the $SU(2)$ group this minimization can be performed analytically, in the $SU(3)$ case the minimization is usually performed *à la* Cabibbo-Marinari, i.e. by iteratively minimizing on the $SU(2)$ subgroups. When this elementary cooling step is subsequently applied to all the lattice links we obtain a cooling step.

Possible alternatives may consist in choosing a different action to be minimized during the smoothing procedure, or in performing a continuous integration of the minimization equations. The latter procedure is known as the gradient flow [49, 50] and has been shown to provide results equivalent to cooling regarding topology [51–54]. Because of its computational simplicity in this work we will thus use cooling, however it will be interesting to consider in future studies also the gradient flow, especially as an independent way to fix the physical scale [50, 55, 56].

Since the topological charge will be measured on smoothed configurations, we can use the simplest discretization of the topological charge density with definite parity [57]

$$q_L(x) = -\frac{1}{29\pi^2} \sum_{\mu\nu\rho\sigma=\pm 1}^{\pm 4} \tilde{\epsilon}_{\mu\nu\rho\sigma} \text{Tr} (U_{x; \mu\nu} U_{x; \rho\sigma}) , \quad (2.4)$$

where $U_{x; \mu\nu}$ is the plaquette located in x and directed along the μ, ν directions. The tensor $\tilde{\epsilon}$ is the completely antisymmetric tensor which coincides with the usual Levi-Civita tensor $\epsilon_{\mu\nu\rho\sigma}$ for positive indices and is defined by $\tilde{\epsilon}_{\mu\nu\rho\sigma} = -\tilde{\epsilon}_{(-\mu)\nu\rho\sigma}$ and antisymmetry for negative indices.

The lattice topological charge $Q_L = \sum_x q_L(x)$ is not in general an integer, although its distribution gets more and more peaked around integer values as the continuum limit is approached. In order to assign to a given configuration an integer value of the topological charge we will follow the prescription introduced in [29]: the topological charge is defined as $Q = \text{round}(\alpha Q_L)$, where ‘round’ denotes the rounding to the closest integer and α is fixed by the minimization of

$$\left\langle (\alpha Q_L - \text{round} [\alpha Q_L])^2 \right\rangle , \quad (2.5)$$

i.e. in such a way that αQ_L is ‘as integer’ as possible. Actually, one could take the non-rounded definition $Q = Q_L$ as well, the only difference being a different convergence of results to the common continuum limit (see ref. [58] for a more detailed discussion on this point). The topological susceptibility is then defined by

$$a^4 \chi = \frac{\langle Q^2 \rangle}{V_4} , \quad (2.6)$$

where $V_4 = N_t N_s^3$ is the four-dimensional volume of the lattice, and the coefficient b_2 has been introduced in eq. (1.6).

We measured the topological charge every 20 cooling steps up to a maximum of 120 steps and verified the stability of the topological quantities under smoothing. Results that will be presented in the following have been obtained by using 100 cooling steps, a number large enough to clearly identify the different topological sectors but for which no significant signals of tunneling to the trivial sector are distinguishable in mean values.

For all data reported in the following we verified that the corresponding time histories were long enough to correctly sample the topological charge. In particular we checked that $\langle Q \rangle$ is compatible with zero and that the topological charge is not frozen. Indeed it is well known that, while approaching the continuum limit, the autocorrelation time of the topological charge increases very steeply until no tunneling events between different sectors happen anymore [28–32]. An example of this behavior can be observed in figure 1, where some time-histories for zero temperature runs are showed, for three different lattice spacings. While the general features of this phenomenon are common to all the lattice actions, the critical value of the lattice spacing at which the charge gets stuck may depend on the specific discretization adopted. In our case we were able to obtain reasonable sampling for lattice spacings down to $a = 0.0572$ fm, while finer lattices (in particular, with $a \lesssim 0.04$ fm) showed severe freezing over thousands of trajectories and had to be discarded.

3 Numerical results

The main purpose of our numerical study is to provide results for the θ -dependence of QCD at finite temperature, in particular above the pseudo-critical temperature, in order to take them as an input for axion phenomenology. However, in order to make sure that our lattice discretization is accurate enough to reproduce the chiral properties of light fermions, we have also performed numerical simulations at zero or low temperature, where results can be compared to reliable analytic predictions.

Indeed, at zero temperature, the full θ dependence of the QCD partition function, can be estimated reliably using chiral Lagrangians [4, 59, 60]. In particular, at leading order in the expansion, χ and b_2 can be written just in terms of m_π , f_π and $z = m_u/m_d$ as

$$\chi^{\text{LO}} = \frac{z}{(1+z)^2} m_\pi^2 f_\pi^2, \quad b_2^{\text{LO}} = -\frac{1}{12} \frac{1+z^3}{(1+z)^3}. \quad (3.1)$$

NLO corrections have been computed in [61–63] and are of the order of percent for physical values of the light quark masses. Using the estimate in [63] we have $\chi^{1/4} = 75.5(5)$ MeV and $b_2 = -0.029(2)$ for $z = 0.48(3)$, while $\chi^{1/4} = 77.8(4)$ MeV and $b_2 = -0.022(1)$ for $z = 1$ used here. Note that for pure SU(3) Yang-Mills these numbers become $\chi_{N_f=0}^{1/4} \approx 180$ MeV [17, 29, 64–68] and¹ $b_{2,N_f=0} = -0.0216(15)$ [58] (see also refs. [29, 68–71] for previous determinations in the literature).

¹It is interesting to note the striking agreement between the ChPT prediction for b_2 in the case of degenerate light flavors and the numerical results obtained for it in the quenched theory [58]. That seems to suggest a similar form of θ -dependence in the low temperature phases of the full and of the quenched theory. At this stage we have no particular explanation for this agreement, which might be considered as a coincidence.




Figure 1. Check for the correct sampling of the topological charge. (Upper left) Topological charge time history and histogram for $a = 0.0824$ fm on a 32^4 lattice. (Upper right) Topological charge time history and histogram for $a = 0.0572$ fm on a 48^4 lattice. (Lower) Topological charge time history for $a = 0.0397$ fm on a 40^4 lattice; in this case the run was stopped earlier because of the visible freezing of the topological charge. Note that the upper histories correspond to approximately the same physical volume: hence the visible shrink in the topological charge distribution is the actual effect of the approach to the continuum limit (see the discussion in section 3).

ChPT provides a prediction at finite temperature as well. In particular we have

$$\begin{aligned}
 \chi_{\text{ChPT}}(T) &= \chi(0) \left(1 - \frac{3 T^2}{2 f_\pi^2} J_1 \left[\frac{T^2}{m_\pi^2} \right] \right), \\
 b_2^{\text{ChPT}}(T) &= b_2(0) \left(1 + \frac{9 m_\pi^2}{2 f_\pi^2} \frac{z(1+z)}{1+z^3} J_2 \left[\frac{T^2}{m_\pi^2} \right] \right),
 \end{aligned}
 \tag{3.2}$$

where the functions J_n are defined in ref. [63]. However, at temperatures around and above T_c , the chiral condensate drops and chiral Lagrangians break down, so that the finite T predictions in eq. (3.2) are expected to fail. In this regime non-perturbative computations based on first principle QCD are mandatory.

3.1 Zero temperature

In table 2 we report our determinations of the topological susceptibility on N_s^4 lattices for several values of N_s and different lattice spacings. The results on the three smallest lattice spacings are also plotted in figure 2.




Figure 2. Volume dependence of the topological susceptibility for different lattice spacings. The lattice extent is reported in physical units, while χ in lattice spacing units. Bands represent best fits to eq. (3.3) for the infinite volume limit extrapolation.

In order to extract the infinite volume limit of χ at fixed lattice spacing, one can either consider only results obtained on the largest available lattices and fit them to a constant, or try to model the behavior of χ with the lattice size using all available data. We have followed both procedures in order to better estimate possible systematics.

The topological susceptibility can be written as the integral of the two point function of the topological charge density; since the η' is the lightest intermediate state that significantly contributes to this two point function (three pions states are OZI suppressed), one expects the leading asymptotic behavior to be

$$\chi_{N_s} \sim \chi_\infty + C e^{-aN_s m_{\eta'}}, \tag{3.3}$$

where C is an unknown constant. This form nicely fits data in the whole available range of aN_s (see figure 2), both when $m_{\eta'}$ is fixed to its physical value and when it is left as a fit parameter. The results obtained using this last procedure (which gives the most conservative estimates) are reported in table 2 and correspond to the entries denoted by $N_s = \infty$. On the other hand, a fit to a constant value works well when using only data coming from lattices with $aN_s > 1.5$ fm, and provides consistent results within errors. This analysis makes also us confident that results obtained on the lattices with $a \simeq 0.1249$ fm and $a \simeq 0.0989$ fm, where a single spatial volume is available with $aN_s > 3$ fm, are not affected by significant finite size effects.

In figure 3 we plot $\chi^{1/4}$, extrapolated to infinite volume, against the square of the lattice spacing, together with the ChPT prediction. Finite cut-off effects are significant, especially for $a \gtrsim 0.1$ fm, meaning that we are not close enough to the continuum limit to reproduce the correct chiral properties of light quarks. In the case of staggered quarks, such lattice artifacts originate mostly from the fact that the full chiral symmetry group is reproduced only in the continuum limit, so that the pion spectrum is composed of a light pseudo-Goldstone boson and other massive states which become degenerate only as




Figure 3. Continuum limit of the topological susceptibility: our final result is $\chi^{1/4} = 73(9)$ MeV. Also shown is the value expected from chiral perturbation theory, denoted by the horizontal dashed line, together with the quenched value. Square points correspond to the combination $\chi_{tc}^{1/4}(a)$ defined in eq. (3.6).

$a \rightarrow 0$. The physical signal vanishes in the chiral limit whereas this is not the case for its discretization effects. This means that it is necessary to work on very fine lattice spacings in order to keep these effects under control, a task which is particularly challenging with present algorithms, because of the freezing of the topological charge.

Despite the large cut-off effects, we can perform the continuum extrapolation of our data. If one considers only the three finest lattice spacings, the finite cut-off effects for this quantity can be parametrized as simple $O(a^2)$ corrections

$$\chi^{1/4}(a) = \chi^{1/4} + Ba^2 \tag{3.4}$$

and a best fit yields $\chi^{1/4} = 73(9)$ MeV, while in order to describe the whole range of available data one must take into account $O(a^4)$ corrections as well, i.e.

$$\chi^{1/4}(a) = \chi^{1/4} + Ba^2 + Ca^4, \tag{3.5}$$

obtaining $\chi^{1/4} = 69(9)$ MeV. Both best fits are reported in figure 3. Therefore we conclude that the continuum extrapolation is already under control with the available lattice data, and is in satisfactory agreement with the predictions of chiral perturbation theory.

In order to further inquire about the reliability of our continuum extrapolation and the importance of the partial breaking of chiral symmetry in the staggered discretization, we studied the combination

$$\chi_{tc}^{1/4}(a) = a\chi^{1/4}(a) \frac{m_\pi^{\text{phys}}}{am_{ngb}(a)}. \tag{3.6}$$

Here $m_{ngb}(a)$ is the mass of one of the non-Goldstone pions, i.e. of a state that becomes massless in the chiral limit only if the continuum limit is performed; since in the continuum

N_s	a [fm]	χa^4
32	0.1249	$8.55(32) \times 10^{-5}$
32	0.0989	$2.22(10) \times 10^{-5}$
12	0.0824	$4.14(20) \times 10^{-6}$
16	0.0824	$6.60(45) \times 10^{-6}$
20	0.0824	$7.17(36) \times 10^{-6}$
24	0.0824	$8.21(70) \times 10^{-6}$
32	0.0824	$7.32(39) \times 10^{-6}$
∞	0.0824	$7.50(40) \times 10^{-6}$
16	0.0707	$1.57(16) \times 10^{-6}$
20	0.0707	$2.06(15) \times 10^{-6}$
24	0.0707	$2.42(16) \times 10^{-6}$
32	0.0707	$2.41(19) \times 10^{-6}$
40	0.0707	$3.29(56) \times 10^{-6}$
∞	0.0707	$2.60(36) \times 10^{-6}$
20	0.0572	$4.34(90) \times 10^{-7}$
24	0.0572	$5.40(76) \times 10^{-7}$
32	0.0572	$4.7(1.2) \times 10^{-7}$
40	0.0572	$6.61(96) \times 10^{-7}$
48	0.0572	$6.88(87) \times 10^{-7}$
∞	0.0572	$6.60(82) \times 10^{-7}$

Table 2. Topological susceptibility in lattice units, measured on several N_s^4 lattices for different values of the lattice spacing. Values denoted by $N_s = \infty$ correspond to the thermodynamical extrapolation and they are obtained by a fit to exponential without fixing $m_{\eta'}$ to its physical value, see eq. (3.3).

$m_{n_{gb}}(a) \rightarrow m_{\pi}^{\text{phys}}$, this ratio converges to $\chi^{1/4}$ as $a \rightarrow 0$. The state with taste structure $\gamma_i \gamma_{\mu}$ was used, whose mass is close to the root mean square value of all the taste states (see ref. [37] figure 2) and the values of $\chi_{tc}^{1/4}(a)$ are shown in figure 3 as square points. It is clear that the combination $\chi_{tc}^{1/4}(a)$ strongly reduces lattice artefacts with respect to $\chi^{1/4}(a)$, moreover a linear fit in a^2 well describes data for all available lattice spacings, giving the result $\chi^{1/4} = 77(3)$. Although a complete study of the systematics affecting $\chi_{tc}^{1/4}(a)$ was not performed (e.g. the dependence of $m_{n_{gb}}(a)$ on the lattice size was not studied, just the largest size was used), this is a strong indication that the dominant source of lattice artefacts in $\chi^{1/4}(a)$ is the chiral symmetry breaking present at finite lattice spacing in the staggered discretization.

3.2 Finite temperature

Finite temperature simulations can in principle be affected by lattice artifacts comparable to those present at $T = 0$. For that reason, we have limited our finite T simulations to the three smallest lattice spacings explored at $T = 0$, i.e. those in the scaling window adopted

N_t	a [fm]	T/T_c	$\chi(T, a)a^4$	$\frac{\chi(T, a)}{\chi(0, a)}$	b_2
16	0.0824	0.964	$6.12(48) \cdot 10^{-6}$	0.817(75)	-0.23(33)
14	0.0824	1.102	$5.25(47) \cdot 10^{-6}$	0.701(70)	-0.191(95)
12	0.0824	1.285	$4.04(11) \cdot 10^{-6}$	0.539(29)	-0.003(25)
10	0.0824	1.542	$2.87(12) \cdot 10^{-6}$	0.383(23)	-0.083(29)
8	0.0824	1.928	$1.296(97) \cdot 10^{-6}$	0.173(15)	-0.065(25)
24	0.0707	0.749	$2.82(36) \cdot 10^{-6}$	1.08(18)	-0.11(12)
16	0.0707	1.124	$1.53(15) \cdot 10^{-6}$	0.589(84)	-0.057(43)
14	0.0707	1.284	$1.385(79) \cdot 10^{-6}$	0.532(62)	-0.019(30)
12	0.0707	1.498	$1.05(11) \cdot 10^{-6}$	0.403(60)	-0.101(21)
10	0.0707	1.798	$7.87(87) \cdot 10^{-7}$	0.302(46)	-0.096(26)
8	0.0707	2.247	$3.41(31) \cdot 10^{-7}$	0.131(18)	-0.077(18)
6	0.0707	2.996	$1.22(21) \cdot 10^{-7}$	0.0469(94)	-0.076(10)
16	0.0572	1.389	$2.92(41) \cdot 10^{-7}$	0.41(16)	-0.049(17)
14	0.0572	1.587	$2.04(32) \cdot 10^{-7}$	0.29(11)	-0.058(16)
12	0.0572	1.852	$1.43(19) \cdot 10^{-7}$	0.202(79)	-0.0626(90)
10	0.0572	2.222	$8.3(1.4) \cdot 10^{-8}$	0.118(48)	-0.0705(74)
8	0.0572	2.777	$4.85(86) \cdot 10^{-8}$	0.068(27)	-0.091(11)
6	0.0572	3.703	$2.51(72) \cdot 10^{-8}$	0.035(16)	-0.0792(12)

Table 3. Finite temperature results; in all cases $N_s = 48$ was used.

for the extrapolation to the continuum limit with only $O(a^2)$ corrections. At fixed lattice spacing the temperature has been varied by changing the temporal extent N_t of the lattice and, in all cases, we have fixed $N_s = 48$. This gives a spatial extent equal or larger than those explored at $T = 0$ and an aspect ratio $N_s/N_t \geq 3$ for all explored values of N_t . The absence of significant finite volume effects has also been verified directly by comparing results with those obtained on $N_t \times 40^3$ lattices. In table 3 we report the numerical values obtained for the topological susceptibility, for the ratio $\chi(T, a)/\chi(T = 0, a)$ (where for $\chi(T = 0, a)$ the infinite volume extrapolation has been taken) and for the cumulant ratio b_2 .

Results for $\chi^{1/4}$ as a function of T/T_c , where $T_c = 155$ MeV, are reported in figure 4 for the three different lattice spacings. The dependence on a is quite strong, as expected from the $T = 0$ case. Inspired by the instanton gas prediction, eq. (1.8), we have performed a fit with the following ansatz

$$\chi^{1/4}(a, T) = A_0(1 + A_1 a^2) \left(\frac{T}{T_c}\right)^{A_2}, \tag{3.7}$$

which also takes into account the dependence on a and nicely describes all data in the range $T > 1.2T_c$ with $\chi^2/\text{dof} \simeq 0.7$. In table 4 we report the best fit values obtained performing the fit in the region $T > T_{\text{cut}}$ for some T_{cut} values; best fit curves are reported in figure 4 as well, together with a band corresponding to the continuum extrapolation.

T_{cut}/T_c	A_0 (MeV)	$A_1(\text{fm}^{-2})$	A_2
1.2	76.0(5.1)	103(17)	-0.671(40)
1.4	80.2(6.9)	100(20)	-0.728(57)
1.6	86(10)	85(24)	-0.750(80)
2.0	83(18)	97(53)	-0.718(14)

Table 4. Best fit values for the coefficients in (3.7) obtained by fitting only data corresponding to temperatures $T > T_{\text{cut}}$.

T_{cut}/T_c	D_0	$D_1(\text{fm}^{-2})$	D_2
1.2	1.17(24)	-2(28)	-2.71(15)
1.4	1.56(39)	-12(30)	-3.02(23)
1.6	1.81(76)	-33(32)	-2.99(39)
2.0	1.8(1.5)	$2(22) \times 10^3$	-2.90(65)

Table 5. Best fit values for the coefficients in (3.8) obtained by fitting only data corresponding to temperatures $T > T_{\text{cut}}$.




Figure 4. Continuum extrapolation of the fourth root topological susceptibility, using the function $\chi^{1/4}(a, T) = A_0(1 + A_1 a^2)(T/T_c)^{A_2}$. Only data corresponding to $T \gtrsim 1.2T_c$ have been used in the fit.

It is remarkable that most of the lattice artifacts disappear when one considers, in place of χ itself, the ratio $\chi(T, a)/\chi(T = 0, a)$, whose dependence on the lattice spacing is indeed quite mild. That is clearly visible in figure 5. Also in this case we have adopted a fit ansatz similar to eq. (3.7)

$$\frac{\chi(a, T)}{\chi(a, T = 0)} = D_0(1 + D_1 a^2) \left(\frac{T}{T_c}\right)^{D_2}, \quad (3.8)$$

which well fits all data in the range $T > 1.2T_c$ with a $\chi^2/\text{dof} \simeq 1$. The best fit curves and the continuum extrapolation are shown in figure 5. The best fit parameters, again for different T_{cut} values, are reported in table 5. It is important to note, in order to assess the




Figure 5. Ratio of the topological susceptibilities $\chi(T)/\chi(T = 0)$, evaluated at fixed lattice spacing. The horizontal solid line describes trivial scaling $\chi(T) = \chi(T = 0)$, while the dashed line is the prediction from finite temperature ChPT and the dashed-dotted line shows the slope predicted by DIGA computation. The band corresponds to the continuum extrapolation using the function $\chi(a, T)/\chi(a, T = 0) = D_0(1 + D_1 a^2)(T/T_c)^{D_2}$, only data corresponding to $T \gtrsim 1.2T_c$ have been used in the fit.




Figure 6. Plot of the exponent of the power-law behavior of $\chi(T)$, extracted from fit to eqs. (3.7) and (3.8) for $T > T_{\text{cut}}$ (see tables 4–5). The value predicted by the dilute instanton gas approximation and perturbation theory is $4A_2, D_2 \sim -8$.

reliability of our continuum limit, that the two different extrapolations, eqs. (3.7) and (3.8), provide perfectly consistent results: that can be appreciated both from figure 6, where we compare the two coefficients D_2 and $4A_2$, and from figure 7, where we directly compare the two continuum extrapolations. As a further check, we have also verified that by fitting $[\chi(a, T)/\chi(a, T = 0)]^{1/4}$ we obtain results in perfect agreement with the ones obtained by fitting $\chi(a, T)/\chi(a, T = 0)$. As an example, using in the fit the values corresponding to




Figure 7. Comparison of the two continuum extrapolations for $\chi(T)^{1/4}$. Continuum extrapolated results for $\chi(T)/\chi(0)$ have been multiplied by the ChPT prediction for $\chi(0)$.

$T > 1.2T_c$ one obtains for the exponent the value $-0.674(38)$, to be compared with $D_2/4$ from the first line of table 5 and A_2 from the first line of table 4. However in the following analysis we will refer to results obtained through the ratio $\chi(T)/\chi(0)$, which is the quantity showing smaller finite cut-off corrections.

Let us now comment on our results for the topological susceptibility. For temperatures below or around T_c , the temperature dependence is quite mild and, for temperatures up to $T \simeq 1.2T_c$, even compatible with the prediction from ChPT, which is reported in figure 5 for comparison. Then, for higher values of T , a sharp power law drop starts. However, it is remarkable that the power law exponent is smaller, by more than a factor two, with respect to the instanton gas computation, eq. (1.8), which predicts $D_2 \simeq -8$ in the case of three light flavors (the dependence on the number of flavor is however quite mild). DIGA is expected to provide the correct result in a region of asymptotically large temperatures, which however seems to be quite far from the range explored in the present study, which goes up to $T \simeq 4T_c$.

This is in sharp contrast with the quenched theory, where the DIGA power law behavior sets in at lower temperatures [13, 22]. In order to allow for a direct comparison, we have reported the DIGA prediction in figure 5, after fixing it by imposing $\chi(T_c) = \chi(0)$ as an overall normalization. As we discuss in the following section, the much milder drop of χ as a function of T has important consequences for axion cosmology.

A determination of the topological susceptibility has been presented in ref. [72], based on the Domain Wall discretization, in the range $T/T_c \sim 0.9-1.2$. Since the data have been produced with different lattice spacings at different temperatures, it is however difficult to compare their results with ours. An extended range of T has been explored in ref. [73] in the presence of twisted mass Wilson fermions, reporting a behavior similar to that observed in this study, although with larger values of the quark masses (corresponding to a pion mass $m_\pi \sim 370$ MeV). The comparison is performed in figure 8, and shows a reasonable




Figure 8. Comparison of our continuum extrapolated $\chi^{1/4}(T)$ (band) with results from ref. [73]. Data for $\chi(T)$ obtained at unphysical quark masses have been rescaled according to the DIGA relation $\chi(T) \sim m_q^2 \sim m_\pi^4$.

agreement if results from ref. [73] are rescaled according to the mass dependence expected from DIGA,² i.e. $\chi(T) \sim m_q^2 \sim m_\pi^4$.

Let us now turn to a discussion of our results for the coefficient b_2 (defined in eq. (1.5)), which is related to the non-gaussianity of the topological charge distribution and gives information about the shape of the θ -dependent part of the free energy density. Data for b_2 are reported in figure 9. For $T < T_c$ this observable is too noisy to give sensible results but a reasonable guess, motivated by what happens in the quenched case and by the ChPT prediction, is that b_2 is almost temperature independent for $T \lesssim T_c$, like the topological susceptibility. In the high temperature region b_2 can instead be measured with reasonable accuracy and, due to the peculiar dependence of the noise on this observable on $\chi\mathcal{V}$, data on the smaller lattice spacing turned out to be significantly more precise than the others (see the discussion in ref. [58]). While data clearly approach the dilute instanton gas prediction $b_2^{\text{DIGA}} = -1/12$ at high temperature, deviations from this value are clearly visible on all the lattice spacings for $T \simeq 1.3 T_c$ and, for the smallest lattice spacing, also up to $T \sim 2.5 T_c$.

This is in striking contrast to what is observed in pure gauge theory, where deviations from $-1/12$ are practically absent for $T \gtrsim 1.15 T_c$, with a precision higher than 10% [13–16]. As discussed in the introduction deviations from b_2^{DIGA} cannot be simply ascribed to a failure of perturbation theory (like e.g. for the behavior of $\chi(T)$) but are instead unambiguous indications of interaction between instantons. Another difference with respect to the quenched case is that in the pure gauge theory (with both gauge groups

²One might wonder why DIGA should work for the mass dependence of χ and not for its dependence on T . A possible explanation is that, while the temperature dependence stems from a perturbative computation, the mass dependence comes from the existence of isolated zero modes in the Dirac operator, i.e. from the very hypothesis of the existence of a dilute gas of instantons, which seems to be verified already at moderately low values of T , see the following discussion on b_2 , but not at $T \sim T_c$.




Figure 9. b_2 evaluated at three lattice spacing. The horizontal solid lines correspond to the value of b_2 at $T = 0$ predicted by ChPT (which is about -0.022) and to the instanton-gas expected value $b_2^{\text{DIGA}} = -1/12$. The dotted-dashed line is the prediction of finite temperature ChPT, while the light blue band is the result of a fit to the smallest lattice spacing data using eq. (3.12) with d_2 independent of T .

SU(N) and G_2 , see [14, 15]) the asymptotic value is approached from below, while in the present case it is approached from above. These peculiar features can be related to a different interaction between instantons mediated by light quarks, as it is clear from the following discussion.

To describe deviations from the instanton gas behavior it is convenient to use the parametrization of $\mathcal{F}(\theta, T)$ introduced in ref. [14]:

$$\mathcal{F}(\theta, T) = \sum_{n>0} c_{2(n-1)}(T) \sin^{2n}(\theta/2) . \tag{3.9}$$

Indeed it is not difficult to show that every even function of θ of period 2π can be written in this form, and the main advantage of this form is that the value of coefficient c_{2n} influences only b_{2j} with $j \geq n$: in particular

$$\begin{aligned} \chi(T) &= c_0(T)/2; & b_2(T) &= -\frac{1}{12} + \frac{c_2(T)}{8\chi(T)}; \\ b_4(T) &= \frac{1}{360} - \frac{c_2(T)}{48\chi(T)} + \frac{c_4(T)}{32\chi(T)} . \end{aligned} \tag{3.10}$$

Since the coefficients c_{2n} parametrize deviations from the instanton gas that manifest themselves only in higher-cumulants of the topological charge, it is natural to interpret eq. (3.9) as a virial expansion, in which the role of the “density” is played by the first coefficient (i.e. the topological susceptibility χ), and to introduce the dimensionless coefficients d_{2n} by

$$c_{2(n-1)}(T) = d_{2(n-1)}(T) \frac{\chi(T)^n}{\chi(T=0)^{n-1}} , \tag{3.11}$$

where $\chi(T = 0)$ was used just as a dimensional normalization and one expects a mild dependence of $d_{2(n-1)}(T)$ on the temperature, since the strongly dependent component

$\chi(T)^n$ have been factorized. To lowest order we thus have

$$b_2(T) = -\frac{1}{12} + \frac{d_2}{8} \frac{\chi(T)}{\chi(T=0)}, \quad (3.12)$$

which nicely describes the b_2 data for the smallest lattice spacing using $d_2 = 0.80(16)$, see figure 9, where the expression $\chi(T)/\chi(T=0) \simeq (T_c/T)^{2.7}$ was used.

In the spirit of a virial expansion interpretation of eq. (3.9), the coefficient d_2 can be considered as the lowest order interaction term between instantons. In particular, a positive value of d_2 corresponds to an attractive potential, which is in sharp contrast with the pure gauge case, where a repulsive, negative value of d_2 is observed. This peculiar difference can be surely interpreted in terms of effective instanton interactions mediated by light quarks, which are likely also responsible for the much slower convergence, with respect to the quenched case, to the DIGA prediction.

4 Implications for axion phenomenology

The big departure of the results for the topological susceptibility at finite temperature from the DIGA prediction has a strong impact on the computation of the axion relic abundance. In particular the model independent contribution from the misalignment mechanism is determined to be [63]

$$\Omega_a^{\text{mis}} = \frac{86}{33} \frac{\Omega_\gamma}{T_\gamma} \frac{n_a^*}{s^*} m_a, \quad (4.1)$$

where Ω_γ and T_γ are the present abundance and temperature of photons while n_a^*/s^* is the ratio between the comoving number density $n_a = \langle m_a a^2 \rangle$ and the entropy density s computed at a late time t_* such that the ratio n_a/s became constant. The number density n_a can be extracted by solving the axion equation of motion

$$\ddot{a} + 3H\dot{a} + V'(a) = 0, \quad (4.2)$$

The temperature (and time) dependence of the Hubble parameter H is determined by the Friedmann equations and the QCD equation of state. The biggest uncertainties come therefore from the temperature dependence of the axion potential $V(a)$. At high temperatures the Hubble friction dominates over the vanishing potential and the field is frozen to its initial value a_0 . As the Universe cools the pull from the potential starts winning over the friction (this happens when $T \approx T_{\text{osc}}$, defined as $m_a(T_{\text{osc}}) \approx 3H(T_{\text{osc}})$) and the axion starts oscillating around the minimum. Shortly after H becomes negligible and the mass term is the leading scale in eq. (4.2). In this regime the approximate WKB solution has the form

$$\begin{aligned} a(t) &\sim A(t) \cos\left(\int_{t_0}^t dt' m_a(t')\right) = \\ &= a(t_0) \left(\frac{m_a(t_0)R^3(t_0)}{m_a(t)R^3(t)}\right)^{1/2} \cos\left(\int_{t_0}^t dt' m_a(t')\right), \end{aligned} \quad (4.3)$$

where $R(t)$ is the cosmic scale factor. Since the energy density is given by $\rho_a \sim m_a^2 A^2/2$, the solution (4.3) implies that what is conserved in the comoving volume is not the energy




Figure 10. Values of the axion decay constant f_a as a function of the initial field value $\theta_0 = a_0/f_a$ such that the axion misalignment contribution matches the full or a tenth of the observed dark matter abundance (red band or dotted green line respectively). When the PQ symmetry is broken only after inflation the axion abundance is reproduced by choosing $\theta_0 \approx 2.2$, i.e. the vertical blue dashed line.

density but $N_a = \rho_a R^3/m_a$, which can be interpreted as the number of axions [5–7]. Through the conservation of the comoving entropy S , it follows that n_a^*/s^* becomes an adiabatic invariant. Hence, it is enough to integrate the equation of motion (4.2) in the small window around the time when $T \approx T_{\text{osc}}$. We integrated numerically eq. (4.2) in the interval between the time when $m_a = H/10$ to that corresponding to $m_a = 2400H$ and extract the ratio n_a^*/s^* when $m_a \sim 300H$, namely a factor a hundred since the oscillation regime begins. The value for T_{osc} varies from T_c to several GeV depending on the axion decay parameter f_a and the temperature dependence of the axion potential. More details about this standard computation can be found for example in [63, 74]. In order to estimate the uncertainty in the results given below we varied the fitting parameters of the topological susceptibility D_2, D_0 and those relative to the QCD equation of state [37] within the quoted statistical and systematic errors.

Given that $b_2(T)$ converges relatively fast to the value predicted by a single cosine potential, we can assume $V(a) = -\chi(T) \cos(a/f_a)$ for $T \gtrsim T_c$. Using the most conservative results for the fit of $\chi(T)$, i.e. $\chi(T)/\chi(0) = (1.8 \pm 1.5)(T_c/T)^{2.90 \pm 0.65}$, in figure 10 we plot the prediction for the parameter f_a as a function of the initial value of the axion field $\theta_0 = a_0/f_a$ assuming that the misalignment axion contribution make up for the whole observed dark matter abundance, $\Omega_{\text{DM}} = 0.259(4)$ [75]. We also plot the case where the axion misalignment contribution accounts only for part (10% for definiteness) of the dark matter abundance.

In some cases the axion field acquires all possible values within the visible horizon, therefore the initial condition to the eq. (4.2) needs to be integrated over. This happens if the PQ symmetry is broken only after inflation or if the PQ symmetric phase is temporarily restored after inflation (e.g. if the Hubble scale during inflation or the maximum

temperature at reheating are at or above the PQ scale f_a). Numerically it corresponds to choosing $\theta_0 \approx 2.2$ and the value of the PQ scale in this case is fixed to:

$$f_a(\theta_0) = (1.00_{-0.26}^{+0.40+0.07} \pm 0.06) \cdot 10^{12} \text{ GeV}, \quad (4.4)$$

where the errors correspond respectively to the uncertainties on the fit parameters D_2 , D_0 , and to the error in the QCD equation of state. Note that, in this case, also other contributions from topological defects are expected to contribute. In case their effects are not negligible, or if axions are not responsible for the whole dark matter abundance, the value above should just be read as an upper bound to the PQ scale.

The value in eq. (4.4) is almost one order of magnitude bigger than the one computed with instantons techniques ($f_a \lesssim 0.2 \cdot 10^{12} \text{ GeV}$), which in fact corresponds to the value we instead get for one tenth of the dark matter abundance.

Some few remarks are in order. While the uncertainties on the axion mass fit we used are not small, the final axion abundance is rather insensitive to them and the prediction for f_a has therefore a good precision. The results above, however, rely on the extrapolation of the axion mass fit formula up to few GeV, where no lattice data is available. In particular for the value of f_a in eq. (4.4) the axion field starts oscillating around $T_{\text{osc}} = 4.3 \text{ GeV}$. An even longer extrapolation is required for $f_a = 1.67 \cdot 10^{11} \text{ GeV}$, corresponding to $\Omega_a^{\text{mis}} = 0.1 \Omega_{\text{DM}}$, where the axion starts oscillating around $T_{\text{osc}} = 7.2 \text{ GeV}$.

Given the stability of the fit in the accessible window of temperatures (see figure 6) big changes in the axion mass behavior are not expected. Still, extending the analysis to even higher temperatures would be extremely useful to control better the extrapolation systematics.

Larger values of f_a corresponds instead to smaller T_{osc} , for example for $f_a = 10^{13} \text{ GeV}$, $T_{\text{osc}} = 2.2 \text{ GeV}$ while for $f_a = 10^{14} \text{ GeV}$, $T_{\text{osc}} = 1.1 \text{ GeV}$. Above these values no extrapolation is required and the corresponding results are free from the extrapolation uncertainties.

5 Conclusions

We studied the topological properties and the θ -dependence for $N_f = 2 + 1$ QCD along a line of constant physics, corresponding to physical quarks masses, and for temperatures up to $4T_c$, where $T_c = 155 \text{ MeV}$ is the pseudo-critical temperature at which chiral symmetry restoration takes place. We explored several lattice spacings, in a range $0.05 - 0.12 \text{ fm}$, in order to perform a continuum extrapolation of our results. Our investigation at even smaller lattice spacings has been hindered by a severe slowing down in the decorrelation of the topological charge.

At zero temperature we observe large cut-off effects for the topological susceptibility. Nevertheless we are able to perform a continuum extrapolation, obtaining from the three finest lattice spacings $\chi^{1/4} = 73(9) \text{ MeV}$, in reasonable agreement with the ChPT prediction in the case of degenerate light flavors, $\chi_{\text{ChPT}}^{1/4} = 77.8(4) \text{ MeV}$.

At finite temperature we observe that cut-off effects are drastically reduced when one considers the ratio $\chi(T)/\chi(T=0)$, which turns out to be the most convenient quantity to perform a continuum extrapolation. The agreement with ChPT persists up to around T_c .

At higher temperatures the topological susceptibility presents the power-law dependence $\chi(T)/\chi(0) = D_0(T/T_c)^{D_2}$, with $D_2 \sim -3$. The exponent of the power-law behavior is however substantially smaller than the DIGA prediction, $D_2 \sim -8$.

Regarding the shape of the θ -dependent part of the free energy density, and in particular the lowest order coefficient b_2 , a visible convergence to the instanton gas prediction is observed in the explored range. With respect to the pure gauge case, the convergence of b_2 to DIGA is slower and the deviation is opposite in sign [14, 15]. This suggests a different interaction between instantons right above T_c , namely repulsive in the quenched case and attractive in full QCD [76].

The deviations from the dilute instanton gas predictions that we found in the present study have a significant impact on axions, resulting in particular in a shift of the axion dark matter window by almost one order of magnitude with respect to estimates based on DIGA. The softer temperature dependence of the topological susceptibility also changes the onset of the axion oscillations, which would now start at a higher temperatures ($T \sim 4$ GeV).

An important point is that this seems an effect directly related to the presence of light fermionic degrees of freedom: indeed, pure gauge computations [13, 22] observe a power-law behavior in good agreement with DIGA in a range of temperatures comparable to those explored in the present study.

One might wonder whether a different power law behavior might set in at temperatures higher than 1 GeV. That claims for future studies extending the range explored by us. The main obstruction to this extension is represented by the freezing of the topological modes at smaller lattice spacings which would be necessary to investigate such temperatures ($a < 0.05$ fm). Such an obstruction could be overcome by the development of new Monte-Carlo algorithms. Proposals in this respect have been made in the past [77, 78] and some new strategies have been put forward quite recently [79, 80]. In view of the exploration of higher temperatures, one should also consider the inclusion of dynamical charm quarks.

Acknowledgments

Numerical simulations have been performed on the Blue Gene/Q Fermi machine at CINECA, based on the agreement between INFN and CINECA (under INFN project NPQCD). Work partially supported by the INFN SUMA Project, by the ERC-2010 DaMESyFla Grant Agreement Number: 267985, by the ERC-2011 NEWPHYSICSHPC Grant Agreement Number: 27975 and by the MIUR (Italy) under a contract PRIN10-11 protocollo 2010YJ2NYW. FN acknowledges financial support from the INFN SUMA project.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R.D. Peccei and H.R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38** (1977) 1440 [[INSPIRE](#)].
- [2] R.D. Peccei and H.R. Quinn, *Constraints Imposed by CP Conservation in the Presence of Instantons*, *Phys. Rev. D* **16** (1977) 1791 [[INSPIRE](#)].
- [3] F. Wilczek, *Problem of Strong p and t Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279 [[INSPIRE](#)].
- [4] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223 [[INSPIRE](#)].
- [5] J. Preskill, M.B. Wise and F. Wilczek, *Cosmology of the Invisible Axion*, *Phys. Lett. B* **120** (1983) 127 [[INSPIRE](#)].
- [6] L.F. Abbott and P. Sikivie, *A Cosmological Bound on the Invisible Axion*, *Phys. Lett. B* **120** (1983) 133 [[INSPIRE](#)].
- [7] M. Dine and W. Fischler, *The Not So Harmless Axion*, *Phys. Lett. B* **120** (1983) 137 [[INSPIRE](#)].
- [8] E. Vicari and H. Panagopoulos, *Theta dependence of $SU(N)$ gauge theories in the presence of a topological term*, *Phys. Rept.* **470** (2009) 93 [[arXiv:0803.1593](#)] [[INSPIRE](#)].
- [9] D.J. Gross, R.D. Pisarski and L.G. Yaffe, *QCD and Instantons at Finite Temperature*, *Rev. Mod. Phys.* **53** (1981) 43 [[INSPIRE](#)].
- [10] T. Schäfer and E.V. Shuryak, *Instantons in QCD*, *Rev. Mod. Phys.* **70** (1998) 323 [[hep-ph/9610451](#)] [[INSPIRE](#)].
- [11] T.R. Morris, D.A. Ross and C.T. Sachrajda, *Higher Order Quantum Corrections in the Presence of an Instanton Background Field*, *Nucl. Phys. B* **255** (1985) 115 [[INSPIRE](#)].
- [12] A. Ringwald and F. Schrempp, *Confronting instanton perturbation theory with QCD lattice results*, *Phys. Lett. B* **459** (1999) 249 [[hep-lat/9903039](#)] [[INSPIRE](#)].
- [13] S. Borsányi et al., *Axion cosmology, lattice QCD and the dilute instanton gas*, *Phys. Lett. B* **752** (2016) 175 [[arXiv:1508.06917](#)] [[INSPIRE](#)].
- [14] C. Bonati, M. D’Elia, H. Panagopoulos and E. Vicari, *Change of θ Dependence in 4D $SU(N)$ Gauge Theories Across the Deconfinement Transition*, *Phys. Rev. Lett.* **110** (2013) 252003 [[arXiv:1301.7640](#)] [[INSPIRE](#)].
- [15] C. Bonati, *Topology and θ dependence in finite temperature G_2 lattice gauge theory*, *JHEP* **03** (2015) 006 [[arXiv:1501.01172](#)] [[INSPIRE](#)].
- [16] G.-Y. Xiong, J.-B. Zhang, Y. Chen, C. Liu, Y.-B. Liu and J.-P. Ma, *Topological susceptibility near T_c in $SU(3)$ gauge theory*, *Phys. Lett. B* **752** (2016) 34 [[arXiv:1508.07704](#)] [[INSPIRE](#)].
- [17] B. Alles, M. D’Elia and A. Di Giacomo, *Topological susceptibility at zero and finite T in $SU(3)$ Yang-Mills theory*, *Nucl. Phys. B* **494** (1997) 281 [*Erratum ibid.* **B 679** (2004) 397] [[hep-lat/9605013](#)] [[INSPIRE](#)].
- [18] B. Alles, M. D’Elia and A. Di Giacomo, *Topological susceptibility in full QCD at zero and finite temperature*, *Phys. Lett. B* **483** (2000) 139 [[hep-lat/0004020](#)] [[INSPIRE](#)].
- [19] C. Gattringer, R. Hoffmann and S. Schaefer, *The Topological susceptibility of $SU(3)$ gauge theory near $T(c)$* , *Phys. Lett. B* **535** (2002) 358 [[hep-lat/0203013](#)] [[INSPIRE](#)].

- [20] B. Lucini, M. Teper and U. Wenger, *Topology of SU(N) gauge theories at $T = 0$ and $T = T(c)$* , *Nucl. Phys. B* **715** (2005) 461 [[hep-lat/0401028](#)] [[INSPIRE](#)].
- [21] L. Del Debbio, H. Panagopoulos and E. Vicari, *Topological susceptibility of SU(N) gauge theories at finite temperature*, *JHEP* **09** (2004) 028 [[hep-th/0407068](#)] [[INSPIRE](#)].
- [22] E. Berkowitz, M.I. Buchoff and E. Rinaldi, *Lattice QCD input for axion cosmology*, *Phys. Rev. D* **92** (2015) 034507 [[arXiv:1505.07455](#)] [[INSPIRE](#)].
- [23] MILC collaboration, A. Bazavov et al., *Topological susceptibility with the asqtad action*, *Phys. Rev. D* **81** (2010) 114501 [[arXiv:1003.5695](#)] [[INSPIRE](#)].
- [24] MILC collaboration, A. Bazavov et al., *Lattice QCD ensembles with four flavors of highly improved staggered quarks*, *Phys. Rev. D* **87** (2013) 054505 [[arXiv:1212.4768](#)] [[INSPIRE](#)].
- [25] ETM collaboration, K. Cichy, E. Garcia-Ramos and K. Jansen, *Topological susceptibility from the twisted mass Dirac operator spectrum*, *JHEP* **02** (2014) 119 [[arXiv:1312.5161](#)] [[INSPIRE](#)].
- [26] ALPHA collaboration, M. Bruno, S. Schaefer and R. Sommer, *Topological susceptibility and the sampling of field space in $N_f = 2$ lattice QCD simulations*, *JHEP* **08** (2014) 150 [[arXiv:1406.5363](#)] [[INSPIRE](#)].
- [27] JLQCD collaboration, H. Fukaya, S. Aoki, G. Cossu, S. Hashimoto, T. Kaneko and J. Noaki, *Topology density correlator on dynamical domain-wall ensembles with nearly frozen topological charge*, *PoS(LATTICE2014)* **323** [[arXiv:1411.1473](#)] [[INSPIRE](#)].
- [28] B. Alles, G. Boyd, M. D’Elia, A. Di Giacomo and E. Vicari, *Hybrid Monte Carlo and topological modes of full QCD*, *Phys. Lett. B* **389** (1996) 107 [[hep-lat/9607049](#)] [[INSPIRE](#)].
- [29] L. Del Debbio, H. Panagopoulos and E. Vicari, *theta dependence of SU(N) gauge theories*, *JHEP* **08** (2002) 044 [[hep-th/0204125](#)] [[INSPIRE](#)].
- [30] L. Del Debbio, G.M. Manca and E. Vicari, *Critical slowing down of topological modes*, *Phys. Lett. B* **594** (2004) 315 [[hep-lat/0403001](#)] [[INSPIRE](#)].
- [31] ALPHA collaboration, S. Schaefer, R. Sommer and F. Virotta, *Critical slowing down and error analysis in lattice QCD simulations*, *Nucl. Phys. B* **845** (2011) 93 [[arXiv:1009.5228](#)] [[INSPIRE](#)].
- [32] R. Kitano and N. Yamada, *Topology in QCD and the axion abundance*, *JHEP* **10** (2015) 136 [[arXiv:1506.00370](#)] [[INSPIRE](#)].
- [33] P. Weisz, *Continuum Limit Improved Lattice Action for Pure Yang-Mills Theory. 1.*, *Nucl. Phys. B* **212** (1983) 1 [[INSPIRE](#)].
- [34] G. Curci, P. Menotti and G. Paffuti, *Symanzik’s Improved Lagrangian for Lattice Gauge Theory*, *Phys. Lett. B* **130** (1983) 205 [Erratum *ibid.* **B 135** (1984) 516] [[INSPIRE](#)].
- [35] C. Morningstar and M.J. Peardon, *Analytic smearing of SU(3) link variables in lattice QCD*, *Phys. Rev. D* **69** (2004) 054501 [[hep-lat/0311018](#)] [[INSPIRE](#)].
- [36] Y. Aoki et al., *The QCD transition temperature: results with physical masses in the continuum limit II*, *JHEP* **06** (2009) 088 [[arXiv:0903.4155](#)] [[INSPIRE](#)].
- [37] S. Borsányi et al., *The QCD equation of state with dynamical quarks*, *JHEP* **11** (2010) 077 [[arXiv:1007.2580](#)] [[INSPIRE](#)].

- [38] S. Borsányi, Z. Fodor, C. Hölbling, S.D. Katz, S. Krieg and K.K. Szabo, *Full result for the QCD equation of state with 2 + 1 flavors*, *Phys. Lett. B* **730** (2014) 99 [[arXiv:1309.5258](#)] [[INSPIRE](#)].
- [39] H. Neuberger, *Exactly massless quarks on the lattice*, *Phys. Lett. B* **417** (1998) 141 [[hep-lat/9707022](#)] [[INSPIRE](#)].
- [40] P. Hasenfratz, V. Laliena and F. Niedermayer, *The Index theorem in QCD with a finite cutoff*, *Phys. Lett. B* **427** (1998) 125 [[hep-lat/9801021](#)] [[INSPIRE](#)].
- [41] M. Lüscher, *Exact chiral symmetry on the lattice and the Ginsparg-Wilson relation*, *Phys. Lett. B* **428** (1998) 342 [[hep-lat/9802011](#)] [[INSPIRE](#)].
- [42] M. Lüscher, *Topological effects in QCD and the problem of short distance singularities*, *Phys. Lett. B* **593** (2004) 296 [[hep-th/0404034](#)] [[INSPIRE](#)].
- [43] L. Giusti and M. Lüscher, *Chiral symmetry breaking and the Banks-Casher relation in lattice QCD with Wilson quarks*, *JHEP* **03** (2009) 013 [[arXiv:0812.3638](#)] [[INSPIRE](#)].
- [44] B. Berg, *Dislocations and Topological Background in the Lattice $O(3)$ σ Model*, *Phys. Lett. B* **104** (1981) 475 [[INSPIRE](#)].
- [45] Y. Iwasaki and T. Yoshie, *Instantons and Topological Charge in Lattice Gauge Theory*, *Phys. Lett. B* **131** (1983) 159 [[INSPIRE](#)].
- [46] S. Itoh, Y. Iwasaki and T. Yoshie, *Stability of Instantons on the Lattice and the Renormalized Trajectory*, *Phys. Lett. B* **147** (1984) 141 [[INSPIRE](#)].
- [47] M. Teper, *Instantons in the Quantized $SU(2)$ Vacuum: A Lattice Monte Carlo Investigation*, *Phys. Lett. B* **162** (1985) 357 [[INSPIRE](#)].
- [48] E.-M. Ilgenfritz, M.L. Laursen, G. Schierholz, M. Müller-Preussker and H. Schiller, *First Evidence for the Existence of Instantons in the Quantized $SU(2)$ Lattice Vacuum*, *Nucl. Phys. B* **268** (1986) 693 [[INSPIRE](#)].
- [49] M. Lüscher, *Trivializing maps, the Wilson flow and the HMC algorithm*, *Commun. Math. Phys.* **293** (2010) 899 [[arXiv:0907.5491](#)] [[INSPIRE](#)].
- [50] M. Lüscher, *Properties and uses of the Wilson flow in lattice QCD*, *JHEP* **08** (2010) 071 [*Erratum ibid.* **1403** (2014) 092] [[arXiv:1006.4518](#)] [[INSPIRE](#)].
- [51] C. Bonati and M. D’Elia, *Comparison of the gradient flow with cooling in $SU(3)$ pure gauge theory*, *Phys. Rev. D* **89** (2014) 105005 [[arXiv:1401.2441](#)] [[INSPIRE](#)].
- [52] K. Cichy et al., *Comparison of different lattice definitions of the topological charge*, *PoS(LATTICE2014)075* [[arXiv:1411.1205](#)] [[INSPIRE](#)].
- [53] Y. Namekawa, *Comparative study of topological charge*, *PoS(LATTICE2014)344* [[arXiv:1501.06295](#)] [[INSPIRE](#)].
- [54] C. Alexandrou, A. Athenodorou and K. Jansen, *Topological charge using cooling and the gradient flow*, *Phys. Rev. D* **92** (2015) 125014 [[arXiv:1509.04259](#)] [[INSPIRE](#)].
- [55] R. Sommer, *Scale setting in lattice QCD*, *PoS(LATTICE 2013)015* [[arXiv:1401.3270](#)] [[INSPIRE](#)].
- [56] Z. Fodor, K. Holland, J. Kuti, S. Mondal, D. Nogradi and C.H. Wong, *The lattice gradient flow at tree-level and its improvement*, *JHEP* **09** (2014) 018 [[arXiv:1406.0827](#)] [[INSPIRE](#)].

- [57] P. Di Vecchia, K. Fabricius, G.C. Rossi and G. Veneziano, *Preliminary Evidence for U(1)-A Breaking in QCD from Lattice Calculations*, *Nucl. Phys. B* **192** (1981) 392 [[INSPIRE](#)].
- [58] C. Bonati, M. D'Elia and A. Scapellato, *θ dependence in SU(3) Yang-Mills theory from analytic continuation*, *Phys. Rev. D* **93** (2016) 025028 [[arXiv:1512.01544](#)] [[INSPIRE](#)].
- [59] P. Di Vecchia and G. Veneziano, *Chiral Dynamics in the Large- N Limit*, *Nucl. Phys. B* **171** (1980) 253 [[INSPIRE](#)].
- [60] H. Leutwyler and A.V. Smilga, *Spectrum of Dirac operator and role of winding number in QCD*, *Phys. Rev. D* **46** (1992) 5607 [[INSPIRE](#)].
- [61] TWQCD collaboration, Y.-Y. Mao and T.-W. Chiu, *Topological Susceptibility to the One-Loop Order in Chiral Perturbation Theory*, *Phys. Rev. D* **80** (2009) 034502 [[arXiv:0903.2146](#)] [[INSPIRE](#)].
- [62] F.-K. Guo and U.-G. Meißner, *Cumulants of the QCD topological charge distribution*, *Phys. Lett. B* **749** (2015) 278 [[arXiv:1506.05487](#)] [[INSPIRE](#)].
- [63] G. Grilli di Cortona, E. Hardy, J.P. Vega and G. Villadoro, *The QCD axion, precisely*, *JHEP* **01** (2016) 034 [[arXiv:1511.02867](#)] [[INSPIRE](#)].
- [64] L. Del Debbio, L. Giusti and C. Pica, *Topological susceptibility in the SU(3) gauge theory*, *Phys. Rev. Lett.* **94** (2005) 032003 [[hep-th/0407052](#)] [[INSPIRE](#)].
- [65] S. Dürr, Z. Fodor, C. Hölbling and T. Kurth, *Precision study of the SU(3) topological susceptibility in the continuum*, *JHEP* **04** (2007) 055 [[hep-lat/0612021](#)] [[INSPIRE](#)].
- [66] M. Lüscher and F. Palombi, *Universality of the topological susceptibility in the SU(3) gauge theory*, *JHEP* **09** (2010) 110 [[arXiv:1008.0732](#)] [[INSPIRE](#)].
- [67] ETM collaboration, K. Cichy, E. Garcia-Ramos, K. Jansen, K. Ottnad and C. Urbach, *Non-perturbative Test of the Witten-Veneziano Formula from Lattice QCD*, *JHEP* **09** (2015) 020 [[arXiv:1504.07954](#)] [[INSPIRE](#)].
- [68] M. Cé, C. Consonni, G.P. Engel and L. Giusti, *Non-Gaussianities in the topological charge distribution of the SU(3) Yang-Mills theory*, *Phys. Rev. D* **92** (2015) 074502 [[arXiv:1506.06052](#)] [[INSPIRE](#)].
- [69] M. D'Elia, *Field theoretical approach to the study of theta dependence in Yang-Mills theories on the lattice*, *Nucl. Phys. B* **661** (2003) 139 [[hep-lat/0302007](#)] [[INSPIRE](#)].
- [70] L. Giusti, S. Petrarca and B. Taglienti, *Theta dependence of the vacuum energy in the SU(3) gauge theory from the lattice*, *Phys. Rev. D* **76** (2007) 094510 [[arXiv:0705.2352](#)] [[INSPIRE](#)].
- [71] H. Panagopoulos and E. Vicari, *The 4D SU(3) gauge theory with an imaginary θ term*, *JHEP* **11** (2011) 119 [[arXiv:1109.6815](#)] [[INSPIRE](#)].
- [72] M.I. Buchoff et al., *QCD chiral transition, U(1)A symmetry and the Dirac spectrum using domain wall fermions*, *Phys. Rev. D* **89** (2014) 054514 [[arXiv:1309.4149](#)] [[INSPIRE](#)].
- [73] A. Trunin, F. Burger, E.-M. Ilgenfritz, M.P. Lombardo and M. Müller-Preussker, *Topological susceptibility from $N_f = 2 + 1 + 1$ lattice QCD at nonzero temperature*, *J. Phys. Conf. Ser.* **668** (2016) 012123 [[arXiv:1510.02265](#)] [[INSPIRE](#)].
- [74] O. Wantz and E.P.S. Shellard, *Axion Cosmology Revisited*, *Phys. Rev. D* **82** (2010) 123508 [[arXiv:0910.1066](#)] [[INSPIRE](#)].
- [75] PLANCK collaboration, P.A.R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, [arXiv:1502.01589](#) [[INSPIRE](#)].

- [76] C.G. Callan Jr., R.F. Dashen and D.J. Gross, *Toward a Theory of the Strong Interactions*, *Phys. Rev. D* **17** (1978) 2717 [[INSPIRE](#)].
- [77] P. de Forcrand, J.E. Hetrick, T. Takaishi and A.J. van der Sijs, *Three topics in the Schwinger model*, *Nucl. Phys. Proc. Suppl.* **63** (1998) 679 [[hep-lat/9709104](#)] [[INSPIRE](#)].
- [78] M. Lüscher and S. Schaefer, *Lattice QCD without topology barriers*, *JHEP* **07** (2011) 036 [[arXiv:1105.4749](#)] [[INSPIRE](#)].
- [79] A. Laio, G. Martinelli and F. Sanfilippo, *Metadynamics Surfing on Topology Barriers: the CP(N-1) Case*, [arXiv:1508.07270](#) [[INSPIRE](#)].
- [80] M.G. Endres, R.C. Brower, W. Detmold, K. Orginos and A.V. Pochinsky, *Multiscale Monte Carlo equilibration: Pure Yang-Mills theory*, *Phys. Rev. D* **92** (2015) 114516 [[arXiv:1510.04675](#)] [[INSPIRE](#)].