

*Tratti maladattivi di personalità,  
ansia e sintomi somatici in adolescenza*

*Maladaptive personality traits,  
anxiety and somatic symptoms in adolescence*

Valentina Spensieri, Simone Amendola

**RASSEGNA DI PSICOLOGIA**

n. 1, vol. XXXVI, 2019  
ISSN PRINT: 1125-5196  
ISSN ONLINE: 1974-4854  
pp. 73-84  
DOI: 10.4458/1965-05



Edizioni Nuova Cultura

The article is protected by copyright and all rights are held exclusively by Sapienza, University of Rome.

Il presente articolo è protetto da copyright e tutti i diritti sono di proprietà della Sapienza, Università degli Studi di Roma.



## Maladaptive personality traits, anxiety and somatic symptoms in adolescence

### Tratti maladattivi di personalità, ansia e sintomi somatici in adolescenza

Valentina Spensieri<sup>a,\*</sup>, Simone Amendola<sup>a</sup>

<sup>a</sup>Department of Clinical and Dynamic Psychology, Sapienza University, Rome, Italy

#### ARTICLE INFO

Submitted: 22 January 2019  
Accepted: 14 March 2019  
DOI: 10.4458/1965-05

#### ABSTRACT

Few studies have explored the link between personality traits and somatic symptoms in adolescence. In contrast, the association between anxiety and somatic symptoms is well-established. This study aimed to evaluate the relationship between maladaptive personality trait domains, anxiety and somatic symptoms. A sample of 303 Italian adolescents (159 males) aged 14 to 17 years were recruited for this study. Participants completed the following self-report measures: the Personality Inventory for DSM-5 Brief Form, the Screen for Child Anxiety-Related Emotional Disorders, and the Children's Somatization Inventory-24. Significant positive correlations emerged between personality trait domains (except for disinhibition), anxiety and somatic symptoms. Hierarchical multiple regression analyses indicated that only psychoticism, among the personality domains, explained unique variance in somatic symptoms, once the role of anxiety and gender was taken into account. This study provides evidence regarding the role of psychoticism, as well as anxiety, in predicting somatic symptoms among healthy adolescents. These findings have important implications for the prevention and clinical management of adolescents who report diverse somatic symptoms. Longitudinal studies are needed to better explore the relationships between these variables in adolescence.

**Keywords:** personality traits; anxiety; somatic symptoms; adolescence.

#### RIASSUNTO

Pochi studi hanno esplorato il legame tra tratti di personalità e sintomi somatici in adolescenza. Al contrario, l'associazione tra ansia e sintomi somatici è stata ben stabilita. L'obiettivo di questo studio è stato quello di indagare le relazioni tra i domini di tratto patologico della personalità, l'ansia e i sintomi somatici.

Un campione di 303 adolescenti italiani (159 maschi) di età compresa tra i 14 e 17 anni ha preso parte a questo studio. I partecipanti hanno completato le seguenti misure di autovalutazione: il Personality Inventory for DSM-5 Brief Form, lo Screen for Child Anxiety-Related Emotional Disorders e il Children's Somatization Inventory-24.

Sono emerse correlazioni positive e significative tra i domini di tratto della personalità (eccetto la Disinibizione), l'ansia e i sintomi somatici. L'analisi di regressione multipla gerarchica ha mostrato che solo lo Psicoticismo, tra i domini della personalità, spiega varianza unica nei sintomi somatici, una volta tenuto conto del ruolo dell'ansia e del genere.

Il presente studio evidenzia il ruolo dello psicoticismo e dell'ansia nel predire i sintomi somatici in adolescenti non clinici. I risultati hanno potenziali implicazioni per la prevenzione e il trattamento degli adolescenti che riferiscono diversi sintomi somatici. Sono necessari studi longitudinali per esplorare le relazioni tra queste variabili in adolescenza.

**Parole chiave:** tratti di personalità; ansia; sintomi somatici; adolescenza.

\*Corresponding author.

Valentina Spensieri  
Department of Dynamic and Clinical  
Psychology, Sapienza University of Rome  
Via degli Apuli 1, 00185, Rome, Italy.  
Email: valentina.spensieri@uniroma1.it



## Introduction

Research on somatic symptomatology has shown that headaches, excessive tiredness, low energy and stomachache are the most frequently self-reported somatic symptoms during adolescence (Cerutti et al., 2017; Essau et al., 2013; Romero-Acosta et al., 2013; Canals et al., 2013). Despite the fact that somatic symptoms are common among adolescents, it is important to underline that they may persist over time, with a negative impact on psychosocial functioning, including poor school attendance and achievement (Bakker et al., 2009; Cerutti et al., 2017b). Furthermore, in children and adolescents, somatic symptoms are frequently associated with anxiety and depressive symptomatology (Campo, 2012; Romero-Acosta et al., 2013; Saps et al., 2009) and predict hospital-based mental health care in adulthood (Bohman et al., 2018).

Egger and colleagues (Egger et al., 1999) revealed that somatic symptoms (e.g., stomachaches, headaches and musculo- skeletal pains) were more prevalent in patients with depression and/or anxiety disorder than in healthy control samples. A prospective cohort study examining the prevalence and impact of pediatric abdominal pain showed that approximately 72% of participants (N=237) reported at least one somatic symptom weekly, and 45% reported at least one gastrointestinal symptom (Saps et al., 2009). Other studies showed that adolescents described one or more somatic symptoms in the last two weeks with percentages ranging from 35 to 52% (Cerutti et al., 2017b; Romero-Acosta et al., 2013) and highlighting an age difference, with adolescents reporting more somatic symptoms than children (Romero-Acosta et al. 2013; Walker et al., 2009). Moreover, girls tend to report more somatic symptoms than boys (Essau et al., 2013; Romero-Acosta et al., 2013; Walker et al., 2009). In particular, Romero-Acosta and colleagues indicated that these gender differences started at 13 years of age and decreased at 15 years (Romero-Acosta et al., 2013).

### *Personality and somatic symptoms*

The Second Edition of the Psychodynamic Diagnostic Manual (PDM-2) is based on the conviction that “all people have personality styles” and includes the notion that “personality issues naturally co-occur with other presenting problems including anxiety, depression, somatic symptoms, addictions, phobias, self-harm, trauma and relationship problems” (Lingiardi & Mc Williams, 2017, p. 17). Research on the association between personality aspects and somatic symptoms is quite limited, specifically during adolescence (Cerutti et al., 2017b; Villanueva Badenes, Prado-Gasco, & Gonzales Barron, 2016). Cerutti and colleagues (2017b) demonstrated that greater difficulty in identifying feelings, a facet of the alexithymia construct, predicted greater functional impairment in children and adolescents through an increase in somatic symptoms. Villanueva Badenes and colleagues (2016) highlighted that higher levels of extraversion and openness and lower levels of consciousness were related to an increased number of somatic complaints in a population of Spanish children. Despite the fact that few studies have explored the link between personality traits and somatic symptoms in childhood and adolescence, a large body of research has explored this association in adults (Neeleman, Nijl, & Ormel, 2004). Specifically, Ode and Robinson (2007) demonstrated an interaction between agreeableness and neuroticism in predicting somatic symptoms among undergraduate students. Specifically, the authors reported that agreeableness plays a significant role in the self-regulation of negative affect, highlighting that the association between neuroticism and somatic symptoms was strong at low levels of agreeableness and absent at high levels of agreeableness (Ode & Robinson, 2007). The findings of a recent longitudinal study (Klinger-König et al., 2018) pointed out interactions between difficulty in identifying feelings, neuroticism and extraversion in predicting physical health symptoms in an adult population. In particular, extraversion decreased the negative impact of neuroticism on somatic symptoms, whereas difficulty in identifying feelings increased it.

The Work Group for Personality and Personality Disorders of the DSM-5 included a proposal of a new model as an “alternative DSM-5 model for personality disorders” in Section III of DSM-5, the section referred to as “Emerging measures and models” (APA, 2013). Two criteria must be met for a personality disorder diagnosis: a) moderate or greater impairment in personality functioning and b) the presence of pathological personality traits. This model emphasizes dimensional personality traits organized into five trait domains, respectively, negative affectivity, detachment, antagonism, disinhibition and psychoticism. These trait domains significantly correlate with the Personality Psychopathology Five domains (Anderson et al., 2013) and with other measures of personality and

psychopathology (Anderson, Sellbom, & Salekin, 2018) and were used in the current study for interpreting our results. Previous research suggested that maladaptive personality traits as measured by the Personality Inventory for DSM-5-Brief Form (PID-5 – BF; APA, 2013) may be associated with emotion dysregulation (Pollock et al., 2016), preferred styles of humor (Zeigler-Hill et al., 2016) as well as persistent and problematic internet use (Gervasi et al., 2017) and opioid dependence (Massaldjieva et al., 2016).

Lugo and colleagues (2019) explored the validity of the DSM-5 personality traits among a Brazilian sample of psychiatric inpatients through the PID-5 highlighting that it represents a valid instrument to discriminate people with severe psychopathological symptoms. In the Italian context, Amendola et al. (2018) investigated the role of a specific maladaptive personality trait (i.e., Psychoticism) in the association between depressive symptoms and internet addiction in a community sample of adolescents, confirming the validity of the alternative DSM-5 model for personality disorders in the examination of internalizing and externalizing symptomatology.

#### *Anxiety and somatic symptoms*

Previous research has indicated that adolescents' self-reported somatic symptoms are related to anxiety both in clinical (Ginsburg, Riddle, & Davies, 2006) and non-clinical samples (Kovacs & Borcsa, 2017; Lavigne, Saps, & Bryant, 2012; Tsao et al., 2009; Muris & Meesters, 2004). Kovacs and Borcsa (2017) showed that anxiety significantly increased the risk for somatic symptoms and restlessness, stomachaches, blushing, palpitations, muscle tension, sweating, and trembling/shaking are the most common symptoms reported in children and adolescents with anxiety disorders (Crawley et al. 2014; Ginsburg et al., 2006). Campo and colleagues (2004) highlighted that anxiety disorders generally preceded abdominal pain. Interestingly, 79% of children and adolescents with recurrent abdominal pain received a diagnosis of anxiety disorder (Campo et al., 2004).

In line with Campo and colleagues, it has been shown that children and adolescents with frequent somatic symptoms are more likely to be diagnosed with anxiety and depression disorders (Domenech-Llaberia et al., 2004; Jellesma et al., 2006). Furthermore, results from recent studies highlighted positive correlations between somatic symptom severity and anxiety disorder severity as well as the degree of the general impairment (Crawley et al., 2014; Sackl-Pammer et al., 2018).

#### *Personality and anxiety*

Research has largely examined whether personality traits as defined by the Big Five model (i.e. extraversion, agreeableness, neuroticism, conscientiousness, openness to experience) represent common predictors of internalizing problems, such as anxiety (Kotov et al., 2010) in adults. Kotov's study (2010) highlights that high neuroticism and low conscientiousness scores were the most powerful traits related to internalizing psychopathology, as well as low extraversion scores, while agreeableness and openness showed weak and equivocal associations (Kotov et al., 2010). Similarly, strong relationships between neuroticism and internalizing symptoms (e.g. depression and anxiety) were also found in adolescent samples, suggesting that these personality traits could reflect the core of internalizing psychopathology (Barbaranelli et al., 2003; Griffith et al., 2010). Furthermore, openness, conscientiousness and extraversion negatively correlated with internalizing symptoms (Barbaranelli et al., 2003).

### **Study Objectives**

In light of the above considerations, there is reason to expect that anxiety plays an important role in the physical health of children and adolescents. It remains unclear whether maladaptive personality trait domains may explain unique variance in somatic symptoms beyond anxiety.

This study aims to explore the relationships between personality trait domains, anxiety and somatic symptoms in Italian schoolchildren. Specifically, we hypothesized that adolescents with higher levels of anxiety would report more somatic symptoms than non-anxious youth.

Furthermore, personality trait domains including negative affect, detachment and disinhibition are hypothesized to be positively related to somatic symptomatology. With regard to the associations between antagonism, psychoticism and somatic symptoms, we had an exploratory purpose. Finally, we expected that personality trait domains would explain additional variance over and above gender and anxiety.

To our knowledge, this is the first study to explore the relationships between personality trait domains as defined by the “Alternative DSM-5 model for personality disorders” and anxiety symptoms in predicting somatic symptoms among a healthy Italian adolescent sample.

## Method

### *Participants*

Three hundred and three adolescents (159 males) aged 14-17 years (Mean age= 14.82, SD= 0.81) were involved in the present study. Participants were recruited in two secondary schools in Italy and all of them completed the entire questionnaire battery. Exclusion criteria for participation in this study were the presence of a diagnosed psychiatric illness, history of significant neurological illness or brain injury and history of recurrent somatic symptoms. All participants were Caucasian. Informed consent was obtained from both participants and their parents before enrolment in the study and anonymity of participants was ensured.

This study was approved by the Ethics Committee of the Department of Dynamic and Clinical Psychology, Sapienza University of Rome (Italy).

### *Measures*

*The Assessment schedule of adolescents' health:* is a questionnaire designed for this study and assesses the health status of participants. Parents were asked to fill-in a questionnaire evaluating the physical and mental health status of their children.

*The Personality Inventory for DSM-5 Brief Form (PID-5-BF; APA, 2013):* is a 25-item self-report personality trait assessment scale. It assesses five maladaptive personality trait domains including negative affect, detachment, antagonism, disinhibition and psychoticism, with each trait domain including 5 items. The PID-5-BF was developed from the PID-5 (Krueger et al., 2012). Each item of the PID-5-BF asks the individual to rate how well the statement describes him or her generally. Each item on the measure is rated on a 4-point scale (i.e., 0=very false or often false; 1=sometimes or somewhat false; 2=sometimes or somewhat true; 3=very true or often true). Higher scores indicate a higher functioning impairment in the investigated domain. Findings from the Italian validation of the instrument suggest that the PID-5 has an adequate internal consistency ( $\alpha = .83$ ) and test-retest reliability (Fossati et al., 2015). Furthermore, the five-factor structure was confirmed. In the present study, the scale had a reliability of  $\alpha = .89$ .

*The Screen for Child Anxiety-Related Emotional Disorders (SCARED; Birmaher et al., 1999):* is a self-report questionnaire assessing anxiety symptoms in children and adolescents. It consists of 41 items rated on a three-point Likert scale, from 0 “Not True or Hardly Ever True” to 2 “Very True or Often True”. The questionnaire includes a scoring matrix in order to indicate the potential presence of general anxiety as well as other specific anxiety disorders. The SCARED was found to have good internal consistency (a coefficient value of approximately .90). Five factors emerged from the factor analysis: panic/somatic, generalized anxiety, separation anxiety, social phobia and school phobia (Birmaher et al., 1999). Additionally, the SCARED showed significant discriminant validity. A cut-off point of 25 is used to indicate the presence of severe anxiety symptomatology. The Italian version of the instrument showed good psychometric properties (Ogliari et al., 2006). In the present study, the SCARED had a reliability of  $\alpha = .87$ .

*The Children's Somatization Inventory-24 (CSI-24; Walker et al., 2009):* a self-report questionnaire evaluating children and adolescents' perceptions of somatic symptoms. It consists of 24 items rated on a 5-point Likert scale, from 0 “Not at all” to 4 “A whole lot”. The CSI-24 was translated into Italian using the translation-back-translation method, with the approval of the author (Cerutti et al., 2017). The total score is obtained by summing the scores given by all of the participant's answers and can vary from 0 to 96. Higher scores indicate greater somatic symptomatology. The scale has good internal consistency ( $\alpha = .88$ ) (Walker et al., 2009). Researchers report results relevant to a single somatization factor and significant correlations with depression, anxiety, functional disability and quality of life, thus corroborating construct validity (Lavigne et al., 2012). In the present study, the CSI-24 demonstrated good internal consistency (Cronbach's  $\alpha$  of .87).

*Data analysis.* Descriptive statistics were performed to evaluate the characteristics of the sample. Analysis of univariate variance (ANOVA) was used to investigate the main differences in personality functioning and maladaptive personality trait domains, anxiety symptoms and somatic symptomatology according to gender.

Participants were divided into two groups, “Anxious group” and “Non-anxious group”, according to the SCARED cut-off of 25 (Birmaher et al., 1999).

Preliminarily, an ANOVA was performed to examine personality and somatic symptom scores in the two different groups of adolescents (Anxious group versus Non-anxious group). Successively, the zero-order correlations between the study variables were calculated in order to analyze the relationships between the constructs explored in the current study.

Finally, a Hierarchical Multiple Regression Analysis was conducted to investigate if personality trait domains explain a statistically significant amount of variance in somatic symptoms (dependent variable) after accounting for all other variables (gender and anxiety). The first model (Model 1) included demographic information such as age. In the next step (Model 2), we added anxiety symptoms. In the final step (Model 3), maladaptive personality trait domains were included.

All data were analyzed using the Statistical Package for Social Science (SPSS) 25.0 for Windows. P values <.05 were considered statistically significant.

## Results

Table 1 shows the descriptive statistics and differences according to gender.

**Table 1 – One-way ANOVA on gender**

	Males Mean ( $\pm$ SD)	Females Mean ( $\pm$ SD)	Total sample Mean ( $\pm$ SD)	F (1,301)	p-value
PID-5-BF	0.61 ( $\pm$ 0.46)	0.67 ( $\pm$ 0.39)	0.64 ( $\pm$ 0.43)	1.26	0.262
Negativeaffect	0.70 ( $\pm$ 0.59)	0.92 ( $\pm$ 0.56)	0.81 ( $\pm$ 0.58)	10.23	0.002**
Detachment	0.51 ( $\pm$ 0.53)	0.60 ( $\pm$ 0.51)	0.55 ( $\pm$ 0.52)	1.99	0.159
Antagonism	0.44 ( $\pm$ 0.47)	0.39 ( $\pm$ 0.42)	0.41 ( $\pm$ 0.45)	0.85	0.357
Disinhibition	0.75 ( $\pm$ 0.66)	0.76 ( $\pm$ 0.55)	0.75 ( $\pm$ 0.61)	0.01	0.920
Psychoticism	0.65 ( $\pm$ 0.59)	0.68 ( $\pm$ 0.50)	0.66 ( $\pm$ 0.55)	0.16	0.687
SCARED	15.28 ( $\pm$ 9.06)	18.65 ( $\pm$ 10.09)	16.88 ( $\pm$ 9.69)	9.38	0.002**
CSI-24	11.33 ( $\pm$ 9.59)	14.69 ( $\pm$ 10.78)	12.93 ( $\pm$ 10.30)	8.25	0.004**

*Note.* \*\*  $p < .01$ ; PID-5-BF= Personality Inventory for DSM-5 Brief Form; SCARED= Screen for Child Anxiety-Related Emotional Disorders; CSI-24= Children’s Somatization Inventory-24

According to the cut-off proposed by Birmaher (Birmaher et al., 1999), 20.1% (n=61) of adolescents reported a severe anxious symptomatology.

Differences between the Anxious group and Non-anxious group are presented in Table 2. Specifically, 26 males and 35 females constitute the Anxious group.

**Table 2 - Differences between Anxious and Non-anxious groups**

	<b>Anxious Group</b> (n=61) Mean ( $\pm$ SD)	<b>Non-anxious group</b> (n=242) Mean ( $\pm$ SD)	F-value (1,301)	P-value
PID-5-BF	0.79 ( $\pm$ 0.39)	0.60 ( $\pm$ 0.43)	9.59	< 0.01
Negative affect	1.06 ( $\pm$ 0.59)	0.74 ( $\pm$ 0.56)	14.62	< 0.001
Detachment	0.76 ( $\pm$ 0.53)	0.50 ( $\pm$ 0.51)	13.43	< 0.001
Antagonism	0.50 ( $\pm$ 0.45)	0.39 ( $\pm$ 0.45)	2.68	0.103
Disinhibition	0.78 ( $\pm$ 0.57)	0.75 ( $\pm$ 0.62)	0.13	0.714
Psychoticism	0.84 ( $\pm$ 0.57)	0.62 ( $\pm$ 0.54)	7.59	< 0.01
CSI-24	19.46 ( $\pm$ 12.90)	11.28 ( $\pm$ 8.82)	34.07	< 0.001

Note. PID-5-BF= Personality Inventory for DSM-5 Brief Form; CSI-24= Children's Somatization Inventory-24.

Table 3 shows the Pearson correlations among personality functioning and personality trait domains, anxiety and somatic symptoms.

**Table 3 – Correlational matrix**

	1	2	3	4	5	6	7	8
1. PID-5-BF	1.00	0.76***	0.85***	0.68***	0.79***	0.84***	0.26***	0.30***
2. Negative affect		1.00	0.63***	0.31***	0.46***	0.52***	0.32***	0.24***
3. Detachment			1.00	0.49***	0.55***	0.66***	0.28***	0.30***
4. Antagonism				1.00	0.43***	0.54***	0.12*	0.16**
5. Disinhibition					1.00	0.57***	0.06	0.16**
6. Psychoticism						1.00	0.24***	0.31***
7. SCARED							1.00	0.39***
8. CSI-24								1.00

Note. \*  $p < .05$ ; \*\*  $p < .01$ ; \*\*\*  $p < .001$ . PID-5-BF= Personality Inventory for DSM-5 BriefForm; SCARED= Screen for Child Anxiety-Related Emotional Disorders; CSI-24= Children's Somatization Inventory-24

#### *Anxiety and maladaptive personality trait domains as predictors of somatic symptoms*

In Table 4, the hierarchical regression analysis with somatic symptoms as dependent variable is presented. In the first step (Model 1), female gender positively predicted somatic symptoms ( $b=0.163$ ,  $p=0.004$ ). The second step (Model 2) indicated that anxiety is a significant predictor of somatic symptoms and, simultaneously, gender resulted non-significant.

Analysis of Model 3 confirmed the importance of anxiety in predicting somatic symptoms and showed that the psychoticism personality domain explained unique variance in somatic symptoms ( $b=0.197$ ,  $p=0.011$ ). Furthermore, gender revealed a tendency towards significance ( $b=0.102$ ,  $p=0.055$ ).

**Table 4 – Hierarchical Multiple Regression Analysis of Predictors of Somatic Symptoms**

<i>Predictor variables</i>	<i>Model 1</i>	<i>Model 2</i>	<i>Model 3</i>
<i>Gender</i>			
Female	0.16 **	0.09	0.10
<i>Anxiety</i>			
SCARED		0.38***	0.31***
<i>Personality trait domains</i>			
Negativeaffect			-0.39
Detachment			0.10
Antagonism			-0.01
Disinhibition			-0.01
Psychoticism			0.20*
R2	0.03	0.16	0.22
R2 change	0.03	0.14	0.06
F change	8.25**	49.41***	4.21**

Note. \* p<.05; \*\* p<.01; \*\*\* p<.001. SCARED= Screen for Child Anxiety-Related Emotional Disorders

### Discussion and conclusions

The primary purpose of this study was to extend our knowledge on the association between anxiety, somatic symptoms and personality trait domains in a sample of Italian adolescents. It has been widely observed that anxious symptoms, together with somatic and depressive symptoms, inevitably interfere with the functioning of adolescents and their adaptive capacities, with negative repercussions on both social and educational benefits, self-esteem and self-efficacy (Mahrer, Montano, & Gold, 2012, Stevanovic, 2012). Anxiety represents one of the most common psychological disorders in childhood and adolescence (Fliet et al., 2019). According to Birmaher's cut-off (Birmaher, et al., 1999), 20.1% of adolescents (n=61) participating in the current study reported a severe anxious symptomatology and fell into the "Anxious Group". An ANOVA analysis showed significant differences between the "Anxious Group" and "Non Anxious Group" in specific personality trait domains (e.g. Negative affect, Detachment and Psychoticism) with the "Anxious Group" scoring higher than "Non Anxious Group". This result is in line with previous research on adult samples, underlining that anxiety plays an important role in mental health (Sadaghiani, 2011; Fielding et al, 2016; Kovacs & Borsca, 2017).

Anxious as well as somatic and depressive manifestations can be traced back to difficulties in affect regulation since unpleasant affective states are expressed through internalizing behaviours, with negative repercussions on the youth's academic and social adaptation, instead of being adequately managed and contained (Parr et al., 2016).

Consistent with earlier cross-sectional studies that have shown how anxiety is associated with an increase risk for somatic symptomatology (David, 2014; Kovacs & Borsca, 2017), our findings suggested higher levels of somatic symptoms in the "Anxious Group" and demonstrated higher degrees of somatic symptomatology in anxious adolescents. A possible explanation may be that anxious youth are more likely than non anxious youth to believe that negative social events happen to them (Halldorsson & Creswell, 2017) and tend to have negative thoughts about themselves. Given that anxious children and adolescents have a higher chance of perceiving and interpreting situations as dangerous, our research findings confirm that severely anxious thoughts activate the body, leading to reactions involving somatic symptoms.

As predicted by our hypothesis, correlational analyses revealed that the total score of the SCARED questionnaire was related to psychological distress, strengthening the notion that higher levels of anxiety are associated with greater somatic symptomatology as well as impairment in personality trait domains. This is in accordance with Liu and colleagues' study (Liu et al., 2018) in which the power of the association between personality, somatic symptoms and psychological distress was explored by using two samples of Australian and Chinese adolescents. Furthermore, our results indicate that only the Disinhibition personality trait was not significantly related to anxiety symptomatology. The Disinhibition personality trait domain of the PID-5 may be interpreted in terms of low conscientiousness, involving irresponsible, impulsive and risk taking behaviors. These features are in contrast with those of anxiety, essentially anxious children and adolescents are more self-conscious, confirming our principal research purpose and supporting the notion that highly self-conscious adolescents are at risk for a variety of internalizing problems, including anxiety (Bowker & Rubin, 2009). Nevertheless, the finding that the Disinhibition personality trait was positively related to somatic symptomatology highlights that risky behaviors are related to physical activation which result in somatic symptomatology.

With respect to the relationship between personality trait domains as defined by the alternative DSM-5 model for personality disorders and anxiety symptoms in predicting somatic symptomatology, our findings emphasized that both anxiety and the Psychoticism personality trait significantly predicted the emergence of somatic symptoms. Psychoticism can be described as a personality trait that involves eccentricity, perceptual problems, misperception of social cues and odd behavior or unusual experiences in behavior (Hopwood et al., 2013). It regards the integration of different functions (e.g., memory, perception) that allow the construction of representations of the external reality. The dysfunctional beliefs that characterize this pathological trait may predispose to the experience of the abnormal thoughts, feelings and behaviors of somatic symptoms. Our results are in keeping with other studies that have demonstrated an association between personality traits and somatic symptoms (Compton et al., 2008; Garcia-Torres et al., 2016, Liu et al., 2018). Interestingly, Liu and colleagues (Liu et al., 2018) found that Psychoticism was related to Australian adolescents' somatic and psychological distress but this link was not significant among Chinese adolescents, indicating that there were significant cross-cultural differences in the association between Psychoticism and somatic distress. Our findings give new contributions to the existing knowledge on the link between maladaptive personality traits and somatic symptoms in adolescence.

Despite the fact that our hypothesis were confirmed by the study's findings, more research is needed to further explore the role of anxiety and psychoticism in predicting somatic symptoms during adolescence, in order to provide useful information for planning preventive interventions for youth.

Overall, important limitations warrant consideration in interpreting our results. First, our sample comprised healthy adolescents, so it is unclear if these findings may be generalized to other populations. Second, this was a cross-sectional study and, consequently, the conclusions drawn should be considered with caution. Third, we used only self-report measures rather than objective assessment which may be less affected by respondent bias. Finally, it is important to note that the effect of Psychoticism other than significant is of modest entity as it explains about only 6% of total variability of somatic complaints. Probably this may be due to the extreme heterogeneity of somatic complaints with respect to Psychoticism manifestation.

In conclusion, the findings of this study highlight that anxiety symptomatology is an important clinical phenomenon in school-age children. The developmental course of anxiety symptoms warrants better understanding in order to increase the possibility of early detection and thus, allow appropriate and adequate interventions that may prevent serious problems in adulthood.

**Author Contributions**

All authors contributed equally to this work.

**Compliance with Ethical Standards****Conflict of interest**

The authors declare that they have no competing interests.

**Funding**

The authors received no financial support for the research, authorship, and/or publication of this article.

**Ethical approval**

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Informed Consent**

Informed consent was obtained from all participants included in the study

## References

- Amendola, S., Spensieri, V., & Cerutti, R. (November 2018). *Does depressive symptomatology moderate the relationship between Problematic internet use and psychoticism domain in adolescence?* Poster presented at 4th International Congress of Clinical and Health Psychology on Children and Adolescents, Palma de Mallorca, Spain.
- American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders. 5th ed.* Washington, DC.
- Anderson, J.L., Sellbom, M., Bagby, R.M., Quilty, L.C., Veltri, C.O., Markon, K.E., & Krueger, R.F. (2013). On the convergence between PSY-5 domains and PID-5 domains and facets: implications for assessment of DSM-5 personality traits. *Assessment*, 20(3), 286-294.
- Anderson, J.L., Sellbom, M., & Salekin, R.T. (2018). Utility of the Personality Inventory for DSM-5-Brief Form (PID-5-BF) in the Measurement of Maladaptive Personality and Psychopathology. *Assessment*, 25(5), 596-607.
- Bakker, R.J., van de Putte, E.M., Kuis, W., & Sinnema, G. (2009). Risk factors for persistent fatigue with significant school absence in children and adolescents. *Pediatrics*, 124(1), e89-95.
- Barbaranelli, C., Caprara, G.V., Rabasca, A., & Pastorelli, C. (2003). A questionnaire for measuring the Big Five in late childhood. *Personality and Individual Differences*, 34, 645–664.
- Birmaher, B., Brent, D.A., Chiappetta, L., Bridge, J., Monga, S., & Baugher, M. (1999). Psychometric properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED): A replication study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(10), 1230–1236.
- Bohman, H., Laftman, S.B., Cleland, N., Lundberg, M., Paaren, A., & Jonsson, U. (2018). Somatic symptoms in adolescence as a predictor of severe mental illness in adulthood: a long-term community-based follow-up study. *Child and Adolescent Psychiatry and Mental Health*, 12:42.
- Bowker, J.C., & Rubin, K.H. (2009). Self-consciousness, friendship quality, and adolescent internalizing problems. *The British journal of developmental psychology*, 27(2), 249-67.
- Campo, J.V. (2012). Annual Research Review - Functional somatic symptoms and associated anxiety and Depression: developmental psychopathology in pediatric practice. *Journal of Child Psychology and Psychiatry*, 53(5), 575-592.
- Campo, J.V., Bridge, J.A., Ehmann, M., Altman, S., Lucas, A., Birmaher, B., Di Lorenzo, C., Iyengar, S., & Brent, D.A. (2004). Recurrent Abdominal Pain, Anxiety, and Depression in Primary Care. *Pediatrics*, 113(4), 817-824
- Crawley, S. A., Caporino, N. E., Birmaher, B., Ginsburg, G., Piacentini, J., Albano, A. M., Sherrill, J., Sakolsky, D., Compton, S. N., Rynn, M., McCracken, J., Gosch, E., Keeton, C., March, J., Walkup, J. T., ...& Kendall, P. C. (2014). Somatic complaints in anxious youth. *Child psychiatry and human development*, 45(4), 398-407.
- Cerutti, R., Spensieri, V., Presaghi, F., Valastro, C., Fontana, A., & Guidetti, V. (2017a). An Exploratory Study on Internet Addiction, Somatic Symptoms and Emotional and Behavioral Functioning in School-Aged Adolescents. *Clinical Neuropsychiatry*, 14 (6), 374-383, 2017
- Cerutti, R., Spensieri, V., Valastro, C., Presaghi, F., Canitano, R., & Guidetti, V. (2017b). A comprehensive approach to understand somatic symptoms and their impact on emotional and psychosocial functioning in children. *PLoS ONE*, 12(2): e0171867.
- Compton, M.T., Carter, T., Kryda, A., Goulding, S.M., & Kaslow, N.J. (2008). The impact of psychoticism on perceived hassles, depression, hostility, and hopelessness in non-psychiatric African Americans. *Psychiatry Research*, 159(1–2), 215-225.
- Domenech-Llaberia, E., Jane, M. C., Canals, J., Ballestri, S., Esparó, G., & Garralda, E. (2004). Parental reports of somatic symptoms in preschool children: Prevalence and associations in a Spanish sample. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43, 598-604.

- Egger, H.L., Costello, E.J., Erkanli, A., & Angold, A. (1999). Somatic complaints and psychopathology in children and adolescents: stomach aches, musculoskeletal pains, and headaches. *Journal of the American Academy of Child and Adolescent Psychiatry* 38(7), 852-860.
- Essau, C.A., Olaya, B., Boszczanin, A., Gilvarry, C., & Bray, D. (2013). Somatic symptoms among children and adolescents in Poland: a confirmatory factor analytic study of the Children Somatization Inventory. *Frontiers in Public Health*, 1:72.
- Fliet, L., Roelofs, J., van Breukelen, G., & Peter Muris (2019). A Longitudinal Study on the Relations Among Fear-Enhancing Parenting, Cognitive Biases, and Anxiety Symptoms in Non-clinical Children. *Child Psychiatry & Human Development*, 1-16.
- Fossati, A., Somma, A., Borroni, S., Markon, K. E., & Krueger, R. F. (2016). The Personality Inventory for DSM-5 Brief Form: Evidence for reliability and construct validity in a sample of community-dwelling Italian adolescents. *Assessment*, 23(6), 683-697.
- Garcia-Torres, F., Alos, F.J., & Perez-Duenas, C. (2016). Commentary: Eysenck personality questionnaire revised psychoticism predicts motivational-somatic symptoms of depression in breast cancer survivors. *Frontiers in Public Health*, 4, 42-42.
- Gervasi, A.M., La Marca, L., Lombardo, E., Mannino, G., Iacolino, C., & Schimmenti, A. (2017). Maladaptive Personality Traits and Internet Addiction Symptoms Among Young Adults: A Study Based on the Alternative DSM-5 Model for Personality Disorders. *Clinical Neuropsychiatry*, 14(1).
- Ginsburg, G.S., Riddle, M.A., & Davies, M. (2006). Somatic Symptoms in Children and Adolescents With Anxiety Disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*, 45(10), 1179-1187.
- Griffith, J.W., Zinbarg, R.E., Craske, M.G., Mineka, S., Rose, R.D., Waters, A.M., ... & Sutton, J.M. (2010). Neuroticism as a common dimension in the internalizing disorders. *Psychological Medicine*, 40, 1125-1136.
- Halldorsson, B., & Creswell, C. (2017). Social anxiety in pre-adolescent children: What do we know about maintenance? *Behaviour Research and Therapy*, 99, 19-36.
- Hopwood, C.J., Schade, N., Krueger, R.F., Wright, A.G.C., & Markon, K.E. (2013). Connecting DSM-5 personality traits and pathological beliefs: Toward a unifying model. *Journal of psychopathology and behavioral assessment*, 35(2), 162-172.
- Jellesma, C. F., Rieffe, C., Terwogt, M. M., & Kneepkens, F.C.M. (2006). Somatic complaints and health care use in children: mod, emotion awareness and sense of coherence. *Social Science and Medicine*, 63, 2640-2648.
- Klinger-König, J., Hertel, J., Terock, J., Völzke, H., Van der Auwera, S., & Grabe, H.J. (2018). Predicting physical and mental health symptoms: Additive and interactive effects of difficulty identifying feelings, neuroticism and extraversion. *Journal of Psychosomatic Research*, 115, 14-23.
- Kovacs, K.I., & Borcsa, M. (2017). The relationship between anxiety, somatic symptoms and hardness in adolescence. *Romanian Journal of Applied Psychology*, 19(2), 42-49.
- Krueger, R.F., Derringer, J., Markon, K.E., Watson, D., & Skodol, A.E. (2012). Initial construction of a Maladaptive Personality Trait Model and Inventory for DSM-5. *Psychological Medicine*, 42(9), 1879-1890.
- Lavigne, J.V., Saps, M., & Bryant, F.B. (2012). Reexamining the factor structure of somatization using the Children's Somatization Inventory (CSI-24) in a Community Sample. *Journal of Pediatric Psychology*, 37(8), 914-924.
- Lingiardi, V., & Mc Williams, N. (2017). *Psychodynamic Diagnostic Manual, Second Edition: PDM-2*. New York: Guilford Press.
- Lipowski, Z. J. (1988). Somatization: the concept and its clinical application. *American Journal of Psychiatry*, 145(11), 1358-1368.
- Liu, Y., Gillespie, N.A., Ye, L., Zhu, G., Duffy, D.L., & Martin, N.G. (2018). The Relationship Between Personality and Somatic and Psychological Distress: A Comparison of Chinese and Australian Adolescents. *Behavior Genetics*, 48(4), 315-322.

- Lugo, V., de Oliveira, S.E.S., Hessel, C.R., Monteiro, R.T., Pasche, N.L., Pavan, G., Motta, L.S., Pacheco, M.A., & Spanemberg, L. (2019). Evaluation of DSM-5 and ICD-11 personality traits using the Personality Inventory for DSM-5 (PID-5) in a Brazilian sample of psychiatric inpatients. *Personality and Mental Health*, 13(1), 24-39.
- Mahrer, N.E., Montano, Z., & Gold, J.I. (2012). Relations between anxiety sensitivity, somatization, and health-related quality of life in children with chronic pain. *Journal of Pediatric Psychology*, 37, 808-816.
- Massaldjieva, R., Georguiev, S., & Hadzhiyska, L. (2016). Maladaptive personality traits in a sample of patients with opioid dependence. *European Health Psychologist*, 18(S), 995.
- Muris, P., & Meesters, C. (2004). Children's somatization symptoms: correlations with trait anxiety, anxiety sensitivity, and learning experiences. *Psychological Reports*, 94, 1269-1275.
- Neeleman, J., Bijl, R., & Ormel, J. (2004). Neuroticism, a central link between somatic and psychiatric morbidity: path analysis of prospective data. *Psychological Medicine*, 34(3), 521-531.
- Parr, N.J., Zeman, J., Braunstein, K., & Price, N. (2016). Peer emotion socialization and somatic complaints in adolescents. *Journal of Adolescence*, 50, 22-30.
- Pollock, N.C., McCabe, G.A., Southard, A.C., & Zeigler-Hill, V. (2016). Pathological personality traits and emotion regulation difficulties. *Personality and Individual Differences*, 95, 168-177.
- Ode, S., & Robinson, M.D. (2007). Agreeableness and the Self-Regulation of Negative Affect: Findings Involving the Neuroticism/Somatic Distress Relationship. *Personality and Individual Differences*, 43(8), 2137-2148.
- Ogliari, A., Citterio, A., Zanoni, A., Fagnani, C., Patriarca, V., Cirrincione, R., Stazi, M.A., Battaglia, M. (2006). Genetic and environmental influences on anxiety dimensions in Italian twins evaluated with the SCARED questionnaire. *Journal of Anxiety Disorders*, 20(6), 760-777.
- Romero-Acosta, K., Canals, J., Hernandez-Martinez, C., Penelo, E., Zolog, T.C., & Domenech-Llaberia, E. (2013). Age and gender differences of somatic symptoms in children and adolescents. *Journal of Mental Health*, 22(1), 33-41.
- Sackl-Pammer, P., Özlü-Erkilic, Z., Jahn, R., Karwautz, A., Pollak, E., Ohmann, S., & Akkaya-Kalayci, T. (2018). Somatic complaints in children and adolescents with social anxiety disorder. *Neuropsychiatrie*, 32, 187-195.
- Saps, M., Seshadri, R., Sztainberg, M., Schaffer, G., Marshall, B.M., & Di Lorenzo, C. (2009). A prospective school-based study of abdominal pain and other common somatic complaints in children. *Journal of Pediatrics*, 154(3), 322-326.
- Shagadiani, N.S.K. (2011). The role of hardiness in decreasing stressors and biological, cognitive and mental reactions. *Procedia- Social and Behavioral Sciences* 30, 2427-2430.
- Stevanovic, D. (2012). Impact of emotional and behavioral symptoms on quality of life in children and adolescents. *Quality of Life Research*, 22(2):333-337.
- Tsao, J.C.I., Allen, L.B., Evans, S., Lu, Q., Myers, C.D., & Zeltzer, L.K. (2009). Anxiety sensitivity and catastrophizing: Associations with pain and somatization in non-clinical children. *Journal of Health Psychology*, 14(8), 1085-1094.
- Villanueva Badenes, L., Prado-Gascó, L., & González Barrón, R. (2016). Emotion awareness, mood and personality as predictors of somatic complaints in children and adults. *Psicothema*, 28(4), 383-388.
- Walker, L. S., Beck, J. E., Garber, J., & Lambert, W. (2009). Children's Somatization Inventory: Psychometric Properties of the Revised Form (CSI-24). *Journal of Pediatric Psychology*, 34(4), 430-440.
- Zeigler-Hill, V., McCabe, G.A., & Vrabel, J.K. (2016). The dark side of humor: DSM-5 pathological personality traits and humor styles. *Europe's journal of psychology*, 12(3), 363.

Finito di stampare nel mese di Marzo 2019  
con tecnologia *print on demand*  
presso il Centro Stampa “Nuova Cultura”  
p.le Aldo Moro n. 5, 00185 Roma  
[www.nuovacultura.it](http://www.nuovacultura.it)  
per ordini: [ordini@nuovacultura.it](mailto:ordini@nuovacultura.it)

[Int\_9788833651965\_a4bnpat\_LM02]