

# Joint distribution of the process and its sojourn time on the positive half-line for pseudo-processes governed by high-order heat equation

Valentina CAMMAROTA\* and Aimé LACHAL<sup>†</sup>

## Abstract

Consider the high-order heat-type equation  $\partial u/\partial t = \pm \partial^N u/\partial x^N$  for an integer  $N > 2$  and introduce the related Markov pseudo-process  $(X(t))_{t \geq 0}$ . In this paper, we study the sojourn time  $T(t)$  in the interval  $[0, +\infty)$  up to a fixed time  $t$  for this pseudo-process. We provide explicit expressions for the joint distribution of the couple  $(T(t), X(t))$ .

**Keywords:** pseudo-process, joint distribution of the process and its sojourn time, Spitzer's identity.

**AMS 2000 Subject Classification:** Primary 60G20; Secondary 60J25, 60K35, 60J05.

---

\*Dipartimento di Statistica, Probabilità e Statistiche Applicate, UNIVERSITY OF ROME 'LA SAPIENZA', P.le A. Moro 5, 00185 Rome, ITALY. *E-mail address:* [valentina.cammarota@uniroma1.it](mailto:valentina.cammarota@uniroma1.it)

<sup>†</sup>Pôle de Mathématiques/Institut Camille Jordan/CNRS UMR5208, Bât. L. de Vinci, INSTITUT NATIONAL DES SCIENCES APPLIQUÉES DE LYON, 20 av. A. Einstein, 69621 Villeurbanne Cedex, FRANCE. *E-mail address:* [aime.lachal@insa-lyon.fr](mailto:aime.lachal@insa-lyon.fr), *Web page:* <http://maths.insa-lyon.fr/~lachal>

# 1 Introduction

Let  $N$  be an integer equal or greater than 2 and  $\kappa_N = (-1)^{1+N/2}$  if  $N$  is even,  $\kappa_N = \pm 1$  if  $N$  is odd. Consider the heat-type equation of order  $N$ :

$$\frac{\partial u}{\partial t} = \kappa_N \frac{\partial^N u}{\partial x^N}. \quad (1.1)$$

For  $N = 2$ , this equation is the classical normalized heat equation and its relationship with linear Brownian motion is of the most well-known. For  $N > 2$ , it is known that no ordinary stochastic process can be associated with this equation. Nevertheless a Markov “pseudo-process” can be constructed by imitating the case  $N = 2$ . This pseudo-process,  $X = (X(t))_{t \geq 0}$  say, is driven by a signed measure as follows. Let  $p(t; x)$  denote the elementary solution of Eq. (1.1), that is,  $p$  solves (1.1) with the initial condition  $p(0; x) = \delta(x)$ . This solution is characterized by its Fourier transform (see, e.g., [13])

$$\int_{-\infty}^{+\infty} e^{i\mu x} p(t; x) dx = e^{\kappa_N t (-i\mu)^N}.$$

The function  $p$  is real, not always positive and its total mass is equal to one:

$$\int_{-\infty}^{+\infty} p(t; x) dx = 1.$$

Moreover, its total absolute value mass  $\rho$  exceeds one:

$$\rho = \int_{-\infty}^{+\infty} |p(t; x)| dx > 1.$$

In fact, if  $N$  is even,  $p$  is symmetric and  $\rho < +\infty$ , and if  $N$  is odd,  $\rho = +\infty$ . The signed function  $p$  is interpreted as the pseudo-probability for  $X$  to lie at a certain location at a certain time. More precisely, for any time  $t > 0$  and any locations  $x, y \in \mathbb{R}$ , one defines

$$\mathbb{P}\{X(t) \in dy | X(0) = x\} / dy = p(t; x - y).$$

Roughly speaking, the distribution of the pseudo-process  $X$  is defined through its finite-dimensional distributions according to the Markov rule: for any  $n > 1$ , any times  $t_1, \dots, t_n$  such that  $0 < t_1 < \dots < t_n$  and any locations  $x, y_1, \dots, y_n \in \mathbb{R}$ ,

$$\mathbb{P}\{X(t_1) \in dy_1, \dots, X(t_n) \in dy_n | X(0) = x\} / dy_1 \dots dy_n = \prod_{i=1}^n p(t_i - t_{i-1}; y_{i-1} - y_i)$$

where  $t_0 = 0$  and  $y_0 = x$ .

This pseudo-process has been studied by several authors: see the references [2] to [4] and the references [8] to [20].

Now, we consider the sojourn time of  $X$  in the interval  $[0, +\infty)$  up to a fixed time  $t$ :

$$T(t) = \int_0^t \mathbb{1}_{[0, +\infty)}(X(s)) ds.$$

The computation of the pseudo-distribution of  $T(t)$  has been done by Beghin, Hochberg, Nikitin, Orsingher and Ragozina in some particular cases (see [2, 4, 9, 16, 20]), and by Krylov and the second author in more general cases (see [10, 11]).

The method adopted therein is the use of the Feynman-Kac functional which leads to certain differential equations. We point out that the pseudo-distribution of  $T(t)$  is actually a genuine probability distribution and in the case where  $N$  is even,  $T(t)$  obeys the famous Paul Lévy’s arcsine law, that is

$$\mathbb{P}\{T(t) \in ds\} / ds = \frac{\mathbb{1}_{(0,t)}(s)}{\pi \sqrt{s(t-s)}}.$$

We also mention that the sojourn time of  $X$  in a small interval  $(-\varepsilon, \varepsilon)$  is used in [3] to define a local time for  $X$  at 0. The evaluation of the pseudo-distribution of the sojourn time  $T(t)$  together with the up-to-date value of the pseudo-process,  $X(t)$ , has been tackled only in the particular cases  $N = 3$  and  $N = 4$

by Beghin, Hochberg, Orsingher and Ragozina (see [2, 4]). Their results have been obtained by solving certain differential equations leading to some linear systems. In [2, 4, 11], the Laplace transform of the sojourn time serves as an intermediate tool for computing the distribution of the up-to-date maximum of  $X$ .

In this paper, our aim is to derive the joint pseudo-distribution of the couple  $(T(t), X(t))$  for any integer  $N$ . Since the Feynman-Kac approach used in [2, 4] leads to very cumbersome calculations, we employ an alternative method based on Spitzer's identity. The idea of using this identity for studying the pseudo-process  $X$  appeared already in [8] and [18]. Since the pseudo-process  $X$  is properly defined only in the case where  $N$  is an even integer, the results we obtain are valid in this case. Throughout the paper, we shall then assume that  $N$  is even. Nevertheless, we formally perform all computations also in the case where  $N$  is odd, even if they are not justified.

The paper is organized as follows.

- In Section 2, we write down the settings that will be used. Actually, the pseudo-process  $X$  is not well defined on the whole half-line  $[0, +\infty)$ . It is properly defined on dyadic times  $k/2^n$ ,  $k, n \in \mathbb{N}$ . So, we introduce ad-hoc definitions for  $X(t)$  and  $T(t)$  as well as for some related pseudo-expectations. For instance, we shall give a meaning to the quantity

$$E(\lambda, \mu, \nu) = \mathbb{E} \left[ \int_0^\infty e^{-\lambda t + i\mu X(t) - \nu T(t)} dt \right]$$

which is interpreted as the 3-parameters Laplace-Fourier transform of  $(T(t), X(t))$ . We also recall in this part some algebraic known results.

- In Section 3, we explicitly compute  $E(\lambda, \mu, \nu)$  with the help of Spitzer's identity. This is Theorem 3.1.
- Sections 4, 5 and 6 are devoted to successively inverting the Laplace-Fourier transform with respect to  $\mu$ ,  $\nu$  and  $\lambda$  respectively. More precisely, in Section 4, we perform the inversion with respect to  $\mu$ ; this yields Theorem 4.1. Next, we perform the inversion with respect to  $\nu$  which gives Theorems 5.1 and 5.2. Finally, we carry out the inversion with respect to  $\lambda$  and the main results of this paper are Theorems 6.2 and 6.3. In each section, we examine the particular cases  $N = 3$  (case of an asymmetric pseudo-process) and  $N = 4$  (case of the biharmonic pseudo-process). For  $N = 2$  (case of rescaled Brownian motion), one can retrieve several classical formulas and we refer the reader to the first draft of this paper [6]. Moreover, our results recover several known formulas concerning the marginal distribution of  $T(t)$ , see also [6].
- The final appendix (Section 7) contains a discussion on Spitzer's identity as well as some technical computations.

## 2 Settings

### 2.1 A first list of settings

In this part, we introduce for each integer  $n$  a step-process  $X_n$  coinciding with the pseudo-process  $X$  on the times  $k/2^n$ ,  $k \in \mathbb{N}$ . Fix  $n \in \mathbb{N}$ . Set, for any  $k \in \mathbb{N}$ ,  $X_{k,n} = X(k/2^n)$  and for any  $t \in [k/2^n, (k+1)/2^n)$ ,  $X(t) = X_{k,n}$ . We can write globally

$$X_n(t) = \sum_{k=0}^{\infty} X_{k,n} \mathbb{1}_{[k/2^n, (k+1)/2^n)}(t).$$

Now, we recall from [13] the definitions of tame functions, functions of discrete observations, and admissible functions associated with the pseudo-process  $X$ . They were introduced by Nishioka [18] in the case  $N = 4$ .

**Definition 2.1.** Fix  $n \in \mathbb{N}$ . A tame function for  $X$  is a function of a finite number  $k$  of observations of the pseudo-process  $X$  at times  $j/2^n$ ,  $1 \leq j \leq k$ , that is a quantity of the form  $F_{k,n} = F(X(1/2^n), \dots, X(k/2^n))$  for a certain  $k$  and a certain bounded Borel function  $F : \mathbb{R}^k \rightarrow \mathbb{C}$ . The "expectation" of  $F_{k,n}$  is defined as

$$\mathbb{E}(F_{k,n}) = \int \dots \int_{\mathbb{R}^k} F(x_1, \dots, x_k) p(1/2^n; x - x_1) \dots p(1/2^n; x_{k-1} - x_k) dx_1 \dots dx_k.$$

**Definition 2.2.** Fix  $n \in \mathbb{N}$ . A function of the discrete observations of  $X$  at times  $k/2^n$ ,  $k \geq 1$ , is a convergent series of tame functions:  $F_{X_n} = \sum_{k=1}^{\infty} F_{k,n}$  where  $F_{k,n}$  is a tame function for all  $k \geq 1$ . Assuming the series  $\sum_{k=1}^{\infty} |\mathbb{E}(F_{k,n})|$  convergent, the “expectation” of  $F_{X_n}$  is defined as

$$\mathbb{E}(F_{X_n}) = \sum_{k=1}^{\infty} \mathbb{E}(F_{k,n}).$$

**Definition 2.3.** An admissible function is a functional  $F_X$  of the pseudo-process  $X$  which is the limit of a sequence  $(F_{X_n})_{n \in \mathbb{N}}$  of functions of discrete observations of  $X$ :  $F_X = \lim_{n \rightarrow \infty} F_{X_n}$ , such that the sequence  $(\mathbb{E}(F_{X_n}))_{n \in \mathbb{N}}$  is convergent. The “expectation” of  $F_X$  is defined as

$$\mathbb{E}(F_X) = \lim_{n \rightarrow \infty} \mathbb{E}(F_{X_n}).$$

In this paper, we are concerned with the sojourn time of  $X$  in  $[0, +\infty)$ :

$$T(t) = \int_0^t \mathbb{1}_{[0, +\infty)}(X(s)) \, ds.$$

In order to give a proper meaning to this quantity, we introduce the similar object related to  $X_n$ :

$$T_n(t) = \int_0^t \mathbb{1}_{[0, +\infty)}(X_n(s)) \, ds.$$

For determining the distribution of  $T_n(t)$ , we compute its 3-parameters Laplace-Fourier transform:

$$E_n(\lambda, \mu, \nu) = \mathbb{E} \left[ \int_0^{\infty} e^{-\lambda t + i\mu X_n(t) - \nu T_n(t)} \, dt \right].$$

In Section 3, we prove that the sequence  $(E_n(\lambda, \mu, \nu))_{n \in \mathbb{N}}$  is convergent and we compute its limit:

$$\lim_{n \rightarrow \infty} E_n(\lambda, \mu, \nu) = E(\lambda, \mu, \nu).$$

Formally,  $E(\lambda, \mu, \nu)$  is interpreted as

$$E(\lambda, \mu, \nu) = \mathbb{E} \left[ \int_0^{\infty} e^{-\lambda t + i\mu X(t) - \nu T(t)} \, dt \right]$$

where the quantity  $\int_0^{\infty} e^{-\lambda t + i\mu X(t) - \nu T(t)} \, dt$  is an admissible function of  $X$ . This computation is performed with the aid of Spitzer’s identity. This latter concerns the classical random walk. Nevertheless, since it hinges on combinatorial arguments, it can be applied to the context of pseudo-processes. We clarify this point in Section 3.

## 2.2 A second list of settings

We introduce some algebraic settings. Let  $\theta_i$ ,  $1 \leq i \leq N$ , be the  $N^{\text{th}}$  roots of  $\kappa_N$  and

$$J = \{i \in \{1, \dots, N\} : \Re \theta_i > 0\}, \quad K = \{i \in \{1, \dots, N\} : \Re \theta_i < 0\}.$$

Of course, the cardinalities of  $J$  and  $K$  sum to  $N$ :  $\#J + \#K = N$ . We state several results related to the  $\theta_i$ ’s which are proved in [11, 13]. We have the elementary equalities

$$\sum_{j \in J} \theta_j + \sum_{k \in K} \theta_k = \sum_{i=1}^N \theta_i = 0, \quad \left( \prod_{j \in J} \theta_j \right) \left( \prod_{k \in K} \theta_k \right) = \prod_{i=1}^N \theta_i = (-1)^{N-1} \kappa_N \quad (2.1)$$

and

$$\prod_{i=1}^N (x - \theta_i) = \prod_{i=1}^N (x - \bar{\theta}_i) = x^N - \kappa_N. \quad (2.2)$$

Moreover, from formula (5.10) in [13],

$$\prod_{k \in K} (x - \theta_k) = \sum_{\ell=0}^{\#K} (-1)^\ell \sigma_\ell x^{\#K-\ell}, \quad (2.3)$$

where  $\sigma_\ell = \sum_{\substack{k_1 < \dots < k_\ell \\ k_1, \dots, k_\ell \in K}} \theta_{k_1} \dots \theta_{k_\ell}$ . We have by Lemma 11 in [11]

$$\sum_{j \in J} \theta_j \prod_{i \in J \setminus \{j\}} \frac{\theta_i x - \theta_j}{\theta_i - \theta_j} = \sum_{j \in J} \theta_j = - \sum_{k \in K} \theta_k = \begin{cases} 1 & \text{if } N \text{ is even,} \\ \frac{1}{2 \sin \frac{\pi}{2N}} = \frac{\cos \frac{\pi}{2N}}{\sin \frac{\pi}{N}} & \text{if } N \text{ is odd.} \end{cases} \quad (2.4)$$

Set  $A_j = \prod_{i \in J \setminus \{j\}} \frac{\theta_i}{\theta_i - \theta_j}$  for  $j \in J$ , and  $B_k = \prod_{i \in K \setminus \{k\}} \frac{\theta_i}{\theta_i - \theta_k}$  for  $k \in K$ . The  $A_j$ 's and  $B_k$ 's solve a Vandermonde system: we have

$$\begin{aligned} \sum_{j \in J} A_j &= \sum_{k \in K} B_k = 1 \\ \sum_{j \in J} A_j \theta_j^m &= 0 \text{ for } 1 \leq m \leq \#J - 1, \quad \sum_{k \in K} B_k \theta_k^m = 0 \text{ for } 1 \leq m \leq \#K - 1. \end{aligned} \quad (2.5)$$

Observing that  $1/\theta_j = \bar{\theta}_j$  for  $j \in J$ , that  $\{\theta_j, j \in J\} = \{\bar{\theta}_j, j \in J\}$  and similarly for the  $\theta_k$ 's,  $k \in K$ , formula (2.11) in [13] gives

$$\sum_{j \in J} \frac{A_j \theta_j}{\theta_j - x} = \sum_{j \in J} \frac{A_j}{1 - \bar{\theta}_j x} = \frac{1}{\prod_{j \in J} (1 - \theta_j x)}, \quad \sum_{k \in K} \frac{B_k \theta_k}{\theta_k - x} = \sum_{k \in K} \frac{B_k}{1 - \bar{\theta}_k x} = \frac{1}{\prod_{k \in K} (1 - \theta_k x)} \quad (2.6)$$

In particular,

$$\sum_{j \in J} \frac{A_j \theta_j}{\theta_j - \theta_k} = \frac{1}{N B_k}, \quad \sum_{k \in K} \frac{B_k \theta_k}{\theta_k - \theta_j} = \frac{1}{N A_j}. \quad (2.7)$$

Set, for any  $m \in \mathbb{Z}$ ,  $\alpha_m = \sum_{j \in J} A_j \theta_j^m$  and  $\beta_m = \sum_{k \in K} B_k \theta_k^m$ . We have, by formula (2.11) of [13],  $\beta_{\#K} = (-1)^{\#K-1} \prod_{k \in K} \theta_k$ . Moreover,  $\beta_{\#K+1} = (-1)^{\#K-1} (\prod_{k \in K} \theta_k) (\sum_{k \in K} \theta_k)$ . The proof of this claim is postponed to Lemma 7.2 in the appendix. We sum up this information and (2.5) into

$$\beta_m = \begin{cases} 1 & \text{if } m = 0, \\ 0 & \text{if } 1 \leq m \leq \#K - 1, \\ (-1)^{\#K-1} \prod_{k \in K} \theta_k & \text{if } m = \#K, \\ (-1)^{\#K-1} (\prod_{k \in K} \theta_k) (\sum_{k \in K} \theta_k) & \text{if } m = \#K + 1, \\ \kappa_N & \text{if } m = N. \end{cases} \quad (2.8)$$

We also have

$$\alpha_{-m} = \sum_{j \in J} \frac{A_j}{\theta_j^m} = \kappa_N \sum_{j \in J} A_j \theta_j^{N-m} = \kappa_N \alpha_{N-m}$$

and then

$$\alpha_{-m} = \begin{cases} 1 & \text{if } m = 0, \\ \kappa_N (-1)^{\#J-1} (\prod_{j \in J} \theta_j) (\sum_{j \in J} \theta_j) & \text{if } m = \#K - 1, \\ \kappa_N (-1)^{\#J-1} \prod_{j \in J} \theta_j & \text{if } m = \#K, \\ 0 & \text{if } \#K + 1 \leq m \leq N - 1, \\ \kappa_N & \text{if } m = N. \end{cases} \quad (2.9)$$

In particular, by (2.1),

$$\alpha_0 \beta_0 = \alpha_{-N} \beta_N = 1, \quad \alpha_{-\#K} \beta_{\#K} = -1, \quad \alpha_{-\#K} \beta_{\#K+1} = \sum_{j \in J} \theta_j, \quad \alpha_{1-\#K} \beta_{\#K} = \sum_{k \in K} \theta_k. \quad (2.10)$$

Concerning the kernel  $p$ , we have from Proposition 1 in [11]

$$p(t; 0) = \begin{cases} \frac{\Gamma(\frac{1}{N})}{N \pi t^{1/N}} & \text{if } N \text{ is even,} \\ \frac{\Gamma(\frac{1}{N}) \cos(\frac{\pi}{2N})}{N \pi t^{1/N}} & \text{if } N \text{ is odd.} \end{cases} \quad (2.11)$$

Proposition 3 in [11] states

$$\mathbb{P}\{X(t) \geq 0\} = \int_0^\infty p(t; -\xi) d\xi = \frac{\#J}{N}, \quad \mathbb{P}\{X(t) \leq 0\} = \int_{-\infty}^0 p(t; -\xi) d\xi = \frac{\#K}{N} \quad (2.12)$$

and formulas (4.7) and (4.8) in [13] yield, for  $\lambda > 0$  and  $\mu \in \mathbb{R}$ ,

$$\int_0^\infty \frac{e^{-\lambda t}}{t} dt \int_{-\infty}^0 (e^{i\mu\xi} - 1) p(t; -\xi) d\xi = \log \left( \prod_{k \in K} \frac{\sqrt[N]{\lambda}}{\sqrt[N]{\lambda} - i\mu\theta_k} \right), \quad (2.13)$$

$$\int_0^\infty \frac{e^{-\lambda t}}{t} dt \int_0^\infty (e^{i\mu\xi} - 1) p(t; -\xi) d\xi = \log \left( \prod_{j \in J} \frac{\sqrt[N]{\lambda}}{\sqrt[N]{\lambda} - i\mu\theta_j} \right).$$

Let us introduce, for  $j \in J$ ,  $m \leq N - 1$  and  $x \geq 0$ ,

$$I_{j,m}(\tau; x) = \frac{Ni}{2\pi} \left( e^{-i\frac{m}{N}\pi} \int_0^\infty \xi^{N-m-1} e^{-\tau\xi^N - \theta_j e^{i\frac{\pi}{N}} x\xi} d\xi - e^{i\frac{m}{N}\pi} \int_0^\infty \xi^{N-m-1} e^{-\tau\xi^N - \theta_j e^{-i\frac{\pi}{N}} x\xi} d\xi \right). \quad (2.14)$$

Formula (5.13) in [13] gives, for  $0 \leq m \leq N - 1$  and  $x \geq 0$ ,

$$\int_0^\infty e^{-\lambda\tau} I_{j,m}(\tau; x) d\tau = \lambda^{-\frac{m}{N}} e^{-\theta_j \sqrt[N]{\lambda} x}. \quad (2.15)$$

We introduce in a very similar manner the functions  $I_{k,m}(\tau; x)$  for  $k \in K$  and  $x \leq 0$ .

**Example 2.1.** Case  $N = 3$ .

• For  $\kappa_3 = +1$ , the third roots of  $\kappa_3$  are  $\theta_1 = 1$ ,  $\theta_2 = e^{i\frac{2\pi}{3}}$ ,  $\theta_3 = e^{-i\frac{2\pi}{3}}$ , and the settings read  $J = \{1\}$ ,  $K = \{2, 3\}$ ,  $A_1 = 1$ ,  $B_2 = \frac{e^{-i\frac{\pi}{6}}}{\sqrt{3}}$ ,  $B_3 = \frac{e^{i\frac{\pi}{6}}}{\sqrt{3}}$ ,  $\alpha_0 = \alpha_{-1} = \alpha_{-2} = 1$ ,  $\beta_0 = 1$ ,  $\beta_{-1} = -1$ . Moreover,

$$I_{1,0}(\tau; x) = \frac{3i}{2\pi} \left( \int_0^\infty \xi^2 e^{-\tau\xi^3 - e^{i\frac{\pi}{3}} x\xi} d\xi - \int_0^\infty \xi^2 e^{-\tau\xi^3 - e^{-i\frac{\pi}{3}} x\xi} d\xi \right).$$

• For  $\kappa_3 = -1$ , the third roots of  $\kappa_3$  are  $\theta_1 = e^{i\frac{\pi}{3}}$ ,  $\theta_2 = e^{-i\frac{\pi}{3}}$ ,  $\theta_3 = -1$ . The settings read  $J = \{1, 2\}$ ,  $K = \{3\}$ ,  $A_1 = \frac{e^{i\frac{\pi}{6}}}{\sqrt{3}}$ ,  $A_2 = \frac{e^{-i\frac{\pi}{6}}}{\sqrt{3}}$ ,  $B_3 = 1$ ,  $\alpha_0 = \alpha_{-1} = 1$ ,  $\beta_0 = \beta_{-2} = 1$ ,  $\beta_{-1} = -1$ . Moreover,

$$I_{1,1}(\tau; x) = \frac{3i}{2\pi} \left( e^{-i\frac{\pi}{3}} \int_0^\infty \xi e^{-\tau\xi^3 - e^{i\frac{2\pi}{3}} x\xi} d\xi - e^{i\frac{\pi}{3}} \int_0^\infty \xi e^{-\tau\xi^3 - x\xi} d\xi \right),$$

$$I_{2,1}(\tau; x) = \frac{3i}{2\pi} \left( e^{-i\frac{\pi}{3}} \int_0^\infty \xi e^{-\tau\xi^3 - x\xi} d\xi - e^{i\frac{\pi}{3}} \int_0^\infty \xi e^{-\tau\xi^3 - e^{-i\frac{2\pi}{3}} x\xi} d\xi \right).$$

Actually, the three functions  $I_{1,0}$ ,  $I_{1,1}$  and  $I_{2,1}$  can be expressed by mean of the Airy function  $\text{Hi}$  defined as  $\text{Hi}(z) = \frac{1}{\pi} \int_0^\infty e^{-\frac{\xi^3}{3} + z\xi} d\xi$  (see, e.g., [1, Chap. 10.4]). Indeed, we easily have by a change of variables, differentiation and integration by parts, for  $\tau > 0$  and  $z \in \mathbb{C}$ ,

$$\int_0^\infty e^{-\tau\xi^3 + z\xi} d\xi = \frac{\pi}{(3\tau)^{4/3}} \text{Hi} \left( \frac{z}{\sqrt[3]{3\tau}} \right),$$

$$\int_0^\infty \xi e^{-\tau\xi^3 + z\xi} d\xi = \frac{\pi}{(3\tau)^{2/3}} \text{Hi}' \left( \frac{z}{\sqrt[3]{3\tau}} \right),$$

$$\int_0^\infty \xi^2 e^{-\tau\xi^3 + z\xi} d\xi = \frac{\pi z}{(3\tau)^{4/3}} \text{Hi} \left( \frac{z}{\sqrt[3]{3\tau}} \right) + \frac{1}{3\tau}.$$

Therefore,

$$I_{1,0}(\tau; x) = \frac{x}{2\sqrt[3]{3}\tau^{4/3}} \left[ e^{i\frac{\pi}{6}} \text{Hi} \left( -\frac{e^{-i\frac{\pi}{3}} x}{\sqrt[3]{3\tau}} \right) + e^{-i\frac{\pi}{6}} \text{Hi} \left( -\frac{e^{i\frac{\pi}{3}} x}{\sqrt[3]{3\tau}} \right) \right], \quad (2.16)$$

$$I_{1,1}(\tau; x) = \frac{\sqrt[3]{3}}{2\tau^{2/3}} \left[ e^{i\frac{\pi}{6}} \text{Hi}' \left( -\frac{e^{i\frac{2\pi}{3}} x}{\sqrt[3]{3\tau}} \right) + e^{-i\frac{\pi}{6}} \text{Hi}' \left( -\frac{x}{\sqrt[3]{3\tau}} \right) \right], \quad (2.17)$$

$$I_{2,1}(\tau; x) = \frac{\sqrt[3]{3}}{2\tau^{2/3}} \left[ e^{i\frac{\pi}{6}} \text{Hi}' \left( -\frac{x}{\sqrt[3]{3\tau}} \right) + e^{-i\frac{\pi}{6}} \text{Hi}' \left( -\frac{e^{-i\frac{2\pi}{3}} x}{\sqrt[3]{3\tau}} \right) \right]. \quad (2.18)$$

**Example 2.2.** Case  $N = 4$ : we have  $\kappa_4 = -1$ . This is the case of the biharmonic pseudo-process. The fourth roots of  $\kappa_4$  are  $\theta_1 = e^{-i\frac{\pi}{4}}$ ,  $\theta_2 = e^{i\frac{\pi}{4}}$ ,  $\theta_3 = e^{i\frac{3\pi}{4}}$ ,  $\theta_4 = e^{-i\frac{3\pi}{4}}$  and the notations read in this case  $J = \{1, 2\}$ ,  $K = \{3, 4\}$ ,  $A_1 = B_3 = \frac{e^{-i\frac{\pi}{4}}}{\sqrt{2}}$ ,  $A_2 = B_4 = \frac{e^{i\frac{\pi}{4}}}{\sqrt{2}}$ ,  $\alpha_0 = \alpha_{-2} = 1$ ,  $\alpha_{-1} = \sqrt{2}$ ,  $\beta_0 = \beta_{-2} = 1$ ,  $\beta_{-1} = -\sqrt{2}$ . Moreover,

$$\begin{aligned} I_{1,1}(\tau; x) &= \frac{2}{\pi} \left( e^{i\frac{\pi}{4}} \int_0^\infty \xi^2 e^{-\tau\xi^4 - x\xi} d\xi + e^{-i\frac{\pi}{4}} \int_0^\infty \xi^2 e^{-\tau\xi^4 + ix\xi} d\xi \right), \\ I_{2,1}(\tau; x) &= \frac{2}{\pi} \left( e^{i\frac{\pi}{4}} \int_0^\infty \xi^2 e^{-\tau\xi^4 - ix\xi} d\xi + e^{-i\frac{\pi}{4}} \int_0^\infty \xi^2 e^{-\tau\xi^4 - x\xi} d\xi \right). \end{aligned} \tag{2.19}$$

### 3 Evaluation of $E(\lambda, \mu, \nu)$

The goal of this section is to evaluate the limit  $E(\lambda, \mu, \nu) = \lim_{n \rightarrow \infty} E_n(\lambda, \mu, \nu)$ . We write  $E_n(\lambda, \mu, \nu) = \mathbb{E}[F_n(\lambda, \mu, \nu)]$  with

$$F_n(\lambda, \mu, \nu) = \int_0^\infty e^{-\lambda t + i\mu X_n(t) - \nu T_n(t)} dt.$$

Let us rewrite the sojourn time  $T_n(t)$  as follows:

$$\begin{aligned} T_n(t) &= \sum_{j=0}^{\lfloor 2^n t \rfloor} \int_{j/2^n}^{(j+1)/2^n} \mathbb{1}_{[0, +\infty)}(X_n(s)) ds - \int_t^{(\lfloor 2^n t \rfloor + 1)/2^n} \mathbb{1}_{[0, +\infty)}(X_n(s)) ds \\ &= \sum_{j=0}^{\lfloor 2^n t \rfloor} \int_{j/2^n}^{(j+1)/2^n} \mathbb{1}_{[0, +\infty)}(X_{j,n}) ds - \int_t^{(\lfloor 2^n t \rfloor + 1)/2^n} \mathbb{1}_{[0, +\infty)}(X_{\lfloor 2^n t \rfloor, n}) ds \\ &= \frac{1}{2^n} \sum_{j=0}^{\lfloor 2^n t \rfloor} \mathbb{1}_{[0, +\infty)}(X_{j,n}) + \left( t - \frac{\lfloor 2^n t \rfloor + 1}{2^n} \right) \mathbb{1}_{[0, +\infty)}(X_{\lfloor 2^n t \rfloor, n}). \end{aligned}$$

Set  $T_{0,n} = 0$  and, for  $k \geq 1$ ,

$$T_{k,n} = \frac{1}{2^n} \sum_{j=1}^k \mathbb{1}_{[0, +\infty)}(X_{j,n}).$$

For  $k \geq 0$  and  $t \in [k/2^n, (k+1)/2^n)$ , we see that

$$T_n(t) = T_{k,n} + \left( t - \frac{k+1}{2^n} \right) \mathbb{1}_{[0, +\infty)}(X_{k,n}) + \frac{1}{2^n}.$$

With this decomposition at hand, we can begin to compute  $F_n(\lambda, \mu, \nu)$ :

$$\begin{aligned} F_n(\lambda, \mu, \nu) &= \int_0^\infty e^{-\lambda t + i\mu X_n(t) - \nu T_n(t)} dt \\ &= \sum_{k=0}^\infty \int_{k/2^n}^{(k+1)/2^n} e^{-\lambda t + i\mu X_{k,n} - \nu T_{k,n} - \frac{\nu}{2^n} + \nu \left( \frac{k+1}{2^n} - t \right) \mathbb{1}_{[0, +\infty)}(X_{k,n})} dt \\ &= e^{-\nu/2^n} \left( \sum_{k=0}^\infty \int_{k/2^n}^{(k+1)/2^n} e^{-\lambda t + \nu \left( \frac{k+1}{2^n} - t \right) \mathbb{1}_{[0, +\infty)}(X_{k,n})} dt \right) e^{i\mu X_{k,n} - \nu T_{k,n}}. \end{aligned}$$

The value of the above integral is

$$\int_{k/2^n}^{(k+1)/2^n} e^{-\lambda t + \nu \left( \frac{k+1}{2^n} - t \right) \mathbb{1}_{[0, +\infty)}(X_{k,n})} dt = e^{-\lambda(k+1)/2^n} \frac{e^{[\lambda + \nu \mathbb{1}_{[0, +\infty)}(X_{k,n})]/2^n} - 1}{\lambda + \nu \mathbb{1}_{[0, +\infty)}(X_{k,n})}.$$

Therefore,

$$F_n(\lambda, \mu, \nu) = \frac{1 - e^{-(\lambda + \nu)/2^n}}{\lambda + \nu} \sum_{k=0}^\infty e^{-\lambda k/2^n + i\mu X_{k,n} - \nu T_{k,n}} \mathbb{1}_{[0, +\infty)}(X_{k,n})$$

$$+ e^{-\nu/2^n} \frac{1 - e^{-\lambda/2^n}}{\lambda} \sum_{k=0}^{\infty} e^{-\lambda k/2^n + i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{(-\infty, 0)}(X_{k,n}).$$

Before applying the expectation to this last expression, we have to check that it defines a function of discrete observations of the pseudo-process  $X$  which satisfies the conditions of Definition 2.2. This fact is stated in the proposition below.

**Proposition 3.1.** *Suppose  $N$  even and fix an integer  $n$ . For any complex  $\lambda$  such that  $\Re(\lambda) > 0$  and any  $\nu > 0$ , the series  $\sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})]$  and  $\sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{(-\infty, 0)}(X_{k,n})]$  are absolutely convergent and their sums are given by*

$$\begin{aligned} \sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})] &= \frac{e^{\nu/2^n} - S_n^+(\lambda, \mu, \nu)}{e^{\nu/2^n} - 1}, \\ \sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{(-\infty, 0)}(X_{k,n})] &= \frac{e^{\nu/2^n} [S_n^-(\lambda, \mu, \nu) - 1]}{e^{\nu/2^n} - 1}, \end{aligned}$$

where

$$\begin{aligned} S_n^+(\lambda, \mu, \nu) &= \exp\left(-\sum_{k=1}^{\infty} \left(1 - e^{-\nu k/2^n}\right) \frac{e^{-\lambda k/2^n}}{k} \mathbb{E}[e^{i\mu X_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})]\right), \\ S_n^-(\lambda, \mu, \nu) &= \exp\left(\sum_{k=1}^{\infty} \left(1 - e^{-\nu k/2^n}\right) \frac{e^{-\lambda k/2^n}}{k} \mathbb{E}[e^{i\mu X_{k,n}} \mathbf{1}_{(-\infty, 0)}(X_{k,n})]\right). \end{aligned}$$

PROOF

• **Step 1.** First, notice that for any  $k \geq 1$ , we have

$$\begin{aligned} & \left| \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})] \right| \\ &= \left| \int \dots \int_{\mathbb{R}^{k-1} \times [0, +\infty)} e^{i\mu x_k - \frac{\nu}{2^n} \sum_{j=1}^k \mathbf{1}_{[0, +\infty)}(x_j)} \mathbb{P}\{X_{1,n} \in dx_1, \dots, X_{k,n} \in dx_k\} \right| \\ &= \left| \int \dots \int_{\mathbb{R}^{k-1} \times [0, +\infty)} e^{i\mu x_k - \frac{\nu}{2^n} \sum_{j=1}^k \mathbf{1}_{[0, +\infty)}(x_j)} p\left(\frac{1}{2^n}; x_1\right) \prod_{j=1}^{k-1} p\left(\frac{1}{2^n}; x_j - x_{j+1}\right) dx_1 \dots dx_k \right| \\ &\leq \int \dots \int_{\mathbb{R}^k} \left| p\left(\frac{1}{2^n}; x_1\right) \prod_{j=1}^{k-1} p\left(\frac{1}{2^n}; x_j - x_{j+1}\right) \right| dx_1 \dots dx_k \\ &= \int \dots \int_{\mathbb{R}^k} \prod_{j=1}^k \left| p\left(\frac{1}{2^n}; y_j\right) \right| dy_1 \dots dy_k = \prod_{j=1}^k \int_{-\infty}^{+\infty} \left| p\left(\frac{1}{2^n}; y_j\right) \right| dy_j = \rho^k. \end{aligned}$$

Hence, we derive the following inequality:

$$\sum_{k=1}^{\infty} \left| e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})] \right| \leq \sum_{k=1}^{\infty} \rho^k \left| e^{-\lambda k/2^n} \right| = \frac{1}{1 - \rho e^{-\Re(\lambda)/2^n}}.$$

We can easily see that this bound holds true also when the factor  $\mathbf{1}_{[0, +\infty)}(X_{k,n})$  is replaced by  $\mathbf{1}_{(-\infty, 0)}(X_{k,n})$ . This shows that the two series of Proposition 3.1 are finite for  $\lambda \in \mathbb{C}$  such that  $\rho e^{-\Re(\lambda)/2^n} < 1$ , that is  $\Re(\lambda) > 2^n \log \rho$ .

• **Step 2.** For  $\lambda \in \mathbb{C}$  such that  $\Re(\lambda) > 2^n \log \rho$ , the Spitzer's identity (7.2) (see Lemma 7.1 in the appendix) gives for the first series of Proposition 3.1

$$\begin{aligned} & \sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})] \\ &= \frac{1}{e^{\nu/2^n} - 1} \left[ e^{\nu/2^n} - \exp\left(-\sum_{k=1}^{\infty} \left(1 - e^{-\nu k/2^n}\right) \frac{e^{-\lambda k/2^n}}{k} \mathbb{E}[e^{i\mu X_{k,n}} \mathbf{1}_{[0, +\infty)}(X_{k,n})]\right) \right]. \end{aligned} \quad (3.1)$$

The right-hand side of (3.1) is an analytic continuation of the Dirichlet series lying in the left-hand side of (3.1), which is defined on the half-plane  $\{\lambda \in \mathbb{C} : \Re(\lambda) > 0\}$ . Moreover, for any  $\varepsilon > 0$ , this continuation is bounded over the half-plane  $\{\lambda \in \mathbb{C} : \Re(\lambda) \geq \varepsilon\}$ . Indeed, we have

$$|\mathbb{E}[e^{i\mu X_{k,n}} \mathbb{1}_{[0,+\infty)}(X_{k,n})]| = \left| \int_0^{+\infty} e^{i\mu\xi} p\left(\frac{k}{2^n}; -\xi\right) d\xi \right| \leq \int_0^{+\infty} \left| p\left(\frac{k}{2^n}; -\xi\right) \right| d\xi < \rho$$

and then

$$\begin{aligned} & \left| \exp\left(-\sum_{k=1}^{\infty} \left(1 - e^{-\nu k/2^n}\right) \frac{e^{-\lambda k/2^n}}{k} \mathbb{E}[e^{i\mu X_{k,n}} \mathbb{1}_{[0,+\infty)}(X_{k,n})]\right)\right| \\ & \leq \exp\left(\rho \sum_{k=1}^{\infty} \frac{e^{-\Re(\lambda)k/2^n}}{k}\right) = \exp\left(-\rho \log(1 - e^{-\Re(\lambda)/2^n})\right) = \frac{1}{(1 - e^{-\Re(\lambda)/2^n})^\rho}. \end{aligned}$$

Therefore, if  $\Re(\lambda) \geq \varepsilon$ ,

$$\left| \exp\left(-\sum_{k=1}^{\infty} \left(1 - e^{-\nu k/2^n}\right) \frac{e^{-\lambda k/2^n}}{k} \mathbb{E}[e^{i\mu X_{k,n}} \mathbb{1}_{[0,+\infty)}(X_{k,n})]\right)\right| \leq \frac{1}{(1 - e^{-\varepsilon/2^n})^\rho}.$$

This proves that the left-hand side of this last inequality is bounded for  $\Re(\lambda) \geq \varepsilon$ . By a lemma of Bohr ([5]), we deduce that the abscissas of convergence, absolute convergence and boundedness of the Dirichlet series  $\sum_{k=0}^{\infty} e^{-\lambda k/2^n} \mathbb{E}[e^{i\mu X_{k,n} - \nu T_{k,n}} \mathbb{1}_{[0,+\infty)}(X_{k,n})]$  are identical. So, this series converges absolutely on the half-plane  $\{\lambda \in \mathbb{C} : \Re(\lambda) > 0\}$  and (3.1) holds on this half-plane. A similar conclusion holds for the second series of Proposition 3.1. The proof is finished. ■

Thanks to Proposition 3.1, we see that the functional  $F_n(\lambda, \mu, \nu)$  is a function of the discrete observations of  $X$  and, by Definition 2.2, its expectation can be computed as follows:

$$\begin{aligned} E_n(\lambda, \mu, \nu) &= \frac{1 - e^{-(\lambda+\nu)/2^n}}{\lambda + \nu} \frac{e^{\nu/2^n} - S_n^+(\lambda, \mu, \nu)}{e^{\nu/2^n} - 1} + \frac{1 - e^{-\lambda/2^n}}{\lambda} \frac{S_n^-(\lambda, \mu, \nu) - 1}{e^{\nu/2^n} - 1} \\ &= \left( \frac{e^{\nu/2^n} (1 - e^{-(\lambda+\nu)/2^n})}{(\lambda + \nu)(e^{\nu/2^n} - 1)} - \frac{1 - e^{-\lambda/2^n}}{\lambda(e^{\nu/2^n} - 1)} \right) \\ & \quad + \frac{1 - e^{-\lambda/2^n}}{\lambda(e^{\nu/2^n} - 1)} S_n^-(\lambda, \mu, \nu) - \frac{1 - e^{-(\lambda+\nu)/2^n}}{(\lambda + \nu)(e^{\nu/2^n} - 1)} S_n^+(\lambda, \mu, \nu). \end{aligned} \quad (3.2)$$

Now, we have to evaluate the limit  $E(\lambda, \mu, \nu)$  of  $E_n(\lambda, \mu, \nu)$  as  $n$  goes toward infinity. It is easy to see that this limit exists; see the proof of Theorem 3.1 below. Formally, we write  $E(\lambda, \mu, \nu) = \mathbb{E}[F(\lambda, \mu, \nu)]$  with

$$F(\lambda, \mu, \nu) = \int_0^{\infty} e^{-\lambda t + i\mu X(t) - \nu T(t)} dt.$$

Then, we can say that the functional  $F(\lambda, \mu, \nu)$  is an admissible function of  $X$  in the sense of Definition 2.3. The value of its expectation  $E(\lambda, \mu, \nu)$  is given in the following theorem.

**Theorem 3.1.** *The 3-parameters Laplace-Fourier transform of the couple  $(T(t), X(t))$  is given by*

$$E(\lambda, \mu, \nu) = \frac{1}{\prod_{j \in J} (\sqrt[\nu]{\lambda + \nu} - i\mu\theta_j) \prod_{k \in K} (\sqrt[\nu]{\lambda} - i\mu\theta_k)}. \quad (3.3)$$

**PROOF**

It is plain that the term lying within the biggest parentheses in the last equality of (3.2) tends to zero as  $n$  goes towards infinity and that the coefficients lying before  $S_n^+(\lambda, \mu, \nu)$  and  $S_n^-(\lambda, \mu, \nu)$  tend to  $1/\nu$ . As a byproduct, we derive at the limit when  $n \rightarrow \infty$ ,

$$E(\lambda, \mu, \nu) = \frac{1}{\nu} [S^-(\lambda, \mu, \nu) - S^+(\lambda, \mu, \nu)] \quad (3.4)$$

where we set

$$S^+(\lambda, \mu, \nu) = \lim_{n \rightarrow \infty} S_n^+(\lambda, \mu, \nu) = \exp\left(-\int_0^{\infty} \mathbb{E}[e^{i\mu X(t)} \mathbb{1}_{[0,+\infty)}(X(t))](1 - e^{-\nu t}) \frac{e^{-\lambda t}}{t} dt\right),$$

$$S^-(\lambda, \mu, \nu) = \lim_{n \rightarrow \infty} S_n^-(\lambda, \mu, \nu) = \exp\left(\int_0^\infty \mathbb{E}[e^{i\mu X(t)} \mathbf{1}_{(-\infty, 0)}(X(t))](1 - e^{-\nu t}) \frac{e^{-\lambda t}}{t} dt\right).$$

We have

$$\begin{aligned} & \int_0^\infty \mathbb{E}[e^{i\mu X(t)} \mathbf{1}_{[0, +\infty)}(X(t))](1 - e^{-\nu t}) \frac{e^{-\lambda t}}{t} dt \\ &= \int_0^\infty \mathbb{E}[(e^{i\mu X(t)} - 1) \mathbf{1}_{[0, +\infty)}(X(t))] \frac{e^{-\lambda t}}{t} dt - \int_0^\infty \mathbb{E}[(e^{i\mu X(t)} - 1) \mathbf{1}_{[0, +\infty)}(X(t))] \frac{e^{-(\lambda+\nu)t}}{t} dt \\ &+ \int_0^\infty \mathbb{P}\{X(t) \geq 0\} \frac{e^{-\lambda t} - e^{-(\lambda+\nu)t}}{t} dt \\ &= \int_0^\infty \frac{e^{-\lambda t}}{t} dt \int_0^\infty (e^{i\mu \xi} - 1) p(t; -\xi) d\xi - \int_0^\infty \frac{e^{-(\lambda+\nu)t}}{t} dt \int_0^\infty (e^{i\mu \xi} - 1) p(t; -\xi) d\xi \\ &+ \mathbb{P}\{X(1) \geq 0\} \int_0^\infty \frac{e^{-\lambda t} - e^{-(\lambda+\nu)t}}{t} dt. \end{aligned}$$

In view of (2.12) and (2.13) and using the elementary equality  $\int_0^\infty \frac{e^{-\lambda t} - e^{-(\lambda+\nu)t}}{t} dt = \log\left(\frac{\lambda+\nu}{\lambda}\right)$ , we have

$$\begin{aligned} & \int_0^\infty \mathbb{E}[e^{i\mu X(t)} \mathbf{1}_{[0, +\infty)}(X(t))](1 - e^{-\nu t}) \frac{e^{-\lambda t}}{t} dt \\ &= \log\left(\prod_{j \in J} \frac{\sqrt[N]{\lambda}}{\sqrt[N]{\lambda} - i\mu\theta_j}\right) - \log\left(\prod_{j \in J} \frac{\sqrt[N]{\lambda + \nu}}{\sqrt[N]{\lambda + \nu} - i\mu\theta_j}\right) + \frac{\#J}{N} \log\left(\frac{\lambda + \nu}{\lambda}\right) = \log\left(\prod_{j \in J} \frac{\sqrt[N]{\lambda + \nu} - i\mu\theta_j}{\sqrt[N]{\lambda} - i\mu\theta_j}\right). \end{aligned}$$

We then deduce the value of  $S^+(\lambda, \mu, \nu)$ . By (2.2),

$$\begin{aligned} S^+(\lambda, \mu, \nu) &= \prod_{j \in J} \frac{\sqrt[N]{\lambda} - i\mu\theta_j}{\sqrt[N]{\lambda + \nu} - i\mu\theta_j} = \frac{\prod_{\ell=1}^N (\sqrt[N]{\lambda} - i\mu\theta_\ell)}{\prod_{j \in J} (\sqrt[N]{\lambda + \nu} - i\mu\theta_j) \prod_{k \in K} (\sqrt[N]{\lambda} - i\mu\theta_k)} \\ &= \frac{\lambda - \kappa_N (i\mu)^N}{\prod_{j \in J} (\sqrt[N]{\lambda + \nu} - i\mu\theta_j) \prod_{k \in K} (\sqrt[N]{\lambda} - i\mu\theta_k)}. \end{aligned} \quad (3.5)$$

Similarly, the value of  $S^-(\lambda, \mu, \nu)$  is given by

$$S^-(\lambda, \mu, \nu) = \prod_{k \in K} \frac{\sqrt[N]{\lambda + \nu} - i\mu\theta_k}{\sqrt[N]{\lambda} - i\mu\theta_k} = \frac{\lambda + \nu - \kappa_N (i\mu)^N}{\prod_{j \in J} (\sqrt[N]{\lambda + \nu} - i\mu\theta_j) \prod_{k \in K} (\sqrt[N]{\lambda} - i\mu\theta_k)}. \quad (3.6)$$

Finally, putting (3.5) and (3.6) into (3.4) immediately leads to (3.3).  $\blacksquare$

**Remark 3.1.** We can rewrite (3.3) as

$$E(\lambda, \mu, \nu) = \frac{1}{\lambda^{\frac{\#K}{N}} (\lambda + \nu)^{\frac{\#J}{N}}} \prod_{j \in J} \frac{\sqrt[N]{\lambda + \nu}}{\sqrt[N]{\lambda + \nu} - i\mu\theta_j} \prod_{k \in K} \frac{\sqrt[N]{\lambda}}{\sqrt[N]{\lambda} - i\mu\theta_k}. \quad (3.7)$$

Actually, this form is more suitable for the inversion of the Laplace-Fourier transform.

In the three next sections, we progressively invert the 3-parameters Laplace-Fourier transform  $E(\lambda, \mu, \nu)$ .

## 4 Inverting with respect to $\mu$

In this part, we invert  $E(\lambda, \mu, \nu)$  given by (3.7) with respect to  $\mu$ .

**Theorem 4.1.** *We have, for  $\lambda, \nu > 0$ ,*

$$\int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx)/dx] dt$$

$$= \begin{cases} \frac{1}{\lambda^{\frac{\#K-1}{N}}(\lambda+\nu)^{\frac{\#J-1}{N}}} \sum_{j \in J} A_j \theta_j \left( \sum_{k \in K} \frac{B_k \theta_k}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda+\nu}} \right) e^{-\theta_j \sqrt[N]{\lambda+\nu} x} & \text{if } x \geq 0, \\ \frac{1}{\lambda^{\frac{\#K-1}{N}}(\lambda+\nu)^{\frac{\#J-1}{N}}} \sum_{k \in K} B_k \theta_k \left( \sum_{j \in J} \frac{A_j \theta_j}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda+\nu}} \right) e^{-\theta_k \sqrt[N]{\lambda} x} & \text{if } x \leq 0. \end{cases} \quad (4.1)$$

PROOF

By (2.6) applied to  $x = i\mu/\sqrt[N]{\lambda+\nu}$  and  $x = i\mu/\sqrt[N]{\lambda}$ , we have

$$\begin{aligned} \prod_{j \in J} \frac{\sqrt[N]{\lambda+\nu}}{\sqrt[N]{\lambda+\nu} - i\mu\theta_j} \prod_{k \in K} \frac{\sqrt[N]{\lambda}}{\sqrt[N]{\lambda} - i\mu\theta_k} &= \prod_{j \in J} \frac{1}{1 - \frac{i\mu}{\sqrt[N]{\lambda+\nu}} \theta_j} \prod_{k \in K} \frac{1}{1 - \frac{i\mu}{\sqrt[N]{\lambda}} \theta_k} \\ &= \sum_{j \in J} \frac{A_j \theta_j}{\theta_j - \frac{i\mu}{\sqrt[N]{\lambda+\nu}}} \sum_{k \in K} \frac{B_k \theta_k}{\theta_k - \frac{i\mu}{\sqrt[N]{\lambda}}} \\ &= \sqrt[N]{\lambda(\lambda+\nu)} \sum_{\substack{j \in J \\ k \in K}} \frac{A_j B_k \theta_j \theta_k}{(\theta_j \sqrt[N]{\lambda+\nu} - i\mu)(\theta_k \sqrt[N]{\lambda} - i\mu)}. \end{aligned}$$

Let us write that

$$\begin{aligned} \frac{1}{(\theta_j \sqrt[N]{\lambda+\nu} - i\mu)(\theta_k \sqrt[N]{\lambda} - i\mu)} &= \frac{1}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda+\nu}} \left( \frac{1}{\theta_j \sqrt[N]{\lambda+\nu} - i\mu} - \frac{1}{\theta_k \sqrt[N]{\lambda} - i\mu} \right) \\ &= \frac{1}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda+\nu}} \left( \int_0^\infty e^{(i\mu - \theta_j \sqrt[N]{\lambda+\nu})x} dx + \int_{-\infty}^0 e^{(i\mu - \theta_k \sqrt[N]{\lambda})x} dx \right). \end{aligned}$$

Therefore, we can rewrite  $E(\lambda, \mu, \nu)$  as

$$\begin{aligned} E(\lambda, \mu, \nu) &= \frac{1}{\lambda^{\frac{\#K-1}{N}}(\lambda+\nu)^{\frac{\#J-1}{N}}} \\ &\quad \times \sum_{\substack{j \in J \\ k \in K}} \frac{A_j B_k \theta_j \theta_k}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda+\nu}} \int_{-\infty}^\infty e^{i\mu x} \left( e^{-\theta_k \sqrt[N]{\lambda} x} \mathbb{1}_{(-\infty, 0]}(x) + e^{-\theta_j \sqrt[N]{\lambda+\nu} x} \mathbb{1}_{[0, \infty)}(x) \right) dx \end{aligned}$$

which is nothing but the Fourier transform with respect to  $\mu$  of the right-hand side of (4.1). ■

**Remark 4.1.** One can observe that formula (24) in [11] involves the density of  $(T(t), X(t))$ , this latter being evaluated at the extremity  $X(t) = 0$  when the starting point is  $x$ . By invoking the duality, we could derive an alternative representation for (4.1). Nevertheless, this representation is not tractable for performing the inversion with respect to  $\nu$ .

**Example 4.1.** For  $N = 3$ , we have two cases to consider. Although this situation is not correctly defined, (4.1) writes formally, with the numerical values of Example 2.1, in the case  $\kappa_3 = 1$ ,

$$\begin{aligned} \int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx)/dx] dt \\ = \begin{cases} \frac{e^{-\sqrt[3]{\lambda+\nu} x}}{\lambda^{2/3} + \sqrt[3]{\lambda(\lambda+\nu)} + (\lambda+\nu)^{2/3}} & \text{if } x \geq 0, \\ \frac{e^{\frac{\sqrt[3]{\lambda}}{2} x} \sqrt{3} \sqrt[3]{\lambda} \cos\left(\frac{\sqrt{3}\sqrt[3]{\lambda}}{2} x\right) - (2\sqrt[3]{\lambda+\nu} + \sqrt[3]{\lambda}) \sin\left(\frac{\sqrt{3}\sqrt[3]{\lambda}}{2} x\right)}{\sqrt{3} \sqrt[3]{\lambda} \left( \lambda^{2/3} + \sqrt[3]{\lambda(\lambda+\nu)} + (\lambda+\nu)^{2/3} \right)} & \text{if } x \leq 0, \end{cases} \end{aligned}$$

and in the case  $\kappa_3 = -1$ ,

$$\int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx)/dx] dt$$

$$= \begin{cases} \frac{e^{-\frac{\sqrt[3]{\lambda+\nu}}{2}x} \sqrt{3} \sqrt[3]{\lambda+\nu} \cos\left(\frac{\sqrt{3}\sqrt[3]{\lambda+\nu}}{2}x\right) + (\sqrt[3]{\lambda+\nu} + 2\sqrt[3]{\lambda}) \sin\left(\frac{\sqrt{3}\sqrt[3]{\lambda+\nu}}{2}x\right)}{\sqrt{3}\sqrt[3]{\lambda+\nu} \lambda^{2/3} + \sqrt[3]{\lambda(\lambda+\nu)} + (\lambda+\nu)^{2/3}} & \text{if } x \geq 0, \\ \frac{e^{\sqrt[3]{\lambda}x}}{\lambda^{2/3} + \sqrt[3]{\lambda(\lambda+\nu)} + (\lambda+\nu)^{2/3}} & \text{if } x \leq 0. \end{cases}$$

**Example 4.2.** For  $N = 4$ , formula (4.1) supplies, with the numerical values of Example 2.2,

$$\int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx)/dx] dt = \begin{cases} \frac{\sqrt{2} e^{-\frac{\sqrt[4]{\lambda+\nu}}{\sqrt{2}}x}}{\sqrt[4]{\lambda+\nu}(\sqrt{\lambda+\nu})\sqrt[4]{\lambda+\nu}} \left[ \sqrt[4]{\lambda+\nu} \cos\left(\frac{\sqrt[4]{\lambda+\nu}}{\sqrt{2}}x\right) + \sqrt[4]{\lambda} \sin\left(\frac{\sqrt[4]{\lambda+\nu}}{\sqrt{2}}x\right) \right] & \text{if } x \geq 0, \\ \frac{\sqrt{2} e^{\frac{\sqrt[4]{\lambda}}{\sqrt{2}}x}}{\sqrt[4]{\lambda}(\sqrt{\lambda+\nu})\sqrt[4]{\lambda+\nu}} \left[ \sqrt[4]{\lambda} \cos\left(\frac{\sqrt[4]{\lambda}}{\sqrt{2}}x\right) - \sqrt[4]{\lambda+\nu} \sin\left(\frac{\sqrt[4]{\lambda}}{\sqrt{2}}x\right) \right] & \text{if } x \leq 0. \end{cases}$$

## 5 Inverting with respect to $\nu$

In this section, we carry out the inversion with respect to the parameter  $\nu$ . The cases  $x \leq 0$  and  $x \geq 0$  lead to results which are not quite analogous. This is due to the asymmetry of our problem. So, we split our analysis into two subsections related to the cases  $x \leq 0$  and  $x \geq 0$ .

### 5.1 The case $x \leq 0$

**Theorem 5.1.** *The Laplace transform with respect to  $t$  of the density of the couple  $(T(t), X(t))$  is given, when  $x \leq 0$ , by*

$$\int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt = -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{m=0}^{\#K} \alpha_{-m}(\lambda s)^{\frac{m}{N}} E_{1, \frac{m+\#J}{N}}(\lambda s) \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k \sqrt[4]{\lambda} x}. \quad (5.1)$$

PROOF

Recall (4.1) in the case  $x \leq 0$ :

$$\int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx)/dx] dt = \frac{1}{\lambda^{\frac{\#K-1}{N}} (\lambda+\nu)^{\frac{\#J-1}{N}}} \sum_{k \in K} B_k \theta_k \left( \sum_{j \in J} \frac{A_j \theta_j}{\theta_k \sqrt[4]{\lambda} - \theta_j \sqrt[4]{\lambda+\nu}} \right) e^{-\theta_k \sqrt[4]{\lambda} x}.$$

We have to invert with respect to  $\nu$  the quantity

$$\frac{1}{(\lambda+\nu)^{\frac{\#J-1}{N}}} \sum_{j \in J} \frac{A_j \theta_j}{\theta_k \sqrt[4]{\lambda} - \theta_j \sqrt[4]{\lambda+\nu}} = - \sum_{j \in J} \frac{A_j}{(\lambda+\nu)^{\frac{\#J-1}{N}} (\sqrt[4]{\lambda+\nu} - \frac{\theta_k}{\theta_j} \sqrt[4]{\lambda})}.$$

By using the following elementary equality, which is valid for  $\alpha > 0$ ,

$$\frac{1}{(\lambda+\nu)^\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty e^{-(\lambda+\nu)s} s^{\alpha-1} ds = \int_0^\infty e^{-\nu s} \left( \frac{s^{\alpha-1} e^{-\lambda s}}{\Gamma(\alpha)} \right) ds,$$

we obtain, for  $|\beta| < \sqrt[4]{\lambda+\nu}$ ,

$$\frac{1}{\sqrt[4]{\lambda+\nu} - \beta} = \frac{1}{\sqrt[4]{\lambda+\nu}} \frac{1}{1 - \frac{\beta}{\sqrt[4]{\lambda+\nu}}} = \sum_{r=0}^{\infty} \frac{\beta^r}{(\lambda+\nu)^{\frac{r+1}{N}}} = \sum_{r=0}^{\infty} \frac{\beta^r}{\Gamma\left(\frac{r+1}{N}\right)} \int_0^\infty e^{-(\lambda+\nu)s} s^{\frac{r+1}{N}-1} ds$$

$$= \int_0^\infty e^{-\nu s} \left( s^{\frac{1}{N}-1} e^{-\lambda s} \sum_{r=0}^\infty \frac{(\beta \sqrt[N]{s})^r}{\Gamma(\frac{r+1}{N})} \right) ds.$$

The sum lying in the last displayed equality can be expressed by means of the Mittag-Leffler function (see [7, Chap. XVIII]):  $E_{a,b}(\xi) = \sum_{r=0}^\infty \frac{\xi^r}{\Gamma(ar+b)}$ . Then,

$$\frac{1}{\sqrt[N]{\lambda + \nu} - \beta} = \int_0^\infty e^{-\nu s} \left( s^{\frac{1}{N}-1} e^{-\lambda s} E_{\frac{1}{N}, \frac{1}{N}}(\beta \sqrt[N]{s}) \right) ds. \quad (5.2)$$

Next, we write

$$\sum_{j \in J} \frac{A_j}{\sqrt[N]{\lambda + \nu} - \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda}} = \int_0^\infty e^{-\nu s} \left[ s^{\frac{1}{N}-1} e^{-\lambda s} \sum_{j \in J} A_j E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda s} \right) \right] ds, \quad (5.3)$$

where

$$\sum_{j \in J} A_j E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda s} \right) = \sum_{j \in J} A_j \sum_{r=0}^\infty \left( \frac{\theta_k}{\theta_j} \right)^r \frac{(\lambda s)^{\frac{r}{N}}}{\Gamma(\frac{r+1}{N})} = \sum_{r=0}^\infty \left( \theta_k^r \sum_{j \in J} \frac{A_j}{\theta_j^r} \right) \frac{(\lambda s)^{\frac{r}{N}}}{\Gamma(\frac{r+1}{N})}.$$

When performing the euclidian division of  $r$  by  $N$ , we can write  $r$  as  $r = \ell N + m$  with  $\ell \geq 0$  and  $0 \leq m \leq N - 1$ . With this, we have  $\theta_j^{-r} = (\theta_j^N)^{-\ell} \theta_j^{-m} = \kappa_N^\ell \theta_j^{-m}$  and  $\theta_k^r = \kappa_N^\ell \theta_k^m$ . Then,

$$\theta_k^r \sum_{j \in J} \frac{A_j}{\theta_j^r} = \theta_k^m \sum_{j \in J} \frac{A_j}{\theta_j^m} = \theta_k^m \alpha_{-m}.$$

Hence, since by (2.9) the  $\alpha_{-m}$ ,  $\#K + 1 \leq m \leq N$ , vanish,

$$\sum_{j \in J} A_j E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda s} \right) = \sum_{\ell=0}^\infty \sum_{m=0}^{\#K} \alpha_{-m} \theta_k^m \frac{(\lambda s)^{\ell + \frac{m}{N}}}{\Gamma(\ell + \frac{m+1}{N})} = \sum_{m=0}^{\#K} \alpha_{-m} \theta_k^m (\lambda s)^{\frac{m}{N}} E_{1, \frac{m+1}{N}}(\lambda s)$$

and (5.3) becomes

$$\sum_{j \in J} \frac{A_j}{\sqrt[N]{\lambda + \nu} - \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda}} = \int_0^\infty e^{-\nu s} \left( s^{\frac{1}{N}-1} e^{-\lambda s} \sum_{m=0}^{\#K} \alpha_{-m} \theta_k^m (\lambda s)^{\frac{m}{N}} E_{1, \frac{m+1}{N}}(\lambda s) \right) ds.$$

As a result, by introducing a convolution product, we obtain

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx) / dx] dt \\ &= -\frac{1}{\lambda^{\frac{\#K-1}{N}}} \sum_{k \in K} B_k \theta_k e^{-\theta_k \sqrt[N]{\lambda} x} \\ & \quad \times \int_0^\infty e^{-\nu s} \left( \int_0^s \frac{\sigma^{\frac{\#J-1}{N}-1} e^{-\lambda \sigma}}{\Gamma(\frac{\#J-1}{N})} \times e^{-\lambda(s-\sigma)} \sum_{m=0}^{\#K} \alpha_{-m} \theta_k^m \lambda^{\frac{m}{N}} (s-\sigma)^{\frac{m+1}{N}-1} E_{1, \frac{m+1}{N}}(\lambda(s-\sigma)) d\sigma \right) ds. \end{aligned}$$

By removing the Laplace transforms with respect to the parameter  $\nu$  of each member of the foregoing equality, we extract

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx)] dt \\ &= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}}} \sum_{m=0}^{\#K} \alpha_{-m} \lambda^{\frac{m}{N}} \left( \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k \sqrt[N]{\lambda} x} \right) \int_0^s \frac{\sigma^{\frac{\#J-1}{N}-1}}{\Gamma(\frac{\#J-1}{N})} (s-\sigma)^{\frac{m+1}{N}-1} E_{1, \frac{m+1}{N}}(\lambda(s-\sigma)) d\sigma. \end{aligned}$$

The integral lying on the right-hand side of the previous equality can be evaluated as follows:

$$\int_0^s \frac{\sigma^{\frac{\#J-1}{N}-1}}{\Gamma(\frac{\#J-1}{N})} (s-\sigma)^{\frac{m+1}{N}-1} E_{1, \frac{m+1}{N}}(\lambda(s-\sigma)) d\sigma = \int_0^s \frac{\sigma^{\frac{\#J-1}{N}-1}}{\Gamma(\frac{\#J-1}{N})} (s-\sigma)^{\frac{m+1}{N}-1} \sum_{\ell=0}^\infty \frac{\lambda^\ell (s-\sigma)^\ell}{\Gamma(\ell + \frac{m+1}{N})} d\sigma$$

$$\begin{aligned}
&= \sum_{\ell=0}^{\infty} \lambda^{\ell} \int_0^s \frac{\sigma^{\frac{\#J-1}{N}-1}}{\Gamma(\frac{\#J-1}{N})} \frac{(s-\sigma)^{\ell+\frac{m+1}{N}-1}}{\Gamma(\ell+\frac{m+1}{N})} d\sigma \\
&= \sum_{\ell=0}^{\infty} \frac{\lambda^{\ell} s^{\ell+\frac{m+\#J}{N}-1}}{\Gamma(\ell+\frac{m+\#J}{N})} = s^{\frac{m+\#J}{N}-1} E_{1, \frac{m+\#J}{N}}(\lambda s)
\end{aligned}$$

from which we deduce (5.1). ■

**Remark 5.1.** An alternative expression for formula (5.1) is for  $x \leq 0$

$$\begin{aligned}
&\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{\substack{j \in J \\ k \in K}} A_j B_k \theta_k E_{\frac{1}{N}, \frac{\#J}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda s} \right) e^{-\theta_k \sqrt[N]{\lambda} x}. \tag{5.4}
\end{aligned}$$

In effect, by (5.1),

$$\begin{aligned}
&\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{\ell=0}^{\infty} \sum_{m=0}^{\#K} \sum_{k \in K} \alpha_{-m} B_k \theta_k^{m+1} \frac{(\lambda s)^{\ell+\frac{m}{N}}}{\Gamma(\ell+\frac{m+\#J}{N})} e^{-\theta_k \sqrt[N]{\lambda} x} \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{\ell=0}^{\infty} \sum_{m=0}^{N-1} \sum_{\substack{j \in J \\ k \in K}} A_j B_k \theta_k \left( \frac{\theta_k}{\theta_j} \right)^m \frac{(\lambda s)^{\ell+\frac{m}{N}}}{\Gamma(\ell+\frac{m+\#J}{N})} e^{-\theta_k \sqrt[N]{\lambda} x}.
\end{aligned}$$

In the last displayed equality, we have extended the sum with respect to  $m$  to the range  $0 \leq m \leq N-1$  because, by (2.9), the  $\alpha_{-m}$ ,  $\#K+1 \leq m \leq N-1$ , vanish. Let us introduce the index  $r = \ell N + m$ . Since  $\left(\frac{\theta_k}{\theta_j}\right)^m = \left(\frac{\theta_k}{\theta_j}\right)^r$ , we have

$$\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt = -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{\substack{j \in J \\ k \in K}} A_j B_k \theta_k \sum_{r=0}^{\infty} \frac{\left(\frac{\theta_k}{\theta_j} \sqrt[N]{\lambda s}\right)^r}{\Gamma\left(\frac{r+\#J}{N}\right)} e^{-\theta_k \sqrt[N]{\lambda} x}$$

which coincide with (5.4).

**Example 5.1.** Case  $N = 3$ . We have formally for  $x \leq 0$ , when  $\kappa_3 = -1$ :

$$\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt = e^{\sqrt[3]{\lambda} x} \left( \frac{e^{-\lambda s}}{\sqrt[3]{s}} E_{1, \frac{2}{3}}(\lambda s) - \sqrt[3]{\lambda} \right)$$

and when  $\kappa_3 = 1$ :

$$\begin{aligned}
&\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= \frac{e^{-\lambda s + \frac{\sqrt[3]{\lambda} x}{2}}}{\sqrt{3} \sqrt[3]{\lambda s^2}} \left[ \sqrt{3} \cos\left(\frac{\sqrt{3} \sqrt[3]{\lambda} x}{2}\right) \left( \sqrt[3]{\lambda s} E_{1, \frac{2}{3}}(\lambda s) - (\lambda s)^{2/3} e^{\lambda s} \right) \right. \\
&\quad \left. + \sin\left(\frac{\sqrt{3} \sqrt[3]{\lambda} x}{2}\right) \left( \sqrt[3]{\lambda s} E_{1, \frac{2}{3}}(\lambda s) + (\lambda s)^{2/3} e^{\lambda s} - 2E_{1, \frac{1}{3}}(\lambda s) \right) \right].
\end{aligned}$$

**Example 5.2.** Case  $N = 4$ . We have, for  $x \geq 0$ ,

$$\begin{aligned}
&\int_0^{\infty} e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= \frac{\sqrt{2} e^{-\lambda s + \frac{\sqrt[4]{\lambda} x}{\sqrt{2}}}}{\sqrt[4]{\lambda} \sqrt{s}} \left[ \cos\left(\frac{\sqrt[4]{\lambda} x}{\sqrt{2}}\right) \left( \sqrt[4]{\lambda s} E_{1, \frac{3}{4}}(\lambda s) - \sqrt{\lambda s} e^{\lambda s} \right) + \sin\left(\frac{\sqrt[4]{\lambda} x}{\sqrt{2}}\right) \left( \sqrt[4]{\lambda s} E_{1, \frac{3}{4}}(\lambda s) - E_{1, \frac{1}{2}}(\lambda s) \right) \right].
\end{aligned}$$

## 5.2 The case $x \geq 0$

**Theorem 5.2.** *The Laplace transform with respect to  $t$  of the density of the couple  $(T(t), X(t))$  is given, when  $x \geq 0$ , by*

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx)] dt \\ &= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}}} \sum_{\substack{j \in J \\ k \in K}} A_j B_k \theta_k \int_0^s \sigma^{\frac{1}{N}-1} E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda \sigma} \right) I_{j, \#J-1}(s - \sigma; x) d\sigma \end{aligned} \quad (5.5)$$

where the function  $I_{j, \#J-1}$  is defined by (2.14).

PROOF

Recall (4.1) in the case  $x \geq 0$ :

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} [\mathbb{E}(e^{-\nu T(t)}, X(t) \in dx) / dx] dt \\ &= \frac{1}{\lambda^{\frac{\#K-1}{N}} (\lambda + \nu)^{\frac{\#J-1}{N}}} \sum_{j \in J} A_j \theta_j \left( \sum_{k \in K} \frac{B_k \theta_k}{\theta_k \sqrt[N]{\lambda} - \theta_j \sqrt[N]{\lambda + \nu}} \right) e^{-\theta_j \sqrt[N]{\lambda + \nu} x}. \end{aligned}$$

We have to invert the quantity  $\frac{e^{-\theta_j \sqrt[N]{\lambda + \nu} x}}{(\lambda + \nu)^{\frac{\#J-1}{N}} (\sqrt[N]{\lambda + \nu} - \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda})}$  with respect to  $\nu$ . Recalling (5.2) and (2.15),

$$\begin{aligned} \frac{1}{\sqrt[N]{\lambda + \nu} - \beta} &= \int_0^\infty e^{-\nu s} \left( s^{\frac{1}{N}-1} e^{-\lambda s} E_{\frac{1}{N}, \frac{1}{N}}(\beta \sqrt[N]{s}) \right) ds, \\ \frac{e^{-\theta_j \sqrt[N]{\lambda + \nu} x}}{(\lambda + \nu)^{\frac{\#J-1}{N}}} &= \int_0^\infty e^{-\nu s} (e^{-\lambda s} I_{j, \#J-1}(s; x)) ds, \end{aligned}$$

we get by convolution

$$\begin{aligned} & \frac{e^{-\theta_j \sqrt[N]{\lambda + \nu} x}}{(\lambda + \nu)^{\frac{\#J-1}{N}} (\sqrt[N]{\lambda + \nu} - \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda})} \\ &= \int_0^\infty e^{-\nu s} \left( \int_0^s \sigma^{\frac{1}{N}-1} e^{-\lambda \sigma} E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda \sigma} \right) \times e^{-\lambda(s-\sigma)} I_{j, \#J-1}(s - \sigma; x) d\sigma \right) ds \\ &= \int_0^\infty e^{-\nu s} \left( e^{-\lambda s} \int_0^s \sigma^{\frac{1}{N}-1} E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda \sigma} \right) I_{j, \#J-1}(s - \sigma; x) d\sigma \right) ds. \end{aligned}$$

This immediately yields (5.5). ■

**Remark 5.2.** Noticing that

$$E_{\frac{1}{N}, \frac{1}{N}} \left( \frac{\theta_k}{\theta_j} \sqrt[N]{\lambda \sigma} \right) = \sum_{r=0}^\infty \frac{\theta_k^r}{\theta_j^r} \frac{(\lambda \sigma)^{\frac{r}{N}}}{\Gamma(\frac{r+1}{N})} = \sum_{\ell=0}^\infty \sum_{m=0}^{N-1} \frac{\theta_k^m}{\theta_j^m} \frac{(\lambda \sigma)^{\ell + \frac{m}{N}}}{\Gamma(\ell + \frac{m+1}{N})} = \sum_{m=0}^{N-1} \frac{\theta_k^m}{\theta_j^m} (\lambda \sigma)^{\frac{m}{N}} E_{1, \frac{m+1}{N}}(\lambda \sigma)$$

and reminding that, from (2.8), the  $\beta_m$ ,  $1 \leq m \leq \#K - 1$ , vanish, we can rewrite (5.5) in the following form. For  $x \geq 0$ ,

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx)] dt \\ &= -e^{-\lambda s} \sum_{m=\#K-1}^{N-1} \left( \sum_{k \in K} B_k \theta_k^{m+1} \right) \lambda^{\frac{m-\#K+1}{N}} \int_0^s \sigma^{\frac{m+1}{N}-1} E_{1, \frac{m+1}{N}}(\lambda \sigma) \left( \sum_{j \in J} \frac{A_j}{\theta_j^m} I_{j, \#J-1}(s - \sigma; x) \right) d\sigma \\ &= -e^{-\lambda s} \sum_{m=\#K}^N \beta_m \lambda^{\frac{m-\#K}{N}} \int_0^s \sigma^{\frac{m}{N}-1} E_{1, \frac{m}{N}}(\lambda \sigma) \Phi_m(s - \sigma; x) d\sigma \end{aligned} \quad (5.6)$$

with  $\Phi_m(\tau; x) = \sum_{j \in J} \frac{A_j}{\theta_j^{m-1}} I_{j, \#J-1}(\tau; x)$ .

**Remark 5.3.** For  $x = 0$ , using formula (5.1) which is valid for  $x \leq 0$ , we get, by (2.8), (2.9) and (2.10),

$$\begin{aligned}
& \int_0^\infty e^{-\lambda t} \mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \Big|_{x=0} dt \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{m=0}^{\#K} \alpha_{-m} \beta_{m+1} (\lambda s)^{\frac{m}{N}} E_{1, \frac{m+\#J}{N}} (\lambda s) \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \left( \alpha_{1-\#K} \beta_{\#K} (\lambda s)^{\frac{\#K-1}{N}} E_{1, 1-\frac{1}{N}} (\lambda s) + \alpha_{-\#K} \beta_{\#K+1} (\lambda s)^{\frac{\#K}{N}} E_{1,1} (\lambda s) \right) \\
&= \frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \left( \sum_{j \in J} \theta_j (\lambda s)^{\frac{\#K-1}{N}} E_{1, 1-\frac{1}{N}} (\lambda s) + \sum_{k \in K} \theta_k (\lambda s)^{\frac{\#K}{N}} e^{\lambda s} \right) \\
&= \left( \sum_{j \in J} \theta_j \right) \frac{e^{-\lambda s}}{\sqrt[N]{s}} \left( E_{1, 1-\frac{1}{N}} (\lambda s) - \sqrt[N]{\lambda s} e^{\lambda s} \right). \tag{5.7}
\end{aligned}$$

On the other hand, with formula (5.6) which is valid for  $x \geq 0$ ,

$$\begin{aligned}
& \int_0^\infty e^{-\lambda t} \mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \Big|_{x=0} dt \\
&= -e^{-\lambda s} \sum_{m=\#K}^N \beta_m \lambda^{\frac{m-\#K}{N}} \int_0^s \sigma^{\frac{m}{N}-1} E_{1, \frac{m}{N}} (\lambda \sigma) \Phi_m(s-\sigma; 0) d\sigma \tag{5.8}
\end{aligned}$$

with

$$\begin{aligned}
\Phi_m(\tau; 0) &= \frac{Ni}{2\pi} \left( \sum_{j \in J} \frac{A_j}{\theta_j^{m-1}} \right) \left( e^{-i \frac{\#J-1}{N} \pi} - e^{i \frac{\#J-1}{N} \pi} \right) \int_0^\infty \xi^{\#K} e^{-\tau \xi^N} d\xi \\
&= \frac{\Gamma(\frac{\#K+1}{N}) \sin(\frac{\#J-1}{N} \pi)}{\pi \tau^{\frac{\#K+1}{N}}} \alpha_{1-m} = \frac{\alpha_{1-m}}{\Gamma(\frac{\#J-1}{N}) \tau^{\frac{\#K+1}{N}}}.
\end{aligned}$$

In view of (2.8), (2.9) and (2.10), we have

$$\begin{aligned}
& \int_0^\infty e^{-\lambda t} \mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \Big|_{x=0} dt \\
&= \frac{e^{-\lambda s}}{\Gamma(\frac{\#J-1}{N})} \left[ \left( \sum_{j \in J} \theta_j \right) \int_0^s \frac{\sigma^{\frac{\#K}{N}-1}}{(s-\sigma)^{\frac{\#K+1}{N}}} E_{1, \frac{\#K}{N}} (\lambda \sigma) d\sigma \right. \\
&\quad \left. + \left( \sum_{k \in K} \theta_k \right) \sqrt[N]{\lambda} \int_0^s \frac{\sigma^{\frac{\#K+1}{N}-1}}{(s-\sigma)^{\frac{\#K+1}{N}}} E_{1, \frac{\#K+1}{N}} (\lambda \sigma) d\sigma \right] \\
&= \left( \sum_{j \in J} \theta_j \right) \frac{e^{-\lambda s}}{\Gamma(\frac{\#J-1}{N})} e^{-\lambda s} \left( \sum_{\ell=0}^\infty \frac{B(\ell + \frac{\#K}{N}, 1 - \frac{\#K+1}{N})}{\Gamma(\ell + \frac{\#K}{N})} \lambda^\ell s^{\ell - \frac{1}{N}} \right. \\
&\quad \left. - \sqrt[N]{\lambda} \sum_{\ell=0}^\infty \frac{B(\ell + \frac{\#K+1}{N}, 1 - \frac{\#K+1}{N})}{\Gamma(\ell + \frac{\#K+1}{N})} (\lambda s)^\ell \right) \\
&= \left( \sum_{j \in J} \theta_j \right) \frac{e^{-\lambda s}}{\sqrt[N]{s}} \left( E_{1, 1-\frac{1}{N}} (\lambda s) - \sqrt[N]{\lambda s} e^{\lambda s} \right).
\end{aligned}$$

Thus, we have checked that the two different formulas (5.7) and (5.8) lead to the same result.

**Example 5.3.** Case  $N = 3$ . For  $x \geq 0$ , (5.5) supplies formally with the numerical values of Example 2.1, when  $\kappa_3 = -1$ ,

$$\int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx)] dt$$

$$\begin{aligned}
&= \frac{e^{-\lambda s}}{\sqrt{3}} \left( e^{\frac{i\pi}{6}} \int_0^s \sigma^{-2/3} E_{\frac{1}{3}, \frac{1}{3}}(-e^{-i\frac{\pi}{3}} \sqrt[3]{\lambda\sigma}) I_{1,1}(s-\sigma; x) d\sigma \right. \\
&\quad \left. + e^{-\frac{i\pi}{6}} \int_0^s \sigma^{-2/3} E_{\frac{1}{3}, \frac{1}{3}}(-e^{i\frac{\pi}{3}} \sqrt[3]{\lambda\sigma}) I_{2,1}(s-\sigma; x) d\sigma \right)
\end{aligned}$$

and when  $\kappa_3 = 1$ ,

$$\begin{aligned}
&\int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= \frac{i e^{-\lambda s}}{\sqrt{3} \sqrt[3]{\lambda}} \left( \int_0^s \sigma^{-2/3} E_{\frac{1}{3}, \frac{1}{3}}(e^{-i\frac{2\pi}{3}} \sqrt[3]{\lambda\sigma}) I_{1,0}(s-\sigma; x) d\sigma \right. \\
&\quad \left. - \int_0^s \sigma^{-2/3} E_{\frac{1}{3}, \frac{1}{3}}(e^{i\frac{2\pi}{3}} \sqrt[3]{\lambda\sigma}) I_{1,0}(s-\sigma; x) d\sigma \right).
\end{aligned}$$

The functions  $I_{1,0}$ ,  $I_{1,1}$  and  $I_{2,1}$  above are respectively given by (2.16), (2.17) and (2.18).

**Example 5.4.** Case  $N = 4$ . For  $x \geq 0$ , (5.5) supplies, with the numerical values of Example 2.2,

$$\begin{aligned}
&\int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= -\frac{e^{-\lambda s}}{2\sqrt[4]{\lambda}} \left( e^{i\frac{\pi}{4}} \int_0^s \sigma^{-3/4} E_{\frac{1}{4}, \frac{1}{4}}(-\sqrt[4]{\lambda\sigma}) I_{1,1}(s-\sigma; x) d\sigma \right. \\
&\quad + e^{-i\frac{3\pi}{4}} \int_0^s \sigma^{-3/4} E_{\frac{1}{4}, \frac{1}{4}}(-i\sqrt[4]{\lambda\sigma}) I_{1,1}(s-\sigma; x) d\sigma \\
&\quad + e^{i\frac{3\pi}{4}} \int_0^s \sigma^{-3/4} E_{\frac{1}{4}, \frac{1}{4}}(i\sqrt[4]{\lambda\sigma}) I_{2,1}(s-\sigma; x) d\sigma \\
&\quad \left. + e^{-i\frac{\pi}{4}} \int_0^s \sigma^{-3/4} E_{\frac{1}{4}, \frac{1}{4}}(-\sqrt[4]{\lambda\sigma}) I_{2,1}(s-\sigma; x) d\sigma \right).
\end{aligned}$$

The functions  $I_{1,1}$  and  $I_{2,1}$  above are given by (2.19).

## 6 Inverting with respect to $\lambda$

In this section, we perform the last inversion in  $F(\lambda, \mu, \nu)$  in order to derive the distribution of the couple  $(T(t), X(t))$ . As in the previous section, we treat separately the two cases  $x \leq 0$  and  $x \geq 0$ .

### 6.1 The case $x \leq 0$

**Theorem 6.1.** *The distribution of the couple  $(T(t), X(t))$  is given, for  $x \leq 0$ , by*

$$\begin{aligned}
&\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/ds dx \\
&= -\frac{Ni}{2\pi} \sum_{m=0}^{\#K} \alpha_{-m} s^{\frac{m-\#K}{N}} \int_0^\infty \xi^{m+\#J} e^{-(t-s)\xi^N} \mathcal{K}_m(x\xi) E_{1, \frac{m+\#J}{N}}(-s\xi^N) d\xi
\end{aligned} \tag{6.1}$$

where

$$\mathcal{K}_m(z) = e^{-i\frac{\#K-m-1}{N}\pi} \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k e^{i\frac{\pi}{N}} z} - e^{i\frac{\#K-m-1}{N}\pi} \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k e^{-i\frac{\pi}{N}} z}.$$

**PROOF**

Assume  $x \leq 0$ . Recalling (5.1), we have

$$\begin{aligned}
&\int_0^\infty e^{-\lambda t} [\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)] dt \\
&= -\frac{e^{-\lambda s}}{\lambda^{\frac{\#K-1}{N}} s^{\frac{\#K}{N}}} \sum_{m=0}^{\#K} \alpha_{-m} (\lambda s)^{\frac{m}{N}} E_{1, \frac{m+\#J}{N}}(\lambda s) \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k \sqrt[N]{\lambda} x}
\end{aligned}$$

$$\begin{aligned}
&= -\sqrt[N]{\lambda} e^{-\lambda s} \sum_{m=0}^{\#K} \alpha_{-m} \sum_{\ell=0}^{\infty} \frac{(\lambda s)^{\ell + \frac{m-\#K}{N}}}{\Gamma(\ell + \frac{m+\#J}{N})} \sum_{k \in K} B_k \theta_k^{m+1} e^{-\theta_k \sqrt[N]{\lambda} x} \\
&= -\sum_{\ell=0}^{\infty} \sum_{m=0}^{\#K} \alpha_{-m} \frac{s^{\ell + \frac{m-\#K}{N}}}{\Gamma(\ell + \frac{m+\#J}{N})} \sum_{k \in K} B_k \theta_k^{m+1} \lambda^{\ell + \frac{m-\#K+1}{N}} e^{-\lambda s - \theta_k \sqrt[N]{\lambda} x}. \tag{6.2}
\end{aligned}$$

We need to invert the quantity  $\lambda^{\ell + \frac{m-\#K+1}{N}} e^{-\lambda s - \theta_k \sqrt[N]{\lambda} x}$  for  $\ell \geq 0$  and  $0 \leq m \leq \#K$  with respect to  $\lambda$ . We intend to use (2.15) which is valid for  $0 \leq m \leq N-1$ . Actually (2.15) holds true also for  $m \leq 0$ ; the proof of this claim is postponed to Lemma 7.3 in the appendix. As a byproduct, for any  $\ell \geq 0$  and  $0 \leq m \leq \#K$ ,

$$\begin{aligned}
\lambda^{\ell + \frac{m-\#K+1}{N}} e^{-\lambda s - \theta_k \sqrt[N]{\lambda} x} &= e^{-\lambda s} \int_0^{\infty} e^{-\lambda u} I_{k, \#K - \ell N - m - 1}(u; x) du \\
&= \int_s^{\infty} e^{-\lambda t} I_{k, \#K - \ell N - m - 1}(t - s; x) dt. \tag{6.3}
\end{aligned}$$

Then, by putting (6.3) into (6.2) and next by eliminating the Laplace transform with respect to  $\lambda$ , we extract

$$\begin{aligned}
&\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \\
&= -\sum_{\ell=0}^{\infty} \sum_{m=0}^{\#K} \alpha_{-m} \frac{s^{\ell + \frac{m-\#K}{N}}}{\Gamma(\ell + \frac{m+\#J}{N})} \sum_{k \in K} B_k \theta_k^{m+1} I_{k, \#K - \ell N - m - 1}(t - s; x) \\
&= -\frac{Ni}{2\pi} \sum_{\ell=0}^{\infty} \sum_{m=0}^{\#K} \alpha_{-m} \frac{s^{\ell + \frac{m-\#K}{N}}}{\Gamma(\ell + \frac{m+\#J}{N})} \\
&\quad \times \sum_{k \in K} B_k \theta_k^{m+1} \left( e^{-i \frac{\#K - \ell N - m - 1}{N} \pi} \int_0^{\infty} \xi^{N - \#K + \ell N + m} e^{-(t-s)\xi^N - \theta_k e^{i \frac{\pi}{N}} x \xi} d\xi \right. \\
&\quad \left. - e^{i \frac{\#K - \ell N - m - 1}{N} \pi} \int_0^{\infty} \xi^{N - \#K + \ell N + m} e^{-(t-s)\xi^N - \theta_k e^{-i \frac{\pi}{N}} x \xi} d\xi \right) \\
&= -\frac{Ni}{2\pi} \sum_{m=0}^{\#K} \alpha_{-m} s^{\frac{m-\#K}{N}} \sum_{k \in K} B_k \theta_k^{m+1} \\
&\quad \times \left( e^{-i \frac{\#K - m - 1}{N} \pi} \int_0^{\infty} \left( \sum_{\ell=0}^{\infty} \frac{(-s\xi^N)^{\ell}}{\Gamma(\ell + \frac{m+\#J}{N})} \right) \xi^{m+\#J} e^{-(t-s)\xi^N - \theta_k e^{i \frac{\pi}{N}} x \xi} d\xi \right. \\
&\quad \left. - e^{i \frac{\#K - m - 1}{N} \pi} \int_0^{\infty} \left( \sum_{\ell=0}^{\infty} \frac{(-s\xi^N)^{\ell}}{\Gamma(\ell + \frac{m+\#J}{N})} \right) \xi^{m+\#J} e^{-(t-s)\xi^N - \theta_k e^{-i \frac{\pi}{N}} x \xi} d\xi \right) \\
&= -\frac{Ni}{2\pi} \sum_{m=0}^{\#K} \alpha_{-m} s^{\frac{m-\#K}{N}} \sum_{k \in K} B_k \theta_k^{m+1} \\
&\quad \times \left( e^{-i \frac{\#K - m - 1}{N} \pi} \int_0^{\infty} \xi^{m+\#J} e^{-(t-s)\xi^N - \theta_k e^{i \frac{\pi}{N}} x \xi} E_{1, \frac{m+\#J}{N}}(-s\xi^N) d\xi \right. \\
&\quad \left. - e^{i \frac{\#K - m - 1}{N} \pi} \int_0^{\infty} \xi^{m+\#J} e^{-(t-s)\xi^N - \theta_k e^{-i \frac{\pi}{N}} x \xi} E_{1, \frac{m+\#J}{N}}(-s\xi^N) d\xi \right).
\end{aligned}$$

The proof of (6.1) is established.  $\blacksquare$

**Remark 6.1.** Let us integrate (6.1) with respect to  $x$  on  $(-\infty, 0]$ . We first compute, by using (2.8),

$$\begin{aligned}
\int_{-\infty}^0 \mathcal{K}_m(x\xi) dx &= -\frac{1}{\xi} \left( \sum_{k \in K} B_k \theta_k^m \right) \left( e^{-i \frac{\#K - m}{N} \pi} - e^{i \frac{\#K - m}{N} \pi} \right) \\
&= \frac{2i}{\xi} \sin\left(\frac{\#K - m}{N} \pi\right) \beta_m = \begin{cases} 0 & \text{if } 1 \leq m \leq \#K, \\ \frac{2i}{\xi} \sin\left(\frac{\#K}{N} \pi\right) & \text{if } m = 0. \end{cases}
\end{aligned}$$

We then obtain

$$\begin{aligned}
\mathbb{P}\{T(t) \in ds, X(t) \leq 0\}/ds &= \frac{N \sin\left(\frac{\#K}{N}\pi\right)}{\pi s^{\frac{\#K}{N}}} \int_0^\infty \xi^{\#J-1} e^{-(t-s)\xi^N} E_{1, \frac{\#J}{N}}(-s\xi^N) d\xi \\
&= \frac{N \sin\left(\frac{\#K}{N}\pi\right)}{\pi s^{\frac{\#K}{N}}} \sum_{\ell=0}^\infty \frac{(-s)^\ell}{\Gamma(\ell + \frac{\#J}{N})} \int_0^\infty \xi^{\ell N + \#J-1} e^{-(t-s)\xi^N} d\xi \\
&= \frac{\sin\left(\frac{\#K}{N}\pi\right)}{\pi s^{\frac{\#K}{N}}(t-s)^{\frac{\#J}{N}}} \sum_{\ell=0}^\infty \left(\frac{s}{t-s}\right)^\ell.
\end{aligned}$$

In the foregoing equality we must assume  $0 < s < t/2$  in order to make convergent the series. From this, we extract

$$\mathbb{P}\{T(t) \in ds, X(t) \leq 0\}/ds = \frac{\sin\left(\frac{\#K}{N}\pi\right)}{\pi t} \left(\frac{t-s}{s}\right)^{\frac{\#K}{N}}. \quad (6.4)$$

We retrieve Theorem 14 of [11].

**Remark 6.2.** Let us evaluate  $\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)$  at  $x = 0$ . For  $0 \leq m \leq \#K$ ,

$$\mathcal{K}_m(0) = e^{-i\frac{\#K-m-1}{N}\pi} \sum_{k \in K} B_k \theta_k^{m+1} - e^{i\frac{\#K-m-1}{N}\pi} \sum_{k \in K} B_k \theta_k^{m+1} = -2i \sin\left(\frac{\#K-m-1}{N}\pi\right) \beta_{m+1}.$$

Observing that  $\sin\left(\frac{\#K-m-1}{N}\pi\right) = 0$  if  $m = \#K - 1$ , in view of (2.8), (2.9) and (2.10), we get

$$\begin{aligned}
\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/ds \Big|_{x=0} &= \frac{N}{\pi} \sin\left(\frac{\pi}{N}\right) \alpha_{-\#K} \beta_{\#K+1} \int_0^\infty \xi^N e^{-(t-s)\xi^N} E_{1,1}(-s\xi^N) d\xi \\
&= \frac{N}{\pi} \sin\left(\frac{\pi}{N}\right) \left(\sum_{j \in J} \theta_j\right) \int_0^\infty \xi^N e^{-t\xi^N} d\xi = \frac{\sin\left(\frac{\pi}{N}\right) \Gamma\left(\frac{1}{N}\right)}{N\pi t^{1+\frac{1}{N}}} \sum_{j \in J} \theta_j.
\end{aligned}$$

Thanks to (2.4) and (2.11), we see that

$$\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/ds \Big|_{x=0} = \frac{1}{t} p(t; 0)$$

and we deduce

$$\mathbb{P}\{T(t) \in ds | X(t) = 0\}/ds = \frac{\mathbb{1}_{(0,t)}(s)}{t},$$

that is,  $(T(t)|X(t) = 0)$  has the uniform law on  $(0, t)$ . This is Theorem 13 of [11].

## 6.2 The case $x \geq 0$

The case  $x \geq 0$  can be related to the case  $x \leq 0$  by using the duality. Let us introduce the dual process  $(X_t^*)_{t \geq 0}$  of  $(X_t)_{t \geq 0}$  defined as  $X_t^* = -X_t$  for any  $t \geq 0$ . It is known that (see [11]):

- If  $N$  is even, the processes  $X$  and  $X^*$  are identical in distribution (because of the symmetry of the heat kernel  $p$ ):  $X^* \stackrel{d}{=} X$ ;
- If  $N$  is odd, we have the equalities in distribution  $(X^+)^* \stackrel{d}{=} X^-$  and  $(X^-)^* \stackrel{d}{=} X^+$  where  $X^+$  is the pseudo-process associated with  $\kappa_N = +1$  and  $X^-$  the one associated with  $\kappa_N = -1$ .

When  $N$  is even, we have  $\{-\theta_j, j \in J\} = \{\theta_k, k \in K\}$ . In this case, for any  $j \in J$ , there exists a unique  $k \in K$  such that  $\theta_j = -\theta_k$  and then

$$A_j = \prod_{i \in J \setminus \{j\}} \frac{\theta_i}{\theta_i - \theta_j} = \prod_{i \in K \setminus \{k\}} \frac{-\theta_i}{-\theta_i + \theta_k} = \prod_{i \in K \setminus \{k\}} \frac{\theta_i}{\theta_i - \theta_k} = B_k$$

and

$$\alpha_m = \sum_{j \in J} A_j \theta_j^m = \sum_{k \in K} B_k (-\theta_k)^m = (-1)^m \beta_m.$$

When  $N$  is odd, we distinguish the roots of  $\kappa_N$  in the cases  $\kappa_N = +1$  and  $\kappa_N = -1$ :

- For  $\kappa_N = +1$ , let  $\theta_i^+$ ,  $1 \leq i \leq N$ , denote the roots of 1 and set  $J^+ = \{i \in \{1, \dots, N\} : \Re(\theta_i^+) > 0\}$  and  $K^+ = \{i \in \{1, \dots, N\} : \Re(\theta_i^+) < 0\}$ ;
- For  $\kappa_N = -1$ , let  $\theta_i^-$ ,  $1 \leq i \leq N$ , denote the roots of  $-1$  and set  $J^- = \{i \in \{1, \dots, N\} : \Re(\theta_i^-) > 0\}$  and  $K^- = \{i \in \{1, \dots, N\} : \Re(\theta_i^-) < 0\}$ .

We have  $\{\theta_j^-, i \in J^-\} = \{-\theta_k^+, k \in K^+\}$  and  $\{\theta_k^-, k \in K^-\} = \{-\theta_j^+, j \in J^+\}$ . In this case, for any  $j \in J^-$ , there exists a unique  $k \in K^+$  such that  $\theta_j^- = -\theta_k^+$  and then

$$A_j^- = \prod_{i \in J^- \setminus \{j\}} \frac{\theta_i^-}{\theta_i^- - \theta_j^-} = \prod_{i \in K^+ \setminus \{k\}} \frac{-\theta_i^+}{-\theta_i^+ - \theta_k^+} = \prod_{i \in K^+ \setminus \{k\}} \frac{\theta_i^+}{\theta_i^+ - \theta_k^+} = B_k^+$$

and similarly  $A_j^+ = B_k^-$ . Moreover, we have

$$\alpha_m^- = \sum_{j \in J^-} A_j^-(\theta_j^-)^m = \sum_{k \in K^+} B_k^+(-\theta_k^+)^m = (-1)^m \sum_{k \in K^+} B_k^+(\theta_k^+)^m = (-1)^m \beta_m^+$$

and similarly  $\alpha_m^+ = (-1)^m \beta_m^-$ .

Now, concerning the connection between sojourn time and duality, we have the following fact. Set

$$\tilde{T}(t) = \int_0^t \mathbf{1}_{(0, +\infty)}(X(u)) du \quad \text{and} \quad T^*(t) = \int_0^t \mathbf{1}_{[0, +\infty)}(X^*(u)) du.$$

Since Spitzer's identity holds true interchanging the closed interval  $[0, +\infty)$  and the open interval  $(0, +\infty)$ , it is easy to see that  $T(t)$  and  $\tilde{T}(t)$  have the same distribution. On the other hand, we have

$$\tilde{T}(t) = \int_0^t \mathbf{1}_{(0, +\infty)}(X(u)) du = \int_0^t \mathbf{1}_{(-\infty, 0)}(X^*(u)) du = \int_0^t [1 - \mathbf{1}_{[0, +\infty)}(X^*(u))] du = t - T^*(t).$$

We then deduce that  $T(t)$  and  $t - T^*(t)$  have the same distribution. Consequently, we can state the lemma below.

**Lemma 6.1.** *The following identity holds:*

$$\mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) = \mathbb{P}\{T^*(t) \in d(t-s), X^*(t) \in d(-x)\} / (ds dx).$$

As a result, the following result ensues.

**Theorem 6.2.** *Assume  $N$  is even. The distribution of  $(T(t), X(t))$  is given, for  $x \geq 0$ , by*

$$\begin{aligned} & \mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \\ &= \frac{Ni}{2\pi} \sum_{m=0}^{\#J} \beta_{-m} (t-s)^{\frac{m-\#J}{N}} \int_0^\infty \xi^{m+\#K} e^{-s\xi^N} \mathcal{J}_m(x\xi) E_{1, \frac{m+\#K}{N}}(-(t-s)\xi^N) d\xi \end{aligned} \quad (6.5)$$

where

$$\mathcal{J}_m(z) = e^{-i\frac{\#J-m-1}{N}\pi} \sum_{j \in J} A_j \theta_j^{m+1} e^{-\theta_j e^{i\frac{\pi}{N}} z} - e^{i\frac{\#J-m-1}{N}\pi} \sum_{j \in J} A_j \theta_j^{m+1} e^{-\theta_j e^{-i\frac{\pi}{N}} z}.$$

**PROOF**

When  $N$  is even, we know that  $X^*$  is identical in distribution to  $X$  and  $(T^*(t), X^*(t))$  is then distributed like  $(T(t), X(t))$ . Thus, by (6.1) and Lemma 6.1, for  $x \geq 0$ ,

$$\begin{aligned} & \mathbb{P}\{T(t) \in ds, X(t) \in dx\} / (ds dx) \\ &= \mathbb{P}\{T(t) \in d(t-s), X(t) \in d(-x)\} / (ds dx) \\ &= -\frac{Ni}{2\pi} \sum_{m=0}^{\#K} \alpha_{-m} (t-s)^{\frac{m-\#K}{N}} \int_0^\infty \xi^{m+\#J} e^{-s\xi^N} \mathcal{K}_m(-x\xi) E_{1, \frac{m+\#J}{N}}(-(t-s)\xi^N) d\xi. \end{aligned}$$

The discussion preceding Lemma 6.1 shows that

$$\mathcal{K}_m(z) = e^{-i\frac{\#J-m-1}{N}\pi} \sum_{j \in J} A_j(-\theta_j)^{m+1} e^{\theta_j e^{i\frac{\pi}{N}} z} - e^{i\frac{\#J-m-1}{N}\pi} \sum_{j \in J} A_j(-\theta_j)^{m+1} e^{\theta_j e^{-i\frac{\pi}{N}} z}.$$

We see that  $\mathcal{K}_m(z) = (-1)^{m+1} \mathcal{J}_m(-z)$  where the function  $\mathcal{J}_m$  is written in Theorem 6.2. Finally, by replacing  $\alpha_{-m}$  by  $(-1)^m \beta_{-m}$  and  $\#J, \#K$  by  $\#K, \#J$  respectively (which actually coincide since  $N$  is even), (6.5) ensues. ■

If  $N$  is odd, although the results are not justified, similar formulas can be stated. We find it interesting to produce them here. We set  $T^\pm(t) = \int_0^t \mathbb{1}_{[0,+\infty)}(X^\pm(u)) du$ .

**Theorem 6.3.** *Suppose that  $N$  is odd. The distribution of  $(T^+(t), X^+(t))$  is given, for  $x \geq 0$ , by*

$$\begin{aligned} & \mathbb{P}\{T^+(t) \in ds, X^+(t) \in dx\} / (ds dx) \\ &= \frac{Ni}{2\pi} \sum_{m=0}^{\#J^+} \beta_{-m}^+(t-s)^{\frac{m-\#J^+}{N}} \int_0^\infty \xi^{m+\#K^+} e^{-s\xi^N} \mathcal{J}_m^+(x\xi) E_{1, \frac{m+\#K^+}{N}}(-(t-s)\xi^N) d\xi \end{aligned} \quad (6.6)$$

where

$$\mathcal{J}_m^+(z) = e^{-i\frac{\#J^+-m-1}{N}\pi} \sum_{j \in J^+} A_j^+(\theta_j^+)^{m+1} e^{-\theta_j^+ e^{i\frac{\pi}{N}} z} - e^{i\frac{\#J^+-m-1}{N}\pi} \sum_{j \in J^+} A_j^+(\theta_j^+)^{m+1} e^{-\theta_j^+ e^{-i\frac{\pi}{N}} z}.$$

PROOF

When  $N$  is odd, we know that  $(X^+)^* \stackrel{d}{=} X^-$  and then  $((T^+)^*(t), (X^+)^*(t)) \stackrel{d}{=} (T^-(t), X^-(t))$ . Thus, by (6.1) and Lemma 6.1, for  $x \geq 0$ ,

$$\begin{aligned} & \mathbb{P}\{T^+(t) \in ds, X^+(t) \in dx\} / (ds dx) \\ &= \mathbb{P}\{T^-(t) \in d(t-s), X^-(t) \in d(-x)\} / (ds dx) \\ &= -\frac{Ni}{2\pi} \sum_{m=0}^{\#K^-} \alpha_{-m}^-(t-s)^{\frac{m-\#K^-}{N}} \int_0^\infty \xi^{m+\#J^-} e^{-s\xi^N} \mathcal{K}_m^-(-x\xi) E_{1, \frac{m+\#J^-}{N}}(-(t-s)\xi^N) d\xi \end{aligned}$$

where

$$\mathcal{K}_m^-(z) = e^{-i\frac{\#K^--m-1}{N}\pi} \sum_{k \in K^-} B_k^-(\theta_k^-)^{m+1} e^{-\theta_k^- e^{i\frac{\pi}{N}} z} - e^{i\frac{\#K^--m-1}{N}\pi} \sum_{k \in K^-} B_k^-(\theta_k^-)^{m+1} e^{-\theta_k^- e^{-i\frac{\pi}{N}} z}.$$

As in the proof of Theorem 6.2, we can write  $\mathcal{K}_m^-(z) = (-1)^{m+1} \mathcal{J}_m^+(-z)$  where the function  $\mathcal{J}_m^+$  is defined in Theorem 6.3. Finally, by replacing  $\alpha_{-m}^-$  by  $(-1)^m \beta_m^+$  and  $\#J^-, \#K^-$  by  $\#K^+, \#J^+$  respectively, (6.6) ensues. ■

Formula (6.6) involves only quantities with associated ‘+’ signs. We have a similar formula for  $X^-$  by changing all ‘+’ into ‘-’. So, we can remove these signs in order to get a unified formula (this is (6.5)) which is valid for even  $N$  and, at least formally, for odd  $N$  without sign.

**Remark 6.3.** Let us integrate (6.5) with respect to  $x$  on  $[0, \infty)$ . We first calculate, recalling that  $\mathcal{J}_m(z) = (-1)^{m+1} \mathcal{K}_m(-z)$  and referring to Remark 6.1,

$$\int_0^\infty \mathcal{J}_m(x\xi) dx = (-1)^{m+1} \int_{-\infty}^0 \mathcal{K}_m(x\xi) dx = \begin{cases} 0 & \text{if } 1 \leq m \leq \#J, \\ -\frac{2i}{\xi} \sin\left(\frac{\#J}{N}\pi\right) & \text{if } m = 0. \end{cases}$$

Then,

$$\begin{aligned} \mathbb{P}\{T(t) \in ds, X(t) \geq 0\} / ds &= \frac{N \sin\left(\frac{\#J}{N}\pi\right)}{\pi(t-s)^{\frac{\#J}{N}}} \int_0^\infty \xi^{\#K-1} e^{-s\xi^N} E_{1, \frac{\#K}{N}}(-(t-s)\xi^N) d\xi \\ &= \frac{N \sin\left(\frac{\#J}{N}\pi\right)}{\pi(t-s)^{\frac{\#J}{N}}} \sum_{\ell=0}^\infty \frac{(-(t-s)^\ell}{\Gamma(\ell + \frac{\#K}{N})} \int_0^\infty \xi^{N\ell + \#K-1} e^{-s\xi^N} d\xi \end{aligned}$$

$$= \frac{\sin\left(\frac{\#J}{N}\pi\right)}{\pi s^{\frac{\#K}{N}}(t-s)^{\frac{\#J}{N}}} \sum_{\ell=0}^{\infty} \left(-\frac{t-s}{s}\right)^{\ell}.$$

In the foregoing equality we must assume  $t/2 < s < t$  in order to make convergent the series. From this, we extract

$$\mathbb{P}\{T(t) \in ds, X(t) \geq 0\}/ds = \frac{\sin\left(\frac{\#J}{N}\pi\right)}{\pi t} \left(\frac{s}{t-s}\right)^{\frac{\#J}{N}} \quad (6.7)$$

and we retrieve Theorem 14 of [11]. By adding (6.4) and (6.7), we obtain the counterpart to the famous Paul Lévy's arc-sine law stated in [11] (Corollary 9):

$$\mathbb{P}\{T(t) \in ds\}/ds = \frac{\sin\left(\frac{\#J}{N}\pi\right)}{\pi} \frac{\mathbf{1}_{(0,t)}(s)}{s^{\frac{\#K}{N}}(t-s)^{\frac{\#J}{N}}}.$$

### 6.3 Examples

In this part, we write out the distribution of the couple  $(T(t), X(t))$  in the cases  $N = 3$  and  $N = 4$ .

**Example 6.1.** Case  $N = 3$ . Let us recall that this case is not fully justified. Nevertheless, we find it interesting to produce the formal corresponding results.

- Suppose  $\kappa_3 = 1$ . Using  $E_{1,1}(-s\xi^3) = e^{-s\xi^3}$  and the values of Example 2.1, (6.1) writes, for  $x \leq 0$ ,

$$\begin{aligned} & \mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx) \\ &= \frac{\sqrt{3}}{2\pi} \left( s^{-2/3} \int_0^{\infty} \xi e^{-(t-s)\xi^3} \tilde{\mathcal{K}}_0(x\xi) E_{1, \frac{1}{3}}(-s\xi^3) d\xi \right. \\ & \quad \left. + s^{-1/3} \int_0^{\infty} \xi^2 e^{-(t-s)\xi^3} \tilde{\mathcal{K}}_1(x\xi) E_{1, \frac{2}{3}}(-s\xi^3) d\xi + \int_0^{\infty} \xi^3 e^{-t\xi^3} \tilde{\mathcal{K}}_2(x\xi) d\xi \right) \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathcal{K}}_0(z) &= -i\sqrt{3}\mathcal{K}_0(z) = e^z - e^{-z/2} \left( \cos \frac{\sqrt{3}z}{2} + \sqrt{3} \sin \frac{\sqrt{3}z}{2} \right), \\ \tilde{\mathcal{K}}_1(z) &= -i\sqrt{3}\mathcal{K}_1(z) = -e^z + e^{-z/2} \left( \cos \frac{\sqrt{3}z}{2} - \sqrt{3} \sin \frac{\sqrt{3}z}{2} \right), \\ \tilde{\mathcal{K}}_2(z) &= -i\sqrt{3}\mathcal{K}_2(z) = e^z + 2e^{-z/2} \cos \frac{\sqrt{3}z}{2}. \end{aligned}$$

For  $x \geq 0$ , (6.6) gives

$$\begin{aligned} & \mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx) \\ &= \frac{3}{2\pi} \left( \frac{1}{\sqrt[3]{t-s}} \int_0^{\infty} \xi^2 e^{-s\xi^3} \tilde{\mathcal{J}}_0(x\xi) E_{1, \frac{2}{3}}(-(t-s)\xi^3) d\xi + \int_0^{\infty} \xi^3 e^{-t\xi^3} \tilde{\mathcal{J}}_1(x\xi) d\xi \right) \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathcal{J}}_0(z) &= i\mathcal{J}_0(z) = 2e^{-z/2} \sin \frac{\sqrt{3}z}{2}, \\ \tilde{\mathcal{J}}_1(z) &= -i\mathcal{J}_1(z) = e^{-z/2} \left( \sqrt{3} \cos \frac{\sqrt{3}z}{2} - \sin \frac{\sqrt{3}z}{2} \right). \end{aligned}$$

- Suppose  $\kappa_3 = -1$ . Likewise, for  $x \leq 0$ ,

$$\begin{aligned} & \mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx) \\ &= \frac{3}{2\pi} \left( \frac{1}{\sqrt[3]{s}} \int_0^{\infty} \xi^2 e^{-(t-s)\xi^3} \tilde{\mathcal{K}}_0(x\xi) E_{1, \frac{2}{3}}(-s\xi^3) d\xi + \int_0^{\infty} \xi^3 e^{-t\xi^3} \tilde{\mathcal{K}}_1(x\xi) d\xi \right) \end{aligned}$$

where

$$\tilde{\mathcal{K}}_0(z) = -i\mathcal{K}_0(z) = -2e^{z/2} \sin \frac{\sqrt{3}z}{2},$$

$$\tilde{\mathcal{K}}_1(z) = -i \mathcal{K}_1(z) = e^{z/2} \left( \sqrt{3} \cos \frac{\sqrt{3}z}{2} + \sin \frac{\sqrt{3}z}{2} \right).$$

For  $x \geq 0$ ,

$\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)$

$$\begin{aligned} &= \frac{\sqrt{3}}{2\pi} \left( (t-s)^{-2/3} \int_0^\infty \xi e^{-s\xi^3} \tilde{\mathcal{J}}_0(x\xi) E_{1, \frac{1}{3}}(-(t-s)\xi^3) d\xi \right. \\ &\quad \left. + (t-s)^{-1/3} \int_0^\infty \xi^2 e^{-s\xi^3} \tilde{\mathcal{J}}_1(x\xi) E_{1, \frac{2}{3}}(-(t-s)\xi^3) d\xi + \int_0^\infty \xi^3 e^{-t\xi^3} \tilde{\mathcal{J}}_2(x\xi) d\xi \right) \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathcal{J}}_0(z) &= i\sqrt{3} \mathcal{J}_0(z) = e^{-z} - e^{z/2} \left( \cos \frac{\sqrt{3}z}{2} - \sqrt{3} \sin \frac{\sqrt{3}z}{2} \right), \\ \tilde{\mathcal{J}}_1(z) &= -i\sqrt{3} \mathcal{J}_1(z) = -e^{-z} + e^{z/2} \left( \cos \frac{\sqrt{3}z}{2} + \sqrt{3} \sin \frac{\sqrt{3}z}{2} \right), \\ \tilde{\mathcal{J}}_2(z) &= i\sqrt{3} \mathcal{J}_2(z) = e^{-z} + 2e^{z/2} \cos \frac{\sqrt{3}z}{2}. \end{aligned}$$

**Example 6.2.** Case  $N = 4$ . Referring to Example 2.2, formula (6.1) writes, for  $x \leq 0$ ,

$\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)$

$$\begin{aligned} &= \frac{2}{\pi} \left( \frac{1}{\sqrt{s}} \int_0^\infty \xi^2 e^{-(t-s)\xi^4} \tilde{\mathcal{K}}_0(x\xi) E_{1, \frac{1}{2}}(-s\xi^4) d\xi \right. \\ &\quad \left. + \frac{\sqrt{2}}{\sqrt[4]{s}} \int_0^\infty \xi^3 e^{-(t-s)\xi^4} \tilde{\mathcal{K}}_1(x\xi) E_{1, \frac{3}{4}}(-s\xi^4) d\xi + \int_0^\infty \xi^4 e^{-t\xi^4} \tilde{\mathcal{K}}_2(x\xi) d\xi \right) \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathcal{K}}_0(z) &= -i \mathcal{K}_0(z) = e^z - \cos z - \sin z, \\ \tilde{\mathcal{K}}_1(z) &= -i \mathcal{K}_1(z) = -e^z + \cos z - \sin z, \\ \tilde{\mathcal{K}}_2(z) &= -i \mathcal{K}_2(z) = e^z + \cos z + \sin z. \end{aligned}$$

For  $x \geq 0$ , (6.5) reads

$\mathbb{P}\{T(t) \in ds, X(t) \in dx\}/(ds dx)$

$$\begin{aligned} &= \frac{2}{\pi} \left( \frac{1}{\sqrt{t-s}} \int_0^\infty \xi^2 e^{-s\xi^4} \tilde{\mathcal{J}}_0(x\xi) E_{1, \frac{1}{2}}(-(t-s)\xi^4) d\xi \right. \\ &\quad \left. + \frac{\sqrt{2}}{\sqrt[4]{t-s}} \int_0^\infty \xi^3 e^{-s\xi^4} \tilde{\mathcal{J}}_1(x\xi) E_{1, \frac{3}{4}}(-(t-s)\xi^4) d\xi + \int_0^\infty \xi^4 e^{-t\xi^4} \tilde{\mathcal{J}}_2(x\xi) d\xi \right) \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathcal{J}}_0(z) &= i \mathcal{J}_0(z) = e^{-z} - \cos z + \sin z, \\ \tilde{\mathcal{J}}_1(z) &= -i \mathcal{J}_1(z) = -e^{-z} + \cos z + \sin z, \\ \tilde{\mathcal{J}}_2(z) &= i \mathcal{J}_2(z) = e^{-z} + \cos z - \sin z. \end{aligned}$$

## 7 Appendix

**Lemma 7.1** (Spitzer). *Let  $(\xi_k)_{k \geq 1}$  be a sequence of independent identically distributed random variables and set  $X_0 = 0$  and  $T_0 = 0$  and, for any  $k \geq 1$ ,*

$$X_k = \xi_1 + \cdots + \xi_k, \quad T_k = \sum_{j=1}^k \mathbb{1}_{[0, +\infty)}(X_j).$$

Then, for  $\mu \in \mathbb{R}$ ,  $\nu > 0$  and  $|z| < 1$ ,

$$\sum_{k=0}^{\infty} \mathbb{E} \left[ e^{i\mu X_k - \nu T_k} \right] z^k = \exp \left( \sum_{k=1}^{\infty} \mathbb{E} \left[ e^{i\mu X_k - \nu k \mathbf{1}_{[0,+\infty)}(X_k)} \right] \frac{z^k}{k} \right), \quad (7.1)$$

$$\sum_{k=0}^{\infty} \mathbb{E} \left[ e^{i\mu X_k - \nu T_k} \mathbf{1}_{[0,+\infty)}(X_k) \right] z^k = \frac{1}{e^\nu - 1} \left[ e^\nu - \exp \left( - \sum_{k=1}^{\infty} (1 - e^{-\nu k}) \mathbb{E} \left[ e^{i\mu X_k} \mathbf{1}_{[0,+\infty)}(X_k) \right] \frac{z^k}{k} \right) \right], \quad (7.2)$$

$$\sum_{k=0}^{\infty} \mathbb{E} \left[ e^{i\mu X_k - \nu T_k} \mathbf{1}_{(-\infty, 0)}(X_k) \right] z^k = \frac{e^\nu}{e^\nu - 1} \left[ \exp \left( \sum_{k=1}^{\infty} (1 - e^{-\nu k}) \mathbb{E} \left[ e^{i\mu X_k} \mathbf{1}_{(-\infty, 0)}(X_k) \right] \frac{z^k}{k} \right) - 1 \right]. \quad (7.3)$$

PROOF

Formula (7.1) is stated in [21] without proof. So, we produce a proof below which is rather similar to one lying in [21] related to the maximum functional of the  $X_k$ 's.

• **Step 1.** Set, for any  $(x_1, \dots, x_n) \in \mathbb{R}^n$  and  $\sigma \in \mathfrak{S}_n$  ( $\mathfrak{S}_n$  being the set of the permutations of  $1, 2, \dots, n$ ),

$$U(x_1, \dots, x_n) = \sum_{k=1}^n \mathbf{1}_{[0, \infty)} \left( \sum_{j=1}^k x_j \right)$$

and

$$V(\sigma; x_1, \dots, x_n) = \sum_{k=1}^{n_\sigma} \#c_k(\sigma) \mathbf{1}_{[0, \infty)} \left( \sum_{j \in c_k(\sigma)} x_j \right).$$

In the definition of  $V$  above, the permutation  $\sigma$  is decomposed into  $n_\sigma$  cycles:  $\sigma = (c_1(\sigma))(c_2(\sigma)) \dots (c_{n_\sigma}(\sigma))$ .

In view of Theorem 2.3 in [21], we have the equality between the two following sets:

$$\{U(\sigma(x_1), \dots, \sigma(x_n)), \sigma \in \mathfrak{S}_n\} = \{V(\sigma; x_1, \dots, x_n), \sigma \in \mathfrak{S}_n\}.$$

We then deduce, for any bounded Borel functions  $\phi$  and  $F$ ,

$$\mathbb{E}[\phi(X_n)F(U(\xi_1, \dots, \xi_n))] = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \mathbb{E} \left[ \phi \left( \sum_{j=1}^n \xi_{\sigma(j)} \right) F(V(\sigma; \xi_1, \dots, \xi_n)) \right].$$

In particular, for  $\phi(x) = e^{i\mu x}$  and  $F(x) = e^{-\nu x}$  (where  $\mu \in \mathbb{R}$  and  $\nu > 0$  are fixed),

$$\begin{aligned} \mathbb{E} \left[ e^{i\mu X_n - \nu \sum_{k=1}^n \mathbf{1}_{[0,+\infty)}(\sum_{j=1}^k \xi_j)} \right] &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \mathbb{E} \left[ \exp \left( i\mu \sum_{k=1}^{n_\sigma} \sum_{j \in c_k(\sigma)} \xi_j - \nu \sum_{k=1}^{n_\sigma} \#c_k(\sigma) \mathbf{1}_{[0, \infty)} \left( \sum_{j \in c_k(\sigma)} \xi_j \right) \right) \right] \\ &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \prod_{k=1}^{n_\sigma} \mathbb{E} \left[ \exp \left( i\mu \sum_{j \in c_k(\sigma)} \xi_j - \nu (\#c_k(\sigma)) \mathbf{1}_{[0, \infty)} \left( \sum_{j \in c_k(\sigma)} \xi_j \right) \right) \right] \\ &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \prod_{k=1}^{n_\sigma} \mathbb{E} \left[ \exp \left( i\mu \sum_{j=1}^{\#c_k(\sigma)} \xi_j - \nu (\#c_k(\sigma)) \mathbf{1}_{[0, \infty)} \left( \sum_{j=1}^{\#c_k(\sigma)} \xi_j \right) \right) \right]. \end{aligned}$$

Denote by  $r_\ell(\sigma)$  the number of cycles of length  $\ell$  in  $\sigma$  for any  $\ell \in \{1, \dots, n\}$ . We have  $r_1(\sigma) + 2r_2(\sigma) + \dots + nr_n(\sigma) = n$ . Then,

$$\begin{aligned} \mathbb{E} \left[ e^{i\mu X_n - \nu T_n} \right] &= \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \prod_{\ell=1}^n \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \right)^{r_\ell(\sigma)} \\ &= \frac{1}{n!} \sum_{\substack{k_1, \dots, k_n \geq 0: \\ k_1 + 2k_2 + \dots + nk_n = n}} N_{k_1, \dots, k_n} \prod_{\ell=1}^n \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \right)^{k_\ell} \end{aligned}$$

where  $N_{k_1, \dots, k_n}$  is the number of the permutations  $\sigma$  of  $n$  objects satisfying  $r_1(\sigma) = k_1, \dots, r_n(\sigma) = k_n$ ; this number is equal to

$$N_{k_1, \dots, k_n} = \frac{n!}{(k_1! 1^{k_1})(k_2! 2^{k_2}) \dots (k_n! n^{k_n})}.$$

Then,

$$\mathbb{E}[e^{i\mu X_n - \nu T_n}] = \sum_{\substack{k_1, \dots, k_n \geq 0: \\ k_1 + 2k_2 + \dots + nk_n = n}} \prod_{\ell=1}^n \frac{1}{k_\ell! \ell^{k_\ell}} \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \right)^{k_\ell}.$$

• **Step 2.** Therefore, the identity between the generating functions follows: for  $|z| < 1$ ,

$$\begin{aligned} \sum_{n=0}^{\infty} \mathbb{E}[e^{i\mu X_n - \nu T_n}] z^n &= \sum_{\substack{n \geq 0, k_1, \dots, k_n \geq 0: \\ k_1 + 2k_2 + \dots + nk_n = n}} \prod_{\ell=1}^n \frac{1}{k_\ell!} \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \frac{z^\ell}{\ell} \right)^{k_\ell} \\ &= \sum_{k_1, k_2, \dots \geq 0} \prod_{\ell=1}^{\infty} \frac{1}{k_\ell!} \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \frac{z^\ell}{\ell} \right)^{k_\ell} \\ &= \prod_{\ell=1}^{\infty} \left[ \sum_{k=1}^{\infty} \frac{1}{k!} \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \frac{z^\ell}{\ell} \right)^k \right] \\ &= \prod_{\ell=1}^{\infty} \exp \left( \mathbb{E} \left[ e^{i\mu X_\ell - \nu \ell \mathbf{1}_{[0, \infty)}(X_\ell)} \right] \frac{z^\ell}{\ell} \right) \\ &= \exp \left( \sum_{n=1}^{\infty} \mathbb{E} \left[ e^{i\mu X_n - \nu n \mathbf{1}_{[0, +\infty)}(X_n)} \right] \frac{z^n}{n} \right). \end{aligned}$$

The proof of (7.1) is finished.

• **Step 3.**

Using the elementary identity  $e^{a\mathbf{1}_A(x)} - 1 = (e^a - 1)\mathbf{1}_A(x)$  and noticing that  $T_k = T_{k-1} + \mathbf{1}_{[0, +\infty)}(X_k)$ , we get for any  $k \geq 1$ ,

$$\mathbb{E}[e^{i\mu X_k - \nu T_k} \mathbf{1}_{[0, +\infty)}(X_k)] = \mathbb{E} \left[ e^{i\mu X_k - \nu T_k} \frac{e^{\nu \mathbf{1}_{[0, +\infty)}(X_k)} - 1}{e^\nu - 1} \right] = \frac{1}{e^\nu - 1} \left[ \mathbb{E}(e^{i\mu X_k - \nu T_{k-1}}) - \mathbb{E}(e^{i\mu X_k - \nu T_k}) \right].$$

Now, since  $X_k = X_{k-1} + \xi_k$  where  $X_{k-1}$  and  $\xi_k$  are independent and  $\xi_k$  have the same distribution as  $\xi_1$ , we have, for  $k \geq 1$ ,

$$\mathbb{E}(e^{i\mu X_k - \nu T_{k-1}}) = \mathbb{E}(e^{i\mu \xi_1}) \mathbb{E}(e^{i\mu X_{k-1} - \nu T_{k-1}}).$$

Therefore,

$$\begin{aligned} \sum_{k=1}^{\infty} \mathbb{E}[e^{i\mu X_k - \nu T_k} \mathbf{1}_{[0, +\infty)}(X_k)] z^k &= \frac{1}{e^\nu - 1} \sum_{k=1}^{\infty} \left( \mathbb{E}[e^{i\mu X_k - \nu T_{k-1}}] - \mathbb{E}[e^{i\mu X_k - \nu T_k}] \right) z^k \\ &= \frac{1}{e^\nu - 1} \left( \mathbb{E}(e^{i\mu \xi_1}) \sum_{k=1}^{\infty} \mathbb{E}[e^{i\mu X_{k-1} - \nu T_{k-1}}] z^k - \sum_{k=1}^{\infty} \mathbb{E}[e^{i\mu X_k - \nu T_k}] z^k \right) \\ &= \frac{1}{e^\nu - 1} \left( (z \mathbb{E}(e^{i\mu \xi_1}) - 1) \sum_{k=0}^{\infty} \mathbb{E}[e^{i\mu X_k - \nu T_k}] z^k + 1 \right). \end{aligned} \quad (7.4)$$

By putting (7.1) into (7.4), we extract

$$\sum_{k=0}^{\infty} \mathbb{E}[e^{i\mu X_k - \nu T_k} \mathbf{1}_{[0, +\infty)}(X_k)] z^k = \frac{1}{e^\nu - 1} \left[ e^\nu - (1 - z \mathbb{E}(e^{i\mu \xi_1})) S(\mu, \nu, z) \right] \quad (7.5)$$

where we set

$$S(\mu, \nu, z) = \exp \left( \sum_{k=1}^{\infty} \mathbb{E} \left[ e^{i\mu X_k - \nu k \mathbf{1}_{[0, +\infty)}(X_k)} \right] \frac{z^k}{k} \right).$$

Next, using the elementary identity  $1 - \zeta = \exp[\log(1 - \zeta)] = \exp[-\sum_{k=1}^{\infty} \zeta^k/k]$  valid for  $|\zeta| < 1$ ,

$$1 - z \mathbb{E}(e^{i\mu\xi_1}) = \exp\left(-\sum_{k=1}^{\infty} [\mathbb{E}(e^{i\mu\xi_1})]^k \frac{z^k}{k}\right) = \exp\left(-\sum_{k=1}^{\infty} \mathbb{E}(e^{i\mu X_k}) \frac{z^k}{k}\right)$$

and then

$$\begin{aligned} (1 - z \mathbb{E}(e^{i\mu\xi_1})) S(\mu, \nu, z) &= \exp\left(\sum_{k=1}^{\infty} \mathbb{E}\left[e^{i\mu X_k - \nu k \mathbb{1}_{[0, +\infty)}(X_k)} - e^{i\mu X_k}\right] \frac{z^k}{k}\right) \\ &= \exp\left(-\sum_{k=1}^{\infty} (1 - e^{-\nu k}) \mathbb{E}[e^{i\mu X_k} \mathbb{1}_{[0, +\infty)}(X_k)] \frac{z^k}{k}\right). \end{aligned} \quad (7.6)$$

Hence, by putting (7.6) into (7.5), formula (7.2) entails.

By subtracting (7.5) from (7.1), we obtain the intermediate representation

$$\sum_{k=0}^{\infty} \mathbb{E}[e^{i\mu X_k - \nu T_k} \mathbb{1}_{(-\infty, 0)}(X_k)] z^k = \frac{1}{e^\nu - 1} [(e^\nu - z \mathbb{E}(e^{i\mu\xi_1})) S(\mu, \nu, z) - e^\nu].$$

By writing, as previously,

$$e^\nu - z \mathbb{E}(e^{i\mu\xi_1}) = e^\nu \exp\left(-\sum_{k=1}^{\infty} \mathbb{E}(e^{i\mu X_k}) \frac{e^{-\nu k} z^k}{k}\right),$$

we find

$$\begin{aligned} (e^\nu - z \mathbb{E}(e^{i\mu\xi_1})) S(\mu, \nu, z) &= e^\nu \exp\left(\sum_{k=1}^{\infty} \mathbb{E}\left[e^{i\mu X_k - \nu k \mathbb{1}_{[0, +\infty)}(X_k)} - e^{i\mu X_k - \nu k}\right] \frac{z^k}{k}\right) \\ &= e^\nu \exp\left(\sum_{k=1}^{\infty} (1 - e^{-\nu k}) \mathbb{E}[e^{i\mu X_k} \mathbb{1}_{(-\infty, 0)}(X_k)] \frac{z^k}{k}\right). \end{aligned}$$

Finally, (7.3) ensues. ■

**Lemma 7.2.** *The following identities hold:*

$$\beta_{\#K} = (-1)^{\#K-1} \prod_{k \in K} \theta_k, \quad \beta_{\#K+1} = (-1)^{\#K-1} \left(\prod_{k \in K} \theta_k\right) \left(\sum_{k \in K} \theta_k\right).$$

PROOF

We label the set  $K$  as  $\{1, 2, 3, \dots, \#K\}$ . By (2.5), we know that the  $B_k$ 's solve a Vandermonde system. Then, by Cramer's formulas, we can write them as fractions of some determinants:  $B_k = V_k/V$  where

$$V = \begin{vmatrix} 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots \\ \theta_1^{\#K-1} & \dots & \theta_{\#K}^{\#K-1} \end{vmatrix} \quad \text{and} \quad V_k = \begin{vmatrix} 1 & \dots & 1 & 1 & 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{k-1} & 0 & \theta_{k+1} & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{k-1}^2 & 0 & \theta_{k+1}^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \theta_1^{\#K-1} & \dots & \theta_{k-1}^{\#K-1} & 0 & \theta_{k+1}^{\#K-1} & \dots & \theta_{\#K}^{\#K-1} \end{vmatrix}.$$

By expanding the determinant  $V_k$  with respect to its  $k^{\text{th}}$  column and next factorizing it suitably, we easily see that

$$V_k = (-1)^{k+1} \begin{vmatrix} \theta_1 & \dots & \theta_{k-1} & \theta_{k+1} & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{k-1}^2 & \theta_{k+1}^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots & \vdots & & \vdots \\ \theta_1^{\#K-1} & \dots & \theta_{k-1}^{\#K-1} & \theta_{k+1}^{\#K-1} & \dots & \theta_{\#K}^{\#K-1} \end{vmatrix}$$

$$= (-1)^{k+1} \frac{\prod_{i \in K} \theta_i}{\theta_k} \begin{vmatrix} 1 & \dots & 1 & 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{k-1} & \theta_{k+1} & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{k-1}^2 & \theta_{k+1}^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots & \vdots & & \vdots \\ \theta_1^{\#K-2} & \dots & \theta_{k-1}^{\#K-2} & \theta_{k+1}^{\#K-2} & \dots & \theta_{\#K}^{\#K-2} \end{vmatrix}.$$

With this at hands, we have

$$\beta_{\#K} = \sum_{k \in K} B_k \theta_k^{\#K} = \frac{\prod_{k \in K} \theta_k}{V} \sum_{k \in K} (-1)^{k+1} \theta_k^{\#K-1} \begin{vmatrix} 1 & \dots & 1 & 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{k-1} & \theta_{k+1} & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{k-1}^2 & \theta_{k+1}^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots & \vdots & & \vdots \\ \theta_1^{\#K-2} & \dots & \theta_{k-1}^{\#K-2} & \theta_{k+1}^{\#K-2} & \dots & \theta_{\#K}^{\#K-2} \end{vmatrix}.$$

We can observe that the sum lying on the above right-hand side is nothing but the expansion of the determinant  $V$  with respect to its last row multiplied by the sign  $(-1)^{\#K-1}$ . This immediately ensues that  $\beta_{\#K} = (-1)^{\#K-1} \prod_{k \in K} \theta_k$ . Similarly,

$$\beta_{\#K+1} = \sum_{k \in K} B_k \theta_k^{\#K+1} = \frac{\prod_{k \in K} \theta_k}{V} \sum_{k \in K} (-1)^{k+1} \theta_k^{\#K} \begin{vmatrix} 1 & \dots & 1 & 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{k-1} & \theta_{k+1} & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{k-1}^2 & \theta_{k+1}^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots & \vdots & & \vdots \\ \theta_1^{\#K-2} & \dots & \theta_{k-1}^{\#K-2} & \theta_{k+1}^{\#K-2} & \dots & \theta_{\#K}^{\#K-2} \end{vmatrix}.$$

The above sum is the expansion with respect to its last row, multiplied by the sign  $(-1)^{\#K-1}$ , of the determinant  $V'$  defined as

$$V' = \begin{vmatrix} 1 & \dots & 1 \\ \theta_1 & \dots & \theta_{\#K} \\ \theta_1^2 & \dots & \theta_{\#K}^2 \\ \vdots & & \vdots \\ \theta_1^{\#K-2} & \dots & \theta_{\#K}^{\#K-2} \\ \theta_1^{\#K} & \dots & \theta_{\#K}^{\#K} \end{vmatrix}.$$

Let  $R_0, R_1, R_2, \dots, R_{\#K-2}, R_{\#K-1}$  denote the rows of  $V'$ . We perform the substitution  $R_{\#K-1} \leftarrow R_{\#K-1} + \sum_{\ell=2}^{\#K} (-1)^\ell \sigma_\ell R_{\#K-\ell}$  where the  $\sigma_\ell$ 's are defined by (2.3). This substitution does not affect the value of  $V'$  and it transforms, e.g., the first term of the last row into

$$\theta_1^{\#K} + \sum_{\ell=2}^{\#K} (-1)^\ell \sigma_\ell \theta_1^{\#K-\ell}.$$

Recall that  $\sigma_\ell = \sum_{1 \leq k_1 < \dots < k_\ell \leq \#K} \theta_{k_1} \dots \theta_{k_\ell}$ . We decompose  $\sigma_\ell$ , by isolating the terms involving  $\theta_1$ , into

$$\theta_1 \sum_{2 \leq k_2 < \dots < k_\ell \leq \#K} \theta_{k_2} \dots \theta_{k_\ell} + \sum_{2 \leq k_1 < k_2 < \dots < k_\ell \leq \#K} \theta_{k_1} \theta_{k_2} \dots \theta_{k_\ell} = \theta_1 \sigma'_{\ell-1} + \sigma'_\ell$$

where we set  $\sigma'_{\#K} = 0$  and  $\sigma'_\ell = \sum_{2 \leq k_1 < k_2 < \dots < k_\ell \leq \#K} \theta_{k_1} \theta_{k_2} \dots \theta_{k_\ell}$ . Therefore, we have

$$\begin{aligned} \theta_1^{\#K} + \sum_{\ell=2}^{\#K} (-1)^\ell \sigma_\ell \theta_1^{\#K-\ell} &= \theta_1^{\#K} + \sum_{\ell=2}^{\#K} (-1)^\ell \sigma'_{\ell-1} \theta_1^{\#K-\ell+1} + \sum_{\ell=2}^{\#K} (-1)^\ell \sigma'_\ell \theta_1^{\#K-\ell} \\ &= \theta_1^{\#K} + \sigma'_1 \theta_1^{\#K-1} = \theta_1^{\#K-1} (\theta_1 + \sigma'_1) = \theta_1^{\#K-1} \left( \sum_{k \in K} \theta_k \right). \end{aligned}$$

The foregoing manipulation works similarly for each term of the last row of  $V'$ . So, we deduce that  $V' = (\sum_{k \in K} \theta_k) V$  and finally  $\beta_{\#K+1} = (-1)^{\#K-1} (\prod_{k \in K} \theta_k) (\sum_{k \in K} \theta_k)$ . ■

**Lemma 7.3.** For any integer  $m \leq N - 1$  and any  $x \geq 0$ ,

$$\int_0^\infty e^{-\lambda u} I_{j,m}(u; x) du = \lambda^{-\frac{m}{N}} e^{-\theta_j \sqrt[N]{\lambda} x}. \quad (2.15)$$

PROOF

This formula is proved in [13] for  $0 \leq m \leq N - 1$ . To prove that it holds true also for negative  $m$ , we directly compute the Laplace transform of  $I_{j,m}(u; x)$ . We have

$$\int_0^\infty e^{-\lambda u} I_{j,m}(u; x) du = \frac{Ni}{2\pi} \left( e^{-i\frac{m}{N}\pi} \int_0^\infty \frac{\xi^{N-m-1}}{\xi^N + \lambda} e^{-\theta_j e^{i\frac{\pi}{N}} x \xi} d\xi - e^{i\frac{m}{N}\pi} \int_0^\infty \frac{\xi^{N-m-1}}{\xi^N + \lambda} e^{-\theta_j e^{-i\frac{\pi}{N}} x \xi} d\xi \right).$$

Let us integrate the function  $H : z \rightarrow \frac{z^{M-1}}{z^N + \lambda} e^{-az}$  for fixed  $a$  and  $M$  such that  $\Re(a) > 0$  and  $M > 0$  on the contour  $\Gamma_R = \{\rho e^{i\varphi} \in \mathbb{C} : \varphi = 0, \rho \in [0, R]\} \cup \{\rho e^{i\varphi} \in \mathbb{C} : \varphi \in (0, -\frac{2\pi}{N}), \rho = R\} \cup \{\rho e^{i\varphi} \in \mathbb{C} : \varphi = -\frac{2\pi}{N}, \rho \in (0, R]\}$ . We get, by residues theorem,

$$\begin{aligned} - \int_0^\infty \frac{z^{M-1}}{z^N + \lambda} e^{-az} dz + e^{-2i\frac{M}{N}\pi} \int_0^\infty \frac{z^{M-1}}{z^N + \lambda} e^{-ae^{-i\frac{2\pi}{N}} z} dz &= 2i\pi \operatorname{Residue}\left(H, \sqrt[N]{\lambda} e^{-i\frac{\pi}{N}}\right) \\ &= \frac{2i\pi}{N} \left(\sqrt[N]{\lambda}\right)^{M-N} e^{-i\frac{M-N}{N}\pi} e^{-a\sqrt[N]{\lambda} e^{-i\frac{\pi}{N}}} \\ &= \frac{2\pi}{Ni} \lambda^{\frac{M}{N}-1} e^{-i\frac{M}{N}\pi} e^{-a\sqrt[N]{\lambda} e^{-i\frac{\pi}{N}}}. \end{aligned}$$

For  $M = N - m$  and  $a = \theta_j e^{i\frac{\pi}{N}} x$ , this yields

$$\int_0^\infty e^{-\lambda u} I_{j,m}(u; x) du = -e^{-i\frac{m}{N}\pi} \lambda^{-\frac{m}{N}} \times (-e^{i\frac{m}{N}\pi}) e^{-\theta_j \sqrt[N]{\lambda} x} = \lambda^{-\frac{m}{N}} e^{-\theta_j \sqrt[N]{\lambda} x}.$$

Hence, (2.15) is valid for  $m \leq N - 1$ . ■

## References

- [1] Abramowitz, M. and Stegun, I.-A.: Handbook of mathematical functions with formulas, graphs, and mathematical tables. Dover Publications, 1992.
- [2] Beghin, L., Hochberg, K.-J. and Orsingher, E.: Conditional maximal distributions of processes related to higher-order heat-type equations. Stochastic Process. Appl. 85 (2000), no. 2, 209–223.
- [3] Beghin, L. and Orsingher, E.: The distribution of the local time for “pseudoprocesses” and its connection with fractional diffusion equations. Stochastic Process. Appl. 115 (2005), 1017–1040.
- [4] Beghin, L., Orsingher, O. and Ragozina, T.: Joint distributions of the maximum and the process for higher-order diffusions. Stochastic Process. Appl. 94 (2001), no. 1, 71–93.
- [5] Bohr, H.: Über die gleichmässige Konvergenz Dirichletscher Reihen. J. für Math. 143 (1913), 203–211.
- [6] Cammarota, V. and Lachal, A.: arXiv:1001.4201 [math.PR].
- [7] Erdélyi, A., Ed.: Higher transcendental functions, vol. III, Robert Krieger Publishing Company, Malabar, Florida, 1981.
- [8] Hochberg, K.-J.: A signed measure on path space related to Wiener measure. Ann. Probab. 6 (1978), no. 3, 433–458.
- [9] Hochberg, K.-J. and Orsingher, E.: The arc-sine law and its analogs for processes governed by signed and complex measures. Stochastic Process. Appl. 52 (1994), no. 2, 273–292.
- [10] Krylov, V. Yu.: Some properties of the distribution corresponding to the equation  $\frac{\partial u}{\partial t} = (-1)^{q+1} \frac{\partial^{2q} u}{\partial^{2q} x}$ . Soviet Math. Dokl. 1 (1960), 760–763.

- [11] Lachal, A.: Distribution of sojourn time, maximum and minimum for pseudo-processes governed by higher-order heat-type equations. *Electron. J. Probab.* 8 (2003), paper no. 20, 1–53.
- [12] Lachal, A.: Joint law of the process and its maximum, first hitting time and place of a half-line for the pseudo-process driven by the equation  $\frac{\partial}{\partial t} = \pm \frac{\partial^N}{\partial x^N}$ . *C. R. Acad. Sci. Paris, Sér. I* 343 (2006), no. 8, 525–530.
- [13] Lachal, A.: First hitting time and place, monopoles and multipoles for pseudo-processes driven by the equation  $\frac{\partial}{\partial t} = \pm \frac{\partial^N}{\partial x^N}$ . *Electron. J. Probab.* 12 (2007), 300–353.
- [14] Lachal, A.: First hitting time and place for the pseudo-process driven by the equation  $\frac{\partial}{\partial t} = \pm \frac{\partial^n}{\partial x^n}$  subject to a linear drift. *Stoch. Proc. Appl.* 118 (2008), 1–27.
- [15] Nakajima, T. and Sato, S.: On the joint distribution of the first hitting time and the first hitting place to the space-time wedge domain of a biharmonic pseudo process. *Tokyo J. Math.* 22 (1999), no. 2, 399–413.
- [16] Nikitin, Ya. Yu. and Orsingher, E.: On sojourn distributions of processes related to some higher-order heat-type equations. *J. Theoret. Probab.* 13 (2000), no. 4, 997–1012.
- [17] Nishioka, K.: Monopole and dipole of a biharmonic pseudo process. *Proc. Japan Acad. Ser. A* 72 (1996), 47–50.
- [18] Nishioka, K.: The first hitting time and place of a half-line by a biharmonic pseudo process. *Japan J. Math.* 23 (1997), 235–280.
- [19] Nishioka, K.: Boundary conditions for one-dimensional biharmonic pseudo process. *Electronic Journal of Probability* 6 (2001), paper no. 13, 1–27.
- [20] Orsingher, E.: Processes governed by signed measures connected with third-order “heat-type” equations. *Lithuanian Math. J.* 31 (1991), no. 2, 220–231.
- [21] Spitzer, F.: A combinatorial lemma and its application to probability theory. *Trans. Amer. Math. Soc.* 82 (1956), 323–339.