

$K \rightarrow \pi$ matrix elements of the chromomagnetic operator on the lattice

M. Constantinou,¹ M. Costa,² R. Frezzotti,³ V. Lubicz,^{4,5} G. Martinelli,⁶ D. Meloni,^{4,5}
H. Panagopoulos,² and S. Simula⁵

(ETM Collaboration)

¹*Department of Physics, Temple University, Philadelphia, Pennsylvania 19122-1801, USA*

²*Department of Physics, University of Cyprus, Nicosia CY-1678, Cyprus*

³*Dipartimento di Fisica, Università di Roma “Tor Vergata” and INFN, Sezione di “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy*

⁴*Dipartimento di Matematica e Fisica, Università Roma Tre, I-00146 Rome, Italy*

⁵*INFN, Sezione di Roma Tre, Via della Vasca Navale 84, I-00146 Rome, Italy*

⁶*Dipartimento di Fisica, Università di Roma “La Sapienza” and INFN,
Sezione di Roma, P.le Aldo Moro 2, I-00185 Rome, Italy*



(Received 9 January 2018; published 2 April 2018)

We present the results of the first lattice QCD calculation of the $K \rightarrow \pi$ matrix elements of the chromomagnetic operator $O_{\text{CM}} = g\bar{s}\sigma_{\mu\nu}G_{\mu\nu}d$, which appears in the effective Hamiltonian describing $\Delta S = 1$ transitions in and beyond the standard model. Having dimension five, the chromomagnetic operator is characterized by a rich pattern of mixing with operators of equal and lower dimensionality. The multiplicative renormalization factor as well as the mixing coefficients with the operators of equal dimension have been computed at one loop in perturbation theory. The power divergent coefficients controlling the mixing with operators of lower dimension have been determined nonperturbatively, by imposing suitable subtraction conditions. The numerical simulations have been carried out using the gauge field configurations produced by the European Twisted Mass Collaboration with $N_f = 2 + 1 + 1$ dynamical quarks at three values of the lattice spacing. Our result for the B parameter of the chromomagnetic operator at the physical pion and kaon point is $B_{\text{CMO}}^{K\pi} = 0.273(69)$, while in the SU(3) chiral limit we obtain $B_{\text{CMO}} = 0.076(23)$. Our findings are significantly smaller than the model-dependent estimate $B_{\text{CMO}} \sim 1-4$, currently used in phenomenological analyses, and improve the uncertainty on this important phenomenological quantity.

DOI: [10.1103/PhysRevD.97.074501](https://doi.org/10.1103/PhysRevD.97.074501)

I. INTRODUCTION

At low energy with respect to the electroweak scale, the standard model (SM) and its possible new physics (NP) extensions are described by an effective Hamiltonian in which the contribution of operators of dimension $d = 4 + n$ are suppressed by n powers of the high-energy (i.e. the electroweak or NP) scale. In the flavor changing $\Delta S = 1$ sector, the effective Hamiltonian contains four operators of dimension $d = 5$ —two electromagnetic (EMO) and two chromomagnetic (CMO) operators. Their contribution to the physical amplitudes is, thus, suppressed by only

one power of the high-energy scale. The $\Delta S = 1$, $d = 5$ effective Hamiltonian has the form

$$H_{\text{eff}}^{\Delta S=1, d=5} = \sum_{i=\pm} (C_{\gamma}^i Q_{\gamma}^i + C_g^i Q_g^i) + \text{H.c.}, \quad (1)$$

where $Q_{\gamma, g}^+$ ($Q_{\gamma, g}^-$) are the parity-even (-odd) EMO and CMO, respectively, defined as

$$Q_{\gamma}^{\pm} = \frac{Q_d e}{16\pi^2} (\bar{s}_L \sigma^{\mu\nu} F_{\mu\nu} d_R \pm \bar{s}_R \sigma^{\mu\nu} F_{\mu\nu} d_L),$$

$$Q_g^{\pm} = \frac{g}{16\pi^2} (\bar{s}_L \sigma^{\mu\nu} G_{\mu\nu} d_R \pm \bar{s}_R \sigma^{\mu\nu} G_{\mu\nu} d_L), \quad (2)$$

with $q_{R,L} = \frac{1}{2}(1 \pm \gamma_5)q$ (for $q = s, d$).

In Fig. 1, we show two examples of Feynman diagrams generating, at low energy, the effective magnetic interactions in the SM and beyond. For illustration of the NP contribution, we have considered the case of SUSY models,

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

in which the $\Delta S = 1$ transition occurs through the exchange of virtual squarks and gluinos and is mediated by the strong interactions.

Note that at least one mass insertion is required in the diagrams, both in the SM and beyond, in order to induce the chirality flip described by the magnetic operators.

A quick inspection of the diagrams of Fig. 1 shows that the Wilson coefficients of the magnetic operators in the SM and NP model are proportional to

$$C_{\gamma,g}^{\text{SM}} \sim \frac{\alpha_W(M_W)}{M_W} \frac{m_s}{M_W}, \quad C_{\gamma,g}^{\text{NP}} \sim \frac{\alpha_s(M_{NP})}{M_{NP}} \delta_{LR}, \quad (3)$$

where M_{NP} represents the typical NP scale, e.g., the gluino mass in the SUSY case, and the factors m_s/M_W and δ_{LR} are generated in the diagrams by the mass insertion. In the SUSY case, for instance, δ_{LR} represents the off-diagonal matrix element of the squark mass matrix normalized to the average squark mass. The transition rate is controlled in the SM by the weak coupling $\alpha_W(M_W)$. This is not generally the case for NP models. In the SUSY transition shown in Fig. 1, for example, the process is mediated by the strong interactions. Therefore, the proportionality of $C_{\gamma,g}^{\text{NP}}$ in Eq. (3) to the strong coupling constant $\alpha_s(M_{NP})$, rather than to the weak coupling as in the SM, compensates in part for the stronger high-energy scale suppression ($M_{NP} > M_W$) in the NP model. Thus, the magnetic interactions receive potentially large contributions from physics beyond the SM.

It is also worth noting that the chirality flipping factor m_s/M_W , which appears in the Wilson coefficients of the magnetic operators in the SM, is of the same size as Λ_{QCD}/M_W , which represents the additional suppression factor of the coefficients of dimension-six operators in the effective Hamiltonian. For this reason, the role of the magnetic operators tends to be marginal in the SM, while it is potentially more relevant for the searches of NP.

The $K \rightarrow \pi$ matrix element of the EMO Q_γ^+ , which is relevant, for instance, for the CP violating part of the rare $K_L \rightarrow \pi^0 \ell^+ \ell^-$ decays [1], has been computed on the lattice both in the quenched [2] and unquenched $N_f = 2$ [3] case. Since the electromagnetic field strength tensor $F_{\mu\nu}$ factorizes out of the hadronic matrix element, the lattice

computation only involves the quark bilinear operator $\bar{s}\sigma^{\mu\nu}d$, and it is relatively straightforward.

Computing the hadronic matrix elements of the CMO Q_g^\pm is, instead, by far more challenging. The main difficulty is represented by the complicated renormalization pattern of the operator, which also involves power divergent mixing with operators of lower dimensionality (see Ref. [4] and Sec. III). The relevant matrix elements with an initial kaon involve one, two or three pions in the final states and are of great phenomenological interest for various processes: the long distance contribution to $K^0 - \bar{K}^0$ mixing [5], $\Delta I = 1/2$, $K \rightarrow \pi\pi$ transitions and ϵ'/ϵ [1], CP violation in $K \rightarrow 3\pi$ decays [6]. These matrix elements are parametrized in terms of suitably defined B parameters:

$$\langle \pi^+ | Q_g^+ | K^+ \rangle = \frac{11}{32\pi^2} \frac{M_K^2 (p_K \cdot p_\pi)}{m_s + m_d} B_{\text{CMO}}^{K\pi}, \quad (4)$$

$$\langle \pi^+ \pi^- | Q_g^- | K^0 \rangle = i \frac{11}{32\pi^2} \frac{M_K^2 M_\pi^2}{f_\pi (m_s + m_d)} B_{\text{CMO}}^{K2\pi}, \quad (5)$$

$$\langle \pi^+ \pi^+ \pi^- | Q_g^+ | K^+ \rangle = - \frac{11}{16\pi^2} \frac{M_K^2 M_\pi^2}{f_\pi^2 (m_s + m_d)} B_{\text{CMO}}^{K3\pi}. \quad (6)$$

At leading order (LO) in SU(3) chiral perturbation theory (ChPT), the CMO has a single representation in terms of pseudo-Goldstone boson fields [7],

$$Q_g^\pm = \frac{11}{256\pi^2} \frac{f_\pi^2 M_K^2}{m_s + m_d} B_{\text{CMO}} [U(D_\mu U^\dagger)(D^\mu U) \pm (D_\mu U^\dagger)(D^\mu U)U^\dagger]_{23}, \quad (7)$$

where the low-energy constant B_{CMO} is estimated to be of order 1 in the chiral quark model of Ref. [7]. Therefore, the three B parameters of Eqs. (4)–(6) are related by SU(3) chiral symmetry, which predicts at LO their equality: $B_{\text{CMO}}^{K\pi} = B_{\text{CMO}}^{K2\pi} = B_{\text{CMO}}^{K3\pi} = B_{\text{CMO}}$. Such an equality is expected to be broken at higher orders in ChPT.

In this work, we evaluate the B parameter appearing in Eq. (4) from the lattice QCD computation of the $\langle \pi | Q_g^+ | K \rangle$ matrix element. We perform numerical simulations by employing the gauge configurations generated by the European Twisted Mass Collaboration (ETMC) with $N_f = 2 + 1 + 1$ dynamical quarks, which include in the sea, besides two light mass-degenerate quarks, also the strange and charm quarks with masses close to their physical values [8–10]. The same gauge ensembles have been used in Ref. [4] to determine nonperturbatively the power divergent mixing coefficients controlling the mixing of the CMO with operators of lower dimension. As for the mixing coefficients with operators of the same dimensionality and for the multiplicative renormalization constant (RC), we adopt the predictions of perturbation theory at one loop obtained in Ref. [4].




FIG. 1. One-loop Feynman diagrams contributing at low energy to the effective magnetic interactions, in the SM (left) and beyond (right). In the latter case we have shown, for illustrative purposes, the case of SUSY models. The crosses denote a mass insertion.

TABLE I. Parameters of the gauge ensembles and the values of the simulated sea and valence quark bare masses (in lattice units) used in this work (see Ref. [10] for details). The values of the kaon mass, M_K , do not correspond to a simulated strange bare quark mass shown in the 8th column, but to the renormalized strange mass interpolated at the physical value, $m_s^{\text{phys}}(\overline{MS}, 2 \text{ GeV}) = 99.6(4.3) \text{ MeV}$, determined in Ref. [10].

Ensemble	β	V/a^4	$a\mu_{\text{sea}} = a\mu_{ud}$	$a\mu_{\sigma}$	$a\mu_{\delta}$	N_{cfg}	$a\mu_s$	$M_{\pi}(\text{MeV})$	$M_K(\text{MeV})$
A30.32	1.90	$32^3 \times 64$	0.0030	0.15	0.19	150	0.0145, 0.0185, 0.0225	275	577
A40.32			0.0040			100		315	588
A50.32			0.0050			150		350	595
A40.24	1.90	$24^3 \times 48$	0.0040	0.15	0.19	150	0.0145, 0.0185, 0.0225	324	594
A60.24			0.0060			150		388	610
A80.24			0.0080			150		438	624
A100.24			0.0100			150		497	650
B25.32	1.95	$32^3 \times 64$	0.0025	0.135	0.170	150	0.0141, 0.0180, 0.0219	259	553
B35.32			0.0035			150		300	562
B55.32			0.0055			150		377	587
B75.32			0.0075			80		437	608
B85.24	1.95	$24^3 \times 48$	0.0085	0.135	0.170	150	0.0141, 0.0180, 0.0219	463	617
D15.48	2.10	$48^3 \times 96$	0.0015	0.12	0.1385	90	0.0118, 0.0151, 0.0184	224	538
D20.48			0.0020			100		255	541
D30.48			0.0030			100		310	554

Preliminary results for the B parameter of the CMO have been presented in Ref. [11], and our final result at the physical pion and kaon point is

$$B_{\text{CMO}}^{K\pi} = 0.273(13)(68) = 0.273(69), \quad (8)$$

where the first error comes from the numerical lattice simulations, while the second error accounts for the perturbative uncertainty in the one-loop determination of the multiplicative renormalization constant. Our result (8) represents the first lattice QCD determination of a matrix element of the CMO. In the SU(3) chiral limit we get $B_{\text{CMO}} = 0.076(23)$. Our findings are significantly smaller than the model-dependent estimate $B_{\text{CMO}} \sim 1\text{--}4$ currently used in phenomenological analyses [1] and improve the uncertainty on this important phenomenological quantity.

The plan of the paper is as follows. In Sec. II, we describe the lattice setup and give the simulation details. In Sec. III, we recall the main results obtained in Ref. [4] on the determination of the power-divergent mixing coefficients needed for the renormalization of the CMO. The mixing subtraction is evaluated in Sec. IV, and it is shown that the renormalized CMO correlator can be determined with a remarkable level of precision. The lattice data for the matrix elements of the renormalized CMO are presented in Sec. V and analyzed in terms of both SU(2) and SU(3) ChPT. Finally, Sec. VI contains our conclusions.

II. SIMULATION DETAILS

Our lattice setup is based on the gauge configurations generated by ETMC with $N_f = 2 + 1 + 1$ dynamical quarks [8,9], adopted in Ref. [10] for the determination of the up, down, strange and charm quark masses, using the

experimental value of the pion decay constant f_{π} to set the lattice scale.¹

The gauge fields are simulated using the Iwasaki gluon action [12], while sea quarks are implemented with the Wilson Twisted Mass Action at maximal twist [13–15]. In order to avoid the mixing of strange and charm quarks induced by lattice artifacts in the unitary twisted mass formulation, we have adopted the mixed action setup described in Ref. [16], where the valence strange quarks are regularized as Osterwalder-Seiler (OS) fermions [17], while the valence up and down quarks have the same action as the sea. The use of different lattice regularisations for the valence and sea quarks of the second generation preserves unitarity in the continuum limit and brings no complications for the operator renormalization pattern in mass-independent schemes, while producing only a modification of discretization effects. Therefore, the uncertainty related to the use of a nonunitary action at finite lattice spacing is incorporated directly in the error due to discretization effects, which will be addressed in Sec. V. Moreover, since we work with both valence and sea quarks at maximal twist, physical observables are guaranteed to be automatically $\mathcal{O}(a)$ improved [15,16].

The details of the ETMC gauge ensembles with $N_f = 2 + 1 + 1$ dynamical quarks are collected in Table I, where the number of the gauge configurations analyzed (N_{cfg}) corresponds to a separation of 20 trajectories. The QCD simulations are carried out at three different values of the inverse bare lattice coupling β , to allow for a controlled

¹Compared to Ref. [10], the number of independent gauge configurations adopted for the ensemble D15.48 has been increased to 90 to improve the statistics.

extrapolation to the continuum limit, and at different lattice volumes. In this work, the up- and down-quark masses are always taken to be degenerate ($m_u = m_d = m_{ud}$) and equal in the sea and valence sectors ($m_{ud}^{\text{sea}} = m_{ud}^{\text{val}} = m_{ud}$). In the light sector, we have simulated quark masses $m_{ud} = \mu_{ud}/Z_P$ in the range $3m_{ud}^{\text{phys}} \lesssim m_{ud} \lesssim 12m_{ud}^{\text{phys}}$, where m_{ud}^{phys} is the physical light-quark mass and $Z_m \equiv Z_P^{-1}$ is the mass renormalization constant (at maximal twist) determined in Ref. [10]. In the strange sector, we have used three values of the valence strange quark mass $m_s = \mu_s/Z_P$ in the range $0.7m_s^{\text{phys}} \lesssim m_s \lesssim 1.2m_s^{\text{phys}}$, where m_s^{phys} is the physical strange quark mass obtained in Ref. [10]. The values of the lattice spacing are $a = \{0.0885(36), 0.0815(30), 0.0619(18)\}$ fm at $\beta = \{1.90, 1.95, 2.10\}$, respectively, the lattice volume goes from $\simeq 2$ to $\simeq 3$ fm and the pion masses, extrapolated to the continuum and infinite volume limits, range from $\simeq 210$ to $\simeq 450$ MeV (see Ref. [10] for further details).

Quark propagators are obtained using the multiple mass solver method [18,19], which allows us to invert the Dirac operator for several quark masses at a relatively low computational cost. The statistical accuracy of the meson correlators is significantly improved by using the ‘‘one-end’’ stochastic method [20], which includes spatial stochastic sources at a single time slice chosen randomly. Statistical errors are evaluated using the jackknife procedure.

III. RENORMALIZATION OF THE CHROMOMAGNETIC OPERATOR

In this section, we briefly review the main results obtained in Ref. [4] for the renormalization of the CMO, whose specific renormalization pattern depends on the details of the lattice regularization, i.e. on the choice of the lattice action. A detailed analysis of the implications of the discrete symmetries of the twisted-mass action was carried out in Ref. [4], showing that the renormalization of the CMO involves the mixing among 13 operators of equal or lower dimensionality,² including also non-gauge-invariant operators vanishing by the equation of motion.

In the case of on-shell matrix elements the mixing simplifies, and the renormalized parity-even CMO can be written as [4]

$$\hat{O}_{\text{CM}} = Z_{\text{CM}} \left[O_{\text{CM}} - \left(\frac{c_{13}}{a^2} + c_2(\mu_s^2 + \mu_d^2) + c_3\mu_s\mu_d \right) S - \frac{c_{12}}{a}(\mu_s + \mu_d)P - c_4 O_4 \right], \quad (9)$$

²The operator mixing pattern considered in Ref. [4] includes one operator (O_6), which is not independent from the other ones, and misses one five-dimensional operator [21], which mixes with the CMO only at two loops in perturbation theory. The results presented in this work are therefore not affected.

where $O_{\text{CM}} \equiv 16\pi^2 Q_g^+ = g\bar{s}\sigma_{\mu\nu}G_{\mu\nu}d$, $S = \bar{s}d$, $P = i\bar{s}\gamma_5 d$ and $O_4 = \square(\bar{s}d)$ are bare operators, valence quarks are taken with the same value of the Wilson r parameter, i.e. $r_s = r_d$ (see Ref. [4]), and μ_s (μ_d) denotes the bare strange (light) quark mass.

Note that the quadratically divergent mixing of the CMO with the scalar density S is common to any regularization, whereas the mixing with the pseudoscalar density P (softened by the proportionality to the quark masses) is peculiar of twisted mass fermions, and it is a consequence of the nonconservation of parity. Moreover, in Eq. (9) the power divergent mixing coefficients c_{12} and c_{13} are scheme and renormalization scale independent [22], while the multiplicative RC Z_{CM} and the coefficients c_i with $i = 2, 3, 4$ depend on both the scheme and the renormalization scale.

It is well known [23] that the determination of power divergent coefficients, controlling the mixing with operators of lower dimension, cannot rely on perturbation theory. The reason is that potential nonanalytic (in g^2) contributions to these coefficients, like those proportional to powers of $(1/a) \exp(-1/(\beta_0 g^2)) \sim \Lambda_{\text{QCD}}$, do not appear in the perturbative expansion. Therefore, while for the present study the multiplicative renormalization factor Z_{CM} and the coefficients c_i with $i = 2, 3, 4$ have been evaluated in perturbation theory at one-loop, the coefficients c_{12} and c_{13} in Eq. (9) have been determined in a nonperturbative way by imposing two suitable subtraction conditions [4]. The first one is that the matrix element of the CMO between external kaon and pion at rest must vanish in the SU(3) chiral limit, namely

$$\frac{1}{Z_{\text{CM}}} \lim_{m_s, m_d \rightarrow 0} \langle \pi | \hat{O}_{\text{CM}} | K \rangle = \lim_{m_s, m_d \rightarrow 0} \langle \pi | O_{\text{CM}} - \frac{c_{13}}{a^2} S | K \rangle = 0, \quad (10)$$

from which the coefficient c_{13} can be determined. The second requirement is the vanishing of the parity violating matrix elements of the CMO up to terms of $\mathcal{O}(a)$, specifically

$$\frac{1}{Z_{\text{CM}}} \langle 0 | \hat{O}_{\text{CM}} | K \rangle = \langle 0 | O_{\text{CM}} - \frac{c_{13}}{a^2} S - \frac{c_{12}}{a}(\mu_s + \mu_d)P | K \rangle = 0, \quad (11)$$

from which the coefficient c_{12} can be calculated once the coefficient c_{13} is determined from Eq. (10).

In Table II, we present the numerical results for the various mixing coefficients, obtained in Ref. [4] at the three values of the inverse coupling β given in Table I. The central values and the errors of the coefficient c_{13} shown in the last column correspond to the averages and the spread of the two nonperturbative determinations corresponding to the choice ‘‘LP’’ given in Table IV of Ref. [4]. For the

TABLE II. Values of the multiplicative renormalization factor Z_{CM} , in the $\overline{\text{MS}}$ scheme at the scale $\mu = 2$ GeV, and of the mixing coefficients c_i in Eq. (9), obtained in Ref. [4]. The results correspond to the three values of the inverse coupling β given in Table I using one-loop perturbation theory, except for c_{12} and c_{13} in the last two columns, which have been obtained nonperturbatively (see text). The perturbative results have been evaluated using the bare coupling $g_0^2 = 6/\beta$ for the power divergent coefficients c_{12} and c_{13} and the boosted coupling $g_P^2 = g_0^2/U_P$ for the other coefficients, where U_P is the average plaquette equal to $\{0.575, 0.585, 0.614\}$ at $\beta = \{1.90, 1.95, 2.10\}$.

β	Z_{CM}	c_2	c_3	c_4	c_{12}	c_{13}	c_{12}	c_{13}
One-loop perturbation theory					Nonperturbative			
1.90	1.781	0.150	0.0	0.0	0.0854	0.962	0.035 (20)	0.89713 (11)
1.95	1.752	0.100	0.0	0.0	0.0832	0.937	0.035 (20)	0.87629 (13)
2.10	1.677	-0.042	0.0	0.0	0.0772	0.870	0.035 (20)	0.81676 (8)

mixing coefficient c_{12} the uncertainty of the nonperturbative results has been found at the level of $\approx 60\%$ [4], and therefore the g^2 -dependence of the nonperturbative determination of c_{12} , shown in the penultimate column, can be safely neglected.

It can be seen that (i) the power divergent coefficient c_{13} has been determined nonperturbatively with a very high level of precision, (ii) the coefficients c_3 and c_4 vanish at one loop, (iii) the coefficient c_2 which starts at $\mathcal{O}(g^2)$ is rather small, and (iv) the multiplicative renormalization factor Z_{CM} receives at one loop a sizable correction ($\sim 70\%$).

In Table II, we also provide the one-loop results for the power divergent coefficients c_{12} and c_{13} . For the latter, the difference between the one-loop and the nonperturbative results is less than 10%. The bulk of the difference is compatible with being a correction of $\mathcal{O}(g^4)$. Thus, genuine nonperturbative contributions to c_{13} are likely to be small, even though a firmer conclusion in this sense would require the calculation of c_{13} at two loops at least.

As far as the coefficient c_{12} is concerned, its size is smaller by (at least) one order of magnitude with respect to c_{13} both perturbatively and nonperturbatively (see Table II). In addition, the corresponding operator is proportional to the first power of the quark masses, with $a(m_s + m_d) \sim 0.02$ in our simulations. For these reasons, the subtraction of the linear divergence in Eq. (9) has a numerically negligible impact on the determination of the CMO matrix elements (see next section).

IV. LATTICE QCD CORRELATORS

In order to evaluate the matrix elements of the renormalized CMO (9), using the values of the mixing coefficients given in Table II, we need to determine the matrix elements of three bare local operators: O_{CM} , $O_S \equiv S$ and $O_P \equiv P$. For the scalar and pseudoscalar densities we adopt the local versions $S = \bar{s}d$ and $P = i\bar{s}\gamma_5 d$, respectively, while in the lattice version of the chromomagnetic operator O_{CM} the gluon tensor $G_{\mu\nu}$ is replaced by its clover discretization $P_{\mu\nu}$, namely, [24]

$$O_{\text{CM}} = g_0 \bar{\psi}_s \sigma_{\mu\nu} P_{\mu\nu} \psi_d, \quad (12)$$

where

$$P_{\mu\nu}(x) \equiv \frac{1}{4a^2} \sum_{j=1}^4 \frac{1}{2ig_0} [P_j(x) - P_j^\dagger(x)], \quad (13)$$

and the sum is over the four plaquettes $P_j(x)$ in the $\mu - \nu$ plane stemming from x and taken in the counterclockwise sense.

The $K \rightarrow \pi$ matrix elements of the bare local operators O_i ($i = \{\text{CM}, S, P\}$) are extracted from the large (Euclidean) time distance behavior of a convenient combination of two- and three-point correlation functions, which for both initial and final mesons at rest are defined as

$$C^{\pi(K)}(t) = \frac{1}{L^3} \sum_{\vec{x}, \vec{z}} \langle P_{\pi(K)}(\vec{x}, t + t_z) P_{\pi(K)}^\dagger(\vec{z}, t_z) \rangle, \quad (14)$$

$$C_i^{K\pi(\pi K)}(t, t') = \frac{1}{L^6} \sum_{\vec{x}, \vec{y}, \vec{z}} \langle P_{\pi(K)}(\vec{x}, t' + t_z) \times O_i(\vec{y}, t + t_z) P_{K(\pi)}^\dagger(\vec{z}, t_z) \rangle, \quad (15)$$

where t' is the time distance between the source and the sink, t is the time distance between the insertion of the operator O_i and the source, while $P_K(x) = i\bar{s}(x)\gamma_5 u(x)$ and $P_\pi(x) = i\bar{d}(x)\gamma_5 u(x)$ are the local interpolating fields of the K and π mesons, respectively. The Wilson parameters r of the two valence quarks in both initial and final mesons are always chosen to have opposite values, i.e., $r_s = r_d = -r_u$, so that the squared meson masses differ from their continuum counterpart only by terms of order $\mathcal{O}(a^2 m_{\text{QCD}})$ [15].

The statistical accuracy of the correlators (14)–(15) is significantly improved by using the all-to-all quark propagators evaluated with the so-called “one-end” stochastic method [20], which includes spatial stochastic sources at a single time slice chosen randomly. Statistical errors are evaluated using the jackknife procedure.

At large time distances two- and three-point correlation functions behave as

$$C^{\pi(K)}(t) \xrightarrow[t \gg a]{} \frac{|Z_{\pi(K)}|^2}{2M_{\pi(K)}} [e^{-M_{\pi(K)}t} + e^{-M_{\pi(K)}(T-t)}], \quad (16)$$

$$C_i^{K\pi(\pi K)}(t, t') \xrightarrow[t \gg a, (t-t) \gg a]{} \frac{Z_{\pi(K)} Z_{K(\pi)}^*}{4M_{\pi} M_K} \langle K(\pi) | O_i | \pi(K) \rangle \times e^{-M_{K(\pi)}t} e^{-M_{\pi(K)}(t-t)}, \quad (17)$$

where $Z_{\pi(K)}$ is the matrix element $\langle 0 | P_{\pi(K)}(0) | \pi(K) \rangle$ and $M_{\pi(K)}$ is the mass of the $\pi(K)$ meson. Both quantities are determined adopting the fitting function (16) in the time interval $[t_{\min}, T/2]$, where t_{\min} is the time distance at which the ground-state starts to dominate the two-point correlator. Explicitly we choose $t_{\min}/a = \{10, 12, 18\}$ at $T/a = \{48, 64, 96\}$ (cf. Table I) for both pion and kaon mesons.

The matrix elements $\langle K | O_i | \pi \rangle$ can be extracted from the time behavior of the following ratios,

$$R_i(t, t') = s_i(t, t') \sqrt{4M_{\pi} M_K} \left| \frac{C_i^{K\pi}(t, t') C_i^{\pi K}(t, t')}{\tilde{C}^{\pi}(t') \tilde{C}^K(t')} \right|, \quad (18)$$

where $s_i(t, t')$ is the sign of correlator $C_i^{K\pi}(t, t')$ and the correlation function $\tilde{C}^{\pi(K)}(t)$ is given by

$$\tilde{C}^{\pi(K)}(t) \equiv \frac{1}{2} \left\{ C^{\pi(K)}(t) + \sqrt{[C^{\pi(K)}(t)]^2 - [C^{\pi(K)}(T/2)]^2} \right\}, \quad (19)$$

which at large time distances behave as

$$\tilde{C}^{\pi(K)}(t) \xrightarrow[t \gg a]{} \frac{Z_{\pi(K)}}{2M_{\pi(K)}} e^{-M_{\pi(K)}t}, \quad (20)$$

i.e. without the backward signal. At large time distances, one has

$$R_i(t, t') \xrightarrow[t \gg a, (t-t) \gg a]{} \langle K | O_i | \pi \rangle, \quad (21)$$

so that the bare matrix elements $\langle K | O_i | \pi \rangle$ can be calculated from the plateau of R_i independently of the matrix elements Z_{π} and Z_K of the interpolating fields. In order to minimize excited state effects, in what follows the source-sink separation is fixed to $t' = T/2$. Therefore, the region of time distances, where both the initial and final ground states dominate leading to the plateau (21), corresponds to $[t_{\min}, T/2 - t_{\min}]$. Such a time interval will be adopted to extract the CMO matrix elements.

The size of the various terms involved in the renormalization of the CMO (see Eq. (22) below) and expressed in lattice units, namely $a^3 R_{\text{CM}}(t, T/2)$, $c_{13} a R_S(t, T/2)$, $c_{12} a (\mu_s + \mu_d) a R_P(t, T/2)$ and $c_2 a^2 (\mu_s^2 + \mu_d^2) a R_S(t, T/2)$, can be inferred from Fig. 2, where the results refer to $\beta = 2.10$ and quark masses $a\mu_d = 0.0020$ and $a\mu_s = 0.0151$

(corresponding to $M_{\pi} \simeq 255$ MeV and $M_K \simeq 520$ MeV). The values adopted for the mixing coefficients c_{12} and c_{13} are the nonperturbative ones, while the value of the coupling entering c_2 is taken from boosted perturbation theory (see Table II). It can be seen that the ratio $a^3 R_{\text{CM}}(t, T/2)$ and the one corresponding to the leading power divergence, $c_{13} a R_S(t, T/2)$, are almost equal, and the subtraction is at the level of 99.7%. The other two terms, i.e., $c_{12} a (\mu_s + \mu_d) a R_P(t, T/2)$ and $c_2 a^2 (\mu_s^2 + \mu_d^2) a R_S(t, T/2)$, are smaller by almost 5 orders of magnitude. The operators $\mu_s \mu_d S$ and $O_4 = \square(\bar{s}d)$ do not




FIG. 2. Time dependence of the various terms contributing to Eq. (22) in lattice units. Upper figure: $a^3 R_{\text{CM}}(t, T/2)$ and $c_{13} a R_S(t, T/2)$. Middle figure: $c_{12} a (\mu_s + \mu_d) a R_P(t, T/2)$ and $c_2 a^2 (\mu_s^2 + \mu_d^2) a R_S(t, T/2)$. Bottom figure: the renormalized ratio $a^3 \hat{R}_{\text{CM}}(t, T/2)$. The results refer to $\beta = 2.10$ and quark masses $a\mu_d = a\mu_{ud} = 0.0020$ and $a\mu_s = 0.0151$, corresponding to $M_{\pi} \simeq 255$ and $M_K \simeq 520$ MeV (ensemble D20.48). The values adopted for the mixing coefficients c_{12} and c_{13} are the nonperturbative ones (see the last two columns of Table II), while the value of the coupling entering c_2 is taken from boosted perturbation theory (see the third column of Table II). In the middle figure, the error of the term $c_{12} a (\mu_s + \mu_d) a R_P(t, T/2)$ is dominated by the large uncertainty of the nonperturbative determination of c_{12} (see the penultimate column of Table II). For the ensemble D20.48 the time interval, where the plateau (21) is expected to occur, is $[t_{\min}, T/2 - t_{\min}]/a = [18, 30]$ (see text) and it is indicated by the vertical dotted lines.

contribute at one loop, since their mixing coefficients vanish (see Table II).

In the lower panel of Fig. 2 we show the renormalized CM ratio $\hat{R}_{\text{CM}}(t, T/2)$, given by

$$\hat{R}_{\text{CM}}(t, T/2) = Z_{\text{CM}} \left\{ R_{\text{CM}}(t, T/2) - \frac{c_{13}}{a^2} R_S(t, T/2) - \frac{c_{12}}{a} (\mu_s + \mu_d) R_P(t, T/2) - c_2 (\mu_s^2 + \mu_d^2) R_S(t, T/2) \right\}. \quad (22)$$

Despite being the outcome of a large numerical subtraction, due almost totally to the power divergent term $(c_{13}/a^2)R_S$, the results for $\hat{R}_{\text{CM}}(t, T/2)$ are clearly different from zero and exhibit plateaux, from which the renormalized CMO matrix element $\langle K | \hat{O}_{\text{CM}} | \pi \rangle$ can be determined quite precisely (see next section). In this respect, we stress that the high level of precision achieved in the nonperturbative determination of c_{13} (see the last column of Table II) plays a crucial role.

Notice that the renormalized CMO ratio $a^3 \hat{R}_{\text{CM}}(t, T/2)$ is 2 orders of magnitude larger than the mixing term $c_2 a^2 (\mu_s^2 + \mu_d^2) a R_S(t, T/2)$. Since c_2 [as well as c_3 and c_4 in Eq. (9)] is known only at one loop in perturbation theory, higher-order corrections and nonperturbative effects might contribute to the renormalized CMO ratio away from the SU(3) chiral limit. However, the smallness of the mixing term $c_2 a^2 (\mu_s^2 + \mu_d^2) a R_S(t, T/2)$ indicates that higher orders and nonperturbative effects are not expected to play a significant role in the determination of the matrix elements of the renormalized CMO within the present statistical uncertainties.

V. MATRIX ELEMENTS OF THE CHROMOMAGNETIC OPERATOR

By having defined the properly renormalized CMO, the matrix elements $\langle K | \hat{O}_{\text{CM}} | \pi \rangle$ can be extracted from the plateaux of the ratio $\hat{R}_{\text{CM}}(t, T/2)$ in the expected time interval $[t_{\text{min}}, T/2 - t_{\text{min}}]$, where t_{min} is the time distance at which excited states have decayed sufficiently from both the source and the sink, namely $t_{\text{min}} = \{10, 12, 18\}$ at $\beta = \{1.90, 1.95, 2.10\}$. As described in Ref. [15], the resulting matrix element $\langle K | \hat{O}_{\text{CM}} | \pi \rangle$ is automatically $\mathcal{O}(a)$ improved.

The B parameter $B_{\text{CMO}}^{K\pi}$, defined in Eq. (4), is easily obtained as

$$B_{\text{CMO}}^{K\pi} = \frac{32\pi^2 m_s + m_d}{11} \frac{\langle K | \hat{O}_{\text{CM}} | \pi \rangle}{M_K^2 M_\pi}. \quad (23)$$

Note that the evaluation of the B parameter (23) does not require the knowledge of the lattice spacing, while it

involves the mass RC Z_m , which in our maximally twisted-mass setup is given by $Z_m = 1/Z_P$, where Z_P is the RC of the pseudoscalar density. For the latter, we adopt the RI'-MOM results obtained in Ref. [10] using the two methods M1 and M2, which differ by $\mathcal{O}(a^2)$ effects.

Besides $B_{\text{CMO}}^{K\pi}$, we have calculated two further B parameters. They correspond to the transitions induced by the CMO in which either the mass m_s of the strange valence quark is taken to be equal to the light-quark mass m_d or the mass m_d of the light valence quark is taken to be equal to the strange quark mass m_s . In both cases, the spectator valence quark, which does not participate in the transition, is always a up-quark with mass m_{ud} [see Eqs. (14) and (15)]. We will refer to the above two transitions as the $\pi\pi$ and KK channels, respectively. Explicitly, one has

$$B_{\text{CMO}}^{\pi\pi} = \frac{32\pi^2 2m_d}{11} \frac{\langle \pi | \hat{O}_{\text{CM}} | \pi \rangle}{M_\pi^2}, \quad (24)$$

$$B_{\text{CMO}}^{KK} = \frac{32\pi^2 m_s + m_d}{11} \frac{\langle K | \hat{O}_{\text{CM}} | K \rangle}{M_K^2}, \quad (25)$$

where $\langle \pi | \hat{O}_{\text{CM}} | \pi \rangle \equiv [\langle K | \hat{O}_{\text{CM}} | \pi \rangle]_{m_s=m_d=m_{ud}}$ and $\langle K | \hat{O}_{\text{CM}} | K \rangle \equiv [\langle K | \hat{O}_{\text{CM}} | \pi \rangle]_{m_d=m_s}$. The $\pi\pi$ and KK channels do not correspond to any physical process, but the set of the three quantities $B_{\text{CMO}}^{\pi\pi}$, $B_{\text{CMO}}^{K\pi}$ and B_{CMO}^{KK} can be analyzed in terms of SU(3) ChPT³

In Fig. 3, we show the results for the B parameters $B_{\text{CMO}}^{\pi\pi}$, $B_{\text{CMO}}^{K\pi}$, and B_{CMO}^{KK} as a function of the renormalized light-quark mass m_{ud} for the ETMC ensembles of Table I in the $\overline{\text{MS}}(2 \text{ GeV})$ scheme. Since we have simulated three values of the strange quark mass around its physical value (see Table I), the results for $B_{\text{CMO}}^{K\pi}$ and B_{CMO}^{KK} , shown in Fig. 3, correspond to a smooth interpolation at $m_s = m_s^{\text{phys}} = 99.6(4.3) \text{ MeV}$, determined in Ref. [10]. It can be seen that, for all the three channels, the results for the B parameters exhibit controllable discretization effects, which are the beneficial consequence of the subtraction of the power-divergent mixings (thanks to the use of maximally twisted Wilson quarks the subtracted CMO matrix elements vanish in the chiral limit even at finite lattice spacing). Moreover, the impact of finite volume effects can be estimated by comparing the results corresponding to the ensembles A40.24 and A40.32, which share common values of the pion mass and the lattice spacing, and differ only by lattice size L . No significant effects are visible within the statistical errors.

As described in Sec. I, the ChPT prediction at LO is that all three B parameters should coincide and be independent

³In Eq. (25), the B parameter B_{CMO}^{KK} is defined in such a way as to guarantee that at LO in SU(3) ChPT the three B parameters $B_{\text{CMO}}^{\pi\pi}$, $B_{\text{CMO}}^{K\pi}$, and B_{CMO}^{KK} are always normalized by the same quark condensate in the chiral limit.




FIG. 3. Values of the B parameters $B_{\text{CMO}}^{\pi\pi}$ (upper), $B_{\text{CMO}}^{K\pi}$ (middle), and B_{CMO}^{KK} (bottom), defined in Eqs. (24), (23), and (25), respectively, as a function of the renormalized light-quark mass m_{ud} in the $\overline{\text{MS}}(2 \text{ GeV})$ scheme. The lattice data for $B_{\text{CMO}}^{K\pi}$ and B_{CMO}^{KK} have been smoothly interpolated to the physical value of the strange quark mass $m_s^{\text{phys}} = 99.6(4.3) \text{ MeV}$ [10]. Note the different scales in the three panels. The dashed, short-dashed and dotted lines correspond to the results of the SU(3)-inspired fit (27) for the three values of the lattice spacing, while the solid lines are the result in the continuum limit. The diamonds represent the values of the B parameters at the physical pion mass and in the continuum limit.

of the light-quark mass. The latter feature is approximately fulfilled only in the case of $\pi \rightarrow \pi$ channel (see upper panel in Fig. 3). Chiral corrections beyond the LO are clearly visible for both the $K\pi$ and KK channels (see middle and bottom panels in Fig. 3), since they exhibit a remarkable dependence on the light-quark mass and deviate strongly from the results corresponding to the $\pi\pi$ channel.

In order to extrapolate to the physical pion mass, we need to take into account at least NLO effects, which however are not known analytically. We observe, in this respect, that the chiral corrections for the CMO matrix element contain also powers of $p_K \cdot p_\pi$. This means that for the B parameter $B_{\text{CMO}}^{K\pi}$, NLO terms proportional not only to M_K^2 and M_π^2 but also to $M_K M_\pi$ should be considered. Note that the quantity $M_K M_\pi \propto m_{ud}^{1/2}$ is nonanalytic in the light-quark mass. In the $\pi\pi$ and KK channels, such a nonanalytic term is not expected.

Thus, we introduce the variables

$$\begin{aligned}\xi_\pi &\equiv \frac{2B_0 m_{ud}}{(4\pi f_0)^2}, \\ \xi_K &\equiv \frac{B_0(m_s + m_{ud})}{(4\pi f_0)^2},\end{aligned}\quad (26)$$

where B_0 and f_0 are LO low-energy constants (LECs) of the SU(3) ChPT [25], and we consider for the generic channel $i \rightarrow j$ with $i(j) = K, \pi$ the following SU(3)-inspired ansatz

$$\begin{aligned}B_{\text{CMO}}^{ij} &= B_{\text{CMO}} \left[1 + \alpha_1 \xi_\pi + \beta_1 (\xi_i + \xi_j) \right. \\ &\quad + \gamma_1 \left(\sqrt{\xi_i} - \sqrt{\xi_j} \right)^2 + \alpha_2 \xi_\pi^2 + \beta_2 (\xi_i + \xi_j)^2 \\ &\quad + \beta_2' (\xi_i - \xi_j)^2 + \gamma_2 \left(\sqrt{\xi_i} - \sqrt{\xi_j} \right)^4 \left. \right] \\ &\quad + a^2 [D_0 + D_1 (\xi_i + \xi_j)],\end{aligned}\quad (27)$$

where B_{CMO} is the LO LEC appearing in Eq. (7) (i.e., the SU(3) chiral limit of the B parameters), while the parameters α_1, β_1 , and γ_1 play the role of NLO LECs, and $\alpha_2, \beta_2, \beta_2'$, and γ_2 are NNLO LECs. In Eq. (27), the two terms proportional to ξ_π and ξ_π^2 are due to the dependence on the mass of the u quark (not involved in the transition) in common for all the channels. Note that the NNLO term proportional to $(\sqrt{\xi_i} - \sqrt{\xi_j})^4$ generates in the $K\pi$ channel nonanalytic terms proportional to $m_{ud}^{1/2}$ and $m_{ud}^{3/2}$.

According to SU(3) ChPT, the LECs are independent of the light and strange quark masses, and therefore the ansatz (27) can be applied to the combined analysis of the $\pi\pi$, $K\pi$, and KK channels. Taking into account that we have simulated three different values of the strange quark mass, the total number of lattice data is 105 (15 points for the $\pi\pi$ channel and 45 points for both $K\pi$ and KK channels) and the number of fitting parameters is 10, whose values have been determined using a χ^2 -minimization procedure based on an uncorrelated χ^2 . The results of the SU(3)-inspired fit (27) are shown in Fig. 3, where it can be seen that the quality of the fit, corresponding to $\chi^2/\text{d.o.f.} \simeq 0.5$, is remarkably good.⁴

After chiral and continuum extrapolations we get at the physical point: $B_{\text{CMO}}^{\pi\pi}|_{\text{phys}} = 0.078(12)$, $B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.279(10)$ and $B_{\text{CMO}}^{KK}|_{\text{phys}} = 0.105(11)$, where the errors are statistical only. The value of the LO LEC B_{CMO} , i.e. the SU(3) chiral limit of the B parameters, is close to the

⁴In Eq. (27) an additional NNLO term proportional to $(\xi_i + \xi_j)(\sqrt{\xi_i} - \sqrt{\xi_j})^2$ can be considered. We have checked that the impact of its inclusion is almost negligible. Moreover, the inclusion of additional discretization terms proportional either to $a^2 \xi_\pi$ or to $a^2 (\sqrt{\xi_i} - \sqrt{\xi_j})^2$ produces no significant effect within the errors.

physical result for the $\pi\pi$ channel, namely $B_{\text{CMO}} = 0.076(14)$. Our results imply that in the $K\pi$ channel the impact of chiral orders higher than the LO corresponds to a strong enhancement factor equal to ≈ 4 . Therefore, since higher orders in ChPT contribute differently in $K \rightarrow \pi$, $K \rightarrow \pi\pi$ and $K \rightarrow \pi\pi\pi$ transitions, the corresponding B parameters are no more simply related to each other beyond the LO.

The exclusion of all the NNLO terms in Eq. (27) (i.e., putting $\alpha_2 = \beta_2 = \beta'_2 = \gamma_2 = 0$) leads to a lower quality fit having $\chi^2/\text{d.o.f.} \simeq 1.5$. This is not surprising, since for a physical strange quark the impact of NNLO terms is expected to be non-negligible.

The result for $B_{\text{CMO}}^{K\pi}|_{\text{phys}}$ is sensitive to the presence of the structures proportional to $(\sqrt{\xi_i} - \sqrt{\xi_j})^2$ and $(\sqrt{\xi_i} - \sqrt{\xi_j})^4$. Putting $\gamma_2 = 0$ in Eq. (27) we obtain $B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.267(10)$ with $\chi^2/\text{d.o.f.} \simeq 0.6$. If all the nonanalytic terms are neglected (i.e., $\gamma_1 = \gamma_2 = 0$), the quality of the corresponding fit deteriorates significantly ($\chi^2/\text{d.o.f.} \simeq 2.8$).

Adopting the SU(3)-inspired fit (27) and averaging the results of the different fits using Eq. (28) of Ref. [10], our result for the B parameter $B_{\text{CMO}}^{K\pi}|_{\text{phys}}$ is

$$\begin{aligned} B_{\text{CMO}}^{K\pi}|_{\text{phys}} &= 0.272(10)_{\text{stat+fit}}(6)_{\text{chir}}(6)_{\text{disc}}(3)_{Z_p} \\ &= 0.272(13), \end{aligned} \quad (28)$$

where

- (i) $()_{\text{stat+fit}}$ indicates the uncertainty induced by both the statistical errors and the fitting procedure itself;
- (ii) $()_{\text{chir}}$ corresponds to the uncertainty related to the chiral extrapolation, obtained using the results corresponding to the inclusion ($\gamma_2 \neq 0$) or the exclusion ($\gamma_2 = 0$) of the NNLO nonanalytic term in Eq. (27);
- (iii) $()_{\text{disc}}$ is the uncertainty related to discretization effects estimated by adding a term proportional to a^4 without any prior;
- (iv) $()_{Z_p}$ is the error induced by the use of the two methods M1 and M2 to obtain the mass RC $Z_m = 1/Z_p$ in Ref. [10].

As a further check, we have also analyzed separately the data for the $K\pi$ channel (see also Ref. [11]) obtained after a smooth interpolation at the physical value of the strange quark mass $m_s^{\text{phys}} = 99.6(4.3)$ MeV [10]. This allows us to adopt the following SU(2)-inspired ansatz,

$$B_{\text{CMO}}^{K\pi} = \alpha + \beta m_{ud} + \gamma m_{ud}^{1/2} + \Delta + Da^2, \quad (29)$$

where the parameters α , β , and γ play the role of $SU(2)$ LECs, while the function Δ includes chiral corrections beyond the NLO. Then, we have performed fits of the 15 lattice points of the $K\pi$ channel (interpolated at $m_s = m_s^{\text{phys}}$) adopting three choices for Δ , namely $\Delta = 0$, $\Delta \propto m_{ud}^{3/2}$ and $\Delta \propto m_{ud}^2$. For all the three choices we obtain a good




FIG. 4. Values of the B parameter $B_{\text{CMO}}^{K\pi}$ as a function of the renormalized light-quark mass m_{ud} in the $\overline{\text{MS}}(2 \text{ GeV})$ scheme. The lattice data have been smoothly interpolated to the physical value of the strange quark mass $m_s^{\text{phys}} = 99.6(4.3)$ MeV [10]. The dashed, short-dashed, and dotted lines correspond to the results of the SU(2)-inspired fit (29) assuming $\Delta = 0$ for the three values of the lattice spacing, while the solid line is the result in the continuum limit. The diamonds represent the values of the B parameters at the physical pion mass and in the continuum limit.

description of the lattice data ($\chi^2/\text{d.o.f.} \simeq 0.4, 0.2$ and 0.2 , respectively). The extrapolation to the physical pion point, in the three cases, yields $B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.275(18)$ ($\Delta = 0$), $B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.340(44)$ ($\Delta \propto m_{ud}^{3/2}$), and $B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.327(51)$ ($\Delta \propto m_{ud}^2$). The results of the fit (29) assuming $\Delta = 0$ are shown in Fig. 4.

Adopting the SU(2)-inspired fit (29) and averaging the results of the different fits using Eq. (28) of Ref. [10], we obtain

$$\begin{aligned} B_{\text{CMO}}^{K\pi}|_{\text{phys}} &= 0.306(37)_{\text{stat+fit}}(44)_{\text{chir}}(16)_{\text{disc}}(7)_{Z_p} \\ &= 0.306(60), \end{aligned} \quad (30)$$

which is consistent with the SU(3) result (28) though with much larger statistical and systematic uncertainties.

Thus, performing a weighted average of the SU(2) and SU(3) results, we quote for $B_{\text{CMO}}^{K\pi}|_{\text{phys}}$ the final value

$$B_{\text{CMO}}^{K\pi}|_{\text{phys}} = 0.273(13)(68)_{PT} = 0.273(69), \quad (31)$$

where the second error accounts for the perturbative uncertainty in the one-loop determination of the multiplicative RC Z_{CM} (see Table II) and it has been estimated to be $\simeq 25\%$ relying on the difference between the values of Z_{CM} obtained with and without boosted perturbation theory. As it can be seen, this error represents the largest source of uncertainty in the determination of the $B_{\text{CMO}}^{K\pi}$ parameter.

Our result (31) represents the first lattice QCD determination of a matrix element of the CMO. In the SU(3) chiral limit we obtain $B_{\text{CMO}} = 0.076(14)(18)_{PT} = 0.076(23)$.

Our findings are significantly smaller than the model-dependent estimate $B_{\text{CMO}} \sim 1-4$ currently adopted in phenomenological analyses [1]. The comparison indicates also that the uncertainty on this important phenomenological quantity has been significantly reduced by lattice QCD. A further drastic improvement in the precision can be achieved by removing the uncertainty due to the use of one-loop perturbation theory for estimating the multiplicative RC Z_{CM} , which is by far the dominating source of uncertainty in our results.

VI. CONCLUSIONS

We have presented the results of the first lattice QCD calculation of the $K \rightarrow \pi$ matrix elements of the chromomagnetic operator $O_{\text{CM}} = g\bar{s}\sigma_{\mu\nu}G_{\mu\nu}d$, which appears in the effective Hamiltonian describing $\Delta S = 1$ transitions in and beyond the standard model.

Having dimension five, the chromomagnetic operator is characterized by a rich pattern of mixing with operators of equal and lower dimensionality. The power divergent coefficients controlling the mixing with operators of lower dimension were determined nonperturbatively in Ref. [4], while the multiplicative renormalization factor as well as the mixing coefficients with the operators of equal dimension have been computed at one-loop in perturbation theory [4]. The precision achieved in the nonperturbative

evaluation of the mixing with the scalar density leads to an extraction of the matrix element of the renormalized chromomagnetic operator with good accuracy.

The numerical simulations have been carried out using the gauge field configurations produced by ETMC with $N_f = 2 + 1 + 1$ dynamical quarks at three values of the lattice spacing. Our result for the B parameter of the chromomagnetic operator at the physical pion and kaon point is $B_{\text{CMO}}^{K\pi} = 0.273(69)$, while in the SU(3) chiral limit we get $B_{\text{CMO}} = 0.076(23)$. Our findings are significantly smaller than the model-dependent estimate $B_{\text{CMO}} \sim 1-4$, currently used in phenomenological analyses, and improve the uncertainty on this important phenomenological quantity.

ACKNOWLEDGMENTS

We warmly thank Jean-Marc Gerard for pointing out the proper chiral normalization of the BCMOKK parameter. We gratefully acknowledge the CPU time provided by PRACE under the Project No. PRA027, ‘‘QCD Simulations for Flavor Physics in the Standard Model and Beyond,’’ on the JUGENE BG/P system at JSC (Germany) and by the agreement between INFN and CINECA under the specific initiative INFN-LQCD123 on the Fermi BG/Q system at CINECA (Italy). V. L., G. M., and S. S. thank MIUR (Italy) for partial support under Contract No. PRIN 2015P5SBHT.

-
- [1] A. J. Buras, G. Colangelo, G. Isidori, A. Romanino, and L. Silvestrini, *Nucl. Phys.* **B566**, 3 (2000).
 - [2] D. Becirevic, V. Lubicz, G. Martinelli, and F. Mescia (SPQcdR Collaboration), *Phys. Lett. B* **501**, 98 (2001).
 - [3] I. Baum, V. Lubicz, G. Martinelli, L. Orifici, and S. Simula, *Phys. Rev. D* **84**, 074503 (2011).
 - [4] M. Constantinou, M. Costa, R. Frezzotti, V. Lubicz, G. Martinelli, D. Meloni, H. Panagopoulos, and S. Simula, *Phys. Rev. D* **92**, 034505 (2015).
 - [5] X. G. He, H. Murayama, S. Pakvasa, and G. Valencia, *Phys. Rev. D* **61**, 071701 (2000).
 - [6] G. D’Ambrosio, G. Isidori, and G. Martinelli, *Phys. Lett. B* **480**, 164 (2000).
 - [7] S. Bertolini, J. O. Eeg, and M. Fabbrichesi, *Nucl. Phys.* **B449**, 197 (1995).
 - [8] R. Baron *et al.* (ETM Collaboration), *J. High Energy Phys.* **06** (2010) 111.
 - [9] R. Baron *et al.* (ETM Collaboration), *Proc. Sci., LATTICE10* (2010) 123.
 - [10] N. Carrasco *et al.* (ETM Collaboration), *Nucl. Phys.* **B887**, 19 (2014).
 - [11] M. Constantinou, M. Costa, R. Frezzotti, V. Lubicz, G. Martinelli, D. Meloni, H. Panagopoulos, and S. Simula, *Proc. Sci. LATTICE*, 2014 (2014) 390, [arXiv:1412.1351](https://arxiv.org/abs/1412.1351).
 - [12] Y. Iwasaki, *Nucl. Phys.* **B258**, 141 (1985).
 - [13] R. Frezzotti, P. A. Grassi, S. Sint, and P. Weisz, *J. High Energy Phys.* **08** (2001) 058.
 - [14] R. Frezzotti and G. C. Rossi, *Nucl. Phys. B, Proc. Suppl.* **128**, 193 (2004).
 - [15] R. Frezzotti and G. C. Rossi, *J. High Energy Phys.* **08** (2004) 007.
 - [16] R. Frezzotti and G. C. Rossi, *J. High Energy Phys.* **10** (2004) 070.
 - [17] K. Osterwalder and E. Seiler, *Ann. Phys. (N.Y.)* **110**, 440 (1978).
 - [18] B. Jegerlehner, [arXiv:hep-lat/9612014](https://arxiv.org/abs/hep-lat/9612014).
 - [19] K. Jansen, M. Papinutto, A. Shindler, C. Urbach, and I. Wetzorke (XLF Collaboration), *J. High Energy Phys.* **09** (2005) 071.
 - [20] C. McNeile and C. Michael (UKQCD Collaboration), *Phys. Rev. D* **73**, 074506 (2006).
 - [21] T. Bhattacharya, V. Cirigliano, R. Gupta, E. Mereghetti, and B. Yoon, *Phys. Rev. D* **92**, 114026 (2015).
 - [22] M. Testa, *J. High Energy Phys.* **04** (1998) 002.
 - [23] L. Maiani, G. Martinelli, and C. T. Sachrajda, *Nucl. Phys.* **B368**, 281 (1992).
 - [24] E. Gabrielli, G. Martinelli, C. Pittori, G. Heatlie, and C. T. Sachrajda, *Nucl. Phys.* **B362**, 475 (1991).
 - [25] S. Aoki *et al.*, *Eur. Phys. J. C* **77**, 112 (2017); See also S. Aoki *et al.*, *Eur. Phys. J. C* **74**, 2890 (2014).