

Comparative anatomy

Plasticity of brain in normal adult some earth-dwelling Anamia and terrestrial Amniota: further review on the trend of seasonal environmental influence on encephalic proliferation, as revealed by immunohistochemistry

Vito Margotta¹, Claudio Chimenti^{2,*}Departments of ¹Animal and Human Biology, and ²Biology and Biotechnology "Charles Darwin", University "La Sapienza", Roma, Italy

Abstract

The immunohistochemical evidence provided by the present authors about the persistence of latent natural brain proliferative potentially in adult vertebrate brain and its response to cyclic seasonal environmental fluctuations (temperature, photoperiod) has been reviewed. These stimuli elicit an otherwise hidden mitotic activity thanks to stem cells still present especially in less high vertebrates like *Triturus carnifex*, *Rana bergeri*, *Podarcis sicula*. The evidence gathered from specimens caught in the wild in spring, summer and autumn has been compared with previous evidence on specimens of newts, frogs and lizards. Signs of proliferation were mainly observed in the typical sites (olfactory bulbs/peduncles, telencephalic hemispheres) occupied by cells in mitotic stand-by. The findings have shown increasing labelling from spring to summer to autumn with minor differences among species, and have confirmed that in adulthood the proliferative therefore the reparative and even the regenerative power of brain cells is highest in urodela (the vertebrates best equipped with quiescent cells), intermediate in anura and lowest in lacertilian Reptiles.

Keywords

Neural stem cells, matrix cells, matrixareas, Amphibians, lacertilian Reptiles.

Review

Since about a half century detailed awareness has been acquired on the plasticity of the brain in some adult heterothermic vertebrates: fresh water, earth-dwelling Anamia and terrestrial Amniota (Margotta and Morelli, 1996).

Among these investigations a thread of research has been devoted to study if in adults the impact of cyclic seasonal environmental fluctuations (consisting in temperature and photoperiod variations) could exert any influence on brain in terms of physiological cell proliferative answer, or unmask a latent spontaneous proliferative power thus making apparent reparative and even regenerative potentialities due to an otherwise hidden mitotic activity of stem cells still present in the adult brain of many vertebrates species, mainly in lower ones.

* Corresponding author. E-mail: claudio.chimenti@uniroma1.it

Such events have been correlated with the persistence in the adult of a stock of brain stem cells which have some own characteristics: morphological (being small and basophilic), physiological (able to proliferate), behavioural (persistence of the tendency to proliferate).

Usually these undifferentiated cells appear as clusters of grouped cells, sometimes layered, in circumscribed areas ("matrix areas", once nicknamed *Matrixzonen*, according to Kirsche, 1967) typically located among the ependyma cells lining each encephalic cavity and in the sub-ependyma layer, and as scattered cells ("matrix cells") within these same areas or elsewhere in cerebral tissues.

The number and size of such stem cells can vary among the different vertebrate groups and species; generally speaking, they appear much more numerous and large in lower than in higher species. In a gradual scale of value regarding the adult fresh water, earth-dwelling Anamnia and heterothermic terrestrial Amniota, such cells in mitotic stand-by appear to be relatively abundant in the urodelan Amphibians (the best provided with these cells among vertebrates), intermediate in the Teleosts, and relatively scarce in the anuran Amphibians and above all in the lacertilian Reptiles.

The number of such sleeping cells can vary according to different encephalic districts and is persistently high in the forebrain (olfactory bulbs/peduncles, telencephalic hemispheres), where they are mainly distributed in characteristic, mirror-like sites: the matrix cells in proximity of the olfactory cavities, the matrix areas at the edge, latero-dorsally and ventrally at the bottom of the sickle-shape, hollow ventricular surface of each telencephalic hemisphere: *zonae germinativae dorsales* and *ventrales*, respectively. In particular, only in lacertilian Reptiles each *zona germinativa dorsalis* appear sub-divisible in two portions, *lateralis* and *medialis*, like in *Lacerta viridis* (Minelli and Del Grande, 1980).

The *zonae ventrales* are the best provided with undifferentiated cells; intra-specific difference can be found among the brain of the earth-dwelling Anamnia and poikilothermal Amniota studied. Matrix cells can be observed here and there in the dien-cephalon. Such cells appear absent from the midbrain, in the *truncus cerebri*, and the *cerebellum* (in both sites, with the only exception of Teleosts in which it is possible to recognize symmetrical, additional areas provided with proliferative potential: the *zonae germinativae caudales* in the midbrain and scattered cells in the cerebellar deep tissue. Sometimes mitotically dormant cells can be found in the *medulla oblongata*.

Both telencephalic *zonae germinativae* are extended antero-posteriorly and active with different time courses: the *zonae dorsales* appear to exhaust their self-maintaining potential earlier than the *zonae ventrales*, which are generally wider and richer in cells (Kirsche, 1967).

All these information has been acquired and expanded through observations on untreated animals and upon experimental intervention, more frequently represented by brain surgery, ablations of encephalic plugs or wider portions sometimes with subsequent hetero- and rarely homo-transplantation (even of the whole brain), and *in vitro* culture of cerebral tissues. Analytical techniques were at first traditional histology, then autoradiography and immunohistochemistry, seldom electron microscopy.

These quiescent cells are remnants of the neural layer which forms in the early embryo and is responsible for the morphogenesis of the central nervous system (Kahle, 1951; Fujita, 1963; Kirsche, 1967), which may explain why the number of such cells decreases during the life of the organism: going from earlier to more advanced

embryonic stages, then through the subsequent larval ones - if present - and eventually into adult life.

The greater part of the information on this subject is the consequence of the exhaustive studies of Kirsche (1967, 1983), who investigated adult non-mammalian vertebrates by traditional histological methods. The studied species ranged from Teleosts to Birds, passing through urodelan and anuran Amphibia and lacertilian Reptiles. This author must be credited for the generation of most data on the features and localization of these putative precursor or stem cells in the adult brain and the recognition that they can be normally silent but are capable of self-reproduction and can start cycling again giving rise to descendants which undergo differentiation into neuronal or glial cells (Kirsche, 1967, 1983).

The persistence also in adult life of such cells in mitotic stand-by, thanks to their proliferative power, sustains both the physiological and experimental proliferative events and explains the reparative and even regenerative potentialities still present in the adult brain of many vertebrate species, especially in the less high ones.

In the last years investigations have been carried on adults of some earth-dwelling Anamnia (frogs and newts) and terrestrial Amniota (lizards) to investigate the spontaneous brain proliferative answer to cyclic seasonal environmental changes (temperature and photoperiod variations). These investigations were performed with the help of immunohistochemistry for Proliferating Cell Nuclear Antigen (PCNA), expressed by proliferating cells (Miyachi et al., 1978; also see Margotta and Chimenti, 2016).

By other authors in the past and by ourselves (Margotta, 2012, 2014a; Margotta and Chimenti, 2016, 2017, 2018, in press, submitted) attention was paid to spring, summer and autumnal seasonal environment, while no attention was devoted to winter season conditions. In the last conditions autoradiographic studies were done by Minelli et al. (1982) on injured and uninjured brain of adult *Rana esculenta*, observing a post-autumnal arrest of encephalic proliferation that reached a nadir in full winter.

Immunohistochemical studies were done by Ramirez et al. (1997) on adult brain-damaged *Podarcis hispanica*, who referred that "...cold (winter) temperature prevented migration of the newly generated immature neurons".

An explanation on the "winter" findings of Minelli et al. (1982) and Ramirez et al. (1997) was proposed by Margotta and Morelli (1997), who, on the basis of the literature, proposed a role for the glial cells in the plasticity of the central nervous system in the adults of several vertebrate species.

Therefore, adult normal brain of *R. bergeri*, *Triturus carnifex*, *Podarcis sicula* - once, respectively, *R. esculenta* (Capula, 2000 a) *T. cristatus carnifex* (Bonifazi, 2000) and *L. viridis* (Capula, 2000 b) - were again investigated by Margotta (2012, 2014a), Chimenti and Margotta (2015) and Margotta and Chimenti (2016, 2017, 2018, in press). Staining was observed in the typical localization of putative stem cells, as "matrix cells" in the olfactory (Figs. 1; 2; 3) and diencephalic districts or as "matrix areas" in the telencephalic hemispheres (Figs. 4; 5; 6) where labelling was in correspondence of the ventricular edges of *zonae germinativae dorsales* (Figs. 4; 5), which in lizards are subdivided, as already said, each in *lateralis* and *medialis* (Fig. 6), and also in relationship with the bottom of the ventricular cavities in *zonae germinativae ventrales* (Figs. 4; 5; 6). Labelled cells were found among the ependymal cells and in the sub-ependymal layer. This pattern emerged also from further immunohistochemical studies on the same earth-dwelling Anamnia (Margotta and Chimenti, 2016, 2017, 2018, in press)




Fig. 1. Drawings (not to scale) of a transverse view of adult normal brain of *Triturus carnifex*. Olfactory bulbs in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, isolated in the ependyma and periependymal grey matter of the olfactory bulbs.




Fig. 2. Drawings (not to scale) of a transverse view of adult normal brain of *Rana bergeri*. Olfactory bulbs in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, isolated in the ependyma and periependymal grey matter of the olfactory bulbs.

and terrestrial Amniota (Margotta and Chimenti, submitted). The findings were more evident in *T. carnifex* (Figs. 1, 4), less so in *R. bergeri* (Figs. 2, 5) and least in *P. sicula* (Figs. 3, 6).

In the investigated species the immunohistochemical signs of proliferation show an ascending trend from spring (Figs. 1a, 2a, 3a, 4a, 5a, 6a), through summer (Figs. 1b, 2b, 3b, 4b, 5b, 6b), to autumn (Figs. 1c, 2c, 3c, 4c, 5c, 6c), with only minor differences linked to the position of each species in the zoological/evolutionary scale.

The present authors' findings in different species and conditions have expanded those of Minelli et al. (1982) and Ramirez et al. (1997) and strongly support the hypothesis proposed by present authors, that the entity of the spontaneous proliferative processes along seasons accounts for the reparative or even regenerative phenomena obtained by previous authors, the extent of which could depend on synergic stimulation by various stressful stimuli: extreme temperature, surgery, other types of trauma. This was also supported by a comparison between the findings of previous authors on adult *R. esculenta* in autumn (Minelli et al. 1982) and on adult *P. hispanica* in summer (Ramirez et al. 1997) and those of present authors on uninjured adult *R. bergeri* (Margotta, 2012; Margotta and Chimenti, 2017, 2018), *P. sicula* (Margotta,




Fig. 3. Drawings (not to scale) of a transverse view of adult normal brain of *Podarcis sicula*. Olfactory bulbs in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, isolated in the ependyma and periependymal grey matter of the olfactory bulbs.




Fig. 4. Drawings (not to scale) of a transverse view of adult normal brain of *Triturus carnifex*. Telencephalic hemispheres in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, clustered in the walls of telencephalic symmetrical ventricles to form the *zonae germinativae dorsales* (arrowheads) and *ventrales* (arrows).

2014a; Margotta and Chimenti, submitted), *T. carnifex* (Margotta and Chimenti, in press) and on adult poikilothermal earth-dwelling and terrestrial vertebrates subjected to cold shock (Chimenti and Margotta, 2013, 2015; Margotta, 2014b, 2015).

It can be said, therefore, that in adults the proliferative and therefore the reparative and even the regenerative power of the brain increases progressively from the lacertilian Reptiles to anuran Amphibians, Teleosts and eventually urodelan Amphibians, due to the respective, available stock of cells in mitotic stand-by of which the brain is especially equipped in urodelans. The latter group of Amphibians is, among vertebrates, that which occupies the most privileged position, since these animals are the richest in undifferentiated cells, therefore those most endowed with proliferative potentiality and hence the best equipped for reparative or even regenerative processes.

Recently, in an immunohistochemistry report (Margotta and Chimenti, 2016) we exposed the influence of seasonal variations alone or coupled with an induced cold shock on spontaneous brain proliferation in adult poikilothermal earth-dwelling Anamnia and terrestrial Amniota; similar results were found in frogs, newts and lizards.




Fig. 5. Drawings (not to scale) of a transverse view of adult normal brain of *Rana bergeri*. Telencephalic hemispheres in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, clustered in the walls of telencephalic symmetrical ventricles to form the *zonae germinativae dorsales* (arrowheads) and *ventrales* (arrows).




Fig. 6. Drawings (not to scale) of a transverse view of adult normal brain of *Podarcis sicula*. Telencephalic hemispheres in specimens caught in the wild in the wild in spring (a), summer (b), autumn (c). The dots represent PCNA immunolabelled matrix cells, clustered in the walls of telencephalic symmetrical ventricles to form the *zonae germinativae latero-dorsales* (arrowheads), *medio-dorsales* (arrowforkeds) and *ventrales* (arrows).

The content of the actual exposition forms with that of the previous reports an *unicum* with which we have attempted to furnish a whole seasonal panorama of the natural cell proliferative trend as consequence of the impact of the cyclic seasonal environmental fluctuations (made of temperature and photoperiodic variations), revealed by the immunoreactive patterns, in the adult brain of heterothermic some vertebrates.

Acknowledgements

This report was supported by a grant from Ministero per l’Istruzione, l’Università e la Ricerca of Italian Republic.

References

- Bonifazi A. (2000) *Triturus carnifex*. In: Bologna M.A., Capula M., Carpaneto G.M. (Eds.) Anfibi e Rettili del Lazio. Fratelli Palombi Editori, Roma. Pp. 42-43.
- Capula M. (2000a) *Rana bergeri* (Günther, 1866) *Rana kl. hispanica* (Bonaparte, 1839). In: Bologna M.A., Capula M., Carpaneto G.M. (Eds.) Anfibi e Rettili del Lazio. Fratelli Palombi Editori, Roma. Pp. 56-57.
- Capula M. (2000b) *Podarcis sicula* (Rafinesque-Schmaltz, 1810). In: Bologna M.A., Capula M., Carpaneto G.M. (Eds.) Anfibi e Rettili del Lazio. Fratelli Palombi Editori, Roma. Pp. 86-87.
- Chimenti C., Margotta V. (2013) Proliferative events experimentally induced by transient cold shock in the brain of adult terrestrial heterothermic vertebrates: preliminary analysis of PCNA expression in *Triturus carnifex*. Ital. J. Anat. Embryol. 118: 105-118
- Chimenti C., Margotta V. (2015) Interaction between spring temperature-photoperiod and experimentally induced transient cold shock influencing proliferative activity in the brain of an adult terrestrial heterothermic vertebrate, *Rana bergeri* (Günther, 1866). Ital. J. Anat. Embryol. 120: 89-98.
- Fujita S. (1963) The matrix cell and cytogenesis in the developing central nervous system. J. Comp. Neur. 120: 37-42.
- Kahle W. (1951) Studien über die Matrixphasen und die örtlichen Reifungsunterschiede im embryonalen menschlichen Gehirn. I. Mitteilung: Die Matrixphasen im allgemeinen: Dtsch. Zschr. Nervenheilk. 166: 273-302.
- Kirsche W. (1967) Über postembryonale Matrixzonen im Gehirn verschiedener Vertebraten und deren Beziehung zur Hirnbauplanlehre. Z. mikrosk.-anat. Forsch. 77: 313-406.
- Kirsche W. (1983) The significance of matrix zones for brain regeneration and brain transplantation with special consideration of lower vertebrates. Chapter 2. In: Wallace R.B., Das G.D. (Eds.), Neural Tissue Transplantation Research. Springer-Verlag, New York Berlin Heidelberg Tokyo. Pp. 65-104.
- Margotta V. (2012) Relationships between seasonal thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Rana bergeri* (Günther, 1866). Ital. J. Anat. Embryol. 117: 45-53.
- Margotta V. (2014a) Relationships between seasonal thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Podarcis sicula*. Ital. J. Anat. Embryol. 119: 29-37.
- Margotta V. (2014b) Proliferative events experimentally induced by a transient cold shock in the brain of adult terrestrial heterothermic vertebrates: preliminary analysis of PCNA expression in *Podarcis sicula*. Ital. J. Anat. Embryol. 119: 81-91.
- Margotta V. (2015) Interaction between autumnal temperature-photoperiod and experimentally induced transient cold shock influences proliferative activity in the brain of an adult terrestrial heterothermic vertebrate, *Rana bergeri*. Ital. J. Anat. Embryol. 120: 192-200.
- Margotta V., Chimenti C. (2016) Plasticity of the central nervous system in adult vertebrates: immunohistochemical report on the effects of seasonal variations alone or coupled with induced cold shock on brain proliferation in fresh water or earth-dwelling Anamnia and heterothermic Amniota. Ital. J. Anat. Embryol. 121: 265-283.

- Margotta V., Chimenti C. (2017) Relationships between seasonal (spring or autumnal) thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Rana bergeri* (Günther, 1986). *Ital. J. Anat. Embryol.* 122: 89-97.
- Margotta V., Chimenti C. (2018) Relationships between summer thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Rana bergeri* (Günther, 1986). *Ital. J. Anat. Embryol.* 123: 100-107.
- Margotta V., Chimenti C. Relationships between seasonal (spring, summer, autumnal) thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Triturus carnifex*. *Ital. J. Anat. Embryol.* (in press).
- Margotta V., Chimenti C. Relationships between seasonal (spring, summer, autumnal) thermal variations and cell proliferation in heterothermic vertebrates, as revealed by PCNA expression in the brain of adult *Podarcis sicula*. *Ital. J. Anat. Embryol.* (submitted).
- Margotta V., Morelli A. (1996) Encephalic matrix areas and post-natal neurogenesis under natural and experimental conditions. *Anim. Biol.* 5: 117-131.
- Margotta V., Morelli A. (1997) Contribution of radial glial cells to neurogenesis and plasticity of central nervous system in adult Vertebrates. *Anim. Biol.* 6: 101-108.
- Minelli G., Del Grande P. (1980) An hypothesis on the influence of the temperature on the telencephalic reparative processes in *Lacerta viridis*. *Z. mikrosk.-anat. Forsch.* 94: 209-216.
- Minelli G., Del Grande P., Franceschini V. (1982) Uptake of 6-H3 thymidine in the normal and regenerating CNS of *Rana esculenta*. *Z. mikrosk.-anat. Forsch.* 96: 201-213.
- Miyachi K., Fritzler M.J., Tan E.M. (1978) Autoantibody to a nuclear antigen in proliferating cells: *J. Immunol.* 121: 2228-2234.
- Ramirez C., Nacher J., Molowny A., Sancez-Sancez F., Irurzun A., Lopez-Garcia C. (1997) Photoperiod-temperature and neuroblast proliferation-migration in the adult lizard cortex. *NeuroReport* 8: 2337-2342.