

Immunometabolic Approaches to Prevent, Detect, And Treat Neonatal Sepsis

Maria Giulia Conti^{1,2}, Asimena Angelidou^{2,3,4}, Joann Diray-Arce^{2,4}, Kinga Smolen^{2,4}, Jessica Lasky-Su^{4,5}, Mario De Curtis¹ and Ofer Levy^{*2,4,6}

¹Department of Maternal and Child Health and Urological Sciences, Sapienza University of Rome, Rome, Italy;

²*Precision Vaccines Program*, Division of Infectious Diseases, Boston Children's Hospital, Boston, MA;

³Division of Newborn Medicine, Boston Children's Hospital, Boston, MA;

⁴Harvard Medical School, Boston, MA;

⁵Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA;

⁶Broad Institute of MIT & Harvard, Cambridge, MA.

Author contributions:

MGC: conception and design, drafting the article; AA: substantial contributions to conception and design, drafting the article, revising the article critically for important intellectual content; JDA: revising the article critically for important intellectual content; KS: revising the article critically for important intellectual content; JLS: drafting the article and revising it critically for important intellectual content; MDC and OL: revising the article critically for important intellectual content, final approval of the version to be published.

Corresponding author*:

Ofer Levy MD, PhD

Precision Vaccines Program

Boston Children's Hospital

Boston, MA 02115

+617-919-2904

ofer.levy@childrens.harvard.edu

Statement of financial support:

No financial assistance was received in support of the article.

Disclosure statement:

OL is a named inventor on several patents relating to vaccine adjuvants.

Category of study:

Review

- Neonatal sepsis is a leading cause of early life mortality and there is an unmet need to address it.
- Emerging evidence indicates that immunometabolism is relevant to the pathophysiology of neonatal sepsis, but much remains to be learned regarding its distinct features and regulation.
- We discuss the interconnection of metabolism and immunity in early life and how distinct immunometabolism may be leveraged to improve prevention, diagnosis, and therapy of neonatal sepsis.

ABSTRACT

The first days of postnatal life are energetically demanding as metabolic functions change dramatically to accommodate drastic environmental and physiologic transitions after birth. It is increasingly appreciated that metabolic pathways are not only crucial for nutrition but also play important roles in regulating inflammation and the host response to infection. Neonatal susceptibility to infection is increased due to a functionally distinct immune response characterized by high reliance on innate immune mechanisms. Interactions between metabolism and the immune response are increasingly recognized as changes in metabolic pathways drive innate immune cell function and activation and consequently host response to pathogens. Moreover, metabolites, such as acetyl-coenzyme A (acetyl-CoA) and succinate have immunoregulatory properties and serve as cofactors for enzymes involved in epigenetic reprogramming or “training” of innate immune cells after an initial infectious exposure. Highly sensitive metabolomic approaches allow us to define alterations in metabolic signatures as they change during ontogeny and as perturbed by immunization or infection thereby linking metabolic pathways to immune cell effector functions. Characterizing the ontogeny of immunometabolism will offer new opportunities to prevent, diagnose and treat neonatal sepsis.

Introduction

Sepsis, leading to life-threatening organ dysfunction due to a dysregulated host response to infection [1], is a major contributor to global morbidity and mortality, particularly in early life [2]. In 2017, the World Health Organization (WHO) estimated 5.4 million deaths in children under five years of age, with newborns (those less than 28 days of life) accounting for nearly 50% of total deaths ([WHO, Child mortality and causes of death, 2017](#)). Those born preterm, comprising 11% of all births globally, are particularly susceptible to infection both early after birth and throughout childhood and at high risk for long-term complications [3].

Defining neonatal sepsis remains challenging, without a clear consensus definition. Indeed, current pediatric definitions for sepsis are not validated in infants [4]. A better understanding of the pathophysiological basis and the molecular and biochemical processes involved in neonatal sepsis is crucial for early diagnosis and intervention.

In early life, response to infection is heavily reliant on innate immunity including anatomical barriers such as skin, intestinal and respiratory mucosae as well as cellular (e.g., macrophages, monocytes, natural killer cells) and soluble (e.g., antimicrobial proteins and peptides and acute-phase reactants) components [5, 6]. Patients with primary innate immunodeficiency (e.g., Interleukin-1 receptor-associated kinase 4 (IRAK-4) deficiency, Myeloid differentiation primary response 88 (MyD88) deficiency, etc.) have a high susceptibility to early life infection [6]. Of note, the innate immune system has adaptive features such that innate immune cells undergo epigenetic reprogramming after stimulation, leading to an altered, potentially enhanced, response after subsequent stimulation with unrelated pathogens, a property termed “innate immune memory” [7].

Key to the regulation of innate immunity is metabolism. Indeed, complex interactions between metabolism and the immune response are increasingly recognized, giving rise to the new field of “immunometabolism” [8]. Metabolic pathways, such as glycolysis, the

tricarboxylic acid (TCA) cycle (also known as Krebs Cycle), fatty acid oxidation (FAO), fatty acid synthesis (FAS) and amino acid pathways, play important roles in the generation of key products responsible for promoting innate immune cell survival or growth, function and activation [8]. The energy cost required to fuel the immune response highlights the crucial role of metabolites and metabolic pathways in sepsis. In addition, certain metabolites such as acetyl-coenzyme A (acetyl-CoA), succinate, nicotinamide adenine dinucleotide (NAD⁺) and α -ketoglutarate can serve as co-factors for epigenetic enzymes, thus potentiating innate immune memory [9]. In the era of systems biology, wherein the entire inventory of molecules of a given class can be measured, the unique dynamic changes of the newborn period can be studied at the molecular level and offer insights into the complex interactions between the host and the insulting pathogen [10]. Sepsis-induced alterations in the epigenome, transcriptome, proteome and metabolome are reflected in changes in the concentration of the small molecules and chemicals in biological fluids and tissues [11], thus opening opportunities for novel approaches to sepsis prevention, diagnosis and treatment.

In this review, we focus on the current understanding of the neonatal energetic and metabolic response to sepsis. We identify certain metabolites as potential diagnostic markers, discuss the use of enhanced nutritional support as a tool to treat sepsis, and identify metabolically active medications as potential ways to manipulate immunometabolism with the goal of preventing and/or treating sepsis.

Metabolic pathophysiology of neonatal sepsis

Immunometabolism is thus far relatively underexplored with respect to neonatal health and disease. In sepsis, host defense to pathogen invasion is characterized by an early pro-inflammatory response, triggered mainly by innate immune system activation, leading to an exaggerated systemic release of cytokines. While aiming at pathogen elimination, this acute

response can cause hypo-tension, cardiovascular dysfunction, tissue damage and multi-organ failure [12]. During the later phases of sepsis, activation of modulatory pathways induces an anti-inflammatory state that contains inflammation and promotes tissue repair, while also limiting host defense; this regulatory mechanism is called innate immune tolerance (minimizing harm from immunopathology) or innate immune paralysis when this progresses toward a dysfunctional state, in which cells are unable to respond properly to stimulation, with consequent increased risk of opportunistic infections [12]. A shift in basic cellular energy metabolism from oxidative phosphorylation and FAO to glycolysis, known as the Warburg Effect, is required during early immune cell activation to provide the necessary rapid energy and metabolic intermediates, while in tolerance and paralysis all major metabolic pathways are impaired [13] [14].

The neonatal immune reaction to infection is markedly different from that of older infants, children and adults [15]: a) neonates demonstrate distinct polarized response to infections, and whole blood transcriptomics in neonates with septic shock demonstrated down-regulation of genes related to innate and adaptive immunity compared to toddlers and older children [16]; b) neonatal sepsis is accompanied by a heightened innate immune cellular response driven by monocytes/macrophages and neutrophils and counter-balanced by inhibitory pathways resulting in a dampened adaptive immune response [15]; c) neonatal innate immunity is distinct: Pattern Recognition Receptors (PRR)-mediated immune responses to most stimuli are generally polarized away from inflammatory/T helper (Th) 1-polarizing cytokines [17-19]. Up-regulation of specific inhibitory signaling genes has been described in neonatal sepsis, likely serving to avoid an excessive inflammatory response [20]. These unique features of the fetal/neonatal immune system serve to avoid energetically costly and potentially harmful immune responses during early life [15, 21].

In the following sections, we will discuss cellular metabolism in innate immune cells at quiescent and activated states (Figure 1) and attempt to elucidate differences between newborns and adults.

Glycolysis and the adenosine system

Glycolysis is a relatively inefficient (producing only two molecules of ATP per unit of glucose), but rapidly initiated pathway for generation of cellular ATP. In acute inflammation, enhanced glycolysis enables immune cells to quickly generate (a) sufficient ATP, the main energy substrate to ensure cell functionality, and (b) biosynthetic intermediates (i.e. serine, glycine, alanine, acetyl-CoA for lipid synthesis) to carry out their specific effector functions [12-14, 22]. ATP serves as a danger-associated molecular pattern (DAMP) for the immune system, which can initiate and prolong immune responses [23]. ATP levels increase significantly during inflammation, hypoxia, or ischemia, as seen in various disease states including preeclampsia and sepsis [24, 25]. To counteract ATP-induced immune effects, cells can hydrolyze ATP into adenosine diphosphate (ADP) and adenosine monophosphate (AMP). AMP can subsequently be broken down into adenosine and phosphate. Adenosine levels rise rapidly in response to systemic inflammation to reduce pro-inflammatory/Th1-polarizing immune responses. Both ATP and adenosine bind to purinergic receptors, so the overall inflammatory effect of ATP on the biological system depends on the balance between ATP and adenosine [26]. Interestingly, levels of adenosine are significantly higher in neonatal compared to adult plasma, contributing to Th2 polarization of Toll like Receptor (TLR)-mediated responses [27, 28]. Comparison of the glycolytic capacity of preterm, term, and adult monocytes at rest demonstrated a severe impairment of glycolysis in preterm monocytes compared to adults, while in monocytes of term newborns, glycolysis was variably affected, reflecting a transitional functional state [22]. Reduced glycolysis was also recently described in

cord blood activated macrophages as a consequence of down-regulation and lack of activity (reduced phosphorylation) of the mammalian target of Rapamycin (mTOR) [29]. The mTOR pathway promotes the metabolic switch toward glycolysis after PRR activation by Pathogen-associated molecular patterns (PAMPs) (i.e. TLRs activation by LPS), and is fundamental for robust cytokine production (i.e. TNF, IL-1 β) and an effective host defense in the acute phase of infection and sepsis [30, 31]. Down-regulation of the mTOR pathway and shutdown of glycolysis is also mediated by alarmins (i.e. S100A8 and S100A9 proteins), which are secreted by neutrophils and activated macrophages and extremely abundant in neonatal serum, thus downregulating responses to microbes and thereby preventing harmful hyper-inflammatory responses [29, 32]. Interestingly in an *in vitro* model of immune paralysis, re-stimulation of immune-tolerant adult human monocytes with LPS revealed a defective ability to mount a Warburg effect with diminished production of lactate and enhanced production of NAD⁺, and was not followed by production of pro-inflammatory cytokines [14]. The same study revealed that tolerant monocytes from adult septic patients have impaired metabolic pathways compared to healthy controls and septic patients with normal cytokine responsiveness (non-tolerant). Similar studies are needed to investigate the neonatal metabolic profile in immune tolerance and paralysis.

TCA cycle

The TCA cycle is based in the mitochondrial matrix and together with oxidative phosphorylation is the major metabolic pathway during the quiescent or non-proliferative cellular stage (e.g., M2 macrophages), efficiently generating ATP for cellular functions. On the other hand, activated innate immune cells (e.g., pro-inflammatory M1 macrophages, cells activated by LPS) experience TCA cycle interruptions to achieve accumulation of intermediary substrates, citrate and succinate, in order to promote their immune functions [33]. While citrate

is necessary for FAS leading to production of inflammatory prostaglandins, succinate is involved in enzymatic activity controlling epigenetic modifications that might lead to a persistent inflammatory state. In a murine model of sepsis, LPS-induced succinate, in bone marrow-derived macrophages, stabilized Hypoxia-inducible factor alpha (HIF- α) interfering with the activity of the enzyme prolyl hydroxylase responsible for its degradation, and leading to an increase of IL-1 β , an important mediator in innate immune signaling during inflammation [34]. Studies are needed to elucidate the role of TCA cycle metabolites in neonatal sepsis.

Lipid metabolism

FAO and FAS have been thought to have opposing roles in immune system functions i.e., FAO is required for anti-inflammatory cytokine production and thus linked with immune tolerance, while FAS seems to positively regulate the generation and function of pro-inflammatory innate immune cells [35]. However, recent literature highlights the role of FAO in macrophage activation, as it also supports inflammasome activation, a molecular complex that processes the secretion of pro-inflammatory cytokines IL-1 β and IL-18 in innate immune cells [36]. Fatty acids serve as source of metabolic energy, and biologically active components of cell membranes [37]. A gene ontology analysis comparing the transcriptome of unstimulated human preterm neonatal, term neonatal and adult monocytes revealed markedly distinct metabolic profiles by age with an overall lower expression of genes related to oxidative phosphorylation and FAO metabolism in addition to lower glycolysis, in preterm monocytes [22]. Studies of host-pathogen interactions using whole blood transcriptomics in culture-positive neonatal sepsis reveal upregulation of genes related to lipid metabolism [20]. Peroxisome proliferator-activated receptor gamma (PPAR γ) is a nuclear receptor involved in lipid metabolism [38] and functionally linked to the inhibition of pro-inflammatory gene expression and to resolution of inflammation [39, 40]; Of note, expression of PPAR γ is

increased in neonatal monocytes [22]. Enhanced lipid metabolism results in increased production of anti-inflammatory cytokines (IL-10, TGF- β), as well as pro-resolving lipid mediators [41], potent anti-inflammatory mediators that regulate the active resolution of inflammation and promotion of wound healing following tissue injury; studies on the role of these mediators in the newborns are still pending.

Amino acid metabolism

Amino acids are substrates for protein synthesis intimately linked to important anabolic cellular signaling pathways, most notably the mTOR pathway. Amino acid metabolism is important in immune function and sepsis [42]. Glutamine metabolism regulates the balance between effector T cells and regulatory T cells (T_{regs}); glutamine deficiency can impair generation and function of Th1 and Th17, whereas T_{regs} generation is not affected [43]. A role for glutamine in the cytotoxic and antimicrobial functions of macrophages has been described for (a) the generation of nitric oxide [44] and (b) induction of IL-1 β by macrophages in response to LPS stimulation [45]. Little is known on the role of amino acid metabolism in the pathogenesis of neonatal sepsis, however certain amino acids, such as glutamine [46] and arginine [47] have been found to be depleted in neonates with sepsis compared to healthy newborns.

Overall, neonatal immunometabolism is apparently programmed to respond immune tolerance, which can result in immune paralysis if not closely regulated.

Metabolomics in neonatal sepsis: a tool to improve diagnosis?

Early diagnosis and intervention remains a challenge for neonatal sepsis [48]. The diagnosis still relies on the clinician's experience due to lack of specificity of clinical signs and symptoms and poor predictive ability of routine laboratory tests, including the "gold standard"

blood culture [48-50]. Combination of technology and biomarker discovery can provide further insights and identify target pathways to improve diagnosis and outcome prediction. Systems biology approaches using technologies that comprehensively measure the inventory of molecules in a given biosample have been introduced to analyze complex datasets of signaling and response pathways to numerous diseases and interventions [51].

Metabolomics comprehensively profiles, in a given biosample, small molecules that are downstream products of enzymatic processes. Metabolic profiling can employ either a) untargeted (global) approaches which measure the broadest range of metabolites in a sample, or b) targeted approaches which provide higher sensitivity and selectivity with known standards of a priori information, aiming to analyze and identify specific metabolites and metabolic pathways of interest [52]. At present, no analytical technology can adequately capture the entirety of a given metabolome. The current challenge for metabolomic studies is the lack of standardization of metabolomic data processing and analyses. One of the strengths of metabolomics is that data can be generated using a variety of biological matrices that are relevant for disease including plasma [53], serum [54-56], saliva [57], exhaled breath condensate [58, 59], urine [46] and fecal samples [60, 61]. Metabolomics has been successfully employed in newborns to identify alterations in plasma steroid and carbohydrate metabolites important for neurodevelopment, rapid cell proliferation and nutrient uptake as well as to characterize molecular ontogeny of plasma across the first week of human life [10].

Sepsis causes disruption in biochemical homeostasis and early detection of metabolite perturbations can predict severity [51]. A case-control study, involving both term and preterm septic human neonates and controls, recently employed urine metabolomics as a tool to identify biomarkers of late-onset neonatal sepsis using non-targeted Nuclear Magnetic Resonance (NMR) spectroscopy and targeted liquid chromatography-tandem mass spectrometry analysis. Septic neonates demonstrated distinct metabolic profiles from non-septic age-matched controls

such as increased amounts of glucose, pyruvate and lactate and lower levels of glutamine and vitamins of the B complex such as riboflavin (vitamin B2) and nicotinamide (vitamin B3) [46]. A similar study on 25 human neonates (9 with diagnosis of sepsis and 16 healthy controls) using NMR and gas-chromatography-mass spectrometry (GC-MS) found increased urine levels of glucose, lactate and acetate, but decreased pentose phosphate pathway metabolites in septic neonates compared to non-septic controls [62]. Metabolomics have also been applied to sepsis-induced acute lung injury, finding differences in distinct plasma metabolites (glutathione, adenosine, phosphatidylserine and sphingomyelin) between septic and healthy adult patients [63]. Changes in the levels of metabolites can also predict sepsis outcome and provide information on responsiveness to therapy. Mass spectrometry-based metabolomics strategies on the plasma metabolome of 21 adult septic shock patients suggest that lipidome alterations play an important role in an individual's response to infection [64]. In a plasma metabolomics and proteomics study involving 1000 adult sepsis patients, FAO was severely impaired in sepsis non-survivors, highlighting fatty acid metabolism as a promising metabolic predictor for survival in sepsis [65].

Thus far much remains to be learned regarding the neonatal metabolome at baseline, during early life ontogeny and during sepsis. Future immunometabolic studies in early life will aid in defining biomarkers of those at high risk of sepsis, and informing new approaches for prophylaxis, diagnosis and treatment of neonatal sepsis.

Possible interventions and new perspectives

Trained immunity and metabolism

Neonates are heavily reliant on their innate immune system for mounting an acute response to infection [6]. Innate immune cells (e.g. macrophages, monocytes, dendritic cells (DCs)) were recently discovered to attain *antigen non-specific* memory capabilities through epigenetic and metabolic reprogramming after subsequent heterologous stimulation [9]. This phenomenon termed as innate immune memory or trained immunity may be manifested as a heightened (training) or attenuated (tolerance) immune response, enabling the host to efficiently and effectively adapt to environmental threats while ensuring cytoprotection [7, 66]. Trained immunity can confer heterologous protection from a variety of subsequent pathogens [67] for a yet undetermined amount of time.

Several metabolites that accumulate in monocytes and macrophages seem to play a role in the induction of epigenetic modulators. The cholesterol synthesis pathway is essential to trained immunity, as statins, inhibitors of the cholesterol synthesis pathway, can prevent the induction of trained immunity [68]. Increased succinate and fumarate levels in activated innate immune cells antagonize histone demethylation, and suppress anti-inflammatory genes, thereby promoting a pro-inflammatory phenotype [34, 69, 70]. Acetyl-CoA, increased in trained monocytes, is an essential substrate for acetylating processes. NAD⁺ is important for epigenetic changes resulting in a switch from glucose to FAO during LPS-induced tolerance [71], which represents a model of sepsis-induced immune paralysis. Understanding trained immunity and targeting cellular metabolic processes, for example via nutritional intervention, may represent a novel approach to optimize immune responsiveness in vulnerable newborns [72].

Impact of microbiome on metabolism in sepsis

The human microbiome, consisting of all microorganisms living in or on the human body, shares a symbiotic relationship with the host and shapes the innate and adaptive immune systems via maintain gut-barrier functions and impacting host metabolism.

Although there is growing evidence that the early life microbiome has important effects on health and disease [73, 74], much remains to be learned regarding the impact of the microbiome in neonatal immune development and risk of neonatal sepsis. The neonatal microbiome undergoes dramatic changes during the first weeks of life influenced by diet, exposure to new microbes, antibiotics and other environmental exposures [75]. Premature or low birth weight neonates often receive antibiotics during hospitalization, shaping the gut microbiome. Antibiotic treatment is associated with reduced gut bacteria diversity and may kill beneficial bacteria allowing multidrug resistant pathogens to dominate the neonate's microbiome [76]. Antibiotics-suppressed gut microbiome affects the whole-body metabolism (including urinary and plasma metabolomes) as demonstrated in antibiotics-treated pigs [77]. Of note, evidence is emerging that microbial metabolites may impact host immune function [78]. Short chain fatty acids (SCFAs), products of bacterial fermentation, have been linked to immune tolerance via (a) inhibition of nuclear factor- κ B (NF- κ B); (b) reduced expression of T cell-activating molecules on antigen-presenting cells [79]; and (c) increased number and function of T_{regs}, including their production of anti-inflammatory cytokines (TGF β and IL-10) [80, 81]. The gene for Free fatty acid receptor 2 (FFAR2), a receptor for gut microbiota-derived SCFAs, was upregulated in human septic newborns [20], suggesting that the metabolic activity of neonatal microbiota influences the innate and adaptive immune response to sepsis and may contribute to an anti-inflammatory state. Disruption of the intestinal microbiome during critical illness, resulting in a "pathobiome", could result in immune suppression and consequent higher risk of sepsis, altering the course and outcome of infection [82, 83]. Further studies on the role of microbiome and related metabolites in neonatal sepsis are needed.

Nutritional support and the role of immunonutrients in preventing neonatal infection

Nutrition plays a key role in the development of multiple systems, including the immune system. In fact, undernutrition is the most common cause of secondary immunodeficiency in the world [84], with malnutrition in the critical early months of development having a profound and long-lasting impact on immunity. Nutrition, both enteral and parenteral, has an important role in the prevention of neonatal sepsis by minimizing nutrient loss and stimulating the maturation of the gastrointestinal tract, one of the largest immune organs of the body [85]. The use of human milk and early enteral nutrition are important for reducing the risk of infections. Indeed, the earlier an infant achieves full enteral nutrition, the lower the risk of late onset sepsis [86]. Human milk is rich in multiple immune-active components, which optimize the intestinal microbiological and metabolic milieu of newborns and protect them from inflammation and infection. Those include immunoglobulins, cytokines and growth factors, medium- and long-chain polyunsaturated free fatty acids, milk oligosaccharides, lactoferrin and lysozyme. The immunomodulatory role of these components is reviewed elsewhere [85]. Human milk provides the gold standard for feeding term infants, while for preterm infants it is routinely supplemented with multi-nutrient fortifiers to provide recommended levels of protein, sodium, chloride, magnesium and other micronutrients necessary to support the hypercatabolic state of prematurity. When exclusive human milk use is not possible, preterm formulas are the main alternative, though likely lacking the immunoprotective effects of human milk, based on observational studies. Harnessing the immunological benefits of maternal breast milk or its beneficial metabolic components might aid in prevention and treatment of neonatal sepsis, especially in vulnerable preterm infants.

Specific essential nutrients such as vitamins and trace minerals are also important to the immune response such as that their deficiency can increase susceptibility to infection. *Vitamin D* (1,25OHD) deficiency has been associated with higher risk of developing sepsis in both human neonatal [87], adult and murine [88] studies. Pretreatment of preterm (<32 weeks

gestational age) neonatal whole blood *in vitro* reversed endotoxin tolerance by enhancing reactive oxygen intermediates production from phagocytes [89]. *Iron* deficiency with or without anemia impairs cell-mediated immune response, intracellular killing of bacteria and secretory IgA responses, increasing susceptibility of infants to infection [90]. Deficits in trace elements are common in both enterally and parenterally-fed infants, with premature infants particularly susceptible to severe acute Zn deficiency states. Zn deficiency causes atrophy of lymphoid tissue, impaired delayed cutaneous hypersensitivity reaction, decreased lymphocyte response to antigens, and impaired chemotaxis of monocytes and polymorphonuclear cells [91]. Serum Zn concentrations of septic adults and pediatric ICU patients are reduced relative to the normal physiological range in healthy age-matched controls [92, 93]. Accordingly, several studies have examined whether Zn supplementation may be helpful in treating sepsis. In a mouse model of sepsis (intraperitoneal fecal-slurry injection with or without Zn supplementation), prophylactic Zn supplementation conferred a significant survival advantage by enhancing pathogen eradication, while simultaneously attenuating potentially detrimental excessive inflammation [94]. It is proposed that Zn redistributes from the blood compartment to the liver to meet the increased metabolic demands of sepsis [95]. Oral Zn supplementation reduced morbidity and mortality in hospitalized very low birth weight neonates in Italy [96], as also demonstrated in a recent meta-analysis [97]. *Selenium* has a role in protecting against oxidative damage and preterm neonates are at risk of selenium deficiency. In a randomized clinical trial conducted in India including 90 preterm infants with <32 weeks of gestation and/or BW \leq 1,500 g, prophylactic selenium supplementation led to significant reductions in late-onset sepsis episodes compared to placebo, though had no effect on overall mortality [98, 99].

Until such time as new recommendations regarding immunonutrient supplementation for at risk including preterm newborns can be made, proven strategies to minimize infection prevention in young infants, such as exclusive use of human milk, standardized enteral feeding

guidelines, avoidance of acid blockade and appropriate minimization of antibiotic exposure should be incorporated in care bundles of term and preterm newborns.

Metabolically active drugs

Infection is associated with tissue infiltration by phagocytic cells such as neutrophils and macrophages and their production of reactive species that contribute to microbicidal activity. An excess of reactive oxygen species (ROS) and reactive nitrogen species is associated with microvascular dysfunction and organ failure. Accordingly, efforts to develop metabolically active agents to prevent or treat sepsis in early life have included development of antioxidants.

Melatonin, an endogenously produced indolamine principally synthesized in the pineal gland, acts as an antioxidant directly by neutralizing ROS and nitrogen oxygen species, and indirectly by stimulating antioxidant enzyme activity, including that of superoxide dismutase, glutathione peroxidase, and glutathione reductase [100]. Supplementation of melatonin and its metabolites have shown effectiveness in human studies performed on several neonatal disorders characterized by massive inflammatory cascade and oxidative injury, including sepsis [88, 101].

Edaravone (3-methyl-1-phenyl-pyrazolin-5-one), a free radical scavenger, exerts multiple antioxidant effects, such as hydroxyl radical scavenging, suppression of hydroxyl-dependent lipid peroxidation, and electron donation to ROS [102]. In a piglet model of neonatal sepsis, edaravone reduced serum total hydroperoxide concentrations 1 hour after cecal ligation and perforation, and nitrite and nitrate levels at 3 and 6 hours in comparison to septic untreated animals, paralleling clinical improvement of septic animals [103]. In the same study, edaravone delayed TNF surge in septic animals and prevented the increase of high mobility group box 1 (HMGB-1), a nuclear transcription factor involved in the systemic inflammatory response.

These findings suggest possible beneficial effects of edaravone on sepsis clinical course in the newborns, but human trials are yet to be performed.

Pentoxifylline (PTX), a methylxanthine derivative with immunomodulating properties, has been used as an adjunct therapy for severe neonatal sepsis. PTX is a phosphodiesterase inhibitor that enhances intracellular cAMP concentrations, inhibits production of TLR-mediated inflammatory cytokines, including TNF- α , IL-6 and Interferon gamma (IFN γ), has beneficial effects on endothelial cell function, and improves microcirculation and tissue perfusion [104]. PTX decreased LPS-induced hyperinflammation in monocytes of preterm infants *in vitro* [105]. Although a recent Cochrane review demonstrated a significant reduction in all-cause mortality of hospitalized septic neonates with adjunct use of PTX [106], a subsequent randomized controlled trial of PTX for preterm infants with late onset sepsis did not demonstrate a significant decrease in neonatal mortality and morbidity [107].

Conclusions

Neonatal sepsis remains a major threat in early life, yet our tools to address it are limited. Growing evidence suggests that the distinct immunometabolism of early life contributes to susceptibility of infection and sepsis risk. On-going research is leveraging powerful new system biology tools to define immunometabolic cellular and molecular signatures and trajectories associated with health and disease [10]. Metabolic agents such as Zn, edaravone and PTX are currently being evaluated as prophylaxis or treatment for neonatal sepsis. Future studies on the ontogeny of early life immunometabolism in health and disease will provide new avenues to prevent, detect and treat neonatal sepsis.

REFERENCES

1. Singer, M., et al., The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, 2016. 315(8): p. 801-10.
2. Collaborators, G.B.D.C.o.D., Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 2018. 392(10159): p. 1736-1788.
3. Srinivasjois, R., et al., Association of Gestational Age at Birth with Reasons for Subsequent Hospitalisation: 18 Years of Follow-Up in a Western Australian Population Study. *PLOS ONE*, 2015. 10(6): p. e0130535.
4. Wynn, J.L., Defining neonatal sepsis. *Curr Opin Pediatr*, 2016. 28(2): p. 135-40.
5. Levy, O., Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nat Rev Immunol*, 2007. 7(5): p. 379-90.
6. Stiehm E R, N.T., Levy O Recognition of immunodeficiency in the first three months of life. . UpToDate Retrieved from <https://www.uptodate.com/>, 2018.
7. Netea, M.G., et al., Trained immunity: A program of innate immune memory in health and disease. *Science*, 2016. 352(6284): p. aaf1098.
8. O'Neill, L., A guide to immunometabolism for immunologists. *Nature Reviews Immunology*, 2016. Sep;16(9):553-65.
9. Arts, R.J.W., et al., Immunometabolic Pathways in BCG-Induced Trained Immunity. *Cell Rep*, 2016. 17(10): p. 2562-2571.

10. Lee, A.H., et al., Dynamic molecular changes during the first week of human life follow a robust developmental trajectory. *Nat Commun*, 2019. 10(1): p. 1092.
11. Evangelatos, N., et al., Metabolomics in Sepsis and Its Impact on Public Health. *Public Health Genomics*, 2017. 20(5): p. 274-285.
12. Angus, D.C. and T. van der Poll, Severe sepsis and septic shock. *N Engl J Med*, 2013. 369(9): p. 840-51.
13. Pearce, E.L. and E.J. Pearce, Metabolic pathways in immune cell activation and quiescence. *Immunity*, 2013. 38(4): p. 633-43.
14. Cheng, S.C., et al., Broad defects in the energy metabolism of leukocytes underlie immunoparalysis in sepsis. *Nat Immunol*, 2016. 17(4): p. 406-13.
15. Ghazal, P., P. Dickinson, and C.L. Smith, Early life response to infection. *Curr Opin Infect Dis*, 2013. 26(3): p. 213-8.
16. Wynn, J.L., et al., The influence of developmental age on the early transcriptomic response of children with septic shock. *Mol Med*, 2011. 17(11-12): p. 1146-56.
17. Kollmann, T.R., et al., Neonatal innate TLR-mediated responses are distinct from those of adults. *J Immunol*, 2009. 183(11): p. 7150-60.
18. Shen, C.M., et al., Development of monocyte Toll-like receptor 2 and Toll-like receptor 4 in preterm newborns during the first few months of life. *Pediatr Res*, 2013. 73(5): p. 685-91.
19. Marchant, E.A., et al., Attenuated innate immune defenses in very premature neonates during the neonatal period. *Pediatr Res*, 2015. 78(5): p. 492-7.
20. Smith, C.L., et al., Identification of a human neonatal immune-metabolic network associated with bacterial infection. *Nat Commun*, 2014. 5: p. 4649.

21. Harbeson, D., et al., Energy Demands of Early Life Drive a Disease Tolerant Phenotype and Dictate Outcome in Neonatal Bacterial Sepsis. *Front Immunol*, 2018. 9: p. 1918.
22. Kan, B., et al., Cellular metabolism constrains innate immune responses in early human ontogeny. *Nat Commun*, 2018. 9(1): p. 4822.
23. Bours, M.J., et al., Adenosine 5'-triphosphate and adenosine as endogenous signaling molecules in immunity and inflammation. *Pharmacol Ther*, 2006. **112**(2): p. 358-404.
24. Idzko, M., D. Ferrari, and H.K. Eltzschig, Nucleotide signalling during inflammation. *Nature*, 2014. 509(7500): p. 310-7.
25. Yegutkin, G.G., Nucleotide- and nucleoside-converting ectoenzymes: Important modulators of purinergic signalling cascade. *Biochim Biophys Acta*, 2008. 1783(5): p. 673-94.
26. Ledderose, C., et al., Purinergic Signaling and the Immune Response in Sepsis: A Review. *Clin Ther*, 2016. 38(5): p. 1054-65.
27. Pettengill, M., et al., Soluble ecto-5'-nucleotidase (5'-NT), alkaline phosphatase, and adenosine deaminase (ADA1) activities in neonatal blood favor elevated extracellular adenosine. *J Biol Chem*, 2013. 288(38): p. 27315-26.
28. Levy, O., et al., The adenosine system selectively inhibits TLR-mediated TNF- α production in the human newborn. *J Immunol*, 2006. 177(3): p. 1956-66.
29. Dreschers, S., et al., Impaired cellular energy metabolism in cord blood macrophages contributes to abortive response toward inflammatory threats. *Nat Commun*, 2019. 10(1): p. 1685.

30. Weichhart, T., M. Hengstschlager, and M. Linke, Regulation of innate immune cell function by mTOR. *Nat Rev Immunol*, 2015. **15**(10): p. 599-614.
31. Everts, B., et al., TLR-driven early glycolytic reprogramming via the kinases TBK1-IKK ϵ supports the anabolic demands of dendritic cell activation. *Nat Immunol*, 2014. **15**(4): p. 323-32.
32. Ulas, T., et al., S100-alarmin-induced innate immune programming protects newborn infants from sepsis. *Nat Immunol*, 2017. **18**(6): p. 622-632.
33. Jha, A.K., et al., Network integration of parallel metabolic and transcriptional data reveals metabolic modules that regulate macrophage polarization. *Immunity*, 2015. **42**(3): p. 419-30.
34. Tannahill, G.M., et al., Succinate is an inflammatory signal that induces IL-1 β through HIF-1 α . *Nature*, 2013. **496**(7444): p. 238-42.
35. Feingold, K.R., et al., Mechanisms of triglyceride accumulation in activated macrophages. *J Leukoc Biol*, 2012. **92**(4): p. 829-39.
36. Moon, J.-S., et al., NOX4-dependent fatty acid oxidation promotes NLRP3 inflammasome activation in macrophages. *Nature Medicine*, 2016. **22**: p. 1002.
37. Innis, S.M., Essential fatty acids in growth and development. *Prog Lipid Res*, 1991. **30**(1): p. 39-103.
38. Varga, T., Z. Czimmerer, and L. Nagy, PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochim Biophys Acta*, 2011. **1812**(8): p. 1007-22.
39. Bouhrel, M.A., et al., PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab*, 2007. **6**(2): p. 137-43.

40. Ricote, M., et al., The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature*, 1998. 391(6662): p. 79-82.
41. Serhan, C.N., Pro-resolving lipid mediators are leads for resolution physiology. *Nature*, 2014. 510(7503): p. 92-101.
42. Kelly, D. and P.E. Wischmeyer, Role of L-glutamine in critical illness: new insights. *Curr Opin Clin Nutr Metab Care*, 2003. 6(2): p. 217-22.
43. Nakaya, M., et al., Inflammatory T cell responses rely on amino acid transporter ASCT2 facilitation of glutamine uptake and mTORC1 kinase activation. *Immunity*, 2014. 40(5): p. 692-705.
44. Murphy, C. and P. Newsholme, Importance of glutamine metabolism in murine macrophages and human monocytes to L-arginine biosynthesis and rates of nitrite or urea production. *Clin Sci (Lond)*, 1998. 95(4): p. 397-407.
45. Wallace, C. and D. Keast, Glutamine and macrophage function. *Metabolism*, 1992. 41(9): p. 1016-20.
46. Sarafidis, K., et al., Urine metabolomics in neonates with late-onset sepsis in a case-control study. *Sci Rep*, 2017. 7: p. 45506.
47. Badurdeen, S., M. Mulongo, and J.A. Berkley, Arginine depletion increases susceptibility to serious infections in preterm newborns. *Pediatric Research*, 2014. 77: p. 290.
48. Gerdes, J.S., Diagnosis and management of bacterial infections in the neonate. *Pediatr Clin North Am*, 2004. 51(4): p. 939-59, viii-ix.
49. Benitz, W.E., Adjunct laboratory tests in the diagnosis of early-onset neonatal sepsis. *Clin Perinatol*, 2010. 37(2): p. 421-38.

50. Hornik, C.P., et al., Early and late onset sepsis in very-low-birth-weight infants from a large group of neonatal intensive care units. *Early Hum Dev*, 2012. 88 Suppl 2: p. S69-74.
51. Skibsted, S., et al., Bench-to-bedside review: future novel diagnostics for sepsis - a systems biology approach. *Crit Care*, 2013. 17(5): p. 231.
52. Johnson, C.H., J. Ivanisevic, and G. Siuzdak, Metabolomics: beyond biomarkers and towards mechanisms. *Nat Rev Mol Cell Biol*, 2016. 17(7): p. 451-9.
53. Vuckovic, D. and J. Pawliszyn, Systematic evaluation of solid-phase microextraction coatings for untargeted metabolomic profiling of biological fluids by liquid chromatography-mass spectrometry. *Anal Chem*, 2011. 83(6): p. 1944-54.
54. OuYang, D., et al., Metabolomic profiling of serum from human pancreatic cancer patients using ¹H NMR spectroscopy and principal component analysis. *Appl Biochem Biotechnol*, 2011. 165(1): p. 148-54.
55. Xuan, J., et al., Metabolomic profiling to identify potential serum biomarkers for schizophrenia and risperidone action. *J Proteome Res*, 2011. 10(12): p. 5433-43.
56. Hasokawa, M., et al., Identification of biomarkers of stent restenosis with serum metabolomic profiling using gas chromatography/mass spectrometry. *Circ J*, 2012. 76(8): p. 1864-73.
57. Alvarez-Sanchez, B., F. Priego-Capote, and M.D. Luque de Castro, Study of sample preparation for metabolomic profiling of human saliva by liquid chromatography-time of flight/mass spectrometry. *J Chromatogr A*, 2012. 1248: p. 178-81.

58. Montuschi, P., et al., NMR spectroscopy metabolomic profiling of exhaled breath condensate in patients with stable and unstable cystic fibrosis. *Thorax*, 2012. 67(3): p. 222-8.
59. Carraro, S., et al., Metabolomics applied to exhaled breath condensate in childhood asthma. *Am J Respir Crit Care Med*, 2007. 175(10): p. 986-90.
60. Chow, J., et al., Fecal metabolomics of healthy breast-fed versus formula-fed infants before and during in vitro batch culture fermentation. *J Proteome Res*, 2014. 13(5): p. 2534-42.
61. Goedert, J.J., et al., Fecal metabolomics: assay performance and association with colorectal cancer. *Carcinogenesis*, 2014. 35(9): p. 2089-96.
62. Fanos, V., et al., Urinary (1)H-NMR and GC-MS metabolomics predicts early and late onset neonatal sepsis. *Early Hum Dev*, 2014. 90 Suppl 1: p. S78-83.
63. Stringer, K.A., et al., Metabolic consequences of sepsis-induced acute lung injury revealed by plasma (1)H-nuclear magnetic resonance quantitative metabolomics and computational analysis. *Am J Physiol Lung Cell Mol Physiol*, 2011. 300(1): p. L4-L11.
64. Cambiaghi, A., et al., Characterization of a metabolomic profile associated with responsiveness to therapy in the acute phase of septic shock. *Sci Rep*, 2017. 7(1): p. 9748.
65. Langley, R.J., et al., An integrated clinico-metabolomic model improves prediction of death in sepsis. *Sci Transl Med*, 2013. 5(195): p. 195ra95.
66. Netea, M.G. and J.W. van der Meer, Trained Immunity: An Ancient Way of Remembering. *Cell Host Microbe*, 2017. 21(3): p. 297-300.

67. Saeed, S., et al., Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science*, 2014. 345(6204): p. 1251086.
68. Bekkering, S., et al., Metabolic Induction of Trained Immunity through the Mevalonate Pathway. *Cell*, 2018. 172(1-2): p. 135-146 e9.
69. Benit, P., et al., Unsuspected task for an old team: succinate, fumarate and other Krebs cycle acids in metabolic remodeling. *Biochim Biophys Acta*, 2014. 1837(8): p. 1330-7.
70. Arts, R.J., et al., Glutaminolysis and Fumarate Accumulation Integrate Immunometabolic and Epigenetic Programs in Trained Immunity. *Cell Metab*, 2016. 24(6): p. 807-819.
71. Liu, T.F., et al., NAD⁺-dependent sirtuin 1 and 6 proteins coordinate a switch from glucose to fatty acid oxidation during the acute inflammatory response. *J Biol Chem*, 2012. 287(31): p. 25758-69.
72. Levy, O. and J.L. Wynn, A prime time for trained immunity: innate immune memory in newborns and infants. *Neonatology*, 2014. 105(2): p. 136-41.
73. Cernada, M., et al., Sepsis in preterm infants causes alterations in mucosal gene expression and microbiota profiles compared to non-septic twins. *Scientific Reports*, 2016. 6(1): p. 25497.
74. Stewart, C., et al., The preterm gut microbiota: changes associated with necrotizing enterocolitis and infection. *Acta Paediatrica*, 2012. 101(11): p. 1121-1127.
75. Gritz, E.C. and V. Bhandari, The human neonatal gut microbiome: a brief review. *Front Pediatr*, 2015. 3: p. 17.

76. Gibson, M.K., et al., Developmental dynamics of the preterm infant gut microbiota and antibiotic resistome. *Nature Microbiology*, 2016. 1(4): p. 16024.
77. Jiang, P., et al., Antibiotic Treatment Preventing Necrotising Enterocolitis Alters Urinary and Plasma Metabolomes in Preterm Pigs. *Journal of Proteome Research*, 2017. 16(10): p. 3547-3557.
78. Rooks, M.G. and W.S. Garrett, Gut microbiota, metabolites and host immunity. *Nat Rev Immunol*, 2016. 16(6): p. 341-52.
79. Vinolo, M.A., et al., Suppressive effect of short-chain fatty acids on production of proinflammatory mediators by neutrophils. *J Nutr Biochem*, 2011. 22(9): p. 849-55.
80. Arpaia, N., et al., Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature*, 2013. 504(7480): p. 451-5.
81. Smith, P.M., et al., The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. *Science*, 2013. 341(6145): p. 569-73.
82. Hotchkiss, R.S., G. Monneret, and D. Payen, Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol*, 2013. 13(12): p. 862-74.
83. Haak, B.W. and W.J. Wiersinga, The role of the gut microbiota in sepsis. *Lancet Gastroenterol Hepatol*, 2017. 2(2): p. 135-143.
84. Black, R.E., et al., Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*, 2008. 371(9608): p. 243-60.
85. Walsh, V. and W. McGuire, Immunonutrition for Preterm Infants. *Neonatology*, 2019. 115(4): p. 398-405.

86. Freitas, B.A., et al., Nutritional therapy and neonatal sepsis. *Rev Bras Ter Intensiva*, 2011. 23(4): p. 492-8.
87. Dhandai, R., et al., Association of vitamin D deficiency with an increased risk of late-onset neonatal sepsis. *Paediatrics and International Child Health*, 2018. 38(3): p. 193-197.
88. Parekh, D., et al., Vitamin D Deficiency in Human and Murine Sepsis*. *Critical Care Medicine*, 2017. 45(2): p. 282-289.
89. Onwuneme, C., et al., Vitamin D enhances reactive oxygen intermediates production in phagocytic cells in term and preterm infants. *Pediatr Res*, 2016. 79(4): p. 654-61.
90. Schlesinger, L. and R. Uauy, Nutrition and neonatal immune function. *Semin Perinatol*, 1991. 15(6): p. 469-77.
91. Loui, A., et al., Nutritional zinc balance in extremely low-birth-weight infants. *J Pediatr Gastroenterol Nutr*, 2001. 32(4): p. 438-42.
92. Besecker, B.Y., et al., A comparison of zinc metabolism, inflammation, and disease severity in critically ill infected and noninfected adults early after intensive care unit admission. *Am J Clin Nutr*, 2011. 93(6): p. 1356-64.
93. Cvijanovich, N.Z., et al., Zinc homeostasis in pediatric critical illness. *Pediatr Crit Care Med*, 2009. 10(1): p. 29-34.
94. Nowak, J.E., et al., Prophylactic zinc supplementation reduces bacterial load and improves survival in a murine model of sepsis. *Pediatr Crit Care Med*, 2012. 13(5): p. e323-9.
95. Alker, W. and H. Haase, Zinc and Sepsis. *Nutrients*, 2018. 10(8).
96. Terrin, G., et al., Zinc supplementation reduces morbidity and mortality in very-low-birth-weight preterm neonates: a hospital-based randomized,

- placebo-controlled trial in an industrialized country. *Am J Clin Nutr*, 2013. 98(6): p. 1468-74.
97. Tang, Z., et al., Efficacy of zinc supplementation for neonatal sepsis: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med*, 2017: p. 1-6.
98. Darlow, B.A. and N.C. Austin, Selenium supplementation to prevent short-term morbidity in preterm neonates. *Cochrane Database Syst Rev*, 2003(4): p. CD003312.
99. Aggarwal, R., et al., Selenium Supplementation for Prevention of Late-Onset Sepsis in Very Low Birth Weight Preterm Neonates. *J Trop Pediatr*, 2016. 62(3): p. 185-93.
100. Gitto, E., et al., Effects of melatonin treatment in septic newborns. *Pediatr Res*, 2001. 50(6): p. 756-60.
101. Gitto, E., et al., Protective role of melatonin in neonatal diseases. *Oxid Med Cell Longev*, 2013. **2013**: p. 980374.
102. Nakamoto, N., et al., A free radical scavenger, edaravone, attenuates steatosis and cell death via reducing inflammatory cytokine production in rat acute liver injury. *Free Radic Res*, 2003. 37(8): p. 849-59.
103. Kato, S., et al., Edaravone, a novel free radical scavenger, reduces high-mobility group box 1 and prolongs survival in a neonatal sepsis model. *Shock*, 2009. 32(6): p. 586-92.
104. Speer, E.M., et al., Pentoxifylline Alone or in Combination with Gentamicin or Vancomycin Inhibits Live Microbe-Induced Proinflammatory Cytokine Production in Human Cord Blood and Cord Blood Monocytes In Vitro. *Antimicrob Agents Chemother*, 2018. 62(12).

105. Schuller, S.S., et al., Pentoxifylline modulates LPS-induced hyperinflammation in monocytes of preterm infants in vitro. *Pediatr Res*, 2017. 82(2): p. 215-225.
106. Pammi, M. and K.N. Haque, Pentoxifylline for treatment of sepsis and necrotizing enterocolitis in neonates. *Cochrane Database Syst Rev*, 2015(3): p. CD004205.
107. Shabaan, A.E., et al., Pentoxifylline therapy for late-onset sepsis in preterm infants: a randomized controlled trial. *Pediatr Infect Dis J*, 2015. 34(6): p. e143-8.

Acknowledgments

We thank Precision Vaccines Program graphic artist Kristin Johnson for optimizing the figure, and Precision Vaccines Program Coordinator, Ms. Diana Vo, for important administrative support.

FIGURE LEGEND

Figure 1. Overview of key immunometabolic pathways in quiescence and sepsis.

Metabolic pathways up-regulated and down-regulated during quiescence, acute-inflammation and tolerance/paralysis in adult monocytes. Distinct metabolic pathways in neonatal monocytes and macrophages that demonstrate down-regulation of glycolysis, oxidative phosphorylation and fatty acid oxidation. During sepsis, neonatal monocytes reveal upregulation of genes related to lipid metabolism but defective glycolysis.