

1 **Cost-efficient sampling methodologies for lake littoral invertebrates in**
2 **compliance with the European Water Framework Directive**

3 Gwendolin Porst^{a,*}, Oliver Miler^a, Louise Donohue^b, Tamara Jurca^b, Francesca Pilotto^{a,c},
4 Mario Brauns^{a,d}, Angelo Solimini^c, Martin Pusch^a

5 ^aLeibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Department of Ecosystem Research,
6 Müggelseedamm 301, 12587 Berlin, Germany

7 ^bSchool of Natural Sciences, Zoology Department, Freshwater Research Group, Trinity College Dublin, Dublin
8 2, Ireland

9 ^cDepartment of Public Health and Infectious Diseases, Sapienza University of Rome, Piazzale Aldo Moro 5,
10 00185 Roma, Italy

11 ^dDepartment River Ecology, Helmholtz Centre for Environmental Research GmbH – UFZ, Brückstr. 3a, 39114
12 Magdeburg, Germany

13 *corresponding author: porst@igb-berlin.de, phone: +49 (0)30 64 181 681, fax: +49 (0)30 64 181 682

14

15

16 **Acknowledgements**

17 This paper is a result of the project WISER (Water bodies in Europe: Integrative Systems to
18 assess Ecological status and Recovery) funded by the European Union under the 7th
19 Framework Programme, Theme 6 (Environment including Climate Change) (contract No.
20 226273), <http://www.wiser.eu>. We thank Daniel Graeber for help with the statistical analysis.

21

22 **Abstract**

23 Lake shores are characterized by a high natural variability, which is increasingly threatened
24 by a multitude of anthropogenic disturbances including morphological alterations to the
25 littoral zone. The European Water Framework Directive (EU WFD) calls for the assessment
26 of lake ecological status by monitoring biological quality elements (BQEs) including benthic
27 macroinvertebrates. To identify cost- and time-efficient sampling strategies for routine lake
28 monitoring, we conducted sampling of littoral invertebrates in 32 lakes located across a
29 European gradient. We compared the efficiency of two sampling methodologies, defined as
30 habitat-specific and pooled composite sampling protocols. Benthic samples were collected
31 from unmodified and morphologically altered shorelines. Variability within macroinvertebrate
32 communities did not differ significantly between sampling protocols across alteration types,
33 lake types and geographical regions. In addition, field composite samples and artificially
34 computed composite samples did not show significant differences in their macroinvertebrate
35 communities, and performed equally well in the calculation of various macroinvertebrate
36 metrics, and in their correlation to a predefined morphological stressor index. We conclude
37 that a benthic invertebrate sampling protocol involving proportional composite sampling
38 represents a time- and cost-efficient method for routine lake monitoring as requested under
39 the EU WFD, and may be applied across various European geographical regions.

40

41 **Key word:** morphological alteration; macroinvertebrates; lake monitoring; method
42 comparison; littoral zone; EU Water Framework Directive.

43

44 **Introduction**

45 The constant increase of anthropogenic disturbances to freshwater ecosystems is threatening
46 their ecological integrity strongly (Carpenter et al., 2007; Strayer & Findlay, 2010; Solimini
47 & Sandin, 2012). While eutrophication and acidification continue to be major threats to
48 European lakes, human modifications of lakeshore zones have only recently been
49 acknowledged as an increasing pressure on their ecological status (Brauns et al., 2007b;
50 Strayer & Findlay, 2010). Lake shores offer habitat for numerous species, dispersal corridors
51 for aquatic fauna and flora, and a variety of ecosystem services such as opportunities for
52 recreation, flood prevention, dissipation of wave energy and preservation of water quality
53 (O'Connor, 1991; Taniguchi et al., 2003; Gabel et al., 2012). Morphological degradation of
54 lakeshores caused *inter alia* by human settlement or industrial development is not only
55 associated with considerable losses in habitat and physical complexity in the lake littoral
56 (Solimini et al., 2006), but also in the above mentioned ecosystem services. Severe effects on
57 lake biotic communities have been demonstrated in detail for littoral fish assemblages
58 (Jennings et al., 1999; Scheuerell & Schindler, 2004) and recently also for benthic
59 invertebrate communities (Brauns et al., 2007b; Porst et al., 2012, in press; Solimini &
60 Sandin, 2012).

61 Littoral benthic invertebrates are a major component of lake ecosystems and their functioning
62 (Wetzel, 2001; Vadeboncoeur et al., 2002) and can be found in their highest diversity in the
63 eulittoral zone which is characterized by its high physical complexity and habitat diversity
64 (Taniguchi et al., 2003; Strayer & Findlay, 2010). This natural habitat diversity offers
65 macroinvertebrates a great variety of ecological niches, protection from foraging predators
66 and refuge from physical disturbance such as wind- or ship-induced waves (O'Connor, 1991;
67 Schneider & Winemiller, 2008; Brauns et al., 2011; Gabel et al., 2012). However, shoreline
68 development is typically accompanied by the loss of important littoral habitats such as

69 emergent or submerged macrophytes, submerged tree roots or coarse woody debris caused by
70 clear cutting of littoral and riparian zones of lakes. Consequently, increasing intensities of
71 shoreline development strongly affect littoral macroinvertebrate communities by reducing
72 littoral invertebrate biodiversity and altering macroinvertebrate community structures at
73 highly modified shorelines (Bänziger, 1995; Brauns et al., 2007b; Porst et al., 2012; McGoff
74 et al., 2013; Pilotto et al., in press)

75 The European Water Framework Directive (EU WFD) (EC, 2000) has acknowledged the
76 influence of increasing morphological alterations on the composition and abundance of biotic
77 communities of European freshwaters. To be in compliance with the requirements of the EU
78 WFD, ecological assessment methods need to be based on biological quality elements (BQEs)
79 including phytoplankton, macrophytes, fish, phytobenthos and benthic invertebrates (EC,
80 2000). The development of assessment tools for the monitoring of ecological integrity of
81 European lakes has so far focused mainly on quantifying the impacts of eutrophication on
82 biotic communities based on phytoplankton(Phillips. et al., 2011; Søndergaard et al., 2011;
83 Mischke et al., 2012), sublittoral and profundal invertebrate abundances and composition
84 (Saether, 1979; Brodersen & Lindegaard, 1999; Langdon et al., 2006). Impacts of
85 anthropogenic shoreline alterations on lake ecological status yet need to be quantified and
86 adequate monitoring programmes developed (EC, 2000). With life-cycles spanning between
87 several months and years and often sedentary aquatic life stages, benthic macroinvertebrate
88 assemblages potentially reflect changes to their physical, chemical and ecological
89 environment over time (Reice & Wohlenberg, 1993; Pinel-Alloul et al., 1996). Benthic
90 macroinvertebrates generally exhibit a strong dependence on the lake littoral and its diversity
91 and will consequently respond to habitat loss (Jurca et al., 2012; Porst et al., 2012; Solimini &
92 Sandin, 2012; Timm & Möls, 2012). Thus, littoral invertebrates can be expected to form a
93 suitable indicator group for the assessment of morphological pressures to lake ecological

94 status as part of routine monitoring programmes (Porst et al., 2012; Solimini & Sandin, 2012;
95 Urbanič et al., 2012).

96 While it has been argued that the high natural variability of littoral habitats and associated
97 macroinvertebrate communities make this organism group unsuitable for assessment purposes
98 (Rasmussen, 1988; Harrison & Hildrew, 1998; Moss et al., 2003), habitat stratification has
99 been identified to overcome the problem of inherent variability of the littoral zone of lakes
100 (Tolonen et al., 2001; Weatherhead & James, 2001; Tolonen & Hämäläinen, 2010). For
101 standardised routine monitoring of lakes, time and cost efficiency are important components
102 which can decide on a monitoring program's feasibility. Assessment methods based on littoral
103 macroinvertebrates typically involve time- and cost-intensive processing and identification of
104 macroinvertebrates in the laboratory, while a comparatively small amount of time and
105 associated expenses have to be spent for collection of samples in the field (Ferraro et al.,
106 1989; Haase et al., 2004; Tolonen & Hämäläinen, 2010; Porst et al., 2012). Habitat-specific
107 sampling regimes, frequently applied for lake monitoring in the past, however, generate
108 considerably higher numbers of macroinvertebrate samples compared to a 'pooled' multi-
109 habitat sampling approach. Consequently, habitat-specific sampling involves a much greater
110 working effort and, thus, potentially accounts for higher associated expenses when compared
111 to a multi-habitat sampling programme. While the stratified sampling regime might improve
112 signal precision by reducing variability within macroinvertebrate samples, the collection of
113 pooled composite macroinvertebrate samples could, thus, offer an alternative time- and cost-
114 effective sampling strategy for routine lake monitoring. So far only a limited number of
115 studies focusing on only a few large oligotrophic and mesotrophic lakes in the Central Baltic
116 region (Schreiber & Brauns, 2010; Porst et al., 2012) and one Mediterranean riverine lake
117 (Mastrantuono et al., in press) have compared the efficiency of habitat-specific and composite
118 sampling techniques for routine assessment of lakes. The suitability of the latter method for

119 routine monitoring purposes has, however, not yet been quantified across a gradient of
120 European lake types.

121 This study aimed at identifying the most suitable sampling methodology for routine
122 monitoring of lake ecological status based on benthic macroinvertebrates in compliance with
123 the requirements of the EU WFD. Based on results from a previous pilot study (Porst et al.,
124 2012) we compared macroinvertebrate samples collected from morphologically altered and
125 unmodified shorelines from a total of 32 lakes located in 3 European countries, with varying
126 trophic status. We tested the adequacy of composite against habitat-specific macroinvertebrate
127 sampling for routine lake monitoring by comparing macroinvertebrate diversity and
128 community structures of unmodified with soft (recreational beaches, grassland) and hard
129 (retaining walls, ripraps) altered shorelines across a trophic and European gradient. Composite
130 sampling comprised pooled proportional sampling of available habitats at a site, while for
131 habitat-specific sampling samples collected from different habitats were kept separate. We
132 hypothesised that pooled composite macroinvertebrate samples would represent a littoral
133 sampling site equally well compared to stratified habitat-specific samples independent of
134 morphological status of a sampling site and are, thus, suitable for routine monitoring of
135 ecological status of European lakes.

136 **Methods**

137 *Invertebrate sampling*

138 Benthic invertebrate samples were collected from 32 lakes in three European
139 countries/geographical regions representing a north-south gradient (Map/Figure 1). In Ireland
140 (North-Western Europe - climate: temperate maritime; topography: lowlands) benthic
141 macroinvertebrates were sampled from 9 lakes in April/May 2009, in Germany (Central
142 Europe - climate: temperate continental; topography: north-eastern lowlands) from 8 lakes in

143 May/April 2010 and in Italy (Southern Europe - Northern Italy: climate: temperate sub-
144 continental; topography: subalpine; Southern Italy: climate: mediterranean; topography:
145 volcanic) from 15 lakes in August-November 2009, with lakes comprising a gradient of total
146 phosphorus (TP range Ireland/North-Western Europe: 8.8 – 80.7 µg/L; TP range
147 Germany/Central Europe: 26.3 – 162.6 µg/L; TP range Italy/Southern Europe: 8 – 130 µg/L).
148 Benthic macroinvertebrate samples were collected from three morphologically differing
149 shoreline types, which were *a priori* classified as ‘soft alteration’ (recreational beaches or
150 riparian clear-cutting/grassland), ‘hard alteration’ (retaining walls and ripraps) and
151 unmodified shorelines. In each study lake three unmodified shoreline sites, three sites with
152 soft alterations and three sites with hard alterations were sampled for benthic
153 macroinvertebrates. Sampling sites comprised a shoreline section of minimum 25 m length
154 and extended to the maximum wadable water depth, generally < 1.2 m. At each sampling site,
155 three habitat-specific samples, ideally from sand, stones and macrophytes plus one composite
156 sample were collected. In cases where not all three habitats were present at a sampling site, a
157 second sample of the dominant habitat at this site was collected. In cases where only one
158 habitat was present, i.e. only sand habitats at recreational beaches, three samples from the
159 same habitat were collected. For habitat-specific samples, macroinvertebrates were collected
160 from an area of 1 m² for each habitat. Composite sampling comprised the collection of
161 macroinvertebrates from different habitats proportional to habitat availability within each
162 sampling site, generally following the method of the AQEM consortium (AQEM Consortium,
163 2002; STAR Consortium, 2003). Sampling of single habitats for habitat-specific and
164 composite sampling generally followed the methods described in Brauns et al. (2007b). In
165 short, samples from stones were collected by brushing off attached macroinvertebrates, while
166 macrophyte and sand habitats were sampled using a hand net (500 µm mesh size). While
167 single habitat samples were kept separate for habitat-specific sampling, macroinvertebrate
168 samples from different habitats were subsequently pooled for the composite sampling

169 approach. All macroinvertebrate samples were preserved in ethanol in the field and processed
170 in the laboratory. Macroinvertebrates were identified to species level, whenever possible,
171 except Chironomidae (subfamily), other Diptera (family), and Oligochaeta (class).

172 *Statistical analysis*

173 Based on findings by McGoff et al. (2013) and Miler et al. (2013), which identified
174 macroinvertebrate communities to differ significantly among geographical regions,
175 macroinvertebrate data were divided into geographical regions for statistical analysis.
176 Initially, we tested whether the habitat configuration at the sampling sites systematically differ
177 with alteration type or ecoregion. Therefore, we conducted a permutational analysis of
178 variance (ANOVA) with number of habitats and proportional availability of habitats as the
179 dependent and alteration type and ecoregion as the independent variables. Permutational
180 ANOVA has the advantage over its classical counterpart that normality and homoscedasticity
181 are not required (Gotelli & Ellison, 2004). The level of significance was calculated with
182 10.000 permutations and the analysis was conducted using the R software (R Core Team,
183 2013). Non-metric multidimensional scaling (NMDS) was used to display similarities in
184 macroinvertebrate community structures between habitat-specific and composite
185 macroinvertebrate samples within different alteration types in each country (PRIMER®
186 version 6, PRIMER-E Ltd, Ivybridge) (Clarke & Warwick, 2001). A two-way nested analysis
187 of similarities with factors 'lake' and 'habitat' (ANOSIM, PRIMER® version 6, PRIMER-E
188 Ltd, Ivybridge) tested for significant differences in macroinvertebrate community structures
189 among habitat and composite samples within alteration types in each country using 9999
190 permutations.

191 To test whether variability of macroinvertebrate community structures within composite
192 samples was significantly different from variability within habitat-specific samples within
193 different alteration types in each ecoregion/country, the homogeneity of dispersion of

194 individual habitats sampled was tested using permutational analysis of multidimensional
195 dispersion with 9999 permutations (PERMDISP, PRIMER® version 6 with PERMANOVA+,
196 PRIMERE Ltd, Ivybridge) (Anderson et al., 2008). Owing to a low number of replicate
197 samples ($n < 3$) the habitat-specific samples from stones at unmodified shoreline sites and from
198 macrophytes at soft alteration sampling sites in Germany could not be included in the
199 ANOSIM or PERMDISP analyses. PERMDISP, furthermore, tested the adequacy of
200 composite samples for monitoring of lake ecological status by comparing the composite
201 samples collected in the field with artificially computed composite samples again within
202 different alteration types and ecoregion/country. To assess the necessity of proportional
203 sampling for the adequate representation of macroinvertebrate communities at a site, artificial
204 composite samples were generated by accumulating single habitat samples once according to
205 their proportional availability at respective sampling sites (proportional artificial composite
206 sample) and again assigning equal weight to each single habitat sample collected at a site
207 (unproportional artificial composite sample). ANOSIM and PERMDISP subsequently tested
208 for differences in macroinvertebrate communities and associated homogeneities of dispersion
209 among collected and proportional and unproportional artificially generated composite samples
210 across different alteration types in each geographical region. NMDS ordinations, ANOSIM
211 and PERMDISP analyses were based on a Bray-Curtis similarity matrix of arcsine-
212 transformed proportional abundance data to account for differences in sampling
213 methodologies.

214 Macroinvertebrate communities can be described for assessment purposes based on ‘metrics’.
215 These are defined as summary measures of parts or processes of a biological system that
216 should change in value along a gradient of anthropogenic impact, i.e. in this case
217 morphological alteration. To test the efficiency of the composite sampling approach for lake
218 assessment based on multimetric indices, 10 invertebrate metrics commonly used for

219 morphological assessment purposes in lakes (Gabriels et al., 2010; Timm & Möls, 2012;
220 Miler et al., 2013) were calculated exemplarily based on macroinvertebrate abundances from
221 proportional and unproportional artificial composite and field composite samples (Table 1).
222 The calculated metrics were subsequently correlated separately with a predefined
223 morphological stressor index using Spearman-Rank correlations. The morphological stressor
224 index was calculated as a mean of variables calculated from Lake Habitat Survey (LHS)
225 parameters (Rowan et al., 2006; Rowan et al., 2008). The stressor index contained the
226 variables ‘Number of habitats’/‘Habitat diversity’, ‘Total PVI’/‘Sum of macrophyte types’,
227 ‘Sum of vegetation cover types’, ‘Sum of Coarse Woody Debris/roots/overhanging
228 vegetation’ (CWD), ‘Pressure index’ and ‘Natural/Artificial dominant land cover type’ and its
229 composition differed between the three geographical regions Germany, Ireland and Italy
230 (Table 2). The development and structure of the morphological stressor index is described in
231 more detail in Miler et al. (2014). Ranges of Spearman-Rank correlation coefficients
232 computed for field composite, proportional and unproportional artificial composite samples
233 were compared using a paired t-test. All metrics were calculated by means of the software
234 program ASTERICS 3.1.1. (www.fliessgewaesserbewertung.de/en) and Spearman-Rank
235 correlations and paired t-tests performed with SAS 9.2 (SAS Institute Inc., Cary, NC, USA.).

236 **Results**

237 *Habitat availability*

238 Habitat diversity as well as proportional availability of habitats varied significantly among
239 alteration types (Permutational ANOVA: $F = 9.97, p < 0.001$; $F = 10.33, p < 0.001$) but not
240 among geographical regions (Permutational ANOVA: $F = 0.96, P > 0.05$; $F = 2.48, p > 0.05$).
241 Similarly, there were no significant interactions between alteration type and ecoregion for
242 habitat diversity ($F = 1.52, p > 0.05$) and proportional habitat availability ($F = 1.25, p > 0.05$).

243 In Germany most dominant habitats found at unmodified sampling sites were sand (n=38;
244 median proportional availability/site 63%, range 14-94%) and macrophytes (n=32; median
245 proportional availability/site 40%, range 30-94%). The only two stone samples collected from
246 unmodified sampling sites in Germany had a median average proportional availability/site of
247 33% (range 6-60%). Soft alteration sampling sites in Germany were dominated by sand
248 habitats with a median proportional availability of 100% (range 60-100%; n=63) while stones
249 accounted for only 16% (range 10-40%) median proportional availability/site when present
250 (n=7). Macrophyte habitats were found only at 2 soft alteration sites representing, however,
251 20 % (range 10-30%) median proportional availability/site. Hard alteration sites were
252 characterized again by sand habitats (n=48) in German lakes with median proportional
253 availability of 90 % (range 30-100%). Stone habitats were found at 7 hard alteration sampling
254 sites and accounted for 30 % (range 5-70%) median proportional availability/site. The only 4
255 macrophyte habitats found at hard alteration sampling sites in German lakes accounted for
256 22.5% (range 5-30%) median proportional availability/site.

257 The most dominant habitat with highest median proportional availability/site at unmodified
258 sampling sites in Ireland were stones (n=40; median proportional availability/site 100%, range
259 33.33-100%). Second highest proportional availability at unmodified sampling sites was
260 found for sand habitats (n=17; median=66.67%, range 42-100%). While a comparatively
261 higher number of macroinvertebrate samples were collected from macrophytes, median
262 proportional availability/site of this habitat accounted for only 33.33% (16%-100%). Number
263 of samples collected from different habitats at soft alteration sampling sites in Irish lakes was
264 relatively equally distributed among habitats (macrophytes n=26; sand n=23; stones n=32) but
265 highest median proportional availability/site was found for stone habitats (median = 100%;
266 range 33.33-100%) followed by sand habitats (median = 94%, range 37-100%) and
267 macrophyte habitats (median = 58.33%, range 12-100%). Hard alteration sampling sites were

268 dominated, however, by stone habitats (n=65) which showed a median proportional
269 availability/site of 100% (range 16-100%). Sand and macrophyte habitats were sampled for
270 macroinvertebrates only from 8 (n=10) and 5 (n=6) sites, respectively and had a median
271 proportional availability/site of 33.33% (range 26-84%) and 33.33% (range 33.33-66.66%),
272 correspondingly at hard alteration sites in Ireland.

273 In Italy macrophytes were the dominant habitat found at unmodified sampling sites (n=80)
274 with a median proportional availability/site of 60% (range 10-100%). Stone and sand habitats
275 accounted for 40 and 30 macroinvertebrate samples, and median proportional
276 availabilities/sites of 60% (range 5-80%) and 40% (range 5-70%), respectively. Soft alteration
277 sampling sites in Italy were characterised by sand habitats (n=106) with median proportional
278 availability/site of 100% (range 40-100%). Stone and macrophyte habitats were represented
279 by 28 and 7 macroinvertebrate samples, respectively, with comparatively lower median
280 proportional availability/site of 70% (range 10-100%) and 30% (range 20-60%),
281 correspondingly. Highest number of samples collected at hard alteration sites in Italy were
282 stone habitat samples (n=71; median proportional availability/site 80%, range 10-100%). Sand
283 habitats accounted for 28 macroinvertebrate samples with median proportional
284 availability/site of 60% (range 30-100%) and macrophytes for 12 macroinvertebrate samples
285 with comparatively low median proportional availability/site of 25% (range 10-100%).

286 *Community composition*

287 NMDS in combination with ANOSIM identified no differences among macroinvertebrate
288 composite and habitat-specific samples at unmodified sampling sites in all countries (Figure
289 2, Table 3). Macroinvertebrate community structures at soft alteration sites varied between
290 composite and stone habitat samples in Germany, and composite and macrophyte habitat
291 samples in Italy (Table 3). No differences in macroinvertebrate community structures were
292 identified among composite and habitat-specific samples in Ireland at soft alteration sampling

293 sites (Figure 2; Table 3). At hard alteration sites NMDS together with ANOSIM identified
294 significant differences in macroinvertebrate community structures only between composite
295 and stone habitat samples in Germany (Table 3). All other habitat-specific samples did not
296 differ from those collected using the composite sampling approach in all countries (Figure 2;
297 Table 3).

298 PERMDISP identified no significant differences in homogeneity of spatial dispersion in
299 macroinvertebrate community structures among composite and habitat-specific samples in all
300 alteration types in Germany. In Ireland, homogeneity of dispersion of macroinvertebrate
301 community structures within composite and habitat-specific samples did not vary significantly
302 from each other at all alteration sites with the exception of composite and stone habitat
303 samples at soft alteration sites (PERMDISP, $t = 2.61$, $P_{(perm)} < 0.05$). In Italy differences in
304 variability in community structures were identified only between composite and sand habitat
305 samples at unmodified and hard alteration sampling sites (PERMDISP, $t = 3.42$ and $t = 4.03$,
306 both $P_{(perm)} < 0.05$ for composite/sand at unmodified and soft alteration sites, respectively).

307 ANOSIM and PERMDISP did not detect significant differences in macroinvertebrate
308 community structures and associated homogeneities of variances between collected and
309 proportional and unproportional artificially generated composite samples, respectively, in all
310 countries and all alteration types (Table 4; PERMDISP, unmodified: $F = 1.305$, $F = 0.152$, F
311 $= 2.788$, hard: $F = 0.1063$, $F = 1.98$, $F = 0.216$, soft: $F = 1.289$, $F = 2.134$, $F = 0.431$,
312 Germany, Ireland and Italy, respectively, all $p > 0.05$).

313 Invertebrate metrics calculated from macroinvertebrate abundances of proportional and
314 unproportional artificial composite and field composite samples performed equally well in
315 correlating with the morphological stressor index (Table 1). Ranges in Spearman-Rank
316 correlations did not differ significantly among different composite sample types (Table 1;
317 paired t-tests, Germany: composite – proportional artificial, $t = 1.49$, $p = 0.1795$, composite –

318 unproportional artificial, $t = 1.60$, $p = 0.1533$; Ireland: composite – proportional artificial, $t =$
319 0.01 , $p = 0.9941$, composite – unproportional artificial, $t = -0.76$, $p = 0.4681$; Italy: composite
320 – proportional artificial, $t = -0.32$, $p = 0.7539$; composite – unproportional artificial, $t = -0.41$,
321 $p = 0.6913$).

322 *Time-effort*

323 Time estimated for the collection and processing of macroinvertebrate samples was assessed
324 in order to compare the efficiency of different sampling methodologies. Collection of German
325 habitat-specific and composite samples in the field accounted for 30 minutes on average each
326 sample. For the sorting of macroinvertebrate habitat-specific samples in the laboratory an
327 experienced worker had to spend 8 h on average per sample. Sorting of German composite
328 macroinvertebrate samples involved 10.3 h on average. Time-effort needed for
329 macroinvertebrate identification, however, was not assessed quantitatively but accounted for
330 the same amount of time on average irrespective of the sampling method used for German
331 samples. In Ireland, field sampling using both sampling protocols also accounted on average
332 for 30 minutes each sample. Sorting of habitat-specific samples in the laboratory involved on
333 average 6 h for an experienced worker while about 10 h had to be spend for sorting of
334 composite samples. Identification of habitat-specific macroinvertebrate samples took on
335 average 4 h and 8 h for composite samples in Ireland. For the collection of macroinvertebrate
336 habitat-specific and composite samples in Italian lakes, an average of 15 minutes was spent
337 per sample in the field. Sorting and identification (no separate estimates available) of
338 macroinvertebrate habitat-specific samples accounted for 7 hours on average each sample
339 while sorting and identification of composite samples took about 11 h per sample. In
340 summary, collection and sorting of macroinvertebrates accounted for 10.8 h using the
341 composite sampling and 25.5 h using the habitat-specific approach in Germany. In Ireland,
342 18.5 h were spent for collection and processing of macroinvertebrates using the composite

343 sampling and 31.5 h with the habitat-specific sampling approach. For the collection and
344 processing of macroinvertebrate samples in Italy, 11.25 h were needed using the composite
345 and 21.75 h with the habitat-specific sampling approach.

346 **Discussion**

347 This study aimed at identifying the most suitable method for routine monitoring of European
348 lakes as required under the EU WFD. The complexity and heterogeneity of littoral habitats
349 has often led to the recommendation of habitat-specific sampling for lake assessment
350 purposes in order to reduce variability within littoral macroinvertebrate samples and
351 consequently improve signal precision (Tolonen et al., 2001; Weatherhead & James, 2001;
352 Brauns et al., 2007a). In accordance with our hypothesis we were able to show that pooled
353 composite benthic macroinvertebrate samples when collected proportional to availability of
354 individual habitats at a morphologically altered or unmodified sampling site, represent
355 individual sampling locations effectively. We were able to corroborate the results from our
356 pilot study (Porst et al., 2012) and to demonstrate that the results apply for a wide range of
357 lake types across a gradient of morphological alterations and a north-south gradient of
358 European geographical regions/countries. Macroinvertebrate community composition of
359 pooled composite samples did not differ significantly from habitat-specific macroinvertebrate
360 samples across differing shoreline types and countries with only a few minor exceptions. In
361 Germany macroinvertebrate stone habitat samples showed significant differences in
362 community composition when compared with composite samples from soft and hard
363 alteration sites. Stone habitats made up only a comparatively small fraction of
364 macroinvertebrate habitats at modified shorelines (both dominated by sand habitat) and
365 consequently only a minor proportion of collected composite samples in Germany. Littoral
366 invertebrate samples collected from macrophyte habitats at soft alteration sites in Italy also
367 varied from composite samples from respective sampling sites. This once again is a result of

368 the comparatively low proportional availability of this habitat at this alteration type in Italy.
369 Macrophyte samples were collected from only few soft alteration sampling sites and
370 represented the lowest proportional availability when compared to the other two habitats at
371 respective morphologically altered sampling locations in Italy.

372 PERMDISP analysis generally revealed no significant differences in homogeneity of
373 dispersion in macroinvertebrate community structures from individual habitats compared with
374 those from pooled composite samples collected at morphologically differing shoreline types
375 across geographical regions. This once again supports the suitability of the collection of
376 pooled macroinvertebrate composite samples for routine lake monitoring as requested under
377 the EU WFD and is in accordance with our preliminary study comparing different sampling
378 methodologies at Lake Werbellin, Germany (Porst et al., 2012). In contrast, Schreiber and
379 Brauns (2010) found variability within habitat-specific macroinvertebrate samples to differ
380 considerably from that of pooled composite samples. The latter study, however, did not
381 account for respective proportional availabilities of individual habitats at each
382 macroinvertebrate sampling location giving each habitat sample equal weight in the
383 computation of artificial pooled samples. This once more emphasizes the importance of the
384 proportional sampling approach for the collection of representative littoral macroinvertebrate
385 samples for the assessment of morphological shoreline alterations as applied in our study.

386 For the assessment of lakes, benthic macroinvertebrate communities collected from single
387 littoral habitats are typically combined into pooled samples in order to obtain a single signal
388 per site. These artificial composite samples also form the basis for the calculation of different
389 macroinvertebrate metrics containing information about certain characteristics or traits of the
390 macroinvertebrate community rather than individual abundances of single species. In our
391 study, proportional and unproportional artificially computed littoral macroinvertebrate
392 composite samples did neither differ significantly in their community structures nor

393 homogeneity of variances in community structures when compared with those of composite
394 samples collected in the field. While variability in macroinvertebrate community structures
395 was generally slightly lower in artificially computed composite samples, the differences were
396 never significant and support the adequacy of the collection of pooled composite
397 macroinvertebrate samples for lake monitoring. Furthermore, proportional composite samples
398 collected in the field proved suitable for use in lake monitoring programmes based on
399 multimetric indices (Hering et al., 2004; Gabriels et al., 2010) for the assessment of lake
400 ecological status. Field composite and proportional and unproportional artificial composite
401 samples performed equally well in the correlation of 10 selected invertebrate metrics typically
402 used for lake morphology assessments with a previously calculated stressor index (Miler et
403 al., 2013). While both artificially computed composite samples showed similar results in the
404 comparison with collected macroinvertebrate composite samples, it should not be concluded
405 that proportional sampling of littoral habitats would not be necessary for obtaining
406 meaningful results in lake assessment programs. In our study habitat proportions in the field
407 generally showed relatively equal distributions among habitats across alteration types and
408 lakes in all countries/geographical regions. We conclude, however, that higher variability in
409 habitat proportions would result in a comparatively less accurate representation of sites using
410 a non-proportional approach as demonstrated in the study by Schreiber and Brauns (2010).

411 Our study demonstrated the suitability of the proportional composite sampling methodology
412 for regular lake monitoring for the generally dominant littoral habitats sand, stones and
413 macrophytes. While these habitats showed highest proportional availabilities across all littoral
414 sampling sites in all three European countries, other macroinvertebrate habitats such as woody
415 debris or roots could also be considered to be included for monitoring purposes. These
416 habitats, which usually account for a fraction of the area of a sampling site only and thus
417 would make up only a small part of respective composite samples, are known to inhabit rare

418 or sensitive macroinvertebrate taxa (Lorenz et al., 2004; Strayer & Findlay, 2010; Porst et al.,
419 2012). The inclusion of disturbance sensitive taxa is required by the EU WFD and metrics
420 describing the percentage or taxa number of disturbance sensitive taxonomic groups are a
421 central part of many macroinvertebrate based multimetric assessment systems (Hering et al.,
422 2004; Lorenz et al., 2004; Hering et al., 2006; Schartau et al., 2008; Gabriels et al., 2010;
423 Timm & Möls, 2012). Our previous study assessing the suitability of the composite sampling
424 method at Lake Werbellin (Porst et al., 2012) already demonstrated the adequacy of the latter
425 sampling method also for the inclusion of these usually comparatively scarcely represented
426 littoral habitats in contrast with the study by Schreiber and Brauns (2010). We recommend the
427 inclusion of additional habitats only if those habitats cover a minimum of 5% area of the
428 sampling site following the AQEM/STAR method for the assessment of streams using benthic
429 invertebrates (AQEM Consortium, 2002; Timm & Möls, 2012) or if assessment is being
430 carried out for conservation purposes rather than basic quality assessment.

431 Time- and cost-effectiveness are important factors for the design and implementation of
432 regular lake monitoring programmes. While usually the largest fraction of time needed for
433 assessment purposes using benthic macroinvertebrates is spent on the processing and
434 identification of samples in the laboratory, the collection of macroinvertebrate samples in the
435 field involves far less time and associated expenses (Ferraro et al., 1989; Haase et al., 2004;
436 Tolonen & Hämäläinen, 2010). In our study the collection of benthic samples in the field
437 using either of the two sampling methods accounted for approximately the same time and
438 made up only a comparatively small amount of total time required for sample processing.
439 Sorting and identification of macroinvertebrate samples in the laboratory was found to be
440 more efficient for individual habitat samples. Total time needed for the collection, sorting and
441 identification of benthic macroinvertebrate samples, however, was about twofold higher for
442 collection, sorting and identification of all habitat samples representing a site. Thus, the

443 working-effort required for the stratified habitat-specific sampling method is considerably
444 higher and consequently accounts for undoubtedly higher associated costs when compared to
445 the suggested composite sampling method.

446 **Conclusions**

447 This study demonstrated that pooled macroinvertebrate composite samples when collected
448 proportionally to habitat availability at a littoral sampling site have the potential of being used
449 in routine monitoring programs for the WFD compliant assessment of European lakes with
450 respect to morphological alterations in the lake littoral. We were able to show that
451 proportional composite samples represent both, morphologically altered as well as unmodified
452 shorelines adequately in terms of macroinvertebrate community compositions across a range
453 of lake types and a European gradient while their processing additionally accounts for
454 considerably less time and associated costs. The results of this study emphasize the
455 importance of applying the proportional sampling approach for the assessment of lake
456 ecological status and support its use as a time and cost effective sampling strategy. While our
457 sampling scheme focused on the three dominant habitats present across the European
458 gradient, the inclusion of additional habitats which might account for only a fraction of the
459 sampling site could be considered for the design of lake assessment programmes beyond the
460 purposes of the EU WFD. In case lake littoral zones are sampled for other purposes, as for
461 identifying effective restoration options for lake littoral habitats, or to survey rare and
462 endangered invertebrate species, we recommend habitat-specific sampling, in order to record
463 habitat specificities of target species.

464 Table 1: Spearman-Rank correlations of metrics calculated from macroinvertebrate
 465 abundances of field composite (CO), proportional (CO1) and unproportional artificial
 466 composite (CO2) samples with the morphological stressor index. Shown are 10 selected
 467 metrics that are typical for morphological assessment methods based on lake invertebrates and
 468 their respective Rho- and p-values.

Metric	CO		CO1		CO2	
	ρ	p	ρ	p	ρ	p
Germany						
ASPT	-0.21	0.084	-0.20	0.088	-0.20	0.088
Margalef Diversity	-0.51	<0.001	-0.48	<0.001	-0.49	<0.001
r/K relationship	0.38	<0.001	0.43	<0.001	0.44	<0.001
Type Lit %	-0.22	0.058	-0.21	0.076	-0.20	0.088
Odonata %	-0.54	<0.001	-0.46	<0.001	-0.46	<0.001
Trichoptera %	-0.40	0.001	-0.41	<0.001	-0.40	<0.004
Diptera %	0.26	0.025	0.11	0.354	0.15	0.199
No. Odonata Taxa	-0.52	<0.001	-0.40	<0.001	-0.40	<0.001
No. Trichoptera Taxa	-0.47	<0.001	-0.42	<0.001	-0.42	<0.001
No. ETO Taxa	-0.42	<0.001	-0.42	<0.001	-0.42	<0.001
Ireland						
ASPT	-0.26	0.020	-0.28	0.013	-0.29	0.009
Margalef Diversity	-0.18	0.118	-0.23	0.039	-0.23	0.043
r/K relationship	-0.11	0.333	0.04	0.738	0.10	0.363
Type Lit %	0.19	0.096	0.18	0.112	0.26	0.022
Odonata %	-0.00	1.000	-0.06	0.628	-0.05	0.658
Trichoptera %	-0.18	0.113	-0.14	0.206	-0.14	0.199
Diptera %	0.11	0.322	0.02	0.866	2	0.884
No. Odonata Taxa	-0.03	0.756	-0.09	0.406	-0.08	0.489
No. Trichoptera Taxa	-0.15	0.174	-0.19	0.094	-0.18	0.109
No. ETO Taxa	-0.25	0.026	-0.23	0.036	-0.23	0.037
Italy						
ASPT	-0.32	<0.001	-0.16	0.053	-0.16	0.053
Margalef Diversity	-0.38	<0.001	-0.42	<0.001	-0.41	<0.001
r/K relationship	0.37	<0.001	0.25	0.006	0.25	0.006
Type Lit %	0.05	0.614	0.17	0.058	0.18	0.042
Odonata %	-0.41	<0.001	-0.48	<0.001	-0.49	<0.001
Trichoptera %	-0.18	0.042	-0.30	<0.001	-0.31	<0.001
Diptera %	0.07	0.418	-0.03	0.731	-0.04	0.678
No. Odonata Taxa	-0.43	<0.001	-0.37	<0.001	-0.37	<0.001
No. Trichoptera Taxa	-0.20	0.028	-0.29	0.001	-0.29	<0.001
No. ETO Taxa	-0.38	<0.001	-0.42	<0.001	-0.42	<0.001

469

470 Table 2: Composition of the morphological stressor index developed for the three
 471 geographical regions Germany, Ireland and Italy.

Stressor Index Component	Geographical region		
	Germany	Ireland	Italy
Number of habitats	X		
Habitat diversity		X	X
Total PVI	X		X
Sum of macrophyte types		X	
Sum of vegetation cover types	X	X	
Sum of CWD/roots/overhanging vegetation	X		X
Pressure index	X	X	X
Natural/artificial dominant land cover type			X

472

473 Table 3: Results from two-way nested ANOSIM analysis comparing benthic macroinvertebrate communities from habitat-specific and composite

474 samples from unmodified, hard and soft alteration sampling sites in different geographical regions/countries.

		Germany		Ireland		Italy		
		<i>Unmodified</i>		<i>Unmodified</i>		<i>Unmodified</i>		
Groups	R-statistic	P %	Groups	R-statistic	P %	Groups	R-statistic	P %
Composite, Macrophytes	-0.027	53.3	Composite, Macrophytes	-0.105	97	Composite, Macrophytes	-0.018	60.6
Composite, Sand	0.071	18.2	Composite, Stones	-0.037	64,5	Composite, Sand	0.087	7.6
			Composite, Sand	-0,054	61,7	Composite, Stones	0.034	24.1
<i>Hard alteration</i>			<i>Hard alteration</i>			<i>Hard alteration</i>		
Groups	R-statistic	P %	Groups	R-statistic	P %	Groups	R-statistic	P %
Composite, Sand	-0.118	94.9	Composite, Stones	-0.077	91,7	Composite, Sand	-0.045	62.1
Composite, Stones	0.562	0.06	Composite, Sand	0,045	30,1	Composite, Stones	-0.027	76
Composite, Macrophytes	-0.013	46.7	Composite, Macrophytes	0,225	11,8	Composite, Macrophytes	0.176	12.8
<i>Soft alteration</i>			<i>Soft alteration</i>			<i>Soft alteration</i>		
Groups	R-statistic	P %	Groups	R-statistic	P %	Groups	R-statistic	P %
Composite, Sand	-0.043	68.4	Composite, Macrophytes	-0.135	95,1	Composite, Macrophytes	0.559	2.2
Composite, Stones	0.73	0.08	Composite, Stones	-0.127	86,7	Composite, Sand	0.045	9.6
			Composite, Sand	0,096	30	Composite, Stones	0.037	32.2

475

- 476 Table 4: Results from two-way nested ANOSIM analysis comparing benthic macroinvertebrate communities from collected composite samples
 477 (CO) with proportional artificial composite (CO1) and unproportional artificial composite (CO2) samples from unmodified, hard and soft alteration
 478 sampling site in different geographical regions/countries.

Germany		Ireland		Italy	
<i>Unmodified</i>		<i>Unmodified</i>		<i>Unmodified</i>	
Groups	R-statistic	Groups	R-statistic	Groups	R-statistic
CO, CO1	-0.087	CO, CO1	-0.09	CO, CO1	0.008
CO, CO2	-0.092	CO, CO2	-0.089	CO, CO2	0.007
CO1, CO2	-0.114	CO1, CO2	-0.097	CO1, CO2	-0.066
	P %		P %		P %
	87.4		97.2		32.7
	89.3		98		33.3
	95.7		99.4		99.8
<i>Hard alteration</i>		<i>Hard alteration</i>		<i>Hard alteration</i>	
Groups	R-statistic	Groups	R-statistic	Groups	R-statistic
CO, CO1	-0.068	CO, CO1	-0.063	CO, CO1	-0.02
CO, CO2	-0.028	CO, CO2	-0.066	CO, CO2	-0.02
CO1, CO2	-0.121	CO1, CO2	-0.095	CO1, CO2	-0.076
	P %		P %		P %
	75.4		84.1		66.4
	55.3		87.1		63.8
	94.8		98.6		99.7
<i>Soft alteration</i>		<i>Soft alteration</i>		<i>Soft alteration</i>	
Groups	R-statistic	Groups	R-statistic	Groups	R-statistic
CO, CO1	-0.062	CO, CO1	-0.063	CO, CO1	0.006
CO, CO2	-0.063	CO, CO2	-0.065	CO, CO2	0.003
CO1, CO2	-0.116	CO1, CO2	-0.11	CO1, CO2	-0.057
	P %		P %		P %
	78.5		81.8		36.3
	77.7		84.2		36.1
	95.5		99.2		99.6

479

480 Figure 1: Map of Europe showing 32 lakes sampled in 3 different European geographical
481 regions.

482 Figure 2: NMDS-plot of macroinvertebrate species arcsine-transformed proportional
483 abundance data from unmodified, soft and hard alteration sampling sites in Germany, Ireland
484 and Italy.

485

486 **References**

- 487 Anderson, M. J., R. N. Gorley & K. R. Clarke, 2008. PERMANOVA+ for PRIMER: Guide to
488 Software and Statistical Methods. PRIMER-E: Plymouth, UK.
- 489 AQEM Consortium, 2002. Manual for the application of the AQEM system. A comprehensive method
490 to assess European streams using benthic macroinvertebrates, developed for the purpose of the
491 Water Framework Directive. Version 1.0. - February 2002. (<http://www.aqem.de>).
- 492 Bänziger, R., 1995. A comparative study of the zoobenthos of eight land-water interfaces (Lake of
493 Geneva). *Hydrobiologia* 300-301: 133-140.
- 494 Brauns, M., X.-F. Garcia, M. T. Pusch & N. Walz, 2007a. Eulittoral macroinvertebrate communities
495 of lowland lakes: discrimination among trophic states. *Freshwater Biology* 52: 1022-1032.
- 496 Brauns, M., X.-F. Garcia, N. Walz & M. T. Pusch, 2007b. Effects of human shoreline development on
497 littoral macroinvertebrates in lowland lakes. *Journal of Applied Ecology* 44: 1138-1144.
- 498 Brauns, M., B. Gücker, C. Wagner, X.-F. Garcia, N. Walz & M. T. Pusch, 2011. Human lakeshore
499 development alters the structure and trophic basis of littoral food webs. *Journal of Applied*
500 *Ecology* 48: 916-925.
- 501 Brodersen, K. P. & C. Lindegaard, 1999. Classification, assessment and trophic reconstruction of
502 Danish lakes using chironomids. *Freshwater Biology* 42: 143-157.
- 503 Carpenter, S. R., B. J. Benson, R. Biggs, J. W. Chipman, J. A. Foley, S. A. Golding, R. B. Hammer, P.
504 C. Hanson, P. T. J. Johnson, A. M. Kamarainen, T. K. Kratz, R. C. Lathrop, K. D. McMahon,
505 B. Provencher, J. A. Rusak, C. T. Solomon, E. H. Stanley, M. G. Turner, M. J. Vander
506 Zanden, C.-H. Wu & H. Yuan, 2007. Understanding Regional Change: A Comparison of Two
507 Lake Districts. *BioScience* 57: 323-335.

508 Clarke, K. R. & R. M. Warwick, 2001. Change in marine communities: an approach to statistical
509 analysis and interpretation, 2nd edition. PRIMER-E; Plymouth.

510 EC, 2000. Council Directive 2000/60/EC establishing a framework for Community action in the field
511 of water policy. Official Journal no L 321/1, 22122000.

512 Ferraro, S. P., F. A. Cole, W. A. Deben & R. C. Swartz, 1989. Power-cost efficiency of 8
513 macrobenthic sampling schemes in Puget Sound, Washington, USA. Canadian Journal of
514 Fisheries and Aquatic Sciences 46: 2157-2165.

515 Gabel, F., X. F. Garcia, I. Schnauder & M. T. Pusch, 2012. Effects of ship-induced waves on littoral
516 benthic invertebrates. Freshwater Biology 57: 2425-2435.

517 Gabriels, W., K. Lock, N. De Pauw & P. L. M. Goethals, 2010. Multimetric Macroinvertebrate Index
518 Flanders (MMIF) for biological assessment of rivers and lakes in Flanders (Belgium).
519 Limnologica - Ecology and Management of Inland Waters 40: 199-207.

520 Gotelli, N. J. & A. M. Ellison, 2004. A Primer of Ecological Statistics. Sinauer Associates, Inc.,
521 Sunderland, MA.

522 Haase, P., S. Lohse, S. Pauls, K. Schindehütte, A. Sundermann, P. Rolauuffs & D. Hering, 2004.
523 Assessing streams in Germany with benthic invertebrates: development of a practical
524 standardised protocol for macroinvertebrate sampling and sorting. Limnologica - Ecology and
525 Management of Inland Waters 34: 349-365.

526 Harrison, S. S. C. & A. A. G. Hildrew, 1998. Patterns in the epilithic community of a lake littoral.
527 Freshwater Biology 39: 477-492.

- 528 Hering, D., C. Feld, O. Moog & T. Ofenböck, 2006. Cook book for the development of a Multimetric
529 Index for biological condition of aquatic ecosystems: Experiences from the European AQEM
530 and STAR projects and related initiatives. *Hydrobiologia* 566: 311-324.
- 531 Hering, D., C. Meier, C. Rawer-Jost, C. K. Feld, R. Biss, A. Zenker, A. Sundermann, S. Lohse & J.
532 Böhmer, 2004. Assessing streams in Germany with benthic invertebrates: selection of
533 candidate metrics. *Limnologica - Ecology and Management of Inland Waters* 34: 398-415.
- 534 Jennings, M. J., M. A. Bozek, G. R. Hatzenbeler, E. E. Emmons & M. D. Staggs, 1999. Cumulative
535 Effects of Incremental Shoreline Habitat Modification on Fish Assemblages in North
536 Temperate Lakes. *North American Journal of Fisheries Management* 19: 18-27.
- 537 Jurca, T., L. Donohue, D. Laketić, S. Radulović & K. Irvine, 2012. Importance of the shoreline
538 diversity features for littoral macroinvertebrate assemblages. *Fundamental and Applied*
539 *Limnology* 180: 175-184.
- 540 Langdon, P. G., Z. Ruiz, K. P. Brodersen & I. D. L. Foster, 2006. Assessing lake eutrophication using
541 chironomids: understanding the nature of community response in different lake types.
542 *Freshwater Biology* 51: 562-577.
- 543 Lorenz, A., D. Hering, C. Feld & P. Rolauuffs, 2004. A new method for assessing the impact of
544 hydromorphological degradation on the macroinvertebrate fauna of five German stream types.
545 *Hydrobiologia* 516: 107-127.
- 546 Mastrantuono, L., F. Pilotto, M. Rossopinti, M. Bazzanti & A. G. Solimini, in press. Response of
547 littoral macroinvertebrates to morphological disturbances in Mediterranean lakes: the case of
548 Lake Piediluco (central Italy). *Fundamental and Applied Limnology*.

549 McGoff, E., J. Aroviita, F. Pilotto, O. Miler, A. G. Solimini, G. Porst, T. Jurca, L. Donohue & L.
550 Sandin, 2013. Assessing the relationship between the Lake Habitat Survey and littoral
551 macroinvertebrate communities in European lakes. *Ecological Indicators* 25: 205-214.

552 Miler, O., G. Porst, E. McGoff, F. Pilotto, L. Donohue, T. Jurca, A. Solimini, L. Sandin, K. Irvine, J.
553 Aroviita, R. Clarke & M. T. Pusch, 2013. Morphological alterations of lake shores in Europe:
554 A multimetric ecological assessment approach using benthic macroinvertebrates. *Ecological*
555 *Indicators* 34: 398-410.

556 Miler, O., G. Porst, E. McGoff, F. Pilotto, L. Donohue, T. Jurca, A. Solimini, L. Sandin, K. Irvine, J.
557 Aroviita, R. Clarke & M. T. Pusch, 2014 (early view). An index of human alteration of lake
558 shore morphology. *Aquatic Conservation: Marine and Freshwater Ecosystems*.

559 Mischke, U., S. Thackeray, M. Dunbar, C. McDonald, L. Carvalho, C. de Hoyos, M. Jarvinen, C.
560 Laplace-Treytore, G. Morabito, B. Skjelbred, A. L. Solheim, B. Brierley & B. Dudley, 2012.
561 Guidance document on sampling, analysis and counting standards for phytoplankton in lakes
562 Project report WISER Deliverable D31-4.

563 Moss, B., D. Stephen, C. Alvarez, E. Becares, W. V. D. Bund, S. E. Collings, E. V. Donk, E. D. Eyto,
564 T. Feldmann, C. Fernández-Aláez, M. Fernández-Aláez, R. J. M. Franken, F. García-Criado,
565 E. M. Gross, M. Gyllström, L.-A. Hansson, K. Irvine, A. Järvalt, J.-P. Jensen, E. Jeppesen, T.
566 Kairesalo, R. Kornijów, T. Krause, H. Künnap, A. Laas, E. Lill, B. Lorens, H. Luup, M. R.
567 Miracle, P. Nöges, T. Nöges, M. Nykänen, I. Ott, W. Peczula, E. T. H. M. Peeters, G. Phillips,
568 S. Romo, V. Russell, J. Salujõe, M. Scheffer, K. Siewertsen, H. Smal, C. Tesch, H. Timm, L.
569 Tuvikene, I. Tonno, T. Virro, E. Vicente & D. Wilson, 2003. The determination of ecological
570 status in shallow lakes - a tested system (ECOFAME) for implementation of the European
571 Water Framework Directive. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13:
572 507-549.

573 O'Connor, N. A., 1991. The effects of habitat complexity on the macroinvertebrates colonising wood
574 substrates in a lowland stream. *Oecologia* 85: 504-512.

575 Phillips., G., G. Morabito, L. Carvalho, A. L. Solheim, B. Skjelbred, J. Moe, T. Andersen, U.
576 Mischke, C. de Hoyos & G. Borics, 2011. Report on lake phytoplankton composition metrics,
577 including a common metric approach for use in intercalibration by all GIGs. Project report
578 WISER Deliverable D31-1.

579 Pilotto, F., M. Bazzanti, V. D. Vito, D. Frosali, F. Livretti, L. Mastrantuono, M. T. Pusch, F. Sena &
580 A. G. Solimini, in press. Relative impacts of morphological alteration to shorelines and
581 eutrophication on littoral macroinvertebrates in Mediterranean lakes. *Freshwater Science*.

582 Pinel-Alloul, B., G. Méthot, L. Lapierre & A. Willsie, 1996. Macroinvertebrate community as a
583 biological indicator of ecological and toxicological factors in Lake Saint-François (Québec).
584 *Environmental Pollution* 91: 65-87.

585 Porst, G., S. Bader, E. Münch & M. Pusch, 2012. Sampling approaches for the assessment of shoreline
586 development based on littoral macroinvertebrates: the case of Lake Werbellin, Germany.
587 *Fundamental and Applied Limnology* 180: 123-131.

588 Porst, G., F. Gabel, S. Lorenz & O. Miler, in press. Implications of hydromorphological alterations to
589 the littoral zone for freshwater ecosystem functioning. *Fundamental and Applied Limnology*.

590 R Core Team, 2013. R: A language and environment for statistical computing. R Foundation for
591 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL [http://www.R-](http://www.R-project.org/)
592 [project.org/](http://www.R-project.org/).

593 Rasmussen, J. B., 1988. Littoral Zoobenthic Biomass in Lakes, and Its Relationship to Physical,
594 Chemical, and Trophic Factors. *Canadian Journal of Fisheries and Aquatic Sciences* 45: 1436-
595 1447.

596 Reice, S. R. & M. Wohlenberg, 1993. Monitoring freshwater benthic macroinvertebrates and benthic
597 processes: measures for assessment of ecosystem health. In Rosenberg, D. M. & V. H. Resh
598 (eds) *Freshwater biomonitoring and benthic macroinvertebrates*. Chapman & Hall, NY.

599 Rowan, J. S., J. Carwardine, R. W. Duck, O. M. Bragg, A. R. Black, M. E. J. Cutler, I. Soutar & P. J.
600 Boon, 2006. Development of a technique for Lake Habitat Survey (LHS) with applications for
601 the European Union Water Framework Directive. *Aquatic Conservation: Marine and*
602 *Freshwater Ecosystems* 16: 637-657.

603 Rowan, J. S., R. W. Duck, J. Carwardine, O. M. Bragg, A. R. Black, M. E. J. Cutler & I. Soutar, 2008.
604 *Lake Habitat Survey in the United Kingdom - Field Survey Guidance Manual*. SNIFFER,
605 Edinburgh.

606 Saether, O. A., 1979. Chironomid communities as water quality indicators. *Ecography* 2: 65-74.

607 Schartau, A., S. J. Moe, L. Sandin, B. McFarland & G. Raddum, 2008. Macroinvertebrate indicators
608 of lake acidification: analysis of monitoring data from UK, Norway and Sweden. *Aquatic*
609 *Ecology* 42: 293-305.

610 Scheuerell, M. D. & D. E. Schindler, 2004. Changes in the Spatial Distribution of Fishes in Lakes
611 Along a Residential Development Gradient. *Ecosystems* 7: 98-106.

612 Schneider, K. & K. Winemiller, 2008. Structural complexity of woody debris patches influences fish
613 and macroinvertebrate species richness in a temperate floodplain-river system. *Hydrobiologia*
614 610: 235-244.

615 Schreiber, J. & M. Brauns, 2010. How much is enough? Adequate sample size for littoral
616 macroinvertebrates in lowland lakes. *Hydrobiologia* 649: 365-373.

617 Solimini, A. G., G. Free, I. Donohue, K. Irvine, M. T. Pusch, B. Rossaro, L. Sandin & A. C. Cardoso,
618 2006. Using benthic Macroinvertebrates to Assess Ecological Status of Lakes: Current
619 Knowledge and Way Forward to Support WFD Implementation. Office for Official
620 Publications of the European Communities, Luxembourg.

621 Solimini, A. G. & L. Sandin, 2012. The importance of spatial variation of benthic invertebrates for the
622 ecological assessment of European lakes. *Fundamental and Applied Limnology* 180: 85-89.

623 Søndergaard, M., S. E. Larsen, T. B. Jørgensen & E. Jeppesen, 2011. Using chlorophyll a and
624 cyanobacteria in the ecological classification of lakes. *Ecological Indicators* 11: 1403-1412.

625 STAR Consortium, 2003. The AQEM sampling method to be applied in STAR. Unpublished report,
626 available at <http://www.eu-star.at/pdf/AqemMacroinvertebrateSamplingProtocol.pdf>.

627 Strayer, D. & S. Findlay, 2010. Ecology of freshwater shore zones. *Aquatic Sciences - Research*
628 *Across Boundaries* 72: 127-163.

629 Taniguchi, H., S. Nakano & M. Tokeshi, 2003. Influences of habitat complexity on the diversity and
630 abundance of epiphytic invertebrates on plants. *Freshwater Biology* 48: 718-728.

631 Timm, H. & T. Möls, 2012. Littoral macroinvertebrates in Estonian lowland lakes: the effects of
632 habitat, season, eutrophication and land use on some metrics of biological quality.
633 *Fundamental and Applied Limnology* 180: 145-156.

634 Tolonen, K. T. & H. Hämäläinen, 2010. Comparison of sampling methods and habitat types for
635 detecting impacts on lake littoral macroinvertebrate assemblages along a gradient of human
636 disturbance. *Fundamental and Applied Limnology* 176: 43-59.

- 637 Tolonen, K. T., H. Hämäläinen, I. J. Holopainen & J. Karjalainen, 2001. Influences of habitat type and
638 environmental variables on littoral macroinvertebrate communities in a large lake system.
639 *Archiv für Hydrobiologie* 152: 39-67.
- 640 Urbanič, G., V. Petkovska & M. Pavlin, 2012. The relationship between littoral benthic invertebrates
641 and lakeshore modification pressure in two alpine lakes. *Fundamental and Applied Limnology*
642 180: 157-173.
- 643 Vadeboncoeur, Y., M. J. Vander Zanden & D. M. Lodge, 2002. Putting the Lake Back Together:
644 Reintegrating Benthic Pathways into Lake Food Web Models. *BioScience* 52: 44-54.
- 645 Weatherhead, M. A. & M. R. James, 2001. Distribution of macroinvertebrates in relation to physical
646 and biological variables in the littoral zone of nine New Zealand lakes. *Hydrobiologia* 462:
647 115-129.
- 648 Wetzel, R. G., 2001. *Limnology. Lake and River Ecosystems*. 3rd Edition. Academic Press.

Figure 1
[Click here to download Figure: Figure 1.JPG](#)




Figure 2
[Click here to download Figure: Figure 2.pdf](#)

