

Xerostomia and Clinical Outcomes in Definitive Intensity Modulated Radiotherapy (IMRT) Versus Three-dimensional Conformal Radiotherapy (3D-CRT) for Head and Neck Squamous Cell Carcinoma: A Meta-analysis

FRANCESCA DE FELICE¹, NICOLA PRANNO², PIERO PAPI², ORLANDO BRUGNOLETTI²,
VINCENZO TOMBOLINI¹ and ANTONELLA POLIMENI²

¹Department of Radiotherapy, Policlinico Umberto I, “Sapienza” University of Rome, Rome, Italy;

²Department of Oral and Maxillofacial Sciences, Policlinico Umberto I,
“Sapienza” University of Rome, Rome, Italy

Abstract. *Background/Aim:* Intensity modulated radiotherapy (IMRT) has been compared with three-dimensional conformal radiotherapy (3D-CRT) in randomized clinical trials for head and neck squamous cell carcinoma (HNSCC). The aim of this meta-analysis was to evaluate the efficacy and toxicity of IMRT and 3D-CRT and identify differences in grade ≥ 2 xerostomia incidence and clinical outcomes. *Materials and Methods:* The preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement was applied. Random-effects models were used. Primary endpoint was xerostomia of grade 2 or worse. Secondary endpoints were overall survival (OS) and loco-regional control (LRC). *Results:* Three randomized clinical trials representing 213 patients were identified. Global, grade ≥ 2 acute xerostomia and late xerostomia at 1 and 2 years after treatment were reduced with the IMRT technique (RR=0.71, 95%CI=0.59-0.86, RR=0.45, 95%CI=0.31-0.65 and RR=0.26, 95%CI=0.15-0.46, respectively). IMRT was not associated with significant OS and LRC improvement compared with 3D-CRT, with OR of 0.70 (95%CI=0.39-1.24; $p=0.22$) and 1.50 (95%CI=0.75-2.98; $p=0.25$). *Conclusion:* This meta-analysis explored the value of IMRT compared to 3D-CRT and confirmed the superiority of IMRT over 3D-CRT in terms of grade ≥ 2 xerostomia rates, but not on clinical outcomes. Its positive impact on tumor control and survival remains to be proven.

Radiation therapy (RT) with or without chemotherapy plays a central role in the treatment of head and neck squamous cell carcinoma (HNSCC), in both definitive and adjuvant setting (1). Intensity modulated RT (IMRT) technique is currently considered standard in RT treatment plans, mainly because of its ability to better limit the dose to adjacent organs at risk over three-dimensional conformal RT (3D-CRT). However, due to the anatomical complexity of the head and neck region, the risk of RT-induced toxicities remains significant and xerostomia still represents a frequent RT-related complication in HNSCC patients (2). However, different studies have demonstrated that IMRT reduces moderate to severe xerostomia onset compared to 3D-CRT treatment (3-9). Whereas its real benefit on clinical outcomes, including overall survival (OS) and loco-regional control (LRC) rates, is still unclear, mostly because these studies were too small to allow any definite conclusion (3-9). Therefore, a pooled analysis of their results is necessary to power IMRT efficacy on clinical outcomes in HNSCC.

The aim of this meta-analysis was to perform a systematic review of the literature, to compare RT-induced xerostomia and clinical outcomes in HNSCC patients treated with IMRT or 3D-CRT and to provide valuable evidence for future research.

Materials and Methods

Data extraction and trials selection. The preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement was followed to perform search strategy and selection processes (10). The meta-analysis included trials without any restrictions on publication date. The last search was carried out on May 2019. Systematic literature electronic search was conducted in Pubmed, Embase and Cochrane central register of controlled trials databases, using the following research criteria: “Head and Neck Neoplasms/radiotherapy”(Mesh) AND (“xerostomia” (MeSH Terms) OR “xerostomia”(All Fields)). The

This article is freely accessible online.

Correspondence to: Francesca De Felice, Department of Radiotherapy, Policlinico Umberto I, “Sapienza” University of Rome, Viale Regina Elena 326, Rome, Italy. Tel: +39 0649973411, Fax: +39 0649973411, e-mail: fradefelice@hotmail.it

Key Words: Radiotherapy, toxicity, xerostomia, head and neck cancer, treatment, survival, multidisciplinary team.

search strategy was linked with the Cochrane highly sensitive search strategy for identifying randomized trials in Pubmed (11). To be eligible, trials needed to compare IMRT and 3D-CRT in definitive HNSCC treatment. Randomized clinical trials, written in English, were included and reference lists of previously published reviews and meta-analysis were explored. Narrative or systematic review articles, retrospective studies, case series, case reports, commentaries, letters to editors and studies involving animals or *in vitro* models were not included. Meeting abstracts were not considered because of the insufficient data provided by the authors.

Two independent reviewers (NP and PP) selected the identified studies based on the title and abstract. If the topic of the study could not be ascertained from its title or abstract, the full-text version was retrieved for evaluation. Disagreement was resolved by a third party (FDF).

Trials were eligible if participants were newly diagnosed, with histologically proven squamous cell carcinoma at study entry. In closer evaluation of potentially eligible articles, when two articles appeared to report results with overlapping data, only the data representing the most recent publication were included in the meta-analysis. Extracted data were recorded into standardized database according to the following parameters: first author's surname, year of publication, sample size of IMRT group and 3D-CRT group, tumor and treatment details, duration of follow-up, xerostomia rates and clinical outcomes.

Focus question. This study attempted to address the following question: does the use of IMRT compared to 3D-CRT reduce the risk of acute and late xerostomia and improve LRC and OS in HNSCC patients?

Outcomes. The intent of the analysis was to evaluate the proportion of patients with grade ≥ 2 acute and late xerostomia. Late xerostomia was assessed at 12 months and 24 months after treatment. We also planned to analyze OS and LRC. Xerostomia was graded using the radiation therapy oncology group and the European organization for research and treatment of cancer (RTOG/EORTC) +/- the late effects of normal tissues subjective-objective management analytic (LENT SOMA) (12). The definition of both OS and LRC was similar across trials. OS and LRC were defined as time from the date of randomization to the defined event using Kaplan-Meier method. The number of events (side effect, death and recurrence), when available, were derived from each study. At least one of these three outcomes should have been assessed and reported in the trial to be included in the present analysis.

Statistical analysis. The grading of recommendations assessment, development and evaluation (GRADE) system was used to rate quality of evidence and grade the strength of recommendations for all included studies (13). Statistical analysis was performed using Review Manager 5.0 (14). The pooled odds ratio (OR) and risk ratio (RR) were calculated using a fixed- or random-effects model. Forest plots were used for graphical representation of each study and pooled analysis. The size of each box represents the weight that the corresponding study exerts in the meta-analysis; confidence intervals (CIs) for each study are displayed as a horizontal line through the box. The pooled OR and RR are symbolized by a solid diamond at the bottom of the forest plot, and the width of the square represents the 95%CI of the OR and RR. OR, RR and 95%CI, for each study were extracted or calculated, based on the published




Figure 1. Flow chart.

studies, according to the methods described by Tierney *et al.* in 2007 (15). A significant two-way *p*-value for comparison was defined as *p*<0.05. Statistical heterogeneity among studies was examined using both the Cochrane Q statistic (significant at *p*<0.1) and the I² value (significant heterogeneity if >50%) (16).


Results

Studies' characteristics. Flowchart of the retrieved studies and their main characteristics are presented in Figure 1 and Table I. Overall, 3 randomized clinical trials, representing 213 patients were included in the final analysis (5, 8-9). Studies included only patients with oral cavity (n=3), oropharynx (n=137), hypopharynx (n=0) and larynx (n=23) cancer, except T1N0 glottic larynx. Patients were randomly assigned to 3D-CRT (n=104) *versus* IMRT (n=109). Except for 23 adjuvant treatments [in Nutting *et al.* trial (5)], all patients received definitive chemoradiotherapy.


Xerostomia. Grade ≥ 2 toxicity analysis mainly demonstrated a benefit in favor of IMRT. The risk of acute (RR=0.71, 95%CI=0.59-0.86), 1-year (RR=0.45, 95%CI=0.31-0.65) and 2-year (RR=0.26, 95%CI=0.15-0.46) xerostomia was consistently and significantly reduced with the IMRT technique. Details are presented in Figure 2. Quality of evidence and strength of recommendation is summarized in Table II.

Survival outcomes. Compared to 3D-CRT, IMRT was not associated with a significant reduction in the risk of death (OR=0.70, 95%CI=0.39-1.24; *p*=0.22). There was no significant difference in LRC between IMRT and 3D-RT (OR=1.50, 95%CI=0.75-2.98; *p*=0.25). The χ^2 tests for heterogeneity of both comparisons showed no significant heterogeneity. Details are shown in Figure 3. The quality of evidence and strength of recommendations for survival outcomes are presented in Table II.

a



b



c



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Figure 2. Forest plot of acute (a), 1-year (b) and 2-year (c) xerostomia profile.

Discussion

Our results indicated that for HNSCC patients IMRT was superior to 3D-CRT in the xerostomia-related toxicity profile, while showed no benefit on OS and LRC rates. Data were statistically homogeneous and results were robust. IMRT was associated with decreased grade ≥2 acute xerostomia and there was a significant difference in late xerostomia between 3D-CRT and IMRT at 1-year and 2-years after treatment. Although the direct comparison

between IMRT and 3D-CRT did not show the superiority of IMRT regarding clinical outcomes, there was a slight increase in local recurrence events in the IMRT group. This higher proportion of events was small but it may have important implications for RT treatment planning. In fact, despite significant progress in the RT technique, the risk of loco-regional failure remains a challenge, mainly related to radio-resistant tumor areas and/or geographic miss (17). In order to improve accuracy in target definition to guarantee good outcomes in terms both of cure and toxicity, fusion of

Table I. Details of the included studies.

Author	ID_study	Population	Patients				Primary end point	Comments
			Total	IMRT	3D-CRT	Median FU		
Ghoshlaskar_2015 (8)	NCT652613	T1-3 N0-2b oral cavity, oropharynx, larynx* or hypopharynx	59	29	30	70 months	Xerostomia G \geq 2	Toxicity was graded using the RTOG scoring system
Gupta_2012 (9)	CTRI/2008/091/000045	T1-T3 N0-2b oropharynx, larynx* or hypopharynx	60	28	32	40 months	Xerostomia G \geq 2	Toxicity was assessed based on the RTOG scoring system
Nutting_2011 (5)	ISRCTN48243537	T1-4 N0-3 oropharynx or hypopharynx	94	47	47	44 months	Xerostomia G \geq 2	Acute side-effects were graded with NCICT. Late RT side-effects were assessed with LENT SOMA and RTOG scoring systems

*Except T1N0 glottic larynx. ID: Identifier; IMRT: intensity modulated radiotherapy; 3D-CRT: three dimensional conformal radiotherapy; FU: follow-up; NCT: number clinical trial; G: grade; RTOG: Radiation Therapy Oncology Group; CTRI: Clinical Trials Registry-India; ISRCTN: International Standard Randomized Controlled Trial; NCICT: National Cancer Institute Common Toxicity Criteria; LENT SOMA: Late Effects of Normal Tissues Subjective-Objective Management Analytic.


functional imaging examination at the contouring stage should be standardized (18).

To minimize indirectness, we restricted the analysis only to those trials that excluded nasopharynx cancer population. This can be explained clinically by nasopharyngeal cancer specific etiology, histology and natural history. It is not surprising that results on survival were not completely in agreement with the previous meta-analysis (4). Both LRC (HR=0.76, 95%CI=0.57-1.01) and OS (HR=0.70, 95%CI=0.57-0.88) benefits of IMRT compared to non-IMRT techniques were clearly driven by the nasopharyngeal series. Authors performed subgroup analysis stratified by primary tumor site and these results were comparable to our findings. There was no significant difference in LRC (HR=1.06, 95%CI=0.71-1.58) and OS (HR=0.85, 95%CI=0.63-1.15; $p=0.29$) between IMRT and non-IMRT (4). Also, xerostomia outcomes were not very different compared to our results, reporting a consistent IMRT benefit in salivary function over conventional techniques (4).


Despite the lower incidence, xerostomia remains a major clinical problem in the IMRT era and parotid-sparing is the recommended method to its prevention (5). But, the importance of the other salivary glands, including both mayor – submandibular and sublingual – and minor – upper/lower lip, buccal mucosa, postero-lateral hard palate, Von Ebner and Weber Blandin Nuhn – glands in quantity, quality and consistency of saliva production should be highlighted (19). In contrast to parotid glands, data on anatomic and dosimetric changes of the other salivary glands are scarce. In general, the

mucinous component is more radio-resistant than serous cells and thus a higher threshold dose is expected for submandibular and minor salivary glands than parotid glands. Reducing submandibular glands mean dose to 39 Gy resulted in gradually improved flow rates and patient-reported dry mouth symptoms, but it was associated to modest rise in the mean doses to surrounding structures, including parotid glands and swallowing structures (20). Probably, to maximize therapeutic gain, radiation dose should be optimized not only to parotids but to all salivary glands. Both major and minor salivary glands-sparing should be the key to preserve salivary function but this benefit could be achieved at the cost of disease control, resulting in increased risk of marginal failures. The best balance between tumor target coverage and salivary glands-sparing approach requires further clinical investigation before any definitive conclusion can be reached. This meta-analysis did not attempt to evaluate the other potential xerostomia-related complications, such as late radiation-associated dysphagia (RAD) and osteoradionecrosis (ORN). Data were not suitable for a cumulative analysis. Obviously late RAD and ORN are also influenced by the direct irradiation of the swallowing apparatus and segments of mandible, respectively. Due to its ballistic characteristics, IMRT exposed large volume of these normal structures to higher doses than previous less conformal treatment, resulting in the highest risk of their damage (21). To reduce this detrimental RT effect, over the years, specific dose-volume parameters have been proposed in addition to the standard maximum dose constraints (22-23). But, at present, these specific constraints have been definitively

a



b



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Figure 3. Forest plot of overall survival (a) and loco-regional control (b).

introduced in daily clinical practice. Optimizing future multimodal treatment for HNSCC requires collection of detailed toxicity data, both clinician-reported and patient-reported outcomes (PROs) (24). A plausible discordance between clinician and patient reporting of toxicity should be considered. For instance, patients in disease remission may have accepted moderate xerostomia, adjusting their expectation accordingly; whereas in other cases the same clinician-reported moderate RT toxicity may drastically impact patient quality of life. PROs inclusion in the rigorous toxicity reporting methods should be useful to provide additional information in decision making process.

In general, this meta-analysis provided additional contribution to published literature data, showing the superiority of IMRT over 3D-CRT in HNSCC patients management. Our main aim was to achieve the highest quality of research. The current meta-analysis was not a literature abstract-based meta-analysis. Conference

proceedings were not considered, due to the lack of statistical quality details and << given the variable evidence of concordance between conference abstracts and their subsequent full-text publications >>, as advised by the agency for healthcare research and quality (AHRQ) (25). All included studies were randomized clinical trials and addressed the same primary end-point and similar methods were used to evaluate xerostomia. The major limitation of the analysis was the relative limited number of trials. But trials' short time accrual period – ranged from 2003 to 2008 – may add homogeneity to results, even though the included trials did not record human papillomavirus (HPV) status. At present HPV evaluation is paramount in HNSCC risk stratification (26). Its positive status discriminates excellent prognosis disease from worst outcome in HPV-negative tumors. Future studies specifically designed and powered to test the benefit of IMRT over 3D-CRT based on HPV stratification and salivary glands-sparing RT approach would provide more

Table II. *Quality of evidence and strength of recommendation: Summary of findings. Intensity modulated radiotherapy (IMRT) compared to three-dimensional conformal radiotherapy (3D-CRT) in head and neck squamous cell carcinoma.*

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with 3D-CRT	Risk with IMRT				
OS assessed with: Kaplan–Meier product limit	40 per 100	32 per 100 (21 to 46)	OR=0.70 (0.39-1.24)	213 (3 RCTs)	⊕⊕⊕○ MODERATE ^{a,b,c}	
LRC assessed with: Kaplan–Meier product limit	17 per 100	24 per 100 (14 to 38)	OR=1.50 (0.75-2.98)	213 (3 RCTs)	⊕⊕⊕○ MODERATE ^{a,b,c}	
Acute xerostomia ≥2 assessed with: Cumulative incidence	76 per 100	54 per 100 (45 to 65)	RR=0.71 (0.59-0.86)	213 (3 RCTs)	⊕⊕⊕○ MODERATE ^{a,b,c}	
1-y xerostomia ≥2 assessed with: Cumulative incidence	59 per 100	27 per 100 (18 to 39)	RR=0.45 (0.31-0.65)	169 (3 RCTs)	⊕⊕⊕○ MODERATE ^{a,b,c}	
2-y xerostomia ≥2 assessed with: Cumulative incidence	55 per 100	14 per 100 (8 to 25)	RR=0.26 (0.15-0.46)	140 (3 RCTs)	⊕⊕⊕○ MODERATE ^{a,b,c}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). CI: Confidence interval; LRC: loco-regional control; OR: odds ratio; OS: overall survival; RR: risk ratio. GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect. ^aMissing outcome data balanced across groups; ^bdetection bias; ^cmay not be downgraded.

conclusive evidence in term of clinical outcomes. We hoped to draw attention to the paucity of good quality data on this topic and improve the quality of future research.

Conclusion

This meta-analysis confirmed the efficacy of IMRT to reduce moderate to severe xerostomia onset, but cannot determine meaningful survival advantages in HNSCC. Prospective randomized trials aimed to compare IMRT *versus* 3D-CRT in an adequate homogeneous HNSCC population, with survival and main RT-related toxicities endpoints are necessary to definitely confirm IMRT value.

Conflicts of Interest

All Authors declare that they have no conflicts of interest that may be relevant to the contents of this manuscript.

Authors' Contributions

FDF, NP, PP have made substantial contributions to the conception of the work, the acquisition and interpretation of data and have drafted the work; OB, VT and AP have substantively revised it. All Authors have approved the submitted version.

References

- 1 National Comprehensive Cancer Network (NCCN). Guidelines Head and Neck Cancers, Version 2.2018. Available at: <http://www.nccn.org> (Last accessed 21/11/2019)
- 2 De Felice F, Musio D, Terenzi V, Valentini V, Cassoni A, Tombolini M, De Vincentiis M and Tombolini V: Treatment improvement and better patient care: which is the most important one in oral cavity cancer? *Radiat Oncol* 9: 263, 2014. PMID: 25479896. DOI: 10.1186/s13014-014-0263-x
- 3 Marta GN, Silva V, de Andrade Carvalho H, de Arruda FF, Hanna SA, Gadia R, da Silva JL, Correa SF, Vita Abreu CE and Riera R: Intensity-modulated radiation therapy for head and neck cancer: systematic review and meta-analysis. *Radiother Oncol* 110(1): 9-15, 2014. PMID: 24332675. DOI: 10.1016/j.radonc.2013.11.010
- 4 Gupta T, Kannan S, Ghosh-Laskar S and Agarwal JP: Systematic review and meta-analyses of intensity-modulated radiation therapy *versus* conventional two-dimensional and/or three-dimensional radiotherapy in curative-intent management of head and neck squamous cell carcinoma. *PLoS One* 13(7): e0200137, 2018. PMID: 29979726. DOI: 10.1371/journal.pone.0200137
- 5 Nutting CM, Morden JP, Harrington KJ, Urbano TG, Bhide SA, Clark C, Miles EA, Miah AB, Newbold K, Tanay M, Adab F, Jefferies SJ, Scrase C, Yap BK, A'Hern RP, Sydenham MA, Emson M, Hall E and PARSPORT trial management group: Parotid-sparing intensity modulated *versus* conventional radiotherapy in head and neck cancer (PARSPORT): a phase 3 multicentre

- randomised controlled trial. *Lancet Oncol* 12(2): 127-136, 2011. PMID: 21236730. DOI: 10.1016/S1470-2045(10)70290-4
- 6 Rosenthal DI, Chambers MS, Fuller CD, Rebueno NC, Garcia J, Kies MS, Morrison WH, Ang KK and Garden AS: Beam path toxicities to non-target structures during intensity-modulated radiation therapy for head and neck cancer. *Int J Radiat Oncol Biol Phys* 72(3): 747-755, 2008. PMID: 18455324. DOI: 10.1016/j.ijrobp.2008.01.012
- 7 Rathod S, Gupta T, Ghosh-Laskar S, Murthy V, Budrukkar A and Agarwal J: Quality-of-life (QOL) outcomes in patients with head and neck squamous cell carcinoma (HNSCC) treated with intensity-modulated radiation therapy (IMRT) compared to three-dimensional conformal radiotherapy (3D-CRT): evidence from a prospective randomized study. *Oral Oncol* 49(6): 634-642, 2013. PMID: 23562564. DOI: 10.1016/j.oraloncology.2013.02.013
- 8 Ghosh-Laskar S, Yathiraj PH, Dutta D, Rangarajan V, Purandare N, Gupta T, Budrukkar A, Murthy V, Kannan S and Agarwal JP: Prospective randomized controlled trial to compare 3-dimensional conformal radiotherapy to intensity-modulated radiotherapy in head and neck squamous cell carcinoma: Long-term results. *Head Neck* 38: E1481-1487, 2016. PMID: 26561342. DOI: 10.1002/hed.24263
- 9 Gupta T, Agarwal J, Jain S, Phurailatpam R, Kannan S, Ghosh-Laskar S, Murthy V, Budrukkar A, Dinshaw K, Prabhaskar K, Chaturvedi P and D'Cruz A: Three-dimensional conformal radiotherapy (3D-CRT) versus intensity modulated radiation therapy (IMRT) in squamous cell carcinoma of the head and neck: a randomized controlled trial. *Radiation Oncol* 104(3): 343-348, 2012. PMID: 22853852. DOI: 10.1016/j.radonc.2012.07.001
- 10 Moher D, Liberati A, Tetzlaff J, Altman DG and PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 62(10): 1006-1012, 2009. PMID: 25554246. DOI: 10.1186/2046-4053-4-1
- 11 Lefebvre C, Manheimer E and Glanville J: Chapter 6: Searching for studies. In: Higgins J, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available at: www.cochrane-handbook.org (Last accessed 21/11/2019)
- 12 Cox JD, Stetz J and Pajak TF: Toxicity criteria of the Radiation Therapy Oncology Group (RTOG) and the European Organization for Research and Treatment of Cancer (EORTC). *Int J Radiat Oncol Biol Phys* 31(5): 1341-1346, 1995. PMID: 7713792. DOI: 10.1016/0360-3016(95)00060-C
- 13 Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, Norris S, Falck-Ytter Y, Glasziou P, DeBeer H, Jaeschke R, Rind D, Meerpohl J, Dahm P and Schünemann HJ: GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 64(4): 383-394, 2011. PMID: 21195583. DOI: 10.1016/j.jclinepi.2010.04.026
- 14 Cochrane. Available at: <http://www.cochrane.org> (Last accessed 21/11/2019)
- 15 Tierney JF, Stewart LA, Ghersi D, Burdett S and Sydes MR: Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials* 8: 16, 2007. PMID: 17555582. DOI: 10.1186/1745-6215-8-16
- 16 Higgins JPT, Thompson SG, Deeks JJ and Altman DG: Measuring inconsistency in meta-analyses. *BMJ* 327: 557-560, 2003. PMID: 12958120. DOI: 10.1136/bmj.327.7414.557
- 17 De Felice F, Thomas C, Barrington S, Pathmanathan A, Lei M and Urbano TG: Analysis of loco-regional failures in head and neck cancer after radical radiation therapy. *Oral Oncol* 51(11): 1051-1055, 2015. PMID: 26506048. DOI: 10.1016/j.oraloncology.2015.08.004
- 18 De Felice F, Musio D, Bulzonetti N, Maghella F and Tombolini V: Target volume delineation based on diffusion-weighted magnetic resonance imaging for locally advanced head and neck cancer. *Anticancer Res* 36(8): 4181-4185, 2016. PMID: 27466529.
- 19 De Felice F, Musio D, Terenzi V, Valentini V, Cassoni A, Tombolini M, De Vincentiis M and Tombolini V: Treatment improvement and better patient care: which is the most important one in oral cavity cancer? *Radiat Oncol* 9: 263, 2014. PMID: 25479896. DOI: 10.1186/s13014-014-0263-x
- 20 Murdoch-Kinch CA, Kim HM, Vineberg KA, Ship JA and Eisbruch A: Dose-effect relationships for the submandibular salivary glands and implications for their sparing by intensity modulated radiotherapy. *Int J Radiat Oncol Biol Phys* 72(2): 373-382, 2008. PMID: 18337023. DOI: 10.1016/j.ijrobp.2007.12.033
- 21 De Felice F, Musio D and Tombolini V: Osteoradionecrosis and intensity modulated radiation therapy: An overview. *Crit Rev Oncol Hematol* 107: 39-43, 2016. PMID: 27823650. DOI: 10.1016/j.critrevonc.2016.08.017
- 22 De Felice F, Thomas C, Patel V, Connor S, Michaelidou A, Sproat C, Kwok J, Burke M, Reilly D, McGurk M, Simo R, Lyons A, Oakley R, Jeannon JP, Lei M and Urbano TG: Osteoradionecrosis following treatment for head and neck cancer and the effect of radiotherapy dosimetry: the Guy's and St Thomas' Head and Neck Cancer Unit experience. *Oral Surg Oral Med Oral Pathol Oral Radiol* 122(1): 28-34, 2016. PMID: 27039003. DOI: 10.1016/j.oooo.2016.01.007
- 23 De Felice F, de Vincentiis M, Luzzi V, Magliulo G, Tombolini M, Ruoppolo G and Polimeni A: Late radiation-associated dysphagia in head and neck cancer patients: evidence, research and management. *Oral Oncol* 77: 125-130, 2018. PMID: 29362118. DOI: 10.1016/j.oraloncology.2017.12.021
- 24 Gilbert A, Ziegler L, Martland M, Davidson S, Efficace F, Sebagn Montefiore D and Velikova G: Systematic review of radiation therapy toxicity reporting in randomized controlled trials of rectal cancer: A comparison of patient-reported outcomes and clinician toxicity reporting. *Int J Radiat Oncol Biol Phys* 92(3): 555-567, 2015. PMID: 26068490. DOI: 10.1016/j.ijrobp.2015.02.021
- 25 Balshem H, Stevens A, Ansari M, Norris S, Kansagara D, Shamliyan T, Chou R, Chung M, Moher D and Dickersin K: Finding grey literature evidence and assessing for outcome and analysis reporting biases when comparing medical interventions: AHRQ and the effective health care program. Agency for healthcare research and Quality, AHRQ methods for effective health care, 2013. PMID: 24404628.
- 26 Lydiatt WM, Patel SG, O'Sullivan B, Brandwein MS, Ridge JA, Migliacci JC, Loomis AM and Shah JP: Head and Neck cancers-major changes in the American Joint Committee on cancer eighth edition cancer staging manual. *CA Cancer J Clin* 67(2): 122-137, 2017. PMID: 28128848. DOI: 10.3322/caac.21389

Received November 16, 2019

Revised November 21, 2019

Accepted November 26, 2019