

## ● REVIEW

# Transcriptional regulation of adult neural stem/progenitor cells: tales from the subventricular zone

Giancarlo Poiana, Roberta Gioia, Serena Sineri, Silvia Cardarelli, Giuseppe Lupo\*, Emanuele Cacci  
Department of Biology and Biotechnology "C. Darwin", Sapienza University of Rome, Rome, Italy

## Abstract

In rodents, well characterized neurogenic niches of the adult brain, such as the subventricular zone of the lateral ventricles and the subgranular zone of the hippocampus, support the maintenance of neural/stem progenitor cells (NSPCs) and the production of new neurons throughout the lifespan. The adult neurogenic process is dependent on the intrinsic gene expression signatures of NSPCs that make them competent for self-renewal and neuronal differentiation. At the same time, it is receptive to regulation by various extracellular signals that allow the modulation of neuronal production and integration into brain circuitries by various physiological stimuli. A drawback of this plasticity is the sensitivity of adult neurogenesis to alterations of the niche environment that can occur due to aging, injury or disease. At the core of the molecular mechanisms regulating neurogenesis, several transcription factors have been identified that maintain NSPC identity and mediate NSPC response to extrinsic cues. Here, we focus on REST, Egr1 and Dbx2 and their roles in adult neurogenesis, especially in the subventricular zone. We review recent work from our and other laboratories implicating these transcription factors in the control of NSPC proliferation and differentiation and in the response of NSPCs to extrinsic influences from the niche. We also discuss how their altered regulation may affect the neurogenic process in the aged and in the diseased brain. Finally, we highlight key open questions that need to be addressed to foster our understanding of the transcriptional mechanisms controlling adult neurogenesis.

**Key Words:** adult neurogenesis; aging; extracellular signaling; gene regulation; neural stem/progenitor cells; transcription factors

\*Correspondence to:

Giuseppe Lupo,  
giuseppe.lupo@uniroma1.it.

orcid:  
0000-0001-7218-4003  
(Giuseppe Lupo)

doi: 10.4103/1673-5374.280301

Received: October 16, 2019

Peer review started: October 24, 2019

Accepted: November 24, 2019

Published online: April 3, 2020

## Introduction

Contrary to the long held "no new neurons after birth" dogma, neurogenesis occurs in adult rodents and possibly other mammals, sustained by pools of neural stem/progenitor cells (NSPCs) that reside in specialized brain niches (Ming and Song, 2011).

In mice, the subventricular zone (SVZ), adjacent to the walls of the lateral ventricles, is the adult neurogenic niche containing the largest number of NSPCs. Specialized astrocytes of the SVZ that contact the lateral ventricle with an apical process extending through the ependymal layer, known as type B cells, have been identified as the neurogenic stem cells of this niche (Fuentelba et al., 2015). These cells can exist in a quiescent or in an activated state. In the latter case, they proliferate to self-renew and/or generate transit amplifying progenitor cells (type C cells or TAPs). Type C cells give rise to neuroblasts (type A cells), which migrate to the olfactory bulb, where they terminally differentiate into distinct types of interneurons that are important for odour discrimination (Kohwi et al., 2007; Brill et al., 2009). Another well described niche is the subgranular zone (SGZ) of the hippocampal dentate gyrus, where NSPCs continuously produce neurons that provide additional plasticity to the adult brain and support several hippocampus-dependent cognitive functions, such as spatial learning and contextual memory capabilities (Gonçalves et al., 2016).

Physical exercise, learning, odour discrimination and enriched environment, among others, are physiological stimuli regulating NSPCs and their progeny in a niche-spe-

cific manner. External stimuli might affect neurogenesis by changing the levels of several extracellular molecules (e.g., neurotransmitters, morphogens, growth factors, cytokines and adhesion/repulsion molecules) in the niche milieu. For example, the use of optogenetics and chemogenetics to activate or inhibit medial septum  $\gamma$ -aminobutyric acid-ergic neurons projecting to the dentate gyrus has indicated that medial septum neuron activity maintains neural stem cell quiescence and controls hippocampal neurogenesis through  $\gamma$ -aminobutyric acid release (Bao et al., 2017). Furthermore, it has been proposed that physical exercise-mediated release of acetylcholine by the septal cholinergic system increases NSPC proliferation in mice, suggesting that acetylcholine may act as an extrinsic signal mediating the positive effect of voluntary running on neurogenesis (Itou et al., 2011).

In rodents, aging also has a major impact on the adult neurogenic process, leading to a reduction of the proliferating NSPC pool and of new neuron production (Lupo et al., 2019). The depletion of proliferating NSPCs in the aged niches may be caused by the progressive consumption of the NSPC reservoir due to terminal differentiation (Encinas et al., 2011), by a decreased ability of quiescent stem cells to undergo activation (Kalamakis et al., 2019), and by age-related changes in different NSPC features, such as their pattern of cell division (symmetric *versus* asymmetric), cell cycle length or survival (Lupo et al., 2019). At the molecular level, the negative regulation of neurogenesis in the aged brain crucially involves age-associated alterations in the niche environment impinging on NSPCs, including the decreased ac-

tivation of signaling pathways promoting NSPC proliferation and neuronal differentiation, such as the epidermal growth factor (EGF) and Wnt pathways, and the increased stimulation of pathways inhibiting neurogenesis, such as transforming growth factor  $\beta$  (TGF- $\beta$ ) signaling and inflammatory pathways (Pineda et al., 2013; Dulken et al., 2019; Kalamakis et al., 2019; Kase et al., 2019). In turn, the extracellular niche modifications due to physiological stimuli or aging induce changes in the gene expression programs controlling NSPC proliferation and differentiation. In this context, transcription factors (TFs) serve as key intracellular hubs in NSPCs, integrating the extrinsic pathways acting in the niche and regulating the intrinsic transcriptional networks driving the neurogenic process.

A number of TFs involved in NSPC regulation, neuronal differentiation and maturation, and synaptic integration have been identified. Several of these TFs are implicated in the control of both adult and embryonic neurogenesis, although their mode of action and expression levels may change at different developmental stages (Götz et al., 2016). In this review, we highlight the regulation of adult neurogenesis by three TFs that we have recently investigated, namely the repressor element 1-silencing transcription factor (REST), Egr1 and Dbx2. We discuss their function in controlling NSPC proliferation and differentiation and the transcriptional networks that they regulate during neurogenesis, with particular focus on the adult SVZ niche. Furthermore, we describe how different extrinsic influences can modulate the activity of REST, Egr1 and Dbx2 in NSPCs or differentiating neurons, suggesting that these TFs might act as regulatory nodes in the molecular pathways controlling neurogenesis in physiological conditions, aging or disease.

## Search Strategy and Selection Criteria

Studies cited in this review were found on the PubMed database, between September and November 2019, using the search terms: adult neurogenesis, neural stem cells, neural progenitors, subventricular zone, transcription factors, REST, Egr1, Dbx2, and various combinations of the above terms.

## The Repressor Element 1-Silencing Transcription Factor Supports Adult Neurogenesis with Cell Type-Specific and Context-Dependent Mechanisms

REST is a zinc finger protein that binds a highly conserved 21–23 bp DNA sequence (REST binding site; RE1 site) and regulates neurogenesis by repressing the transcription of neuronal specific genes (Chong et al., 1995; Schoenherr and Anderson, 1995). Reducing REST activity reconfigures chromatin accessibility, allowing neuronal gene expression and the direct conversion of non-neuronal cells (e.g., astrocytes or fibroblasts) into neurons (Masserdotti et al., 2015; Drouin-Ouellet et al., 2017). With the advent of genome-wide analyses, several thousand RE1 sites linked to genes that encode both neuronal and non-neuronal proteins have been

identified (Bruce et al., 2004; Mortazavi et al., 2006; Johnson et al., 2008). Binding of REST at the RE1 sites allows the recruitment of CoREST and mSin3A, and the assembly of a repressor complex including histone deacetylases, histone methyltransferases and the histone demethylase LSD1. As part of this complex, REST controls gene expression through an epigenetic remodelling mechanism (Ballas et al., 2005).

REST functions as an intrinsic factor controlling neural development. Its levels are elevated in neural progenitors, where REST represses neuronal genes, and decrease when neural progenitors differentiate into neurons (Chen et al., 1998; Ballas et al., 2005; Mandel et al., 2011; Yang et al., 2012; Nechiporuk et al., 2016). Transient overexpression of REST during the neurogenic phase of brain development blocked progenitor migration and neuronal differentiation, indicating that REST downregulation is crucial to allow neurogenesis (Mandel et al., 2011). In addition, REST keeps neuronal genes protected from DNA damage in apical cortical neural progenitors during the S phase of the cell cycle, avoiding accumulation of chromosomal abnormalities that would either commit cells to apoptosis and result in microcephaly or, in combination with the loss of p53, trigger the formation of brain tumors (Nechiporuk et al., 2016). These observations highlight the importance of properly regulating the timing and the levels of REST activity during brain development.

REST is also expressed in adult NSPCs and its inactivation affects neurogenesis, leading to NSPC depletion and decreased granule neuron generation in the adult hippocampus (Kuwabara et al., 2004; Gao et al., 2011; Mukherjee et al., 2016). To better dissect its role in specific NSPC subpopulations, REST was deleted in quiescent hippocampal neural stem cells (qNSC) by injections of a Cre-p2A-mCherry lentivirus expressed under the control of the human glial fibrillary acidic protein promoter into REST fl/fl mice, finding that REST prevented qNSC activation. The conditional inactivation of REST in the TAP population caused its decrease and the activation of the neuronal differentiation program. These findings indicate that REST acts cell autonomously to support the maintenance of both the qNSC and the TAP pools by preventing their progression along the neurogenic lineage (Mukherjee et al., 2016).

Genome-wide ChIP-seq analysis of REST binding sites, combined with RNA-seq profiling of genes deregulated by REST knockdown, showed that the majority of the REST targets were neuronal genes, suggesting that their REST-dependent repression may be crucial to preserve the qNSC and TAP populations (Mukherjee et al., 2016). It would be interesting to overexpress neuronal REST target genes in qNSCs or TAPs to verify whether this is sufficient to promote their lineage progression. Notably, several of the REST-controlled neuronal genes were differentially expressed in qNSCs and TAPs, indicating that the transcriptional response of REST target genes is cell type-dependent and stage-dependent, as previously proposed (Bruce et al., 2004; Hohl and Thiel, 2005; Johnson et al., 2008; Noh et al., 2012; Lu et al., 2014; McClelland et al., 2014; Mukherjee et al., 2016). Since REST levels are similar in qNSCs and TAPs, the differential regu-


lation of neuronal target genes in these cell populations may involve context-dependent interactions of REST with other TFs (Mukherjee et al., 2016).

This analysis also identified unique REST targets involved in ribosome biogenesis (e.g., the genes coding for nucleophosmin 1 (*Npm1*) and ribosomal protein L4 (*Rpl4*) and cell cycle regulation (e.g., *Mms22l*), whose overexpression either alone or in combination stimulated proliferation in quiescent cells. These data suggest that REST-regulated genes implicated in ribosome biogenesis and cell cycle are important players in controlling the transition between quiescent and activated stem cells (Mukherjee et al., 2016). They also strengthen the idea that REST is a multifaceted protein with context-dependent and cell type-specific roles, highlighting the importance of studying REST in different cell types, and contexts, to fully unveil its function in adult neurogenesis.

Along with other groups, we have investigated the role of REST in the SVZ by means of *in vitro* NSPC cultures. SVZ-derived NSPCs in which REST function was impaired generated a higher percentage of neurons when induced to differentiate. Notably, these cells showed reduced neurosphere formation and increased neuronal differentiation in comparison with control cultures even when maintained in culture media supplemented with growth factors, a condition suitable for NSPC expansion and the maintenance of multipotency (Gao et al., 2011; Soldati et al., 2015). Altogether, these data indicate that REST/NRSF is required for the self-renewal of adult SVZ NSPCs and that REST directly controls the neurogenic lineage program *in vitro* (Figure 1). This conclusion is supported by the observation that NSPCs isolated from the SVZ of 3 week-old REST/NRSF<sup>loxP</sup> mice and then infected with an adenovirus expressing Cre/GFP showed activation of REST target genes and reduced binding of REST at the RE1 sequence of proneural genes, such as

*Ascl1* and *NeuroD1*. *NeuroD1* upregulation following REST inactivation correlated with increased histone H4 acetylation, increased H3K4me2 and decreased H3K27me3. This is in accordance with the role of REST in the recruitment of histone methylases/demethylases and REST-dependent acetylation/deacetylation (Gao et al., 2011; Mukherjee et al., 2016).

We identified REST target genes in SVZ NSPCs by combining a transcriptomic analysis of REST deficient cells together with a ChIP-seq profiling of REST binding sites throughout the genome. These analyses provided a comprehensive REST-regulated gene expression resource and indicated that REST direct target genes were significantly enriched for pathways associated with nervous system development/function and disease (Soldati et al., 2015). Of note, a few genes identified as REST targets in our study (e.g., *Bcl11B*, *Cels3*, *BMP6*) had been previously annotated as stem cell enriched genes by a transcriptome analysis performed on NSPCs freshly isolated from the SVZ of adult mice by means of FACS-based purification (Beckervordersandforth et al., 2010). Among the genes directly regulated by REST, we found that the expression of a bone morphogenetic protein (BMP) family member, *BMP6*, was increased both at the mRNA and protein level upon REST knockdown. BMPs are important regulators of neurogenesis and gliogenesis (Panchision and McKay, 2002). In addition, BMP pathway activation is implicated in the control of neural stem cell quiescence in the adult hippocampus (Mira et al., 2010). We showed that administration of exogenous BMP6, at a concentration comparable to that released by NSPCs upon REST knockdown, elicited an anti-proliferative and pro-differentiative effect on NSPCs, suggesting that REST-mediated control of BMP signaling may regulate NSPC properties in the SVZ (Soldati et al., 2015).



**Figure 1 Proposed model of the REST-dependent control of NSPC proliferation and neuron differentiation/survival.**

Decreased levels of REST activity in NSPCs derepress the transcription of genes associated with the inhibition of NSPC proliferation and the stimulation of differentiation, thus promoting NSPC progression along the neurogenic lineage. In differentiated neurons, REST levels and/or cellular localization change as a result of different physiological or pathological events. These alterations of REST activity may exert neurotoxic or neuroprotective effects, with context-dependent mechanisms. NSPCs: Neural stem/progenitor cells; REST: repressor element 1-silencing transcription factor.

In other studies, secretory genes have been identified as a class of REST target genes (Kim et al., 2015). Among them, SCG2 encodes for Secretogranin II, which has been proposed to act as an extrinsic molecule mediating the effects of REST knockdown in promoting neuronal differentiation and enhanced neuritic outgrowth in hippocampal NSPCs. Notably, these effects were observed both in the cells transfected with a shREST-mCherry plasmid and in the non-transfected cells within the same cultures, suggesting that REST controls neuronal differentiation in a paracrine manner. Gain and loss of function experiments allowed the authors to identify SCG2 as part of a non-cell autonomous signaling pathway, downstream of REST, which is necessary for neuronal differentiation in hippocampal NSPCs. In future work, it will be interesting to explore whether similar signals also act in the adult SVZ both *in vitro* and *in vivo*.

To date, there are few published *in vivo* studies addressing REST function in the SVZ. One of them performed a conditional knockout of REST in NSPCs using Sox1-Cre/+; Rest<sup>2lox/2lox</sup> mice (Aoki et al., 2012). The authors showed that neurogenesis was affected neither in the adult SVZ, nor in the dentate gyrus, at variance with a previous study reporting altered neurogenesis after the conditional deletion of REST in the adult hippocampus of Nestin-CreERT2/R26R-YFP mice injected with tamoxifen (Gao et al., 2011). This discrepancy might be due to the different strategies used to delete REST *in vivo*. REST knockout in Sox1-Cre/+; Rest<sup>2lox/2lox</sup> adult mice starts at an early embryonic stage, since Sox1 expression begins at embryonic day 7.5–8.5 in the neural tube, whereas abrogation of REST function in Nestin-CreERT2/R26R-YFP mice was induced in the adult brain. Compensatory mechanisms activated by the early embryonic deletion of REST may mask its role in the adult brain and explain the lack of an obvious neurogenic phenotype. Notably, it has been reported that REST knockout animals carrying a deletion of exon 2 or 4 still produce the C-terminal or the N-terminal REST repressor domains (Gao et al., 2011; Aoki et al., 2012; Mukherjee et al., 2016), which might retain repressive functions (Andrés et al., 1999; Naruse et al., 1999; Grimes et al., 2000; Roopra et al., 2000; Ballas et al., 2001), questioning the conclusion that REST function is dispensable for adult neurogenesis (Aoki et al., 2012). The use of classical knockout REST models, complemented with mice containing a conditional gene trap cassette in an intron of the REST gene that terminates transcription upstream of the initiator codon (Nechiporuk et al., 2016), might help to unravel the role of REST in adult neurogenesis.

It will also be important to shed light on the molecular mechanisms regulating REST expression and function under both physiological and pathological conditions. REST is also expressed, although at low levels, in specific neuronal subpopulations at embryonic stages (Nechiporuk et al., 2016), during postnatal development (Rodenas-Ruano et al., 2012), and in the adult brain (Kuwabara et al., 2004; Gao et al., 2011), suggesting that REST function may be required even in differentiated neurons (Yeo et al., 2009; Rodenas-Ruano et al., 2012). Accordingly, mice conditionally overexpressing

REST in cells expressing dopamine receptor 2, including striatal neurons, showed impaired spontaneous locomotion and displayed some of the alteration associated with human Huntington's disease (Lu et al., 2018). Importantly, the expression of REST and of its target genes is altered in post-mitotic neuronal subpopulations in pathological conditions, such as ischemia (Calderone et al., 2003; Noh et al., 2012; Kaneko et al., 2014), seizures (Palm et al., 1998), Huntington's disease (Zuccato et al., 2003), and Alzheimer's disease (AD) (Lu et al., 2014) (**Figure 1**). Elevated REST levels have been associated with striatal neurotoxicity in Huntington's disease. Huntingtin mutations destroy REST interactions and retention in the cytoplasm, favouring REST aberrant accumulation in the nucleus and the repression of specific target genes, such as the one coding for brain-derived neurotrophic factor, a neurotrophin essential for striatal neurons survival (Zuccato et al., 2003). *Vice versa*, loss of nuclear REST occurs in AD patients and AD animal models. In this context, cognitive impairment and neurodegeneration have been correlated with the derepression of REST target genes involved in cell death (e.g., BAX and DAXX) and AD (e.g., Presenilin2). Compared to control neurons, REST-deficient neurons treated with A $\beta$ <sub>42</sub> showed increased degeneration and cell death, which could be rescued by lentiviral transduction of REST. Moreover, REST overexpression in the SH-SY5Y human neural cell line decreased cell death upon oxidative stress. These observations suggest that, in differentiated neurons, REST may exert a neuroprotective function (Lu et al., 2014).

Although an impairment of adult neurogenesis has been associated with several neurodegenerative diseases (Steiner et al., 2006), there are very few studies addressing the effects of changes in REST function on the response of NSPCs to brain insults. In a recent *in vitro* study on NSPCs derived from sporadic AD patients or in pluripotent stem cells carrying an AD-associated APO4 allele, reduced nuclear REST localization and decreased REST-RE1 binding have been described (Meyer et al., 2019). The transcriptomic analysis of NSPC models of AD showed modifications in REST regulated gene networks, loss of self-renewal capability, premature neuronal differentiation, accelerated synapse formation and increased electrical excitability, as compared with control cells, indicating that REST dysfunction emerges in NSPCs and persists in neurons (Meyer et al., 2019). Since REST knockout triggers a depletion of the NSPC pool (Gao et al., 2011; Mukherjee et al., 2016), it is tempting to speculate that an impaired neurogenesis due to an altered REST function may contribute to the cognitive decline associated with AD.

Finally, high levels of REST are associated with enhanced proliferation in several neural cell tumours. In particular, elevated REST expression has been detected in medulloblastoma, neuroblastoma and glioblastoma and its inhibition promotes cell cycle exit and differentiation of cancer cells (Su et al., 2006; Singh et al., 2011; Conti et al., 2012). These data suggest that REST plays a key role in the self-renewal of both normal NSPCs and malignant neural cancer cells, thus representing a candidate target for therapeutic interventions in

multiple neuropathological contexts.

## **Egr1 and Epidermal Growth Factor Signaling in the Regulation of Adult Neural Stem/Progenitor Cell Proliferation and Differentiation**

*Egr1* is an immediate early gene defined as an “early growth response gene” (Sukhatme et al., 1988), also known under other names and acronyms, such as *Krox24*, *Zif268*, *Tis8* and *NGFI-A*. It encodes for a zinc finger TF that binds to the DNA sequence GCG(G/T)GGGCG (Crosby et al., 1991), which is present at the promoter of a large number of genes involved in a wide variety of physiological and pathological cell processes.


The modulation of its activity levels depends on several extracellular signals, such as tumour necrosis factor  $\alpha$ , TGF- $\beta$ , EGF and platelet-derived growth factor A and B (Khachigian et al., 1995; Liu et al., 1996; Yao et al., 1997; Kaufmann and Thiel, 2001). Intracellular Ca<sup>2+</sup>-dependent pathways also regulate *Egr1* expression and its biological functions (Thiel et al., 2010). *Egr1* function is also modulated by intracellular co-activators, such as CREB-binding protein and p300 (Silverman et al., 1998), and co-repressors, such as NGFI-A binding protein 1 and 2, which bind to a specific *Egr1* domain and block the activation of its target genes (Thiel et al., 2000). In turn, *Egr1* regulates several tumor suppressors, such as TGF- $\beta$ 1, PTEN, p53 and fibronectin (Baron et al., 2006). *Egr1* protein levels, when detectable, are generally low and depend on the analyzed cell type and developmental stage; they may rapidly increase or decrease in physiological and pathological or stressful states, possibly as a consequence of a rapid turnover. *Egr1* activity is also dependent on its phosphorylation state, which is regulated through the phosphorylation of *Egr1* Ser<sup>26</sup> by the extracellular signal-regulated kinase-1 (ERK) (Gregg and Fraizer, 2011; Santiago et al., 2019). In particular, a role for EGF and its receptor (EGFR) in controlling *Egr1* phosphorylation via the ERK pathway has been well characterized (Alagappan et al., 2013).

In the nervous system, *Egr1* expression and activity has been demonstrated in the adult hippocampus, where it is involved in synaptic plasticity, learning, long term potentiation and memory consolidation (Jones et al., 2001; Cheval et al., 2012). *Egr1* role in the control of cell proliferation has been described in several cell types, including astrocytes (Biesiada et al., 1996). As demonstrated in PC12 pheochromocytoma cells (Levkovitz and Baraban, 2001), and in a mouse neuroblastoma cell line (N18TG2) (Salani et al., 2009), *Egr1* also regulates neuronal differentiation. Several *Egr1* transcriptional targets have been identified in PC12 cells, used as an *in vitro* model of neuronal differentiation (Adams et al., 2017). Nerve growth factor (NGF) signaling via the NGF receptor TrkA promotes PC12 differentiation (Segal and Greenberg, 1996), whereas EGF stimulates PC12 proliferation (Huff et al., 1981). Both NGF and EGF activate the ERK signaling pathway, but a short (30–60 minutes) EGF stimulation promotes proliferation, whereas NGF treatment induces

a longer lasting stimulation (4–6 hours) of ERK signaling, that ultimately leads to differentiation. In this context, *Egr1* cooperates with CREB and AP-1 to activate genes involved in the NGF-dependent differentiation of PC12 cells. This cohort of *Egr1* targets includes several genes implicated in neural development, such as the gene encoding for p35, which is necessary for Cdk5 activation and neurite outgrowth in response to NGF; *Vgf* and *Hbegf*, which encode for neurotrophic factors; *Kdm6b/Jmjd3* and *Kctd11*, which contribute to the differentiation of stem cells to neurons; *Sema6a*, which is involved in axon guidance; and *Arc*, which plays a role in synaptic plasticity, learning and memory (Adams et al., 2017). It will be important to investigate whether these genes are also regulated by *Egr1* in differentiating NSPCs in the *in vivo* neurogenic niches.

We have studied the role of *Egr1* in controlling the proliferation and differentiation of adult mouse SVZ NSPCs using *in vitro* culture systems, in which SVZ NSPCs can be maintained in a proliferative state by treatment with basic fibroblast growth factor and EGF. In these conditions, *Egr1* expression is upregulated in NSPCs and supports their proliferation. The presence of EGF is specifically required to maintain elevated *Egr1* transcription levels in NSPCs and sustain their proliferation via EGF receptor and ERK pathway activation. In contrast, EGF removal from the culture medium promotes *Egr1* downregulation in NSPCs, as well as their cell cycle exit and differentiation towards neuronal and glial fates. We found that EGF removal causes the downregulation of the genes coding for cyclin D1, D2 and D3, and the upregulation of p27, coding for a cell cycle inhibitor. *Egr1* overexpression in NSPCs has the opposite effect, causing an increase in the expression levels of cell cycle regulators and a decrease in those of cell cycle inhibitors. Of note, *Egr1* overexpression rescues the cell proliferation decrease observed after EGF removal, suggesting that this TF may play a significant role in supporting the proliferation of adult SVZ NSPCs downstream of the EGF-EGFR signaling pathway (Cera et al., 2018; **Figure 2**).

Other studies have investigated the role played by *Egr1* in adult SVZ NSPCs *in vivo*. Using a mouse model of recovery from an ischemic injury, it was found that *Egr1* expression is upregulated, and its nuclear levels increase, in SVZ NSPCs of mice recovering after a treatment that simulates hypoxic conditions. Moreover, upon hypoxia/ischemia, *EGFR* mRNA and protein levels increase, promoting the recruitment of NSPCs in the cell cycle, a reduction in their transition time through the cell cycle and an expansion of the NSPC pool (Alagappan et al., 2013). It was therefore important to determine whether there is a causal relationship between the concomitant increase in the expression levels of both *Egr1* and EGFR and the expansion of the SVZ NSPC population following ischemia. In support of this hypothesis, *Egr1* binding to the EGFR promoter region has been detected following ischemic injury, suggesting that *Egr1* can stimulate NSPC proliferation in the SVZ through the positive regulation of EGFR expression, at least in this experimental paradigm (Alagappan et al., 2013). Furthermore, it has been demonstrated



**Figure 2 Proposed model of the regulation of SVZ NSPC proliferation by the antagonistic activities of *Dbx2* and *Egr1*.** EGFR signaling may promote NSPC proliferation by repressing *Dbx2* and by activating *Egr1* expression. In turn, these TFs may exert opposite effects on the expression of genes coding for cell cycle activators and inhibitors and for other regulators of NSPC proliferation. According to this model, changes in the levels of EGFR signaling may be instrumental, acting via *Dbx2* and *Egr1*, to the decrease in NSPC proliferation associated with the transition towards quiescent or differentiating states or with aging. EGFR: Epidermal growth factor receptor; NSPCs: neural stem/progenitor cells; SVZ: subventricular zone; TFs: transcription factors.

that *Egr1* overexpression in several cancer cell lines causes *Egr1* binding to the EGFR promoter and enhances EGFR expression (Nishi et al., 2002).

Another study recently reported the ability of quercetin, a flavonoid with a strong antioxidant activity that stimulates adult hippocampal neurogenesis (Tchantchou et al., 2009), to increase the number of proliferating NSPCs and *Egr1* mRNA levels in the adult dentate gyrus of a rat model of AD (Karimipour et al., 2019). This suggests the interesting possibility that *Egr1* may be involved in mediating the positive effects of quercetin on hippocampal neurogenesis, and, more in general, regulate the neurogenic process also in the SGZ niche.

In the developing brain, *Egr1* expression pattern has been mapped and its mRNA has been detected in neural progenitors and neurons both in the embryonic and in the neonatal rat brain, but not in radial glial cells or in mature glial cell types (Wells et al., 2011). This result is in agreement with above mentioned studies (Alagappan et al., 2013), which found that *Egr1* upregulation in the adult SVZ following ischemic injury takes place in the neural progenitor population, but not in astrocytes or in astrocyte-like cells. Although it seems unlikely that *Egr1* role changes drastically when comparing the neurogenic process in the developing or in the adult brain, it remains to be investigated whether there are specific functions of this TF that are peculiar to either the developing or the adult stages.

### The *Dbx2* Homeobox Gene in the Regulation of Adult Neural Stem/Progenitor Cell Proliferation during Aging

Over the last 3 years, several studies have employed next generation sequencing methodologies to investigate the ge-

nome-wide molecular changes that take place during SVZ aging, especially in NSPCs. To this aim, different laboratories have compared the transcriptomic signatures of single SVZ cells (Shi et al., 2018; Dulken et al., 2019; Kalamakis et al., 2019), sorted SVZ NSPC bulk populations (Leeman et al., 2018; Kalamakis et al., 2019), SVZ NSPC *in vitro* cultures (Lupo et al., 2018; Shi et al., 2018), or dissected SVZ tissues (Apostolopoulou et al., 2017; Shi et al., 2018) obtained from mice at different ages between 2 and 24 months old. In one of these studies, we have described the genome-wide profiling of mRNA, DNA methylation and histone H3 lysine 4/27 trimethylation levels in NSPCs of the 3 and the 18 months old mouse SVZ that were expanded for two passages in non-adherent culture conditions (Lupo et al., 2018). Among the genes displaying age-associated transcriptional and/or epigenetic changes in this analysis, we focused on the homeobox gene *Dbx2*, given the pivotal roles that homeodomain TFs play in neural development (Lupo et al., 2000; Wigle and Eisenstat, 2008). *Dbx2*, in particular, is involved in the specification of interneuron fates in the developing spinal cord together with its close homolog *Dbx1* (Pierani et al., 1999). When compared with young adult samples, aged NSPC cultures showed enhanced *Dbx2* transcript levels, which were also detected in freshly sorted NSPCs from the SVZ of adult and aged mice (Lupo et al., 2018). These results are supported by independent transcriptomic datasets of freshly sorted SVZ NSPCs, or freshly dissected SVZ tissues, from young adult and aged mice, reporting increased *Dbx2* mRNA levels in aged samples (Apostolopoulou et al., 2017; Kalamakis et al., 2019).

Although the global transcriptional changes taking place during NSPC aging have been partially elucidated by the above described studies, the molecular drivers of these changes remain unclear. Several extracellular signaling



pathways, such as the EGFR, TGF- $\beta$ , mammalian target of rapamycin and Wnt pathways, are known to be differentially modulated in the young adult and in the aged SVZ neurogenic niche and they could potentially determine the age-related gene expression changes detected in NSPCs (Paliouras et al., 2012; Pineda et al., 2013; Zhu et al., 2014; Lupo et al., 2019). We found that *Dbx2* mRNA levels increased when young adult SVZ NSPCs were cultured for 24 hours without EGF, suggesting that EGFR signaling might negatively regulate *Dbx2* expression (Lupo et al., 2018). Since the levels of EGFR, of its TGF- $\alpha$  ligand and of its intracellular mediator ERK have been shown to be downregulated in the aged SVZ (Enwere et al., 2004; Kase et al., 2019), there might be a causative link between the decrease of EGFR signaling levels and the enhanced *Dbx2* mRNA levels detected in the aged SVZ. Further experiments specifically addressing the epistatic relationships between EGFR signaling and *Dbx2* need to be performed. Moreover, the age-related increase of *Dbx2* expression levels may result from an increase in its transcription and/or its transcript stability, with the latter potentially involving microRNAs. Since microRNAs are implicated in NSPC regulation (Cacci et al., 2017) and brain aging (Danka Mohammed et al., 2017), it would be worth to address their potential role in regulating *Dbx2* expression in NSPCs.

To investigate whether *Dbx2* upregulation in aged NSPCs may be involved in promoting the phenotypic changes associated with NSPC aging, we overexpressed this gene in young adult NSPCs by producing transgenic NSPC cultures with constitutive, or doxycycline-inducible expression of a *Dbx2* transgene. These assays showed a reduction in NSPC culture growth upon constitutive or acute overexpression of *Dbx2*, as exemplified by the smaller size of neurospheres obtained by non-adherent culture of NSPCs overexpressing *Dbx2* in comparison with control NSPCs. Elevated *Dbx2* expression levels did not cause major effects on cell viability or cell differentiation in young adult NSPCs cultured in proliferating conditions (Lupo et al., 2018). This suggests that the growth phenotype of *Dbx2* overexpressing cultures may result from altered NSPC proliferation dynamics, which might be related to the previously reported changes in the transition between activated and quiescent states and/or in the cell cycle length of aged NSPCs (Daynac et al., 2016; Apostolopoulou et al., 2017; Bast et al., 2018; Kalamakis et al., 2019). Further experiments are required to pinpoint the alterations of NSPC proliferation caused by *Dbx2* overexpression, and to investigate whether the inhibition of *Dbx2* function may improve the proliferative capacity of aged NSPCs.

Since *Dbx2* encodes for a TF, the phenotypic effects observed in NSPCs undergoing elevated expression of this gene are expected to result from the dysregulation of *Dbx2*-dependent transcriptional networks. By performing gene expression analyses in NSPCs overexpressing *Dbx2*, we have identified a few putative targets of its transcriptional activity (Lupo et al., 2018; **Figure 2**). In particular, *Dbx2* may negatively regulate the homeobox genes *Prrx1* and *Gsx1*, since they were both downregulated in SVZ NSPCs overexpressing *Dbx2*. *In vivo*, *Prrx1* is co-expressed in NSPCs of the SVZ

and the SGZ together with the Sox2 TF, and these proteins may bind each other according to biochemical assays. When retroviral vectors were employed *in vivo* to constitutively express *Prrx1* in NSPCs of the adult mouse SGZ, NSPCs were forced to remain in an undifferentiated state and their progression to neuronal fates was impaired. Conversely, *Prrx1* knockdown in NSPC cultures from the whole mouse forebrain caused a depletion of proliferating cells (Shimozaki et al., 2013). These results suggest a functional link between the downregulation of *Prrx1* and the growth phenotype caused by *Dbx2* overexpression, although the specific role of *Prrx1* in the regulation of the SVZ niche needs to be investigated. Expression of *Gsx1* has been described in the mouse embryonic ventral telencephalon, at the level of NSPCs transiting between the ventricular Zone and the SVZ during a period of active neurogenesis at midgestation (Pei et al., 2011). This spatio-temporal pattern suggests a potential role for *Gsx1* in the regulation of embryonic NSPC proliferation and differentiation, which appears to be supported by the analysis of transgenic or mutant mouse embryos with *Gsx1* gain or loss of function. *Gsx1* activity, however, is largely redundant with that of the closely related homolog *Gsx2*, and the individual roles of these TFs remain unclear (Pei et al., 2011; Chapman et al., 2018). Furthermore, *Gsx2* is also implicated in the adult mouse SVZ niche, where it modulates NSPC progression towards neuronal differentiation in physiological conditions and in response to injury (López-Juárez et al., 2013), but the role of *Gsx1* in adult neurogenesis is still unknown. *Igf2bp2*, coding for a RNA binding protein, is another gene downregulated in NSPCs overexpressing *Dbx2*. Similar to *Gsx1/2*, *Igf2bp2* role during embryonic neurogenesis overlaps with that of the close homolog *Igf2bp1*. Both of them are involved in regulating NSPC proliferation and differentiation in the developing cerebral cortex, but *Igf2bp1* expression declines postnatally and is undetectable in the young adult SVZ (Fujii et al., 2013; Nishino et al., 2013). There is no available report to date on the function of *Igf2bp1* in adult neurogenesis.

We also found genes that were upregulated upon *Dbx2* overexpression, such as those coding for the cell cycle inhibitor p21 and for Dll4, a Notch ligand implicated in the modulation of NSPC proliferation and differentiation in the adult SVZ niche (Bicker et al., 2017; Lin et al., 2019), thus suggesting that *Dbx2* may activate or repress the transcription of different genes. Other studies, however, indicate that *Dbx2* functions as a transcriptional repressor. The N-terminal regions of *Dbx1* and *Dbx2* contain short aminoacidic stretches sharing sequence similarity with the transcriptional repressor domain of the *Drosophila* Engrailed homeoprotein (Muhr et al., 2001; Karaz et al., 2016). Both *Dbx1* and *Dbx2* could bind the *Drosophila* Groucho transcriptional co-repressor *in vitro* (Muhr et al., 2001), and exhibited transcriptional repressor activity in luciferase reporter assays performed in COS-7 or HEK293 cells (Ma et al., 2011; Karaz et al., 2016). Furthermore, misexpression of a chimaeric protein containing the *Dbx2* homeodomain fused to the Engrailed repressor domain in the developing chick spinal cord could mimic some of the

effects of full-length Dbx2 misexpression, whereas a deletion mutant lacking the Engrailed repressor homologous sequence or a chimaeric protein containing the Dbx2 homeodomain fused to the transcriptional activator domain of VP16 were ineffective (Muhr et al., 2001). These observations raise the question of how Dbx2 overexpression leads to the increased transcript levels of certain genes, such as *p21* and *Dll4*. One possibility is that these genes are negatively regulated by a repressor, itself inhibited by *Dbx2*. Of note, *Gsx1* and *Gsx2* contain putative Engrailed homologous motifs, similar to Dbx proteins, suggesting that they too may function as transcriptional repressor (Muhr et al., 2001).

Taken together, the observations described in this section suggest the possible involvement of Dbx2 in the molecular mechanisms underlying the age-associated changes of NSPCs in the adult mouse SVZ. As schematized in **Figure 2**, decreased activation of EGFR signaling during SVZ aging may lead to an opposite increase in Dbx2 expression levels in NSPCs of the aged niche. In turn, as a result of its enhanced levels, Dbx2 may directly downregulate a subset of downstream genes including *Prrx1*, *Gsx1* and *Igf2bp2* by acting as a transcriptional repressor, whereas other genes such as *p21* and *Dll4* may be upregulated through an indirect mechanism (dashed lines in **Figure 2**). By modulating a cohort of genes implicated in the regulation of NSPC proliferation, elevated levels of Dbx2 in aged NSPCs may contribute to the decreased proliferative activity and the reduced neurogenic output of the aged SVZ niche. Bearing in mind that the aging process of the adult neurogenic niches entails several complex intrinsic and extrinsic mechanisms (Lupo et al., 2019), in future work it will be interesting to investigate Dbx2 function and regulation in SVZ NSPCs in more detail, including the genome-wide identification of its transcriptional targets and binding sites, the assessment of the effects of its functional inactivation and the elucidation of the upstream molecular pathways controlling its expression in the young adult and in the aged SVZ niche.

## Conclusions and Future Perspectives

As discussed in this review, a substantial body of data has been collected on the role played by the TFs studied in our and other laboratories during adult neurogenesis. One major feature of the evidences presented here is that the activity of REST, Egr1 and Dbx2 in the adult neurogenic niches is modulated by extrinsic mechanisms associated with physiological stimuli, aging, injury or disease. Furthermore, these TFs and at least some of the extracellular pathways modulating their activity, such as EGF signaling, are able to affect NSPC proliferation and differentiation. Therefore, REST, Egr1 and Dbx2 may be involved both in the intrinsic molecular networks governing NSPC self-renewal and neurogenic potential and, crucially, also in NSPC responses to the extrinsic influences toning up or down neurogenesis in physiological or pathological conditions. Within this intriguing scenario, several unresolved issues remain to be addressed in follow up studies.

Among these TFs, the role of REST in adult neurogenesis has been investigated in much more detail, revealing its

complex ability to regulate several events spanning the whole neurogenic process from qNSCs till neuronal maturation and activity. The molecular mechanisms by which this TF can perform distinct roles in different cell types and contexts is a major avenue for future investigations, along with a deeper characterization of the molecular pathways modulating its activity in response to various extrinsic stimuli. Moreover, although neuronal REST appears to have a protective role during physiological aging and the dysregulation of REST-dependent pathways in neurons may play a role in pathological aging (Lu et al., 2014), REST contribution to age-associated alterations in NSPCs remains surprisingly unaddressed. In the case of Egr1, there are several hints that it may be involved in normal or pathological brain aging processes, since changes in its expression levels have been detected in the hippocampus of aged AD human subjects (compared to adult and aged non-AD subjects) (Lanke et al., 2018), in the hippocampus of aged rats following spatial behaviour (compared to adult rats in the same behaviour conditions) (Penner et al., 2016), and in the cortex of aged mice (compared to young adult mice) (Rosa et al., 2018). As for REST, it is not known whether Egr1 may be implicated in aging processes in the adult neurogenic niches. Finally, our knowledge of the function of Dbx2 in adult neurogenesis remains very limited and much more work needs to be done to address it, especially using *in vivo* experimental models.

In future studies, it will also be important to map the possible interactions between REST, Egr1 and Dbx2 within the bigger picture of the global gene expression programs driving the neurogenic process. Based on their potential modulation by EGF signaling and their shared ability to affect NSPC proliferation, it is tempting to speculate that these TFs may functionally interact, directly or indirectly, in adult NSPCs. Supporting this hypothesis, we previously showed that Egr1 can regulate REST expression in neuroblastoma cells (Salani et al., 2009). The potential interplay among REST, Egr1 and Dbx2 will need to be investigated both by functional assays and by comparing the genome-wide transcriptional networks regulated by them. Finally, given the potential differences between adult neurogenesis in humans and rodents (Sanai et al., 2011; Sorrells et al., 2018), it would also be crucial to compare their expression profiles in the NSPC niches of these organisms and investigate their function using human *in vitro* models of the neurogenic process.

**Author contributions:** GP, GL and EC wrote the initial manuscript draft. RG, SS and SC critically revised the manuscript. All authors approved the final manuscript draft.

**Conflicts of interest:** The authors declare no conflicts of interest.

**Financial support:** None.

**Copyright license agreement:** The Copyright License Agreement has been signed by all authors before publication.

**Plagiarism check:** Checked twice by iThenticate.

**Peer review:** Externally peer reviewed.

**Open access statement:** This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Open peer reviewers:** Gregory Wohl Kirschen, Stony Brook Univer-



sity School of Medicine, USA; Jason H. Huang, Baylor Scott & White Health, USA.

**Additional file:** Open peer review reports 1 and 2.

## References

- Adams KW, Kletsov S, Lamm RJ, Elman JS, Mullenbrock S, Cooper GM (2017) Role for Egr1 in the transcriptional program associated with neuronal differentiation of PC12 cells. *PLoS One* 12:e0170076.
- Alagappan D, Balan M, Jiang Y, Cohen RB, Kottenko SV, Levison SW (2013) Egr-1 is a critical regulator of EGF-receptor-mediated expansion of subventricular zone neural stem cells and progenitors during recovery from hypoxia-hypoglycemia. *ASN Neuro* 5:183-193.
- Andrés ME, Burger C, Peral-Rubio MJ, Battaglioli E, Anderson ME, Grimes J, Dallman J, Ballas N, Mandel G (1999) CoREST: a functional corepressor required for regulation of neural-specific gene expression. *Proc Natl Acad Sci U S A* 96:9873-9878.
- Aoki H, Hara A, Era T, Kunisada T, Yamada Y (2012) Genetic ablation of Rest leads to in vitro-specific derepression of neuronal genes during neurogenesis. *Dev Camb Engl* 139:667-677.
- Apostolopoulou M, Kiehl TR, Winter M, Cardenas De La Hoz E, Boles NC, Bjornsson CS, Zuloaga KL, Goderie SK, Wang Y, Cohen AR, Temple S (2017) Non-monotonic changes in progenitor cell behavior and gene expression during aging of the adult V-SVZ neural stem cell niche. *Stem Cell Rep* 9:1931-1947.
- Ballas N, Battaglioli E, Atouf F, Andres ME, Chenoweth J, Anderson ME, Burger C, Moniwa M, Davie JR, Bowers WJ, Federoff HJ, Rose DW, Rosenfeld MG, Brehm P, Mandel G (2001) Regulation of neuronal traits by a novel transcriptional complex. *Neuron* 31:353-365.
- Ballas N, Grunseich C, Lu DD, Speh JC, Mandel G (2005) REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* 121:645-657.
- Bao H, Asrican B, Li W, Gu B, Wen Z, Lim SA, Haniff J, Ramakrishnan C, Deisseroth K, Philpot B, Song J (2017) Long-range GABAergic inputs regulate neural stem cell quiescence and control adult hippocampal neurogenesis. *Cell Stem Cell* 21:604-617.
- Baron V, Adamson ED, Calogero A, Ragona G, Mercola D (2006) The transcription factor Egr1 is a direct regulator of multiple tumor suppressors including TGFbeta1, PTEN, p53, and fibronectin. *Cancer Gene Ther* 13:115-124.
- Bast L, Calzolari F, Strasser MK, Hasenauer J, Theis FJ, Ninkovic J, Marr C (2018) Increasing neural stem cell division asymmetry and quiescence are predicted to contribute to the age-related decline in neurogenesis. *Cell Rep* 25:3231-3240.
- Beckervordersandforth R, Tripathi P, Ninkovic J, Bayam E, Lepier A, Stempfhuber B, Kirchhoff F, Hirrlinger J, Haslinger A, Lie DC, Beckers J, Yoder B, Irmeler M, Götz M (2010) In vivo fate mapping and expression analysis reveals molecular hallmarks of prospectively isolated adult neural stem cells. *Cell Stem Cell* 7:744-758.
- Bicker F, Vasic V, Horta G, Ortega F, Nolte H, Kavayanifar A, Keller S, Stankovic ND, Irmeler PN, Benedetto R, Lutz B, Bäuerle T, Hartwig J, Baumgart J, Krüger M, Radyushkin K, Alberi L, Berninger B, Schmidt MHH (2017) Neurovascular EGFL7 regulates adult neurogenesis in the subventricular zone and thereby affects olfactory perception. *Nat Commun* 8:15922.
- Biesiada E, Razandi M, Levin ER (1996) Egr-1 activates basic fibroblast growth factor transcription. Mechanistic implications for astrocyte proliferation. *J Biol Chem* 271:18576-18581.
- Brill MS, Ninkovic J, Wimpenny E, Hodge RD, Ozen I, Yang R, Lepier A, Gascón S, Erdelyi F, Szabo G, Parras C, Guillemot F, Frotscher M, Berninger B, Hevner RF, Raineteau O, Götz M (2009) Adult generation of glutamatergic olfactory bulb interneurons. *Nat Neurosci* 12:1524-1533.
- Bruce AW, Donaldson IJ, Wood IC, Yerbury SA, Sadowski MI, Chapman M, Göttgens B, Buckley NJ (2004) Genome-wide analysis of repressor element 1 silencing transcription factor/neuron-restrictive silencing factor (REST/NRSF) target genes. *Proc Natl Acad Sci U S A* 101:10458-10463.
- Cacci E, Negri R, Biagioni S, Lupo G (2017) Histone methylation and microRNA-dependent regulation of epigenetic activities in neural progenitor self-renewal and differentiation. *Curr Top Med Chem* 17:794-807.
- Calderone A, Jover T, Noh K, Tanaka H, Yokota H, Lin Y, Grooms SY, Regis R, Bennett MVL, Zukin RS (2003) Ischemic insults derepress the gene silencer REST in neurons destined to die. *J Neurosci Off J Soc Neurosci* 23:2112-2121.
- Cera AA, Cacci E, Toselli C, Cardarelli S, Bernardi A, Gioia R, Giorgi M, Poiana G, Biagioni S (2018) Egr-1 maintains NSC proliferation and its overexpression counteracts cell cycle exit triggered by the withdrawal of epidermal growth factor. *Dev Neurosci* 40:223-233.
- Chapman H, Riesenberger A, Ehrman LA, Kohli V, Nardini D, Nakafuku M, Campbell K, Waclaw RR (2018) Gsx transcription factors control neuronal versus glial specification in ventricular zone progenitors of the mouse lateral ganglionic eminence. *Dev Biol* 442:115-126.
- Chen ZF, Paquette AJ, Anderson DJ (1998) NRSE/REST is required in vivo for repression of multiple neuronal target genes during embryogenesis. *Nat Genet* 20:136-142.
- Cheval H, Chagneau C, Levasseur G, Veyrac A, Faucon-Bigué N, Laroche S, Davis S (2012) Distinctive features of Egr transcription factor regulation and DNA binding activity in CA1 of the hippocampus in synaptic plasticity and consolidation and reconsolidation of fear memory. *Hippocampus* 22:631-642.
- Chong JA, Tapia-Ramírez J, Kim S, Toledo-Aral JJ, Zheng Y, Boutros MC, Altshuler YM, Frohman MA, Kraner SD, Mandel G (1995) REST: a mammalian silencer protein that restricts sodium channel gene expression to neurons. *Cell* 80:949-957.
- Conti L, Crisafulli L, Caldera V, Tortoreto M, Brilli E, Conforti P, Zunino F, Magrassi L, Schiffer D, Cattaneo E (2012) REST controls self-renewal and tumorigenic competence of human glioblastoma cells. *PLoS One* 7:e38486.
- Crosby SD, Puetz JJ, Simburger KS, Fahrner TJ, Milbrandt J (1991) The early response gene NGFI-C encodes a zinc finger transcriptional activator and is a member of the GCGGGGGCG (GSG) element-binding protein family. *Mol Cell Biol* 11:3835-3841.
- Danka Mohammed CP, Park JS, Nam HG, Kim K (2017) MicroRNAs in brain aging. *Mech Ageing Dev* 168:3-9.
- Daynac M, Morizur L, Chicheportiche A, Mouthon MA, Boussin FD (2016) Age-related neurogenesis decline in the subventricular zone is associated with specific cell cycle regulation changes in activated neural stem cells. *Sci Rep* 6:21505.
- Drouin-Ouellet J, Lau S, Brattås PL, Rylander Ottosson D, Pircs K, Grassi DA, Collins LM, Vuono R, Andersson Sjöland A, Westergren-Thorsson G, Graff C, Minthon L, Toresson H, Barker RA, Jakobsson J, Parmar M (2017) REST suppression mediates neural conversion of adult human fibroblasts via microRNA-dependent and -independent pathways. *EMBO Mol Med* 9:1117-1131.
- Dulken BW, Buckley MT, Navarro Negredo P, Saligrama N, Cayrol R, Leeman DS, George BM, Boutet SC, Hebestreit K, Pluvinage JV, Wyss-Coray T, Weissman IL, Vogel H, Davis MM, Brunet A (2019) Single-cell analysis reveals T cell infiltration in old neurogenic niches. *Nature* 571:205-210.
- Encinas JM, Michurina TV, Peunova N, Park JH, Tordo J, Peterson DA, Fishell G, Koulakov A, Enikolopov G (2011) Division-coupled astrocytic differentiation and age-related depletion of neural stem cells in the adult hippocampus. *Cell Stem Cell* 8:566-579.
- Enwere E, Shingo T, Gregg C, Fujikawa H, Ohta S, Weiss S (2004) Aging results in reduced epidermal growth factor receptor signaling, diminished olfactory neurogenesis, and deficits in fine olfactory discrimination. *J Neurosci Off J Soc Neurosci* 24:8354-8365.
- Fuentealba LC, Rompani SB, Parraguez JL, Obernier K, Romero R, Cepko CL, Alvarez-Buylla A (2015) Embryonic origin of postnatal neural stem cells. *Cell* 161:1644-1655.
- Fujii Y, Kishi Y, Gotoh Y (2013) IMP2 regulates differentiation potentials of mouse neocortical neural precursor cells. *Genes Cells Devoted Mol Cell Mech* 18:79-89.
- Gao Z, Ure K, Ding P, Nashaat M, Yuan L, Ma J, Hammer RE, Hsieh J (2011) The master negative regulator REST/NRSF controls adult neurogenesis by restraining the neurogenic program in quiescent stem cells. *J Neurosci Off J Soc Neurosci* 31:9772-9786.
- Gonçalves JT, Schafer ST, Gage FH (2016) Adult neurogenesis in the hippocampus: from stem cells to behavior. *Cell* 167:897-914.
- Götz M, Nakafuku M, Petrik D (2016) Neurogenesis in the developing and adult brain-similarities and key differences. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a018853.
- Gregg J, Fraizer G (2011) Transcriptional regulation of EGR1 by EGF and the ERK signaling pathway in prostate cancer cells. *Genes Cancer* 2:900-909.
- Grimes JA, Nielsen SJ, Battaglioli E, Miska EA, Speh JC, Berry DL, Atouf F, Holdener BC, Mandel G, Kouzarides T (2000) The co-repressor mSin3A is a functional component of the REST-CoREST repressor complex. *J Biol Chem* 275:9461-9467.

- Hohl M, Thiel G (2005) Cell type-specific regulation of RE-1 silencing transcription factor (REST) target genes. *Eur J Neurosci* 22:2216-2230.
- Huff K, End D, Guroff G (1981) Nerve growth factor-induced alteration in the response of PC12 pheochromocytoma cells to epidermal growth factor. *J Cell Biol* 88:189-198.
- Itou Y, Nochi R, Kuribayashi H, Saito Y, Hisatsune T (2011) Cholinergic activation of hippocampal neural stem cells in aged dentate gyrus. *Hippocampus* 21:446-459.
- Johnson R, Teh CH, Kunarso G, Wong KY, Srinivasan G, Cooper ML, Volta M, Chan SS, Lipovich L, Pollard SM, Karuturi RKM, Wei C, Buckley NJ, Stanton LW (2008) REST regulates distinct transcriptional networks in embryonic and neural stem cells. *PLoS Biol* 6:e256.
- Jones MW, Errington ML, French PJ, Fine A, Bliss TV, Garel S, Charnay P, Bozon B, Laroche S, Davis S (2001) A requirement for the immediate early gene *Zif268* in the expression of late LTP and long-term memories. *Nat Neurosci* 4:289-296.
- Kalamakis G, Brüne D, Ravichandran S, Bolz J, Fan W, Ziebell F, Stiehl T, Catalá-Martinez F, Kupke J, Zhao S, Llorens-Bobadilla E, Bauer K, Limpert S, Berger B, Christen U, Schmezer P, Mallm JP, Berninger B, Anders S, Del Sol A, et al. (2019) Quiescence modulates stem cell maintenance and regenerative capacity in the aging brain. *Cell* 176:1407-1419.
- Kaneko N, Hwang JY, Gertner M, Pontarelli F, Zukin RS (2014) Casein kinase 1 suppresses activation of REST in insulted hippocampal neurons and halts ischemia-induced neuronal death. *J Neurosci Off J Soc Neurosci* 34:6030-6039.
- Karaz S, Courgeon M, Lepetit H, Bruno E, Pannone R, Tarallo A, Thouzé F, Kerner P, Vervoort M, Causeret F, Pierani A, D'Onofrio G (2016) Neuronal fate specification by the *Dbx1* transcription factor is linked to the evolutionary acquisition of a novel functional domain. *EvoDevo* 7:18.
- Karimipour M, Rahbarghazi R, Tayefi H, Shimia M, Ghanadian M, Mahmoudi J, Bagheri HS (2019) Quercetin promotes learning and memory performance concomitantly with neural stem/progenitor cell proliferation and neurogenesis in the adult rat dentate gyrus. *Int J Dev Neurosci Off J Int Soc Dev Neurosci* 74:18-26.
- Kase Y, Otsu K, Shimazaki T, Okano H (2019) Involvement of p38 in age-related decline in adult neurogenesis via modulation of Wnt signaling. *Stem Cell Rep* 12:1313-1328.
- Kaufmann K, Thiel G (2001) Epidermal growth factor and platelet-derived growth factor induce expression of *Egr-1*, a zinc finger transcription factor, in human malignant glioma cells. *J Neurol Sci* 189:83-91.
- Khachigian LM, Williams AJ, Collins T (1995) Interplay of *Sp1* and *Egr-1* in the proximal platelet-derived growth factor A-chain promoter in cultured vascular endothelial cells. *J Biol Chem* 270:27679-27686.
- Kim HJ, Denli AM, Wright R, Baul TD, Clemenson GD, Morcos AS, Zhao C, Schafer ST, Gage FH, Kagalwala MN (2015) REST regulates non-cell-autonomous neuronal differentiation and maturation of neural progenitor cells via secretogranin II. *J Neurosci Off J Soc Neurosci* 35:14872-14884.
- Kohwi M, Petryniak MA, Long JE, Ekker M, Obata K, Yanagawa Y, Rubenstein JLR, Alvarez-Buylla A (2007) A subpopulation of olfactory bulb GABAergic interneurons is derived from *Emx1*- and *Dlx5/6*-expressing progenitors. *J Neurosci Off J Soc Neurosci* 27:6878-6891.
- Kuwabara T, Hsieh J, Nakashima K, Taira K, Gage FH (2004) A small modulatory dsRNA specifies the fate of adult neural stem cells. *Cell* 116:779-793.
- Lanke V, Moolamalla STR, Roy D, Vinod PK (2018) Integrative analysis of hippocampus gene expression profiles identifies network alterations in aging and Alzheimer's disease. *Front Aging Neurosci* 10:153.
- Leeman DS, Hebestreit K, Ruetz T, Webb AE, McKay A, Pollina EA, Dulken BW, Zhao X, Yeo RW, Ho TT, Mahmoudi S, Devarajan K, Passegué E, Rando TA, Frydman J, Brunet A (2018) Lysosome activation clears aggregates and enhances quiescent neural stem cell activation during aging. *Science* 359:1277-1283.
- Levkovitz Y, Baraban JM (2001) A dominant negative inhibitor of the *Egr* family of transcription regulatory factors suppresses cerebellar granule cell apoptosis by blocking *c-Jun* activation. *J Neurosci Off J Soc Neurosci* 21:5893-5901.
- Lin R, Cai J, Kenyon L, Iozzo R, Rosenwasser R, Iacovitti L (2019) Systemic factors trigger vasculature cells to drive notch signaling and neurogenesis in neural stem cells in the adult brain. *Stem Cells Dayt Ohio* 37:395-406.
- Liu C, Adamson E, Mercola D (1996) Transcription factor *EGR-1* suppresses the growth and transformation of human HT-1080 fibrosarcoma cells by induction of transforming growth factor beta 1. *Proc Natl Acad Sci U S A* 93:11831-11836.
- López-Juárez A, Howard J, Ullom K, Howard L, Grande A, Pardo A, Waclaw R, Sun YY, Yang D, Kuan CY, Campbell K, Nakafuku M (2013) *Gsx2* controls region-specific activation of neural stem cells and injury-induced neurogenesis in the adult subventricular zone. *Genes Dev* 27:1272-1287.
- Lu L, Marisetty A, Liu B, Kamal MM, Gumin J, Veo B, Cai Y, Kassem DH, Weng C, Maynard ME, Hood KN, Fuller GN, Pan ZZ, Cykowski MD, Dash PK, Majumder S (2018) REST overexpression in mice causes deficits in spontaneous locomotion. *Sci Rep* 8:12083.
- Lu T, Aron L, Zullo J, Pan Y, Kim H, Chen Y, Yang TH, Kim HM, Drake D, Liu XS, Bennett DA, Colaiacovo MP, Yankner BA (2014) REST and stress resistance in ageing and Alzheimer's disease. *Nature* 507:448-454.
- Lupo G, Andreazzoli M, Gestri G, Liu Y, He RQ, Barsacchi G (2000) Homeobox genes in the genetic control of eye development. *Int J Dev Biol* 44:627-636.
- Lupo G, Gioia R, Nisi PS, Biagioni S, Cacci E (2019) Molecular mechanisms of neurogenic aging in the adult mouse subventricular zone. *J Exp Neurosci* 13:1179069519829040.
- Lupo G, Nisi PS, Esteve P, Paul YL, Novo CL, Sidders B, Khan MA, Biagioni S, Liu HK, Bovolenta P, Cacci E, Rugg-Gunn PJ (2018) Molecular profiling of aged neural progenitors identifies *Dbx2* as a candidate regulator of age-associated neurogenic decline. *Aging Cell* 17:e12745.
- Ma P, Zhao S, Zeng W, Yang Q, Li C, Lv X, Zhou Q, Mao B (2011) *Xenopus Dbx2* is involved in primary neurogenesis and early neural plate patterning. *Biochem Biophys Res Commun* 412:170-174.
- Mandel G, Fiondella CG, Covey MV, Lu DD, Loturco JJ, Ballas N (2011) Repressor element 1 silencing transcription factor (REST) controls radial migration and temporal neuronal specification during neocortical development. *Proc Natl Acad Sci U S A* 108:16789-16794.
- Masserdotti G, Gillotin S, Sutor B, Drechsel D, Irmeler M, Jørgensen HF, Sass S, Theis FJ, Beckers J, Berninger B, Guillemot F, Götz M (2015) Transcriptional mechanisms of proneural factors and REST in regulating neuronal reprogramming of astrocytes. *Cell Stem Cell* 17:74-88.
- McClelland S, Brennan GP, Dubé C, Rajpara S, Iyer S, Richichi C, Bernard C, Baram TZ (2014) The transcription factor NRSF contributes to epileptogenesis by selective repression of a subset of target genes. *eLife* 3:e01267.
- Meyer K, Feldman HM, Lu T, Drake D, Lim ET, Ling KH, Bishop NA, Pan Y, Seo J, Lin YT, Su SC, Church GM, Tsai LH, Yankner BA (2019) REST and neural gene network dysregulation in iPSC models of Alzheimer's disease. *Cell Rep* 26:1112-1127.
- Ming GL, Song H (2011) Adult neurogenesis in the mammalian brain: significant answers and significant questions. *Neuron* 70:687-702.
- Mira H, Andreu Z, Suh H, Lie DC, Jessberger S, Consiglio A, San Emeterio J, Hortigüela R, Marqués-Torrejón MA, Nakashima K, Colak D, Götz M, Fariñas I, Gage FH (2010) Signaling through *BMPRI-IA* regulates quiescence and long-term activity of neural stem cells in the adult hippocampus. *Cell Stem Cell* 7:78-89.
- Mortazavi A, Leeper Thompson EC, Garcia ST, Myers RM, Wold B (2006) Comparative genomics modeling of the NRSF/REST repressor network: from single conserved sites to genome-wide repertoire. *Genome Res* 16:1208-1221.
- Muhr J, Andersson E, Persson M, Jessell TM, Ericson J (2001) Groucho-mediated transcriptional repression establishes progenitor cell pattern and neuronal fate in the ventral neural tube. *Cell* 104:861-873.
- Mukherjee S, Brulet R, Zhang L, Hsieh J (2016) REST regulation of gene networks in adult neural stem cells. *Nat Commun* 7:13360.
- Naruse Y, Aoki T, Kojima T, Mori N (1999) Neural restrictive silencer factor recruits *mSin3* and histone deacetylase complex to repress neuron-specific target genes. *Proc Natl Acad Sci U S A* 96:13691-13696.

- Nechiporuk T, McGann J, Mullendorff K, Hsieh J, Wurst W, Floss T, Mandel G (2016) The REST remodeling complex protects genomic integrity during embryonic neurogenesis. *eLife* 5:e09584.
- Nishi H, Nishi KH, Johnson AC (2002) Early Growth Response-1 gene mediates up-regulation of epidermal growth factor receptor expression during hypoxia. *Cancer Res* 62:827-834.
- Nishino J, Kim S, Zhu Y, Zhu H, Morrison SJ (2013) A network of heterochronic genes including Imp1 regulates temporal changes in stem cell properties. *eLife* 2:e00924.
- Noh KM, Hwang JY, Follenzi A, Athanasiadou R, Miyawaki T, Grealley JM, Bennett MVL, Zukin RS (2012) Repressor element-1 silencing transcription factor (REST)-dependent epigenetic remodeling is critical to ischemia-induced neuronal death. *Proc Natl Acad Sci U S A* 109:E962-971.
- Paliouras GN, Hamilton LK, Aumont A, Joppé SE, Barnabé-Heider F, Fernandes KJL (2012) Mammalian target of rapamycin signaling is a key regulator of the transit-amplifying progenitor pool in the adult and aging forebrain. *J Neurosci Off J Soc Neurosci* 32:15012-15026.
- Palm K, Belluardo N, Metsis M, Timmusk T (1998) Neuronal expression of zinc finger transcription factor REST/NRSF/XBR gene. *J Neurosci Off J Soc Neurosci* 18:1280-1296.
- Panchision DM, McKay RDG (2002) The control of neural stem cells by morphogenic signals. *Curr Opin Genet Dev* 12:478-487.
- Pei Z, Wang B, Chen G, Nagao M, Nakafuku M, Campbell K (2011) Homeobox genes *Gsx1* and *Gsx2* differentially regulate telencephalic progenitor maturation. *Proc Natl Acad Sci U S A* 108:1675-1680.
- Penner MR, Parrish RR, Hoang LT, Roth TL, Lubin FD, Barnes CA (2016) Age-related changes in *Egr1* transcription and DNA methylation within the hippocampus. *Hippocampus* 26:1008-1020.
- Pierani A, Brenner-Morton S, Chiang C, Jessell TM (1999) A sonic hedgehog-independent, retinoid-activated pathway of neurogenesis in the ventral spinal cord. *Cell* 97:903-915.
- Pineda JR, Daynac M, Chicheportiche A, Cebrian-Silla A, Sii Felice K, Garcia-Verdugo JM, Boussin FD, Mouthon MA (2013) Vascular-derived TGF- $\beta$  increases in the stem cell niche and perturbs neurogenesis during aging and following irradiation in the adult mouse brain. *EMBO Mol Med* 5:548-562.
- Rodenas-Ruano A, Chávez AE, Cossio MJ, Castillo PE, Zukin RS (2012) REST-dependent epigenetic remodeling promotes the developmental switch in synaptic NMDA receptors. *Nat Neurosci* 15:1382-1390.
- Roopra A, Sharling L, Wood IC, Briggs T, Bachfischer U, Paquette AJ, Buckley NJ (2000) Transcriptional repression by neuron-restrictive silencer factor is mediated via the Sin3-histone deacetylase complex. *Mol Cell Biol* 20:2147-2157.
- Rosa P, Zerbinati C, Crestini A, Canudas AM, Ragona G, Confaloni A, Iuliano L, Calogero A (2018) Heme oxygenase-1 and brain oxysterols metabolism are linked to *Egr-1* expression in aged mice cortex, but not in hippocampus. *Front Aging Neurosci* 10:363.
- Salani M, Anelli T, Tocco GA, Lucarini E, Mozzetta C, Poiana G, Tata AM, Biagioni S (2009) Acetylcholine-induced neuronal differentiation: muscarinic receptor activation regulates *EGR-1* and REST expression in neuroblastoma cells. *J Neurochem* 108:821-834.
- Sanai N, Nguyen T, Ihrie RA, Mirzadeh Z, Tsai HH, Wong M, Gupta N, Berger MS, Huang E, Garcia-Verdugo JM, Rowitch DH, Alvarez-Buylla A (2011) Corridors of migrating neurons in the human brain and their decline during infancy. *Nature* 478:382-386.
- Santiago FS, Sanchez-Guerrero E, Zhang G, Zhong L, Raftery MJ, Khachigian LM (2019) Extracellular signal-regulated kinase-1 phosphorylates early growth response-1 at serine 26. *Biochem Biophys Res Commun* 510:345-351.
- Schoenherr CJ, Anderson DJ (1995) The neuron-restrictive silencer factor (NRSF): a coordinate repressor of multiple neuron-specific genes. *Science* 267:1360-1363.
- Segal RA, Greenberg ME (1996) Intracellular signaling pathways activated by neurotrophic factors. *Annu Rev Neurosci* 19:463-489.
- Shi Z, Geng Y, Liu J, Zhang H, Zhou L, Lin Q, Yu J, Zhang K, Liu J, Gao X, Zhang C, Yao Y, Zhang C, Sun YE (2018) Single-cell transcriptomics reveals gene signatures and alterations associated with aging in distinct neural stem/progenitor cell subpopulations. *Protein Cell* 9:351-364.
- Shimozaki K, Clemenson GD, Gage FH (2013) Paired related homeobox protein 1 is a regulator of stemness in adult neural stem/progenitor cells. *J Neurosci Off J Soc Neurosci* 33:4066-4075.
- Silverman ES, Du J, Williams AJ, Wadgaonkar R, Drazen JM, Collins T (1998) cAMP-response-element-binding-protein-binding protein (CBP) and p300 are transcriptional co-activators of early growth response factor-1 (*Egr-1*). *Biochem J* 336 (Pt 1):183-189.
- Singh A, Rokes C, Gireud M, Fletcher S, Baumgartner J, Fuller G, Stewart J, Zage P, Gopalakrishnan V (2011) Retinoic acid induces REST degradation and neuronal differentiation by modulating the expression of SCF( $\beta$ -TRCP) in neuroblastoma cells. *Cancer* 117:5189-5202.
- Soldati C, Caramanica P, Burney MJ, Toselli C, Bithell A, Augusti-Tocco G, Stanton LW, Biagioni S, Buckley NJ, Cacci E (2015) RE1 silencing transcription factor/neuron-restrictive silencing factor regulates expansion of adult mouse subventricular zone-derived neural stem/progenitor cells in vitro. *J Neurosci Res* 93:1203-1214.
- Sorrells SF, Paredes MF, Cebrian-Silla A, Sandoval K, Qi D, Kelley KW, James D, Mayer S, Chang J, Auguste KI, Chang EF, Gutierrez AJ, Kriegstein AR, Mathern GW, Oldham MC, Huang EJ, Garcia-Verdugo JM, Yang Z, Alvarez-Buylla A (2018) Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555:377-381.
- Steiner B, Wolf S, Kempermann G (2006) Adult neurogenesis and neurodegenerative disease. *Regen Med* 1:15-28.
- Su X, Gopalakrishnan V, Stearns D, Aldape K, Lang FF, Fuller G, Snyder E, Eberhart CG, Majumder S (2006) Abnormal expression of REST/NRSF and *Myc* in neural stem/progenitor cells causes cerebellar tumors by blocking neuronal differentiation. *Mol Cell Biol* 26:1666-1678.
- Sukhatme VP, Cao XM, Chang LC, Tsai-Morris CH, Stamenkovich D, Ferreira PC, Cohen DR, Edwards SA, Shows TB, Curran T (1988) A zinc finger-encoding gene coregulated with *c-fos* during growth and differentiation, and after cellular depolarization. *Cell* 53:37-43.
- Tchantchou F, Lacor PN, Cao Z, Lao L, Hou Y, Cui C, Klein WL, Luo Y (2009) Stimulation of neurogenesis and synaptogenesis by bilobalide and quercetin via common final pathway in hippocampal neurons. *J Alzheimers Dis* 18:787-798.
- Thiel G, Kaufmann K, Magin A, Lietz M, Bach K, Cramer M (2000) The human transcriptional repressor protein NAB1: expression and biological activity. *Biochim Biophys Acta* 1493:289-301.
- Thiel G, Mayer SI, Müller I, Stefano L, Rössler OG (2010) *Egr-1-A* Ca(2+)-regulated transcription factor. *Cell Calcium* 47:397-403.
- Wells T, Rough K, Carter DA (2011) Transcription mapping of embryonic rat brain reveals *EGR-1* induction in *SOX2* neural progenitor cells. *Front Mol Neurosci* 4:6.
- Wigle JT, Eisenstat DD (2008) Homeobox genes in vertebrate forebrain development and disease. *Clin Genet* 73:212-226.
- Yang YJ, Baltus AE, Mathew RS, Murphy EA, Evrony GD, Gonzalez DM, Wang EP, Marshall-Walker CA, Barry BJ, Murn J, Tatarakis A, Mahajan MA, Samuels HH, Shi Y, Golden JA, Mahajnah M, Shenhav R, Walsh CA (2012) Microcephaly gene links trithorax and REST/NRSF to control neural stem cell proliferation and differentiation. *Cell* 151:1097-1112.
- Yao J, Mackman N, Edgington TS, Fan ST (1997) Lipopolysaccharide induction of the tumor necrosis factor-alpha promoter in human monocytic cells. Regulation by *Egr-1*, *c-Jun*, and *NF-kappaB* transcription factors. *J Biol Chem* 272:17795-17801.
- Yeo M, Berglund K, Augustine G, Liedtke W (2009) Novel repression of *Kcc2* transcription by REST-RE-1 controls developmental switch in neuronal chloride. *J Neurosci Off J Soc Neurosci* 29:14652-14662.
- Zhu Y, Demidov ON, Goh AM, Virshup DM, Lane DP, Bulavin DV (2014) Phosphatase WIP1 regulates adult neurogenesis and WNT signaling during aging. *J Clin Invest* 124:3263-3273.
- Zuccato C, Tartari M, Crotti A, Goffredo D, Valenza M, Conti L, Cataudella T, Leavitt BR, Hayden MR, Timmusk T, Rigamonti D, Cattaneo E (2003) Huntingtin interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes. *Nat Genet* 35:76-83.