REGULAR FUNCTIONS ON SPHERICAL NILPOTENT ORBITS IN COMPLEX SYMMETRIC PAIRS: CLASSICAL HERMITIAN CASES

PAOLO BRAVI, JACOPO GANDINI

ABSTRACT. Given a classical semisimple complex algebraic group G and a symmetric pair (G, K) of Hermitian type, we study the closures of the spherical nilpotent K-orbits in the isotropy representation of K. We show that all such orbit closures are normal and describe the K-module structure of their ring of regular functions.

INTRODUCTION

Let G be a connected semisimple complex algebraic group, and let K be the fixed point subgroup of an algebraic involution θ of G.

The Lie algebra \mathfrak{g} of G splits into the sum of eigenspaces of θ ,

 $\mathfrak{g}=\mathfrak{k}\oplus\mathfrak{p},$

where the Lie algebra \mathfrak{k} of K is the eigenspace of eigenvalue 1, and \mathfrak{p} is the eigenspace of eigenvalue -1. The latter is called the isotropy representation of K.

In the present paper, we continue the systematic study initiated in [4] of the spherical nilpotent K-orbits in \mathfrak{p} , classified by King [13].

Here we treat the classical symmetric pairs (G, K) of Hermitian type: in particular, K is a maximal Levi subgroup of G, and $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$ splits into the sum of two simple K-modules dual to each other. Under this assumption, we prove that all spherical nilpotent orbit closures are normal (Theorem 4.3), and we compute the K-module structure of their coordinate rings.

In Appendix A we report the list of the spherical nilpotent K-orbits in \mathfrak{p} for all classical symmetric pairs $(\mathfrak{g}, \mathfrak{k})$ of Hermitian type. In the list, every orbit is labelled with the corresponding signed partition, [11].

For every orbit we provide an explicit description of a representative $e \in \mathfrak{p}$, as an element of a normal triple $\{h, e, f\}$, and the centralizer of e, which we denote by K_e . All these data can be deduced from [13].

Then we provide the Luna spherical system associated with $N_K(K_e)$, the normalizer of K_e in K, which is a wonderful subgroup of K.

Let us write $e = e_1 + e_2$ with $e_1 \in \mathfrak{p}_1$ and $e_2 \in \mathfrak{p}_2$. If e_1 and e_2 are both non-zero, then the orbit Ke is a *bicone*, the normalizer of K_e is the common stabilizer of the

²⁰¹⁰ Mathematics Subject Classification. Primary 14M27; Secondary 20G05.

Key words and phrases. Nilpotent orbits; symmetric spaces; spherical varieties.

lines $[e_1] \in \mathbb{P}(\mathfrak{p}_1)$ and $[e_2] \in \mathbb{P}(\mathfrak{p}_2)$, and $N_K(K_e)/K_e$ is a 2-dimensional complex torus.

The Luna spherical systems are used to deduce the normality of the K-orbit closures, and to compute the K-module structure of the corresponding coordinate rings.

In the tables of Appendix B we summarize the results of our computations. In Tables 1–6 we describe the K-module structure of $\mathbb{C}[\overline{Ke}]$ by giving a set of generators of its weight semigroup $\Gamma(\overline{Ke})$ (that is, the semigroup of the highest weights occurring in $\mathbb{C}[\overline{Ke}]$). Tables 7–11 contain the Luna spherical system of $N_K(K_e)$.

In Section 1 we adapt the criterion of normality for spherical cones used in our previous papers [6] and [4] to the case of a spherical multicone. In Section 2 we compute the Luna spherical systems of the normalizers $N_K(K_e)$. In Section 3 we show that the multiplication of sections of globally generated line bundles on the corresponding wonderful varieties is surjective. In Section 4 we deduce our results on normality and semigroups.

Acknowledgments. We would like to thank Andrea Maffei for his help in this project and, in particular, for sharing with us his ideas which led to Section 1. We would like to thank also the anonymous referee for some useful remarks.

Notation. As in our previous paper, simple roots of irreducible root systems are denoted by $\alpha_1, \alpha_2, \ldots$ and enumerated as in Bourbaki, when belonging to different irreducible components they are denoted by $\alpha_1, \alpha_2, \ldots, \alpha'_1, \alpha'_2, \ldots, \alpha''_1, \alpha''_2, \ldots$, and so on. When G (resp. K, T, \ldots) is an algebraic group, we will denote the associated Lie algebra by the corresponding fraktur character \mathfrak{g} (resp. $\mathfrak{k}, \mathfrak{t}, \ldots$). If moreover X is a G-variety (resp. a K-variety) and $x \in X$, we will denote the stabilizer of x by G_x (resp. K_x).

1. A CRITERION FOR THE NORMALITY OF A SPHERICAL MULTICONE

In this section, G will denote a connected reductive complex algebraic group (possibly not semisimple). By generalizing an argument due to C. De Concini in [12], in this section we will give a criterion (Theorem 1.2) to test the normality of a spherical multicone. We will apply it in Section 4 to study the normality of closures of spherical K-orbits in \mathfrak{p} , where K is a symmetric subgroup of a semisimple group and \mathfrak{p} is the isotropy representation of K.

Fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$. Let $\Pi = \{\lambda_1, \ldots, \lambda_m\} \subset \mathcal{X}(T)$ be a finite set of dominant weights and denote $V_{\Pi} = \bigoplus_{i=1}^m V(\lambda_i)$. For all $i = 1, \ldots, m$ we denote by $\hat{\pi}_i \colon V \longrightarrow V(\lambda_i)$ the corresponding projection.

Let $e \in V_{\Pi}$ and suppose that Ge is spherical. We will assume that $\hat{\pi}_i(e) \neq 0$ for all $i = 1, \ldots, m$. Under this assumption, we have a well-defined equivariant map $\pi_i \colon Ge \longrightarrow \mathbb{P}(V(\lambda_i))$ for all i, hence diagonally we get an equivariant map

$$\pi\colon Ge\longrightarrow \mathbb{P}(V(\lambda_1))\times\ldots\times\mathbb{P}(V(\lambda_m)).$$

We say that a closed subvariety $Z \subset V_{\Pi}$ is a *multicone* (w.r.t. the given decomposition of V_{Π}) if, for all $z \in Z$ and $(\xi_1, \ldots, \xi_m) \in \mathbb{C}^m$, it holds

$$\xi_1 \hat{\pi}_1(z) + \ldots + \xi_m \hat{\pi}_m(z) \in Z_{\cdot}$$

Given a spherical orbit $Ge \subset V_{\Pi}$, we will define a wonderful G-variety X endowed with a map

$$\phi \colon X \longrightarrow \mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m))$$

which is birational on its image and which identifies Π with a set of globally generated line bundles on X. If moreover $\overline{Ge} \subset V_{\Pi}$ is a multicone, we will establish a combinatorial criterion for \overline{Ge} to be normal in terms of Π , regarded as a subset of Pic(X).

Proposition 1.1. Let $Ge \subset V_{\Pi}$ be a spherical orbit, then $G_{\pi(e)} = N_G(G_e)$.

Proof. Write $e = \sum_{i=1}^{m} e_i$ with $e_i \in V(\lambda_i) \subset V_{\Pi}$, and notice that $G_e = G_{e_1} \cap \ldots \cap G_{e_m}$. Being spherical, for all $i = 1, \ldots, m$ the subgroup G_e fixes pointwise a unique line in $V(\lambda_i)$, namely $[e_i]$. If $g \in N_G(G_e)$, it follows that $ge_i \in V(\lambda_i)^{G_e}$ for all $i = 1, \ldots, m$, hence $g \in G_{[e_1]} \cap \ldots \cap G_{[e_m]} = G_{\pi(e)}$. Therefore $N_G(G_e) \subset G_{\pi(e)}$. Similarly, if $g \in G_{\pi(e)}$, then $ge_i \in V(\lambda_i)^{G_e}$ for all $i = 1, \ldots, m$, hence $ge \in V_{\Pi}^{G_e}$ and $g^{-1}hge = e$ for all $h \in G_e$, that is $g \in N_G(G_e)$. This shows the equality $G_{\pi(e)} = N_G(G_e)$.

Given a spherical subgroup $H \subset G$, it follows by a theorem of F. Knop that the homogeneous space $G/N_G(H)$ admits a wonderful compactification (see [16, Corollary 6.5] or [7, Proposition 2.4]). Notice that in general the normalizer of a spherical subgroup is not self-normalizing (see e.g. [2, Example 4]). The spherical homogeneous spaces occurring in our cases 2.4 and 4.4 give other examples of this kind.

Let us now fix some notation and recall some general facts on line bundles and their sections on a wonderful variety X, see e.g. [6, Section 1] and the references therein. Let Δ be the set of colors of X (that is, the *B*-stable prime divisors of Xwhich are not *G*-stable) and if $D \in \mathbb{Z}\Delta$ denote by $\mathcal{L}_D \in \operatorname{Pic}(X)$ the corresponding line bundle. Recall that $\operatorname{Pic}(X)$ is a free Abelian group with basis the line bundles \mathcal{L}_D with $D \in \Delta$, and that, under the indentification $\operatorname{Pic}(X) = \mathbb{Z}\Delta$, the semigroup of globally generated line bundles is identified with $\mathbb{N}\Delta$. Given $\mathcal{L}, \mathcal{L}' \in \operatorname{Pic}(X)$ we denote by

 $m_{\mathcal{L},\mathcal{L}'}: \Gamma(X,\mathcal{L}) \otimes \Gamma(X,\mathcal{L}') \longrightarrow \Gamma(X,\mathcal{L} \otimes \mathcal{L}')$

the multiplication of sections. If $\mathcal{L} = \mathcal{L}_D$ and $\mathcal{L}' = \mathcal{L}_E$ with $D, E \in \mathbb{Z}\Delta$, we will also denote $m_{\mathcal{L},\mathcal{L}'}$ by $m_{D,E}$.

Let $Y \subset X$ be the unique closed *G*-orbit and let $y_0 \in Y$ be the unique B^- -fixed point, where B^- denotes the opposite Borel subgroup of *B*. Recall the set of the spherical roots of *X*, defined by

$$\Sigma = \{T \text{-weights in } T_{u_0} X / T_{u_0} Y \}.$$

Then there is a bijective correspondence between spherical roots and G-stable divisors in X, which allows to regard $\mathbb{Z}\Sigma$ as a sublattice of $\operatorname{Pic}(X) = \mathbb{Z}\Delta$. This defines a partial order \leq_{Σ} on $\mathbb{N}\Delta$ as follows: if $D, E \in \mathbb{N}\Delta$, then $D \leq_{\Sigma} E$ if and only if $E - D \in \mathbb{N}\Sigma$. We say that $E \in \mathbb{N}\Delta$ is *minuscule* w.r.t. \leq_{Σ} if there exists no $F \in \mathbb{N}\Delta$ such that $F \leq_{\Sigma} E$ and $F \neq E$. Notice that $\Gamma(X, \mathcal{L}_E)$ is an irreducible *G*-module if and only if *E* is minuscule in $\mathbb{N}\Delta$ w.r.t. \leq_{Σ} . Given $F \in \mathbb{N}\Delta$, let indeed $V_F \subset \Gamma(X, \mathcal{L}_F)$ be the *G*-module generated by a canonical section, and given $\gamma \in \mathbb{Z}\Sigma$ (regarded as a sublattice of $\mathbb{Z}\Delta$) let $s^{\gamma} \in \Gamma(X, \mathcal{L}_{\gamma})^G$. Then for $E \in \mathbb{N}\Delta$ we have the following decomposition into simple *G*-modules

(1.1)
$$\Gamma(X, \mathcal{L}_E) = \bigoplus_{F \leqslant_{\Sigma} E} s^{E-F} V_F.$$

Going back to our setting, suppose that $Ge \subset V_{\Pi}$ is a spherical orbit. Then $G/G_{\pi(e)}$ admits a wonderful compactification X. Notice that the center Z(G) of G acts trivially on $G/G_{\pi(e)}$, hence it acts trivially on X as well.

For all $i = 1, \ldots, m$, let $\phi_i \colon G/G_{\pi(e)} \longrightarrow \mathbb{P}(V(\lambda_i))$ be the restriction of the projection defined on $\mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m))$. By the theory of spherical embeddings (see in particular [15, Theorem 5.1]), any dominant morphism $G/G_{\pi(e)} \longrightarrow Y$ to a complete spherical *G*-variety *Y* with a unique closed orbit can be extended to an equivariant morphism $X \longrightarrow Y$. This in particular applies to the closure of $\phi_i(G/G_{\pi(e)})$, since $\mathbb{P}(V(\lambda_i))$ contains a unique *B*-fixed point that is contained in every closed *G*-orbit. Thus ϕ_i extends to an equivariant morphism $\phi_i \colon X \longrightarrow \mathbb{P}(V(\lambda_i))$, that we still denote by the same symbol by abuse of notation. Mapping *X* diagonally to $\mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m))$ via $\phi = (\phi_1, \ldots, \phi_m)$, we get then an equivariant morphism

$$\phi \colon X \longrightarrow \mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m)).$$

which extends the inclusion $G/G_{\pi(e)} \longrightarrow \pi(Ge) \subset \mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m)).$

For all i = 1, ..., m, let $\mathcal{L}_i \in \operatorname{Pic}(X)$ be the globally generated line bundle defined by setting $\mathcal{L}_i = \phi_i^* \mathcal{O}(1)$, and let $D_i \in \mathbb{N}\Delta$ be the unique *B*-stable divisor such that $\mathcal{L}_{D_i} = \mathcal{L}_i$. We denote

$$\Delta_{\Pi}(e) = \{D_1, \dots, D_m\}.$$

For every $(d_1, \ldots, d_m) \in \mathbb{N}^m$, according with equation (1.1) the *G*-module $\Gamma(X, \mathcal{L}_1^{d_1} \otimes \ldots \otimes \mathcal{L}_m^{d_m})$ decomposes into the direct sum of the simple modules with highest weights of the form

(1.2)
$$d_1\lambda_1^* + \ldots + d_m\lambda_m^* - (d_1D_1 + \ldots + d_mD_m - F)$$

where $F \in \mathbb{N}\Delta$ and $F \leq_{\Sigma} d_1 D_1 + \ldots + d_m D_m$. Notice that we denote by λ^* the highest weight of $V(\lambda)^*$, the dual of the simple module of highest weight λ .

We will prove the following.

Theorem 1.2. Suppose that the orbit Ge is spherical and its closure $\overline{Ge} \subset V_{\Pi}$ is a multicone, and assume that the multiplication of sections $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all globally generated $\mathcal{L}, \mathcal{L}' \in \operatorname{Pic}(X)$. Then $\overline{Ge} \subset V$ is normal if and only if every $D \in \Delta_{\Pi}(e)$ is minuscule in $\mathbb{N}\Delta$.

Denote by A(Ge) the coordinate ring of \overline{Ge} and by $A_d(Ge)$ its component of degree d, then A(Ge) is generated by $A_1(Ge) = \bigoplus_{\lambda \in \Pi} V(\lambda)^*$. Consider the multigraded ring

$$\widetilde{A}(Ge) = \bigoplus_{(d_1,\dots,d_m)\in\mathbb{N}^m} \Gamma(X,\mathcal{L}_1^{d_1}\otimes\dots\otimes\mathcal{L}_m^{d_m}).$$

and denote $\widetilde{Ge} = \operatorname{Spec} \widetilde{A}(Ge)$. Setting

$$\widetilde{A}_d(Ge) = \bigoplus_{d_1 + \ldots + d_m = d} \Gamma(X, \mathcal{L}_1^{d_1} \otimes \ldots \otimes \mathcal{L}_m^{d_m})$$

we get a canonical inclusion $A_d(Ge) \subset \widetilde{A}_d(Ge)$: indeed for all $i = 1, \ldots, m$ the *G*-module $V(\lambda_i)^*$ is canonically identified with a submodule in $\Gamma(X, \mathcal{L}_i) \subset \widetilde{A}_1(Ge)$. This makes canonically A(Ge) a subring of $\widetilde{A}(Ge)$, and we get a projection

$$p\colon \widetilde{Ge} \longrightarrow \overline{Ge}$$

Theorem 1.2 is a consequence of the following description of the normalization of \overline{Ge} .

Theorem 1.3. Let Ge be spherical and $\overline{Ge} \subset V_{\Pi}$ be a multicone, and suppose that the multiplication of sections $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all $\mathcal{L},\mathcal{L}' \in \operatorname{Pic}(X)$. Then $p: \widetilde{Ge} \longrightarrow \overline{Ge}$ is the normalization map.

Remark 1.4. In [5] a standard monomial theory for the Cox ring of a wonderful variety was constructed. By making use of such a tool, reasoning as in [5, Proposition 5.2] one can actually show that \widetilde{Ge} has rational singularities, hence it is in particular normal. We will give anyway a direct proof of the normality of \widetilde{Ge} by using an easy geometric argument (see also [6, Section 7]).

Proof of Theorem 1.2. By Theorem 1.3 it follows that \overline{Ge} is normal if and only if $A(Ge) = \widetilde{A}(Ge)$. Since the multiplication of sections $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all globally generated $\mathcal{L}, \mathcal{L}' \in \operatorname{Pic}(X)$, it follows that $\widetilde{A}(Ge)$ is generated by its degree one component $\widetilde{A}_1(Ge)$. Therefore $A(Ge) = \widetilde{A}(Ge)$ if and only if $A_1(Ge) = \widetilde{A}_1(Ge)$, if and only if $\Gamma(X, \mathcal{L}_i) = V(\lambda_i)^*$ for all $i = 1, \ldots, m$. Being $\mathcal{L}_i = \mathcal{L}_{D_i}$, by the description of the irreducible components of $\Gamma(X, \mathcal{L}_{D_i})$ (see e.g. [6, Proposition 1.1]), this is equivalent to the fact that every D_i is minuscule in $\mathbb{N}\Delta$.

1.1. **Proof of Theorem 1.3.** We will split the proof of Theorem 1.3 in several propositions. The following shows that $\widetilde{A}(Ge)$ is a normal ring.

Suppose that Y is a normal variety and let $\mathcal{M}_1, \ldots, \mathcal{M}_k \in \operatorname{Pic}(Y)$, we denote

$$A(\mathcal{M}_1,\ldots,\mathcal{M}_k) = \bigoplus_{(d_1,\ldots,d_k)\in\mathbb{N}^k} \Gamma(Y,\mathcal{M}_1^{d_1}\otimes\ldots\otimes\mathcal{M}_k^{d_k})$$

and define correspondingly a sheaf of \mathcal{O}_Y -algebras on Y by setting

$$\mathcal{A}(\mathcal{M}_1,\ldots,\mathcal{M}_k) = \bigoplus_{(d_1,\ldots,d_k)\in\mathbb{N}^k} \mathcal{M}_1^{d_1}\otimes\ldots\otimes\mathcal{M}_k^{d_k}.$$

Proposition 1.5. Suppose that Y is a normal variety and let $\mathcal{M}_1, \ldots, \mathcal{M}_k \in \operatorname{Pic}(Y)$. Then $A(\mathcal{M}_1, \ldots, \mathcal{M}_k)$ is a normal ring.

Proof. Notice that $A(\mathcal{M}_1, \ldots, \mathcal{M}_k)$ is a domain since Y is irreducible. Suppose first that Y is affine and that $\mathcal{M}_i \simeq \mathcal{O}_Y$ for all $i = 1, \ldots, k$. Then $A(\mathcal{M}_1, \ldots, \mathcal{M}_k) = \mathbb{C}[Y][s_1, \ldots, s_k]$ is a polynomial ring with coefficients in $\mathbb{C}[Y]$, hence it is normal since Y is so.

Consider now the general case. Let $s_1, s_2 \in A(\mathcal{M}_1, \ldots, \mathcal{M}_k)$ and suppose that s_1/s_2 is integral over $A(\mathcal{M}_1, \ldots, \mathcal{M}_k)$. Let $U \subset Y$ be an affine open subset such that $\mathcal{M}_i|_U$ is trivial for all i. Then $(s_1/s_2)|_U$ is integral over $A(\mathcal{M}_1|_U, \ldots, \mathcal{M}_k|_U)$, thus it belongs to $A(\mathcal{M}_1|_U, \ldots, \mathcal{M}_k|_U)$. The claim follows since we can cover Y with affine open subsets $U \subset Y$ such that $\mathcal{M}_i|_U$ is trivial for all $i = 1, \ldots, k$. \Box

In particular we have

$$A(Ge) = A(\mathcal{L}_1, \dots, \mathcal{L}_m) = \Gamma(X, \mathcal{A}(\mathcal{L}_1, \dots, \mathcal{L}_m)).$$

Notice that, for all i = 1, ..., m, the map $\phi_i \colon X \longrightarrow \mathbb{P}(V(\lambda_i))$ factors through $\mathbb{P}(\Gamma(X, \mathcal{L}_i)^*)$. Therefore the map $\phi = (\phi_1, ..., \phi_m)$ factors as follows

$$X \xrightarrow{\widetilde{\phi}} \mathbb{P}(\Gamma(X, \mathcal{L}_1)^*) \times \ldots \times \mathbb{P}(\Gamma(X, \mathcal{L}_m)^*) \xrightarrow{\psi} \mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m)),$$

where $\psi = (\psi_1, \dots, \psi_m)$ is the canonical projection.

Consider the multiplication map


$$\bigoplus_{n\geq 0} \mathsf{S}^n\big(\widetilde{A}_1(Ge)\big) \longrightarrow A(\mathcal{L}_1,\ldots,\mathcal{L}_m):$$

it is surjective by the assumption of Theorem 1.3, and its kernel coincides with the homogeneous ideal of $\tilde{\phi}(X)$. It follows that \tilde{Ge} is the multicone over $\tilde{\phi}(X)$, whereas \overline{Ge} is by assumption the multicone over $\phi(X)$.

By Proposition 1.1 the map $\phi: X \longrightarrow \phi(X)$ is birational. Therefore the restriction of π induces a birational map $\tilde{\phi}(X) \longrightarrow \phi(X)$, and taking the affine multicones it follows that $p: \widetilde{Ge} \longrightarrow \overline{Ge}$ is birational as well. Therefore to conclude the proof of Theorem 1.3 we are left to show that $\widetilde{A}(Ge)$ is integral over A(Ge). The argument of the following proof is due to M. Brion (see [9, Proposition 2.1]).

Proposition 1.6. $\widehat{A}(Ge)$ is an integral extension of A(Ge).

Proof. Denote $Z = \mathbb{P}(V(\lambda_1)) \times \ldots \times \mathbb{P}(V(\lambda_m))$, for $i = 1, \ldots, m$ let $p_i \colon Z \longrightarrow \mathbb{P}(V(\lambda_i))$ be the projection. Denote $\mathcal{A}_Z = \mathcal{A}(p_1^*\mathcal{O}(1), \ldots, p_m^*\mathcal{O}(1))$, a sheaf of \mathcal{O}_Z -algebras, and set $L_Z = \operatorname{Spec} \Gamma(Z, \mathcal{A}_Z)$. Similarly denote $\mathcal{A}_X = \mathcal{A}(\mathcal{L}_1, \ldots, \mathcal{L}_m)$, a sheaf of \mathcal{O}_X -algebras, and set $L_X = \operatorname{Spec} \Gamma(X, \mathcal{A}_X)$. Then we have a pullback diagram



Notice that $\widetilde{A}(Ge) = \Gamma(L_X, \mathcal{O}_{L_X}) = \Gamma(L_Z, \overline{\phi}_* \mathcal{O}_{L_X})$, whereas A(Ge) is the image of the natural morphism $\Gamma(L_Z, \mathcal{O}_{L_Z}) \longrightarrow \Gamma(L_Z, \overline{\phi}_* \mathcal{O}_{L_X})$. Notice that $\overline{\phi}$ is projective, so that $\overline{\phi}_* \mathcal{O}_{L_X}$ is a coherent sheaf on L_Z . Hence $\Gamma(L_Z, \overline{\phi}_* \mathcal{O}_{L_X})$ is a finitely generated $\Gamma(L_Z, \mathcal{O}_{L_Z})$ -module, or equivalently $\widetilde{A}(Ge)$ is a finitely generated A(Ge)-module.

2. The spherical systems

For the notation and the generalities about wonderful subgroups and Luna spherical systems we refer to our previous paper [4, Section 1].

In this section we compute the wonderful subgroups of K associated with the spherical systems given in the tables of Appendix B. Since, as one can check case-by-case, they correspond up to conjugation to the normalizers of the centralizers $N_K(K_e)$ of the representatives e given in Appendix A, we obtain the following.

Proposition 2.1. The Luna spherical systems of the wonderful K-varieties associated to the spherical nilpotent K-orbits in \mathfrak{p} , for the classical symmetric pairs of Hermitian type, are those given in Tables 7–11.

We start with the spherical systems whose set Σ is empty:

- Table 7: 1.1 (r = 1), 1.2 (r = 1), 1.3 (r = s = 1), 1.6 (r = s = 0 and q = 1), 1.7 (r = s = 0 and p = 1);
- Table 8: 2.1, 2.2;
- Table 9: 3.1 (r = 1), 3.2 (r = 1), 3.3 (r = s = 1);
- Table 10: 4.1, 4.2;
- Table 11: 5.1 (r = 1), 5.2 (r = 1), 5.3 (r = s = 1).

Here it is immediate to see that the parabolic subgroup Q of K given in Appendix A is the wonderful subgroup associated with the corresponding spherical K-system.

For the remaining spherical systems, let M be the Levi subgroup of K corresponding with supp Σ . We will just compute the wonderful subgroup H of M associated with the M-spherical system obtained from the given K-spherical system by localization in supp Σ . It is then immediate to deduce from H which is the wonderful subgroup K associated to the given K-spherical system and check that it is equal, up to conjugation, to the normalizer of K_e given in the corresponding case of Appendix A.

2.1. Symmetric cases. From the following K-spherical systems, by localizing in $\sup \Sigma$, we get the M-spherical systems of symmetric subgroups of M. Since these are well-known, we just provide a reference:

- in the cases 1.1 (r > 1), 1.2 (r > 1), 1.3 (r > 1 or s > 1) of Table 7 we get the case 2 of [7], or the direct product of two of them;
- in the cases 1.6 (r = s = 0 and q > 1), 1.7 (r = s = 0 and p > 1) of Table 7, 2.4 of Table 8, 4.4 of Table 10, 5.4 of Table 11 we get the case 3 of [7];
- in the case 2.3 of Table 8 we get the case 9 of [7];
- in the cases 3.1 (r > 1), 3.2 (r > 1), 3.3 (r > 1 or s > 1) of Table 9 we get the case 5 of [7], or the direct product of two of them;
- in the case 4.3 of Table 10 we get the case 15 of [7];
- in the cases 5.1 (r > 1), 5.2 (r > 1), 5.3 (r > 1 or s > 1) of Table 11 we get the case 6 of [7], or the direct product of two of them.

2.2. Other reductive cases. From the K-spherical systems 1.4 and 1.5 of Table 7, by localizing in supp Σ , we get the M-spherical system of a wonderful reductive (but not symmetric) subgroup of M: the case 43 of [7].

2.3. Morphisms of type \mathcal{L} . In the remaining cases, 1.6 (r + s > 0) and 1.7 (r + s > 0) of Table 7, the spherical K-system (S^{p}, Σ, A) admits a distinguished set of colors Δ' such that the corresponding quotient


 $(S^{\rm p}/\Delta', \Sigma/\Delta', A/\Delta')$

is the spherical system of a wonderful K-variety which is obtained by parabolic induction from a wonderful K_h -variety. Such distinguished set of colors Δ' may be not minimal, let us describe it in the case 1.6, the other one is similar. We assume r and s both to be non-zero and r + s < q - 1, in the other cases the description is similar but simpler.

Under this assumption, localizing the K-spherical system of the case 1.6 in supp Σ , we obtain the following spherical system, which we label as $a^{y}(r, r) + a(t) + a^{y}(s, s)$ (here t = q - r - s - 1) for a group M of semisimple type $A_r \times A_{r+s+t} \times A_s$:

$$\begin{split} S^{\mathbf{p}} &= \{\alpha'_{r+2}, \dots, \alpha'_{r+t-1}\},\\ \Sigma &= \{\alpha_{1}, \dots, \alpha_{r}, \ \alpha'_{1}, \dots, \alpha'_{r}, \ \alpha'_{r+1} + \dots + \alpha'_{r+t}, \ \alpha'_{r+t+1}, \dots, \alpha'_{r+t+s}, \ \alpha''_{1}, \dots, \alpha''_{s})\}\\ \Delta &= \{D_{1}, \dots, D_{2r+1}, \ D_{2r+2}, D_{2r+3}, \ D_{2r+4}, \dots, D_{2r+2s+4}\},\\ \text{and full Cartan pairing} \\ \alpha_{1} &= D_{1} + D_{2} - D_{3}, \\ \alpha_{i} &= -D_{2i-2} + D_{2i-1} + D_{2i} - D_{2i+1} \text{ for } 2 \leqslant i \leqslant r, \\ \alpha'_{i} &= -D_{2i-1} + D_{2i} + D_{2i+1} - D_{2i+2} \text{ for } 1 \leqslant i \leqslant r, \\ \alpha'_{r+1} + \dots + \alpha'_{r+t} &= -D_{2r+1} + D_{2r+2} + D_{2r+3} - D_{2r+4}, \\ \alpha'_{r+t+i} &= -D_{2r+2i+1} + D_{2r+2i+2} + D_{2r+2i+3} - D_{2r+2i+4} \text{ for } 1 \leqslant i \leqslant s, \\ \alpha''_{i} &= -D_{2r+2i+2} + D_{2r+2i+3} + D_{2r+2i+4} - D_{2r+2i+5} \text{ for } 1 \leqslant i \leqslant s-1, \\ \alpha''_{s} &= -D_{2r+2s+2} + D_{2r+2s+3} + D_{2r+2s+4}. \end{split}$$

Notice that we have re-enumerated the simple roots of M to simplify the notation. If t > 1 the Luna diagram is as follows.




Let us consider the following subsets of colors:

$$\Delta_1 = \{ D_{2i} : 1 \leq i \leq r \}, \Delta_2 = \{ D_{2r+2i+3} : 1 \leq i \leq s \}$$

Notice that Δ_1 (resp. Δ_2) consists of the colors represented by circles above vertices on the left (resp. right) hand side of the diagram. Both are minimal distinguished with quotient of higher defect, see [4, Section 1.5.5]. The quotient by $\Delta' = \Delta_1 \cup \Delta_2$ is as follows.

$$\Sigma/\Delta' = \{\alpha_2 + \alpha'_1, \dots, \alpha_r + \alpha'_{r-1}, \alpha'_{r+1} + \dots + \alpha'_{r+t}, \alpha'_{r+t+2} + \alpha''_1, \dots, \alpha'_{r+t+s} + \alpha''_{s-1}\}$$

This spherical system corresponds to a subgroup of M which is a parabolic induction of a symmetric subgroup, that is, it can be decomposed as LP^{u} (a Levi decomposition) where $P = L_P P^u$ is the parabolic subgroup of M corresponding to $\operatorname{supp}(\Sigma/\Delta')$ and L is a symmetric subgroup of L_P corresponding to



More explicitly, L is the normalizer of $SL(r) \times GL(t) \times SL(s)$ in L_P of semisimple type $A_r \times A_r \times A_t \times A_s \times A_s$, where SL(r) is skew-diagonal in $SL(r) \times SL(r)$ and SL(s) is skew-diagonal in $SL(s) \times SL(s)$.

The wonderful subgroup H associated with the above spherical M-system $a^{y}(r, r) + a(t) + a^{y}(s, s)$ can therefore be taken as $L_{H}H^{u}$, with $L_{H} \subset L$ and $H^{u} \subset P^{u}$, where Lie $P^{u}/\text{Lie }H^{u}$ is the direct sum of two simple L_{H} -modules while L_{H} and L differ only by their connected center.

The codimension of L_H in L is equal to the increase in defect, which in this case is equal to 2.

The unipotent radical Lie H^{u} is as follows. There exist $W_{0,1}$ and $W_{1,1}$ L_{H} -submodules of Lie P^{u} of dimension r, isomorphic as L_{H} -modules but not as L-modules. Analogously there exist $W_{0,2}$ and $W_{1,2}$ L_{H} -submodules of Lie P^{u} of dimension s, isomorphic as L_{H} -modules but not as L-modules. Denoting by V the L_{H} -complement of $W_{0,1} \oplus W_{1,1} \oplus W_{0,2} \oplus W_{1,2}$ in Lie P^{u} , as L_{H} -module we have

$$\operatorname{Lie} H_1^{\mathrm{u}} = W_1 \oplus W_2 \oplus V$$

where W_1 is a simple L_H -submodule of $W_{0,1} \oplus W_{1,1}$ which projects non-trivially on both summands, and analogously W_2 is a simple L_H -submodule of $W_{0,2} \oplus W_{1,2}$ which projects non-trivially on both summands.

Remark 2.2. Notice that the given subsets of colors Δ_1 and Δ_2 decompose the above spherical system in the sense of [8, Definition 2.2.2], hence the corresponding wonderful variety is a non-trivial wonderful fiber product.

3. Projective normality

In this section we prove the following result, that will be used to study the normality of the spherical nilpotent K-orbit closures in \mathfrak{p} . Reasoning as in Section 1, recall that every spherical nilpotent K-orbit in \mathfrak{p} determines naturally a wonderful K-variety.

Theorem 3.1. Let $(\mathfrak{g}, \mathfrak{k})$ be a classical symmetric pair of Hermitian type, let $\mathcal{O} \subset \mathfrak{p}$ be a spherical nilpotent K-orbit and let X be the wonderful K-variety associated to \mathcal{O} . Then the multiplication of sections $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}' \in \operatorname{Pic}(X)$.

Some generalities on line bundles on wonderful varieties and their sections have already been recalled in Section 1, we keep on using the notation introduced in [6, Section 1].

3.1. General reductions. The surjectivity of the multiplication of sections of globally generated line bundles on a wonderful variety is known for some important families, namely for the *wonderful symmetric varieties* (see [10]), for the *wonderful*

model varieties (of simply-connected type) and for the wonderful comodel varieties (see [6]).

We will essentially reduce the proof of Theorem 3.1 to the known cases recalled above, by making use of the operations of localization, quotient and parabolic induction on spherical systems, which provide us with some reduction steps in the study of the multiplication maps. We recall such reductions, which are proved in [4] and [6], and then we will apply them to the cases under consideration.

Lemma 3.2 ([4, Lemma 2.4]). Let X be a wonderful variety and let $X' \subset X$ be a wonderful subvariety. If $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all globally generated $\mathcal{L},\mathcal{L}' \in$ $\operatorname{Pic}(X)$, then $m_{\mathcal{L},\mathcal{L}'}$ is surjective for all globally generated $\mathcal{L},\mathcal{L}' \in \operatorname{Pic}(X')$.

Lemma 3.3 ([6, Corollary 1.4]). Let X be a wonderful variety with set of colors Δ , let X' be a quotient of X by a distinguished subset $\Delta_0 \subset \Delta$ with set of colors Δ' and identify Δ' with $\Delta \setminus \Delta_0$. If $D \in \mathbb{N}\Delta$ and $\operatorname{supp}(D) \cap \Delta_0 = \emptyset$ and if $\mathcal{L}_D \in \operatorname{Pic}(X)$ and $\mathcal{L}'_D \in \operatorname{Pic}(X')$ are the line bundles corresponding to D regarded as an element in $\mathbb{N}\Delta$ and in $\mathbb{N}\Delta'$, then $\Gamma(X, \mathcal{L}_D) = \Gamma(X', \mathcal{L}'_D)$.

In particular, if $m_{D,E}$ is surjective for all $D, E \in \mathbb{N}\Delta$, then $m_{D',E'}$ is surjective for all $D', E' \in \mathbb{N}\Delta'$.

Lemma 3.4 ([6, Proposition 1.6]). Let X be a wonderful variety and suppose that X is the parabolic induction of a wonderful variety X'. Then for all $\mathcal{L}, \mathcal{L}'$ in $\operatorname{Pic}(X)$ the multiplication $m_{\mathcal{L},\mathcal{L}'}$ is surjective if and only if the multiplication $m_{\mathcal{L}|X'}, \mathcal{L}'|_{X'}$ is surjective.

We now apply previous reductions to our cases. In particular, we will show that to prove Theorem 3.1 it is enough to prove the surjectivity of the multiplication just for the following basic case, labelled $a^{x}(1,1,1)$, since the other basic cases have already been treated in [10] and [6]:

Let $\mathcal{O} \subset \mathfrak{p}$ be a spherical nilpotent *K*-orbit as in Theorem 3.1 and let *X* be the corresponding wonderful variety. When *X* is a flag variety, or equivalently $\Sigma = \emptyset$, the surjectivity of the multiplication is trivial.

By Lemma 3.4, the surjectivity of the multiplication on X is reduced to the surjectivity of the multiplication on Z, the localization of X at the subset $\operatorname{supp} \Sigma \subset S$. These localizations are described in Section 2.

In the cases


1.6 (r = s = 0 and q > 1), 1.7 (r = s = 0 and p > 1), 2.3, 2.4, 4.3, 4.4, 5.4,

treated in Section 2.1, the wonderful variety Z is a rank one wonderful variety which is homogeneous under its automorphism group (see [1] for a description of these varieties). Therefore in these cases Z is a flag variety, and the surjectivity of the multiplication is trivial. In the remaining cases of Section 2.1 the wonderful variety Z is the wonderful compactification of an adjoint symmetric variety, and the surjectivity of the multiplication holds thanks to [10].

In the cases 1.4 and 1.5, treated in Section 2.2, we get the wonderful variety Z with spherical system $a^{x}(1,1,1)$.


In the cases of Section 2.3, Z is the wonderful variety with spherical system $a^{y}(r, r) + a(t) + a^{y}(s, s)$. The surjectivity of the multiplication in this case can be reduced to the surjectivity of the multiplication for a comodel wonderful variety, which is known by [6, Theorem 5.2].

Let indeed Z' be the comodel wonderful variety of cotype $A_{2(r+t+s)}$, which is the wonderful variety with the following spherical system for a group G of semisimple type $A_{r+t+s+1} \times A_{r+t+s}$, where the set Σ of spherical roots is equal to the set of simple roots of G.



Consider the wonderful subvariety of Z' associated to $\Sigma \setminus \{\alpha_{r+1}, \ldots, \alpha_{r+t+1}\},\$

then the set of colors $\{D_{\alpha'_{s+1}}^+, D_{\alpha'_{s+2}}^\pm, \dots, D_{\alpha'_{s+t-1}}^\pm, D_{\alpha'_{s+t}}^-\}$ is distinguished, and the corresponding quotient is a parabolic induction of Z.



Therefore the surjectivity of the multiplication of Z follows from that of Z' thanks to Lemmas 3.2, 3.3 and 3.4.

3.2. Projective normality of $a^{x}(1,1,1)$. Consider X for the semisimple group $G = SL(2) \times SL(2) \times SL(2)$ with spherical system $a^{x}(1,1,1)$.

The spherical roots of X coincide with the simple roots, and we enumerate them as follows: $\sigma_1 = \alpha$, $\sigma_2 = \alpha'$, $\sigma_3 = \alpha''$. Moreover we enumerate the colors of X in the following way: $D_1 = D_{\alpha''}^+$, $D_2 = D_{\alpha}^-$, $D_3 = D_{\alpha}^+$. Then the spherical roots are expressed in terms of colors as follows:

$$\begin{aligned} \sigma_1 &= -D_1 + D_2 + D_3, \\ \sigma_2 &= D_1 - D_2 + D_3, \\ \sigma_3 &= D_1 + D_2 - D_3. \end{aligned}$$

Notice that the restriction $\omega : \operatorname{Pic}(X) \to \Lambda$ is injective in this case (see e.g. [20, Lemma 30.24]). Indeed, by identifying Λ^+ with \mathbb{N}^3 we have

$$\begin{aligned} \omega(\mathcal{L}_{D_1}) &= (0, 1, 1), \quad \omega(\mathcal{L}_{D_2}) &= (1, 0, 1), \quad \omega(\mathcal{L}_{D_3}) &= (1, 1, 0), \\ \omega(\mathcal{L}_{\sigma_1}) &= (2, 0, 0), \quad \omega(\mathcal{L}_{\sigma_2}) &= (0, 2, 0), \quad \omega(\mathcal{L}_{\sigma_3}) &= (0, 0, 2). \end{aligned}$$

Thus every $\mathcal{L} \in \operatorname{Pic}(X)$ is uniquely determined by the corresponding triple $\omega(\mathcal{L}) \in \mathbb{Z}^3$.

The surjectivity of the multiplication of sections of globally generated line bundles can be proved fairly easily in this case by making use of the machinery of low triples developed in [6], namely by classifying the covering differences of the partial ordered set $(\mathbb{N}\Delta, \leq_{\Sigma})$ and then studying the fundamental low triples. However, in order to increase the accessibility to the non-experts, we will write down a proof by more elementary language, only involving representation theoretic considerations. In any case, even though we do not mention low triples and covering differences, they are intrinsically hidden inside the arguments that appear in the present proof.

Let $H \subset G$ be the generic stabilizer of X. Identifying sections of line bundles on X with functions on G which are semiinvariant with respect to the right action of H, for every globally generated line bundle $\mathcal{L} \in \operatorname{Pic}(X)$ we can regard the space of sections $\Gamma(X, \mathcal{L})$ as a G-stable submodule of $\mathbb{C}[G]^{(H)}$.

If $m \in \mathbb{N}$, denote by V(m) the simple SL(2)-module of highest weight m, and if $\underline{m} = (m, m', m'') \in \mathbb{N}^3$ we denote $V(\underline{m}) = V(m) \otimes V(m') \otimes V(m'')$, regarded as a G-module.

Let

$$\mathbf{T} = \{ (m, m', m'') \in \mathbb{N}^3 : V(m'') \subset V(m) \otimes V(m') \}$$

be the tensor semigroup of SL(2). Then by the Clebsch-Gordan rule we have

$$\mathbf{T} = \{ (m, m', m'') \in \mathbb{N}^3 : m + m' + m'' \in 2\mathbb{N} \text{ and } |m - m'| \leq m'' \leq m + m' \}.$$

Notice that T is stable under permutations: indeed

$$V(m'') \subset V(m) \otimes V(m') \iff \left(V(m)^* \otimes V(m')^* \otimes V(m'')\right)^{\operatorname{diag}\operatorname{SL}(2)} \neq 0,$$

the latter conditon is independent on the order of the triple (m, m', m'') since $V(m) \simeq V(m)^*$ and $V(m') \simeq V(m')^*$ are self-dual modules.

In our case, we can apply the previous discussion as follows. Indeed, by [7, Case 43] we can assume that

$$H = (Z_{\mathrm{SL}(2)} \times Z_{\mathrm{SL}(2)} \times Z_{\mathrm{SL}(2)}) \operatorname{diag} \mathrm{SL}(2).$$

Since $V(\underline{m}) \simeq V(\underline{m})^*$ is self dual, we have then

$$\mathbb{C}[G]^{(H)} = \mathbb{C}[G]^{\operatorname{diag}\operatorname{SL}(2)} \simeq \bigoplus_{\underline{m} \in \mathbb{N}^3} V(\underline{m}) \otimes V(\underline{m})^{\operatorname{diag}\operatorname{SL}(2)} = \bigoplus_{\underline{m} \in \operatorname{T}} V(\underline{m}).$$

By the *G*-equivariant isomorphism $\mathbb{C}[G]^{(H)} \simeq \bigoplus_{D \in \mathbb{N}\Delta} V_D$, as a consequence of the previous description we see that the map ω identifies the semigroup of globally generated line bundles on *X* with the tensor semigroup T.

The *G*-equivariant embedding $V(\underline{m}) \longrightarrow \mathbb{C}[G]^{\operatorname{diag SL}(2)}$ can be more explicitly expressed in terms of matrix coefficients as follows. Fix for all $k \in \mathbb{N}$ an equivariant isomorphism $v \longmapsto \psi_v$ between V(k) and $V(k)^*$.

If $\underline{m} = (m, m', m'') \in T$, by the Clebsch-Gordan rule V(m'') always occurs with multiplicity one in $V(m) \otimes V(m')$, thus we have an equivariant projection

$$\pi_{m,m'}^{m''} \colon V(m) \otimes V(m') \longrightarrow V(m'')$$

which is uniquely determined up to a scalar factor.

For any simple tensor $v \otimes v' \otimes v'' \in V(m) \otimes V(m') \otimes V(m'')$ we can define a matrix coefficient $f_{v \otimes v' \otimes v''} \in \mathbb{C}[G]$ by setting, for all $(g, g', g'') \in G$,

$$f_{v\otimes v'\otimes v''}(g,g',g'')=\langle \pi_{m,m'}^{m''}(gv\otimes g'v'),g''\psi_{v''}\rangle,$$

Notice that $f_{v \otimes v' \otimes v''} \in \mathbb{C}[G]^{\operatorname{diag SL}(2)}$. Thus extending linearly we get an embedding of $V(\underline{m})$ in $\mathbb{C}[G]^{\operatorname{diag SL}(2)}$.

Define a partial order on T, by setting $(m, m', m'') \ge (n, n', n'')$ if the differences m-n, m'-n', m''-n'' are all nonnegative even numbers. This is the partial order on T induced by the partial order \leq_{Σ} on $\mathbb{N}\Delta$ via the isomorphism $T \simeq \mathbb{N}\Delta$.

For all $\underline{m} \in \mathbb{T}$, define a submodule of $\mathbb{C}[G]^{\text{diag SL}(2)}$ by setting

$$\Gamma(\underline{m}) = \bigoplus_{\underline{n} \leq \underline{m}} V(\underline{n}).$$

If $\underline{m} \in T$ and $\mathcal{L}_{\underline{m}} \in \operatorname{Pic}(X)$ is the corresponding globally generated line bundle on X, then by (1.1) we have a G-equivariant isomorphism

$$\Gamma(X, \mathcal{L}_m) \simeq \Gamma(\underline{m}).$$

If moreover $\underline{n} \in \mathbf{T}$, then we have a commutative diagram

where the upper arrow is the multiplication of sections and the lower arrow is the multiplication in the invariant ring $\mathbb{C}[G]^{\operatorname{diag SL}(2)}$.

Lemma 3.5. Let $\underline{k} = (k, k', k'')$, $\underline{m} = (m, m', m'')$, $\underline{n} = (n, n', n'') \in T$. Suppose that (m, n, k), (m', n', k'), $(m'', n'', k'') \in T$. If V(k'') occurs with multiplicity one inside $V(m) \otimes V(m') \otimes V(n) \otimes V(n')$, then $V(\underline{k}) \subset V(\underline{m}) \cdot V(\underline{n})$.

Proof. Consider the diagram

$$V(m) \otimes V(n) \otimes V(m') \otimes V(n') \xrightarrow{\pi_{m,n}^{k} \otimes \pi_{m',n'}^{k'}} V(k) \otimes V(k') \xrightarrow{\pi_{m,n'}^{m''} \otimes \pi_{n,n'}^{n''}} V(k) \otimes V(k') \xrightarrow{\pi_{m'',n''}^{k''}} V(k') \xrightarrow{\pi_{m'',n''}^{k''}} V(k')$$

Then $\pi_{k,k'}^{k''} \circ (\pi_{m,n}^k \otimes \pi_{m',n'}^{k'})$ and $\pi_{m'',n''}^{k'''} \circ (\pi_{m,m'}^{m''} \otimes \pi_{n,n'}^{n''})$ are both non-zero and equivariant for the action of SL(2). Since V(k'') occurs with multiplicity one inside $V(m) \otimes V(m') \otimes V(n) \otimes V(n')$ these two projections must be proportional, hence up to a renormalization we may assume that

$$\pi_{k,k'}^{k''} \circ (\pi_{m,n}^k \otimes \pi_{m',n'}^{k'}) = \pi_{m'',n''}^{k''} \circ (\pi_{m,m'}^{m''} \otimes \pi_{n,n'}^{n''}).$$

Let now $v \otimes v' \otimes v'' \in V(\underline{m})$ and $w \otimes w' \otimes w'' \in V(\underline{n})$ be simple tensors and consider the product of the corresponding matrix coefficients.

Then we have

$$(f_{v\otimes v'\otimes v''}f_{w\otimes w'\otimes w''})(g,g',g'') =$$

= $\langle (\pi_{m,m'}^{m''}\otimes \pi_{n,n'}^{n''})(g.(v\otimes w)\otimes g'.(v'\otimes w')), g''.(\psi_{v''}\otimes \psi_{w''})\rangle$

Since $V(k'') \subset V(m'') \otimes V(n'')$, we can find a nonzero element $u'' = \sum_h v''_h \otimes w''_h \in V(k'')$, for suitable elements $v''_h \in V(m'')$ and $w''_h \in V(n'')$. Then, up to a nonzero scalar factor, we have

$$\begin{split} \sum_{h} \left(f_{v \otimes v' \otimes v''_{h}} f_{w \otimes w' \otimes w''_{h}} \right) (g, g', g'') &= \\ &= \langle (\pi_{m,m'}^{m''} \otimes \pi_{n,n'}^{n''}) \left(g(v \otimes w) \otimes g'.(v' \otimes w') \right), \, g''.(\sum_{h} \psi_{v''_{h}} \otimes \psi_{w''_{h}}) \rangle = \\ &= \langle \pi_{m'',n''}^{k''} \circ (\pi_{m,m'}^{m''} \otimes \pi_{n,n'}^{n''}) \left(g.(v \otimes w) \otimes g'.(v' \otimes w') \right), \, g''.\psi_{u''} \rangle = \\ &= \langle \pi_{k,k'}^{k''} \circ (\pi_{m,n}^{k} \otimes \pi_{m',n'}^{k'}) \left(g.(v \otimes w) \otimes g'.(v' \otimes w') \right), \, g''.\psi_{u''} \rangle = \\ &= \langle \pi_{k,k'}^{k''} \left(g\pi_{m,n}^{k} (v \otimes w) \otimes g' \pi_{m',n'}^{k'} (v' \otimes w') \right), \, g'''\psi_{u''} \rangle, \end{split}$$

which is the matrix coefficient of

$$\pi^k_{m,n}(v\otimes w)\otimes\pi^{k'}_{m',n'}(v'\otimes w')\otimes u''\in V(k)\otimes V(k')\otimes V(k'').$$

Since $V(k) \subset V(m) \otimes V(n)$ and $V(k') \subset V(m') \otimes V(n')$, we can find nonzero elements $u = \sum_i v_i \otimes w_i \in V(m) \otimes V(n)$ and $u' = \sum_j v'_j \otimes w'_j \in V(m') \otimes V(n')$ with $u \in V(k)$ and $u' \in V(k')$, thus

$$\sum_{i,j,h} \left(f_{v_i \otimes v'_j \otimes v''_h} f_{w_i \otimes w'_j \otimes w''_h} \right) (g,g',g'') = \langle \pi_{k,k'}^{k''} (gu \otimes g'u'), g'' \psi_{u''} \rangle$$

is the matrix coefficient of a nonzero element in $V(\underline{k})$, and the claim follows. \Box

Remark 3.6. Notice that the multiplication in $\mathbb{C}[G]^{\text{diag SL}(2)}$ is degenerate in the sense of [6, Proposition 9.1], that is, the inclusion

$$V(\underline{m}) \cdot V(\underline{n}) \subset \bigoplus_{\underline{k} \in \mathcal{T} \ : \ V(\underline{k}) \subset V(\underline{m}) \otimes V(\underline{n})} V(\underline{k})$$

is not necessarily an equality for all \underline{m} and \underline{n} in T. For example, for $\underline{k} = (2, 2, 2)$ and $\underline{m} = (1, 1, 2), V(\underline{k})$ occurs in the tensor product $V(\underline{m})^{\otimes 2}$. However $V(\underline{k})$ does not occur in the symmetric product $S^2V(\underline{m})$, thus $V(\underline{k}) \not\subset V(\underline{m})^2$ inside $\mathbb{C}[G]^{\text{diag SL}(2)}$.

In terms of matrix coefficients, for $\underline{k} = (k, k', k'')$, $\underline{m} = (m, m', m'')$, $\underline{n} = (n, n', n'')$ in T with $V(\underline{k}) \subset V(\underline{m}) \otimes V(\underline{n})$, it can be easily shown that

$$V(\underline{k}) \subset V(\underline{m}) \cdot V(\underline{n}) \quad \iff \pi_{m'',n''}^{k''} \circ \left(\pi_{m,m'}^{m''} \otimes \pi_{n,n'}^{n''}\right) \circ \left(\iota_k^{m,n} \otimes \iota_{k'}^{m',n'}\right) \neq 0$$

where $\iota_k^{m,n}$ and $\iota_{k'}^{m',n'}$ denote the equivariant injections $V(k) \longrightarrow V(m) \otimes V(n)$ and $V(k') \longrightarrow V(m') \otimes V(n')$, respectively.

Proposition 3.7. Let \underline{m} and $\underline{n} \in T$, then $\Gamma(\underline{m}) \cdot \Gamma(\underline{n}) = \Gamma(\underline{m} + \underline{n})$.

14

Proof. We have to show that, for all $\underline{k} \leq \underline{m} + \underline{n}$, it holds $V(\underline{k}) \subset \Gamma(\underline{m}) \cdot \Gamma(\underline{n})$. We proceed by induction on the sum m + n + m' + n' + m'' + n''. Let $\underline{k} \in T$ with $\underline{k} \leq \underline{m} + \underline{n}$.

Suppose first that $\underline{k} = \underline{m} + \underline{n}$. If $v_{\underline{m}} \in V(\underline{m})$ and $v_{\underline{n}} \in V(\underline{n})$ are highest weight vectors, then their product $v_{\underline{m}}v_{\underline{n}}$ is a highest weight vector of weight $\underline{m} + \underline{n}$, thus $V(\underline{m} + \underline{n}) \subset V(\underline{m}) \cdot V(\underline{n}) \subset \Gamma(\underline{m}) \cdot \Gamma(\underline{n})$. Therefore we will assume that $\underline{k} < \underline{m} + \underline{n}$. Case 1. Suppose that both m and n have zero entries. Then m and n have a

unique zero entry, and up to a permutation of the coordinates we may assume that $\underline{m} = (m, 0, m)$ and $\underline{n''} > 0$. Thus we have either $\underline{n} = (n, 0, n)$ or $\underline{n} = (0, n', n')$.

Suppose that $\underline{n} = (n, 0, n)$. Then it must be $\underline{k} = (k, 0, k)$ with k < m + n, thus $\underline{k} \leq \underline{m} + \underline{n} - (2, 0, 2)$. If $\underline{m} = \underline{n} = (1, 0, 1)$, then $\underline{k} = (0, 0, 0)$ and $V(\underline{k}) \subset V(\underline{m}) \cdot V(\underline{n})$ by Lemma 3.5. Therefore we can assume that either $m \geq 2$ or $n \geq 2$. Suppose that $m \geq 2$, the other case is similar: then $(m - 2, 0, m - 2) \in T$, and by induction

$$V(\underline{k}) \subset \Gamma(m-2, 0, m-2) \cdot \Gamma(\underline{n}) \subset \Gamma(\underline{m}) \cdot \Gamma(\underline{n})$$

Suppose now that $\underline{n} = (0, n', n')$. Then $\underline{k} < (m, n', m+n')$, and since $k'' \leq k+k' \leq m+n'$ it must be k'' < m+n'. If k+k' = m+n' or if m=n'=1, then $\underline{k} = (m, n', k'')$ and by Lemma 3.5 we get $V(\underline{k}) \subset V(\underline{m}) \cdot V(\underline{n})$. Thus we can assume that k+k' < m+n', and that either $m \geq 2$ or $n' \geq 2$. Suppose we are in the first case, the other one is similar. Then $(m-2, 0, m-2) \in T$ and $\underline{k} \leq (m-2, m-2, 0) + \underline{n}$, and the claim follows again by the induction.

Case 2. Suppose that k'' < m'' + n'' and that either m'' > |m - m'| or n'' > |n - n'|. We can assume that we are in the first case. Then $(m, m', m'' - 2) \in \mathbb{T}$ and $\underline{k} \leq (m, m', m'' - 2) + \underline{n}$, thus by induction

$$V(\underline{k}) \subset \Gamma(m, m', m'' - 2) \cdot \Gamma(\underline{n}) \subset \Gamma(\underline{m}) \cdot \Gamma(\underline{n}).$$

Notice that this covers the case $\underline{k} = (0, 0, 0)$. Indeed by Case 1 we can assume that either \underline{m} or \underline{n} has no zero entry. Therefore, up to a permutation of the coordinates, we always have either m'' > |m - m'| or n'' > |n - n'|.

Case 3. Suppose that $\underline{k} \neq (0, 0, 0)$. Up to a permutation of the coordinates we can assume that k'' < k + k'.

Assume that m'' = m + m', and n'' = n + n'. By Case 1 and by symmetry we can also assume that all the entries of \underline{m} are nonzero. Then $(m, m', m'' - 2) \in \mathbb{T}$. Moreover by assumption we have $k'' < k + k' \leq m + n + m' + n' = m'' + n''$. Therefore $\underline{k} \leq (m, m', m'' - 2) + \underline{n}$, and by induction

$$V(\underline{k}) \subset \Gamma(m, m', m'' - 2) \cdot \Gamma(\underline{n}) \subset \Gamma(\underline{m}) \cdot \Gamma(\underline{n}).$$

Assume now that either m'' < m + m' or n'' < n + n'. We can assume that we are in the first case. Then (m - 1, m' - 1, m'') and (k - 1, k' - 1, k'') both belong to T and $(k - 1, k' - 1, k'') < (m - 1, m' - 1, m'') + \underline{n}$, thus by induction

$$V(k-1, k-1, k'') \subset \Gamma(m-1, m'-1, m'') \cdot \Gamma(\underline{n}).$$

Therefore

$$V(\underline{k}) \subset V(1,1,0) \cdot V(k-1,k'-1,k'') \subset V(1,1,0) \cdot \Gamma(m-1,m'-1,m'') \cdot \Gamma(\underline{n}),$$

and the claim follows because $V(1,1,0) \cdot \Gamma(m-1,m'-1,m'') \subset \Gamma(\underline{m}).$

Corollary 3.8. The multiplication $m_{D,E}$ is surjective for all $D, E \in \mathbb{N}\Delta$.

4. Normality and weight semigroups

Let $K \subset G$ be a Hermitian symmetric subgroup. As it is well known (see e.g. [19, Section 5.5]), this is equivalent to require that K is the Levi factor of a parabolic subgroup Q of G with Abelian unipotent radical. This implies that $Q \subset G$ is a maximal parabolic subgroup (we assume G to be almost simple).

Recall that we have fixed a maximal torus T in K and a Borel subgroup B of K containing T. We denote by $\mathcal{X}(T)$ the weight lattice of T. Recall the decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$.

The torus T is also a maximal torus in G. We can choose a Borel subgroup of G containing B, the Borel subgroup of K, and contained in Q.

If Q_{-} denotes the opposite parabolic subgroup, then we get the K-module decomposition

$$\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$$

where \mathfrak{p}_1 (resp. \mathfrak{p}_2) is the Lie algebra of the unipotent radical of Q (resp. Q_-). Notice that \mathfrak{p}_1 and \mathfrak{p}_2 are irreducible K-modules, dual to each other. More precisely, if $\theta_G \in \mathcal{X}(T)$ denotes the highest root of G (w.r.t. the above choice of a Borel subgroup in G), then $\mathfrak{p}_1 = V(\theta_G)$ is the irreducible K-module of highest weight θ_G , and $\mathfrak{p}_2 = V(\theta_G)^*$ is the irreducible K-module of lowest weight $-\theta_G$.

We denote by Z_K the identity component of the center of K, and by \mathfrak{z}_K its Lie algebra. Since G/K is a Hermitian symmetric space, dim $Z_K = 1$.

Proposition 4.1. Z_K acts non-trivially on \mathfrak{p}_1 and \mathfrak{p}_2 .

Proof. Let $z(\xi)$ ($\xi \in \mathbb{C}^*$) be a parametrization of $Z_K \simeq \mathbb{C}^*$. Since $\mathfrak{p}_2 \simeq \mathfrak{p}_1^*$, it follows that $z(\xi).e = \xi^m e_1 + \xi^{-m} e_2$ with $m \in \mathbb{Z}$. Suppose that m = 0: then Z_K acts trivially on $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$, hence it acts trivially on \mathfrak{g} since it acts trivially on \mathfrak{k} . Therefore Z_K is in the center of G, which is absurd since G is semisimple. \Box

It follows that \mathfrak{p}_1 and \mathfrak{p}_2 are not isomorphic as representations of K. Let χ be the character of Z_K acting on \mathfrak{p}_1 . By making use of the classification of the standard parabolic subgroups of G with Abelian unipotent radical, we can describe this character explicitly.

Indeed, let $S_G = \{\alpha_1, \ldots, \alpha_n\}$ be the set of simple roots of G, and denote by $[\theta_G : \alpha_i]$ the coefficient of α_i in θ_G . Then a standard parabolic subgroup $Q \subset G$ has an Abelian unipotent radical if and only if it is maximal, corresponding to a root $\alpha_p \in S_G$ such that $[\theta_G : \alpha_p] = 1$. In the following list we give all the simple roots (of irreducible root systems of classical type) with this property:

- (1) If G is of type $A_n: \alpha_1, \ldots, \alpha_n;$
- (2) If G is of type B_n : α_1 ;
- (3) If G is of type C_n : α_n ;
- (4) If G is of type D_n : $\alpha_1, \alpha_{n-1}, \alpha_n$.

Let $\mathfrak{t}_G \subset \mathfrak{g}$ be the Cartan subalgebra generated by the fundamental coweights $\omega_1^{\vee}, \ldots, \omega_n^{\vee}$, and let $\mathfrak{t}_K^{\mathrm{ss}} \subset \mathfrak{t}_G$ be the subalgebra generated by the simple coroots of

K. Since K is the Levi subgroup of G defined by the set of simple roots $S_G \setminus \{\alpha_p\}$, it follows that \mathfrak{z}_K , which is by definition the annihilator of $(\mathfrak{t}_K^{ss})^*$ in \mathfrak{t}_G , is generated by the fundamental coweight ω_p^{\vee} .

On the other hand, assuming G simply connected, we have that the cocharacter lattice $\mathcal{X}(T)^{\vee}$ is equal to the coroot lattice $\mathbb{Z}S^{\vee}$, therefore

$$\mathcal{X}(Z_K)^{\vee} = \mathfrak{z}_K \cap \mathbb{Z}S^{\vee}$$

is generated by $m\omega_p^{\vee}$ where $m \in \mathbb{N}$ is the minimum such that $m\omega_p^{\vee} \in \mathbb{Z}S^{\vee}$. We list the value of m here below for all the possible cases of G and α_p :

- (A_n, α_p) : $m = (n + 1)/\gcd(p, n + 1)$; - (B_n, α_1) , (C_n, α_n) , (D_n, α_1) : m = 2; - (D_n, α_{n-1}) , (D_n, α_n) : m = 2 if n is even, m = 4 if n is odd.

If $z(\xi)$ is the 1-parameter-subgroup in Z_K given by $m\omega_p^{\vee}$, it follows then

$$\chi(z(\xi)) = \xi^{m\theta_G(\omega_p^{\vee})} = \xi^m$$

In the following proposition we will show that if $Ke \subset \mathfrak{p}$ is a nilpotent orbit, then $\overline{Ke} \subset \mathfrak{p}$ is a bicone with respect to the decomposition $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$. This will allow us to study the normality of \overline{Ke} by making use of Theorem 1.2.

Proposition 4.2. Write $e = e_1 + e_2$ with $e_1 \in \mathfrak{p}_1$ and $e_2 \in \mathfrak{p}_2$. Then $\xi_1 e_1 + \xi_2 e_2 \in Ke$ for all $\xi_1, \xi_2 \in \mathbb{C}^*$.

Proof. Let $\{e, f, h\}$ be a normal \mathfrak{sl}_2 -triple containing e, then $h \in K$ and [h, e] = 2e. If $t(\xi) = \exp(\xi h)$ ($\xi \in \mathbb{C}^*$) is the one parameter subgroup of K obtained exponentiating the line generated by h, it follows that $t(\xi).e = \xi e$, namely $t(\xi).e_1 = \xi e_1$ and $t(\xi).e_2 = \xi e_2$.

On the other hand, by the previous proposition, the connected component Z_K of the center of K acts non-trivially on \mathfrak{p}_1 , therefore we can take a parametrization $z(\xi)$ of Z ($\xi \in \mathbb{C}^*$) such that $z(\xi).e = \xi^m e_1 + \xi^{-m} e_2$ with $m \neq 0$. It follows that every combination of e_1 and e_2 with non-zero coefficients can be written in the form $t(\xi)z(\xi').e$ for some $\xi, \xi' \in \mathbb{C}^*$.

Let X be the wonderful compactification of $K/N_K(K_e)$, and denote by Σ and by Δ its set of spherical roots and its set of colors. For i = 1, 2, let $\pi_i \colon \mathfrak{p} \longrightarrow \mathfrak{p}_i$ be the projections corresponding to the decomposition $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$. If $\pi_i(e) \neq 0$, we denote by $D_{\mathfrak{p}_i} \in \mathbb{N}\Delta$ the element such that $\mathfrak{p}_i = V_{D_{\mathfrak{p}_i}}^*$, and the image of \overline{Ke} in $\mathbb{P}(\mathfrak{p}_i)$ coincides with the image of the corresponding map $\phi_{D_{\mathfrak{p}_i}} \colon X \longrightarrow \mathbb{P}(\mathfrak{p}_i)$. In particular, if e projects non-trivially both on \mathfrak{p}_1 and \mathfrak{p}_2 , then the image of \overline{Ke} in $\mathbb{P}(\mathfrak{p}_1) \times \mathbb{P}(\mathfrak{p}_2)$ coincides with the image of X mapped diagonally via $\phi_{D_{\mathfrak{p}_1}}$ and $\phi_{D_{\mathfrak{p}_2}}$. For convenience we also set $D_{\mathfrak{p}_i} = 0$ if $\pi_i(e) = 0$, and we denote

$$\Delta_{\mathfrak{p}}(e) = \{D_{\mathfrak{p}_1}, D_{\mathfrak{p}_2}\}.$$

By Theorem 3.1 the multiplication of sections of globally generated line bundles on the wonderful compactification of $K/N_K(K_e)$ is surjective, hence by Theorem 1.3 it follows that $\overline{Ke} \subset \mathfrak{p}$ is normal if and only if every $D \in \Delta_{\mathfrak{p}}(e)$ is minuscule in $\mathbb{N}\Delta$ with respect to the partial order \leq_{Σ} , or zero. Below we will see that this condition is always fulfilled, hence we get the following. **Theorem 4.3.** Let $(\mathfrak{g}, \mathfrak{k})$ be a classical symmetric pair of Hermitian type and let $Ke \subset \mathfrak{p}$ be a spherical nilpotent orbit. Then \overline{Ke} is normal.

Remark 4.4. The normality of \overline{Ge} is well known and may be deduced from [17]. In particular, if $(\mathfrak{g}, \mathfrak{k})$ is a classical symmetric pair of Hermitian type and Ke is a spherical nilpotent orbit in \mathfrak{p} , then \overline{Ge} is always normal.

Let us denote by $\Gamma(X)$ the weight semigroup of a K-spherical variety X,

$$\Gamma(X) = \{\lambda \in \mathcal{X}(T) : \operatorname{Hom}(\mathbb{C}[X], V(\lambda)) \neq 0\}.$$

Denoting for i = 1, 2 the highest weight of \mathfrak{p}_i^* as a *G*-module by λ_i^* , the previous theorem together with Theorem 1.3 imply that $\Gamma(\overline{Ke})$ consists of the weights

$$n_1\lambda_1^* + n_2\lambda_2^* - (n_1D_{\mathfrak{p}_1} + n_2D_{\mathfrak{p}_2} - E)$$

for $(n_1, n_2) \in \mathbb{N}^2$, $E \in \mathbb{N}\Delta$ with $E \leq_{\Sigma} n_1 D_{\mathfrak{p}_1} + n_2 D_{\mathfrak{p}_2}$, see equation (1.2).

Beyond showing the normality of \overline{Ke} , we obtain the weight semigroups $\Gamma(\overline{Ke})$ by computing the corresponding semigroups

$$\Gamma_{\Delta_{\mathfrak{p}}(e)} = \{ (n_1, n_2, E) \in \mathbb{N}^2 \times \mathbb{N}\Delta : E \leq_{\Sigma} n_1 D_{\mathfrak{p}_1} + n_2 D_{\mathfrak{p}_2} \}.$$

The generators of the weight semigroup $\Gamma(\overline{Ke}) = \Gamma(\widetilde{Ke})$ are given in Tables 1–6, in Appendix B. In the same tables we also provide the codimension of $\overline{Ke} \smallsetminus Ke$ in \overline{Ke} . Notice that, if \overline{Ke} is normal and the codimension of $\overline{Ke} \smallsetminus Ke$ in \overline{Ke} is greater than 1, then $\mathbb{C}[\overline{Ke}] = \mathbb{C}[Ke]$, so that the weight semigroup of Ke actually coincides with $\Gamma(\overline{Ke})$.

We now report the details of the computation of the semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}$. We omit the cases where X is a flag variety or a parabolic induction of a wonderful symmetric variety (see Section 2): in these cases the combinatorics of the spherical systems is easier. By [14], the normality of \overline{Ke} was already known in all these cases, since they all satisfy $\operatorname{ht}_{\mathfrak{p}}(e) = 2$ (see Appendix B). Some of the corresponding weight semigroups $\Gamma(\overline{Ke})$ were obtained in [3] by using different techniques.

Remark 4.5. In [18], for the complex symmetric pair $(SL(p+q), S(GL(p) \times GL(q))))$, K. Nishiyama gave a description of the coordinate rings of the closures of some special spherical orbits, those which can be obtained as *theta lift in the stable range*. Actually, in that symmetric pair, the only spherical orbits which are not theta lifts in the stable range correspond to the following cases: 1.4, 1.5, 1.6 (r + s = q - 1)and 1.7 (r + s = p - 1).

Notation. For all $E = \sum_{D \in \Delta} k_D D \in \mathbb{Z}\Delta$, define its positive part $E^+ = \sum_{k_D > 0} k_D D$ and its height $\operatorname{ht}(E) = \sum_{D \in \Delta} k_D$.

4.1. Cases 1.4 and 1.5. We consider the case 1.4, the other one is analogous. Let $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$ be the set of spherical roots and $\Delta = \{D_1, D_2, D_3, D_4, D_5\}$ the set of colors of X, where we denote

 $\sigma_1 = \alpha', \quad \sigma_2 = \alpha_1, \quad \sigma_3 = \alpha_{p-1}$

$$D_1 = D_{\alpha'}^+, \quad D_2 = D_{\alpha'}^-, \quad D_3 = D_{\alpha_1}^+, \quad D_4 = D_{\alpha_{p-2}}, \quad D_5 = D_{\alpha_2}$$
(if $p = 4, D_4 = D_5$).

We have in this case $\mathfrak{p}_1 = V(\omega_1 + \omega' + \chi)$ and $\mathfrak{p}_2 = V(\omega_{p-1} + \omega' - \chi)$, $D_{\mathfrak{p}_1} = D_1$ and $D_{\mathfrak{p}_2} = D_2$. One can easily see that every covering difference $\gamma \in \mathbb{N}\Sigma$ is a simple root or the sum of two simple roots, thus satisfies $\operatorname{ht}(\gamma^+) = 2$. In particular every $D \in \mathbb{N}\Delta$ is minuscule, therefore Theorem 1.2 implies that \overline{Ke} is normal.

Proposition 4.6. The semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ is generated by

 $(1,0,D_1), (0,1,D_2), (1,1,D_3), (2,0,D_4), (0,2,D_5).$

Remark 4.7. Notice that if $D_{\mathfrak{p}_1}$ and $D_{\mathfrak{p}_2}$ are two distinct elements of Δ in order to compute generators for $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ it is actually enough to compute generators for the semigroup

$$\Gamma_{\Delta_{\mathfrak{p}}(e)}^{\Sigma} = \left\{ \gamma \in \mathbb{N}\Sigma : \operatorname{supp}(\gamma^{+}) \subset \{D_{\mathfrak{p}_{1}}, D_{\mathfrak{p}_{2}}\} \right\},\$$

which is the image of the homomorphism $\Gamma_{\Delta_{\mathfrak{p}}(e)} \longrightarrow \mathbb{N}\Sigma$ defined by $(n_1, n_2, E) \mapsto n_1 D_{\mathfrak{p}_1} + n_2 D_{\mathfrak{p}_2} - E$. For every generator $\gamma \in \Gamma_{\Delta_{\mathfrak{p}}(e)}^{\Sigma}$ there exists a unique minimal triple mapping to γ , and $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ is generated by such triples together with $(1, 0, D_{\mathfrak{p}_1})$ and $(0, 1, D_{\mathfrak{p}_2})$.

Proof of Proposition 4.6. Let us show that $\Gamma^{\Sigma}_{\Delta_n(e)}$ is generated by

$$D_1 + D_2 - D_3 = \sigma_1, \ 2D_1 - D_4 = \sigma_1 + \sigma_3, \ 2D_2 - D_5 = \sigma_1 + \sigma_2.$$

Indeed, these are generators of the semigroup

 $\{a_1\sigma_1 + a_2\sigma_2 + a_3\sigma_3 \in \mathbb{N}\Sigma : a_1 \geqslant a_2 + a_3\}$

and the condition $a_1 \ge a_2 + a_3$ is just equivalent to requiring the non-positivity of the coefficient of D_3 in $a_1\sigma_1 + a_2\sigma_2 + a_3\sigma_3$ (written as an element of $\mathbb{Z}\Delta$), which is equal to $-a_1 + a_2 + a_3$.

4.2. Cases 1.6 and 1.7. We consider the case 1.6, the other one is analogous.

4.2.1. We assume first r + s < q - 1, the case r + s = q - 1 will be treated below, separately.

For $i = 1, \ldots, r$, we denote $\sigma_{2i-1}^1 = \alpha_{p-i}$ and $\sigma_{2i}^1 = \alpha'_i$. Similarly, for $i = 1, \ldots, s$, we denote $\sigma_{2i-1}^2 = \alpha_i$ and $\sigma_{2i}^2 = \alpha'_{q-i}$. Finally, we denote $\tau = \alpha'_{r+1} + \ldots + \alpha'_{q-s-1}$. Then

$$\Sigma = \{\sigma_1^1, \dots, \sigma_{2r}^1, \sigma_1^2, \dots, \sigma_{2s}^2, \tau\}$$

For the set of colors we introduce the following notation. For all $h \leq 2r + 2$, set

$$D_{h}^{1} = \begin{cases} D_{\alpha_{p-i}}^{-} & \text{if } h = 2i - 1, \text{ for } i \leqslant r \\ D_{\alpha_{p-i}}^{+} & \text{if } h = 2i, \text{ for } i \leqslant r \\ D_{\alpha'_{r}}^{-} & \text{if } h = 2r + 1 \\ D_{\alpha_{p-r-1}}^{-} & \text{if } h = 2r + 2 \end{cases}$$

For all $h \leq 2s + 2$, set

$$D_{h}^{2} = \begin{cases} D_{\alpha_{i}}^{-} & \text{if } h = 2i - 1, \text{ for } i \leqslant s \\ D_{\alpha_{i}}^{+} & \text{if } h = 2i, \text{ for } i \leqslant s \\ D_{\alpha_{i}'}^{-} & \text{if } h = 2s + 1 \\ D_{\alpha_{s+1}}^{-} & \text{if } h = 2s + 2 \end{cases}$$

Notice that if p = r + s + 1 then $D_{2r+2}^1 = D_{2s+2}^2$.

We also set

$$D_{2r+3}^{1} = \begin{cases} D_{\alpha'_{s+1}} & \text{if } r+s < q-2 \\ D_{\tau}^{+} & \text{if } r+s = q-2 \end{cases}, \quad D_{2s+3}^{2} = \begin{cases} D_{\alpha'_{q-s-1}} & \text{if } r+s < q-2 \\ D_{\tau}^{-} & \text{if } r+s = q-2 \end{cases}$$

(if r + s = q - 2, the spherical root τ is equal to a simple root, $\tau = \alpha'_{r+1} = \alpha'_{q-s-1}$, we assume $c(D^+_{\tau}, \alpha'_r) = c(D^-_{\tau}, \alpha'_{q-s}) = -1$).

Therefore

$$\Delta = \{ D_1^1, \dots, D_{2r+3}^1, D_1^2, \dots, D_{2s+3}^2 \}.$$

Let us suppose r, s > 0. We have $D_{\mathfrak{p}_1} = D_2^1$ and $D_{\mathfrak{p}_2} = D_2^2$. As explained at the end of Section 3.1, X is a parabolic induction of a quotient of a localization of a comodel wonderful variety of cotype A, therefore by [6, Proposition 3.2] every covering difference $\gamma \in \mathbb{N}\Sigma$ satisfies $\operatorname{ht}(\gamma^+) = 2$. In particular every element $D \in \Delta$ is minuscule, therefore Theorem 1.2 implies that \overline{Ke} is normal.

For notational purposes, set $r_1 = r$ and $r_2 = s$. For k = 1, 2 and $h \leq 2r_k + 2$, we denote

$$\tilde{D}_{h}^{k} = \begin{cases} D_{h}^{k} & \text{if } h < 2r_{k} + 1\\ D_{h}^{k} + D_{h+1}^{k} & \text{if } h = 2r_{k} + 1, 2r_{k} + 2 \end{cases}$$

Proposition 4.8. The semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ is generated by the elements $(i, 0, \tilde{D}_{2i}^1)$ for $i \leq r_1 + 1$, $(0, j, \tilde{D}_{2j}^2)$ for $j \leq r_2 + 1$ and $(i, j, \tilde{D}_{2i-1}^1 + \tilde{D}_{2j-1}^2)$ for $i \leq r_1 + 1$, $j \leq r_2 + 1$.

Proof. As noticed in Remark 4.7, it is enough to compute generators for the semigroup

$$\Gamma^{\Sigma}_{\Delta_{\mathfrak{p}}(e)} = \left\{ \gamma \in \mathbb{N}\Sigma : \operatorname{supp}(\gamma^{+}) \subset \{D_{2}^{1}, D_{2}^{2}\} \right\}.$$

Notice that, for k = 1, 2 and $i = 2, \ldots, r_k + 1$, it holds

$$\sigma_1^k + \ldots + \sigma_{2i-2}^k = D_2^k + \tilde{D}_{2i-2}^k - \tilde{D}_{2i}^k.$$

Therefore,

$$\gamma_i^k := \sum_{u=1}^{i-1} (i-u)(\sigma_{2u-1}^k + \sigma_{2u}^k)$$

is equal to $iD_2^k - \tilde{D}_{2i}^k$.

Notice also that, for $i \leq r_1 + 1$ and $j \leq r_2 + 1$,

$$\sum_{u=i}^{r_1} \sigma_{2u}^1 + \sum_{v=j}^{r_2} \sigma_{2v}^2 + \tau = \tilde{D}_{2i}^1 + \tilde{D}_{2j}^2 - \tilde{D}_{2i-1}^1 - \tilde{D}_{2j-1}^2$$

Therefore,

$$\gamma_{i,j} := \sum_{u=1}^{i-1} (i-u)(\sigma_{2u-1}^1 + \sigma_{2u}^1) + \sum_{u=i}^{r_1} \sigma_{2u}^1 + \sum_{v=1}^{j-1} (j-v)(\sigma_{2v-1}^2 + \sigma_{2v}^2) + \sum_{v=j}^{r_2} \sigma_{2v}^2 + \tau$$

is equal to $iD_2^1 + jD_2^2 - \tilde{D}_{2i-1}^1 - \tilde{D}_{2j-1}^2$.

We claim that the semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}^{\Sigma}$ is generated by the elements of the form γ_i^k , for k = 1, 2 and $2 \leq i \leq r_k + 1$, and $\gamma_{i,j}$, for $1 \leq i \leq r_1 + 1$ and $1 \leq j \leq r_2 + 1$.

20

Let us write $\gamma = \sum_{h=1}^{2r_1} a_h^1 \sigma_h^1 + \sum_{h=1}^{2r_2} a_h^2 \sigma_h^2 + b\tau$ as an element of N Σ . Let us denote by d_h^k the coefficient of D_h^k in γ (written as element of $\mathbb{Z}\Delta$), for $h \neq 2r_k + 2$ there is no ambiguity. We have

$$d_1^k = a_1^k - a_2^k, \quad d_h^k = -a_{h-2}^k + a_{h-1}^k + a_h^k - a_{h+1}^k \ (3 \le h \le 2r_k - 1),$$

$$d_{2r_k}^k = -a_{2r_k-2}^k + a_{2r_k-1}^k + a_{2r_k}^k, \quad d_{2r_k+1}^k = -a_{2r_k-1}^k + a_{2r_k}^k - b, \quad d_{2r_k+3}^k = -a_{2r_k}^k + b_{2r_k+3}^k + b_{2r_k+3}^k - b_{2r_k+3}^k = -a_{2r_k}^k + b_{2r_k+3}^k - b_{2r_k+3$$

Furthermore, every spherical root lies in the lattice generated by $\tilde{D}_{1}^{k}, \ldots, \tilde{D}_{2r_{k}+2}^{k}$, with $k \in \{1, 2\}$. Denoting by \tilde{d}_{h}^{k} the coefficient of \tilde{D}_{h}^{k} in γ , we have $\tilde{d}_{2r_{k}+1}^{k} = d_{2r_{k}+1}^{k}$ and $\tilde{d}_{2r_{k}+2}^{k} = d_{2r_{k}+3}^{k}$, with $k \in \{1, 2\}$.

Assume $\operatorname{supp}(\gamma^+) \subset \{D_2^1, D_2^2\}$, then the coefficients \tilde{d}_h^k are non-positive for $h \neq 2$. Let us write γ as a combination with non-negative integer coefficients of the γ_i^k $(k = 1, 2 \text{ and } 2 \leq i \leq r_k + 1)$ and the $\gamma_{i,j}$ $(1 \leq i \leq r_1 + 1 \text{ and } 1 \leq j \leq r_2 + 1)$. We have

$$\sum_{i=1}^{r_1+1} \tilde{d}_{2i-1}^1 = -b = \sum_{j=1}^{r_2+1} \tilde{d}_{2j-1}^2.$$

Therefore, there exist integers $c_{i,j} \leq 0$ (for $i \leq r_1 + 1$ and $j \leq r_2 + 1$) such that $\sum_j c_{i,j} = \tilde{d}_{2i-1}^1$ and $\sum_i c_{i,j} = \tilde{d}_{2j-1}^2$. Indeed, for k = 1, 2 and $i \leq r_k$, we can set $n_i^k = -\sum_{u=1}^i \tilde{d}_{2u-1}^k$ and take

$$-c_{i,j} = \operatorname{card}\{n \in \mathbb{N} \mid n_{i-1}^1 < n \leq n_i^1 \text{ and } n_{j-1}^2 < n \leq n_j^2\}.$$

We claim that γ is equal to

$$\sum_{i=2}^{r_1+1} -\tilde{d}_{2i}^1 \gamma_i^1 + \sum_{j=2}^{r_2+1} -\tilde{d}_{2j}^2 \gamma_j^2 + \sum_{i=1}^{r_1+1} \sum_{j=1}^{r_2+1} -c_{i,j} \gamma_{i,j}.$$

Indeed, the coefficient of σ_{2i-1}^1 in the above expression is equal to

$$\sum_{u=i+1}^{r_1+1} -\tilde{d}_{2u}^1(u-i) + \sum_{u=i+1}^{r_1+1} \sum_{v=1}^{r_2+1} -c_{u,v}(u-i)$$
$$= \sum_{u=i+1}^{r_1+1} -(\tilde{d}_{2u}^1 + \tilde{d}_{2u-1}^1)(u-i)$$
$$= a_{2i-1}^1.$$

The coefficient of σ_{2i}^1 is equal to

$$\begin{split} &\sum_{u=i+1}^{r_1+1} -\tilde{d}_{2u}^1(u-i) + \sum_{u=1}^{i} \sum_{v=1}^{r_2+1} -c_{u,v} + \sum_{u=i+1}^{r_1+1} \sum_{v=1}^{r_2+1} -c_{u,v}(u-i) \\ &= \sum_{u=1}^{i} -\tilde{d}_{2u-1}^1 + \sum_{u=i+1}^{r_1+1} -(\tilde{d}_{2u}^1 + \tilde{d}_{2u-1}^1)(u-i) \\ &= a_{2i}^1. \end{split}$$

Analogously, the same holds for σ_h^2 , for any h. It remains the coefficient of τ , which is equal to

$$\sum_{i=1}^{r_1+1} \sum_{j=1}^{r_2+1} -c_{i,j} = b.$$

The case r = s = 0 is a parabolic induction of a wonderful symmetric variety. We are left with the case r > 0 and s = 0 (the other one, r = 0 and s > 0, is analogous). Let us keep the same notation as above, notice that there exists no D_1^2 and we have

$$\Delta = \{D_1^1, \dots, D_{2r+3}^1\} \cup \{D_2^2, D_3^2\}.$$

In this case, $D_{\mathfrak{p}_1} = D_2^1$ and $D_{\mathfrak{p}_2} = \tilde{D}_2^2 = D_2^2 + D_3^2$. Both are minuscule, and \overline{Ke} is normal.

The description of the $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ given in the above proposition remains valid. The proof is slightly simpler: every spherical root lies in the lattice generated by $\tilde{D}_1^1, \ldots, \tilde{D}_{2r+2}^1$ and $\tilde{D}_1^2, \tilde{D}_2^2$ which are still linearly independent, denoting by \tilde{d}_h^k the coefficient of \tilde{D}_h^k in γ , the semigroup

$$\left\{\gamma \in \mathbb{N}\Sigma : \tilde{d}_h^1 \leqslant 0 \ \forall \ h \neq 2\right\}$$

is generated by the elements of the form γ_i^1 , for $2 \leq i \leq r+1$, and $\gamma_{i,1}$, for $1 \leq i \leq r+1$.

4.2.2. We now consider the case r + s = q - 1.

Let us keep the same notation as above, as far as possible. Indeed, there exists no $\tau,$ so we have

$$\Sigma = \{\sigma_1^1, \dots, \sigma_{2r}^1, \sigma_1^2, \dots, \sigma_{2s}^2\}$$

and

$$\Delta = \{D_1^1, \dots, D_{2r+2}^1, D_1^2, \dots, D_{2s+2}^2\}$$

Let us suppose r, s > 0. We have $D_{\mathfrak{p}_1} = D_2^1$ and $D_{\mathfrak{p}_2} = D_2^2$, which as in previous case are minuscule. Therefore, \overline{Ke} is normal.

For convenience we also define $D_{2r+3}^1 = D_{2s+1}^2$ and $D_{2s+3}^2 = D_{2r+1}^1$. As in previous case, set $r_1 = r$ and $r_2 = s$. If k = 1, 2 and $h \leq 2r_k + 2$, denote

$$\tilde{D}_{h}^{k} = \begin{cases} D_{h}^{k} & \text{if } h < 2r_{k} + 1\\ D_{h}^{k} + D_{h+1}^{k} & \text{if } h = 2r_{k} + 1, 2r_{k} + 2 \end{cases}$$

Notice that if p = q+1 then $D_{2r_1+2}^1 = D_{2r_2+2}^2$, thus $\tilde{D}_{2r_1+2}^1 = \tilde{D}_{2r_2+1}^2$ and $\tilde{D}_{2r_2+2}^2 = \tilde{D}_{2r_1+1}^1$.

Proposition 4.9. The semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ is generated by the elements $(i, 0, \tilde{D}_{2i}^1)$ for $i \leq r_1 + 1$, $(0, j, \tilde{D}_{2j}^2)$ for $j \leq r_2 + 1$ and $(i, j, \tilde{D}_{2i-1}^1 + \tilde{D}_{2j-1}^2)$ for $i \leq r_1 + 1$, $j \leq r_2 + 1$ with $i + j < r_1 + r_2 + 2$.

Proof. We follow the line of the proof of the previous proposition. As in that case, it is enough to compute generators for $\Gamma_{\Delta_n}^{\Sigma}(e)$.

For k = 1, 2 and $i = 2, \ldots, r_k + 1$, we have

$$\gamma_i^k := \sum_{u=1}^{i-1} (i-u)(\sigma_{2u-1}^k + \sigma_{2u}^k) = iD_2^k - \tilde{D}_{2i}^k.$$

22

For $i \leq r_1 + 1$ and $j \leq r_2 + 1$ with $i + j < r_1 + r_2 + 2$, we have

$$\begin{aligned} \gamma_{i,j} &:= \sum_{u=1}^{i-1} (i-u)(\sigma_{2u-1}^1 + \sigma_{2u}^1) + \sum_{u=i}^{r_1} \sigma_{2u}^1 + \sum_{v=1}^{j-1} (j-v)(\sigma_{2v-1}^2 + \sigma_{2v}^2) + \sum_{v=j}^{r_2} \sigma_{2v}^2 \\ &= iD_2^1 + jD_2^2 - \tilde{D}_{2i-1}^1 - \tilde{D}_{2j-1}^2. \end{aligned}$$

Let us prove that $\Gamma_{\Delta_{\mathfrak{p}}(e)}^{\Sigma}$ is generated by the elements of the form γ_i^k , for k = 1, 2and $2 \leq i \leq r_k + 1$, and $\gamma_{i,j}$, for $1 \leq i \leq r_1 + 1$ and $1 \leq j \leq r_2 + 1$ with $i+j < r_1 + r_2 + 2$.

Let us write $\gamma = \sum_{h=1}^{2r_1} a_h^1 \sigma_h^1 + \sum_{h=1}^{2r_2} a_h^2 \sigma_h^2 \in \mathbb{N}\Sigma$ and denote by d_h^k the coefficient of D_h^k in γ , for $h < 2r_k + 2$. We have

$$\begin{aligned} d_1^k &= a_1^k - a_2^k, \quad d_h^k = -a_{h-2}^k + a_{h-1}^k + a_h^k - a_{h+1}^k \ (3 \leqslant h \leqslant 2r_k - 1), \\ d_{2r_k}^k &= -a_{2r_k-2}^k + a_{2r_k-1}^k + a_{2r_k}^k, \\ d_{2r_1+1}^1 &= -a_{2r_1-1}^1 + a_{2r_1}^1 - a_{2r_2}^2, \quad d_{2r_2+1}^2 = -a_{2r_2-1}^2 + a_{2r_2}^2 - a_{2r_1}^1. \end{aligned}$$

Assume supp $(\gamma^+) \subset \{D_2^1, D_2^2\}$, then the coefficients d_h^k are non-positive for $h \neq 2$. We have

$$\sum_{i=1}^{r_k} d_{2i-1}^k = a_{2r_k-1}^k - a_{2r_k}^k, \quad \sum_{i=1}^{r_1+1} d_{2i-1}^1 = -a_{2r_2}^2, \quad \sum_{j=1}^{r_2+1} d_{2j-1}^2 = -a_{2r_1}^1,$$

and moreover

$$d_{2r_1+1}^1 + d_{2r_2+1}^2 = -a_{2r_1-1}^1 - a_{2r_2-1}^2.$$

Therefore, there exist non-positive integers c^1 , c^2 and $c_{i,j}$, for $i \leq r_1 + 1$ and $j \leq r_2 + 1$ with $i + j < r_1 + r_2 + 2$, such that

$$\sum_{j=1}^{r_2+1} c_{i,j} = d_{2i-1}^1 \ \forall \ i \leqslant r_1, \qquad \sum_{i=1}^{r_1+1} c_{i,j} = d_{2j-1}^2 \ \forall \ j \leqslant r_2,$$
$$c^1 + \sum_{i=1}^{r_1} c_{i,r_2+1} = d_{2r_2+1}^2, \qquad c^2 + \sum_{j=1}^{r_2} c_{r_1+1,j} = d_{2r_1+1}^1,$$
$$c^1 + \sum_{j=1}^{r_2} c_{r_1+1,j} = -a_{2r_1-1}^1, \quad \text{and} \quad c^2 + \sum_{i=1}^{r_1} c_{i,r_2+1} = -a_{2r_2-1}^2.$$

Indeed, if we assume (without loss of generality) $a_{2r_1}^1 \leq a_{2r_2}^2$, we can take $c^1 = a_{2r_1}^1 - a_{2r_2}^2 - b$ and $c^2 = -b$ where $b = \min(a_{2r_1-1}^1, a_{2r_2-1}^2 - a_{2r_2}^2 + a_{2r_1}^1)$. For the $c_{i,j}$'s one can do as in the proof of the previous proposition.

We claim that γ is equal to

$$\left(\sum_{i=2}^{r_1} -d_{2i}^1 \gamma_i^1\right) - c^1 \gamma_{r_1+1}^1 + \left(\sum_{j=2}^{r_2} -d_{2j}^2 \gamma_j^2\right) - c^2 \gamma_{r_2+1}^2 + \sum_{\substack{1 \leqslant i \leqslant r_1+1\\1 \leqslant j \leqslant r_2+1\\i+j \leqslant r_1+r_2+2}} -c_{i,j} \gamma_{i,j}.$$

Indeed, the coefficient of σ^1_{2i-1} in the above expression is equal to

$$\left(\sum_{u=i+1}^{r_1} -d_{2u}^1(u-i)\right) - c^1(r_1+1-i) + \sum_{\substack{i+1 \le u \le r_1+1 \\ 1 \le v \le r_2+1 \\ u+v < r_1+r_2+2}} -c_{u,v}(u-i)$$
$$= \left(\sum_{u=i+1}^{r_1} -(d_{2u}^1 + d_{2u-1}^1)(u-i)\right) + a_{2r_1-1}^1(r_1+1-i)$$
$$= a_{2i-1}^1.$$

The coefficient of σ_{2i}^1 is equal to

$$\left(\sum_{u=i+1}^{r_1} -d_{2u}^1(u-i)\right) - c^1(r_1+1-i) + \left(\sum_{u=1}^{i}\sum_{v=1}^{r_2+1} -c_{u,v}\right) + \sum_{\substack{i+1 \le u \le r_1+1 \\ 1 \le v \le r_2+1 \\ u+v < r_1+r_2+2}} -c_{u,v}(u-i)$$

$$= \left(\sum_{u=1}^{i} -d_{2u-1}^1\right) + \left(\sum_{u=i+1}^{r_1} -(d_{2u}^1+d_{2u-1}^1)(u-i)\right) + a_{2r_1-1}^1(r_1+1-i)$$

$$= a_{2i}^1.$$

The same holds for σ_h^2 , for any h.

The case r = s = 0 corresponds to a flag variety. We are left with the case r > 0 and s = 0 (the other one, r = 0 and s > 0, is analogous). Keeping the same notation as above, there exists no D_1^2 and we have

$$\Delta = \{D_1^1, \dots, D_{2r+2}^1\} \cup \{D_2^2\}.$$

In this case, $D_{\mathfrak{p}_1} = D_2^1$ and $D_{\mathfrak{p}_2} = \tilde{D}_2^2 = D_2^2 + D_{2r+1}^1$. Both are minuscule, and \overline{Ke} is normal.

The description of the semigroup $\Gamma_{\Delta_{\mathfrak{p}}(e)}$ remains the same: denoting by d_h^k the coefficient of D_h^k in γ , for $h \leq 2r$, the semigroup

$$\left\{\gamma \in \mathbb{N}\Sigma \,:\, d_h^1 \leqslant 0 \,\,\forall \,\, h \neq 2\right\}$$

is generated by the elements of the form γ_i^1 , for $2 \leq i \leq r+1$, and $\gamma_{i,1}$, for $1 \leq i \leq r$.

Appendix A. List of spherical nilpotent K-orbits in \mathfrak{p} in the classical Hermitian cases

Here we report the list of the spherical nilpotent K-orbits in \mathfrak{p} for all symmetric pairs $(\mathfrak{g}, \mathfrak{k})$ of classical Hermitian type.

Every (complex) K-orbit in \mathfrak{p} is labelled with the signed partition of the corresponding real nilpotent orbit. In each case we provide a normal triple $\{h, e, f\}$, with e a representative of the orbit.

We denote by Q the parabolic subgroup of K whose Lie algebra is equal to

$$\operatorname{Lie} Q = \bigoplus_{i \geqslant 0} \mathfrak{k}(i),$$

where $\mathfrak{k}(i)$ is the adh-eigenspace in \mathfrak{k} of eigenvalue *i*.

We describe the centralizer of h, denoted by K_h or by L, which is a Levi subgroup of Q. Let Q^u be the unipotent radical of Q. Then we describe the centralizer of e, denoted by K_e . A Levi subgroup of K_e is always given by L_e , the centralizer of ein L. The unipotent radical of K_e is explicitly described as L_e -submodule of Q^u .

1.
$$SL(p+q)/S(GL(p) \times GL(q))$$
.

$$\begin{split} &K = \mathrm{S}(\mathrm{GL}(p) \times \mathrm{GL}(q)), \, p, q \geqslant 2, \, \mathfrak{p} = V(\omega_1 + \omega'_{q-1}) \oplus V(\omega_{p-1} + \omega'_1) \text{ as } K^{\mathrm{ss}}\text{-module.} \\ &\mathrm{If} \; p = 1 \; \mathrm{and} \; q \geqslant 2, \, \mathfrak{p} = V(\omega'_{q-1}) \oplus V(\omega'_1). \; \mathrm{If} \; p \geqslant 2 \; \mathrm{and} \; q = 1, \, \mathfrak{p} = V(\omega_1) \oplus V(\omega_{p-1}). \\ &\mathrm{If} \; p = q = 1, \, \mathfrak{p} = V(0) \oplus V(0). \end{split}$$

Let us fix a basis e_1, \ldots, e_p of \mathbb{C}^p and denote by $\varphi_1, \ldots, \varphi_p$ the dual basis of $(\mathbb{C}^p)^*$. Similarly, let us fix a basis e'_1, \ldots, e'_q of \mathbb{C}^q and denote by $\varphi'_1, \ldots, \varphi'_q$ the dual basis of $(\mathbb{C}^q)^*$. Then $K = \mathrm{S}(\mathrm{GL}(\mathbb{C}^p) \times \mathrm{GL}(\mathbb{C}^q))$ and

$$\mathfrak{p} = \left(\mathbb{C}^p \otimes (\mathbb{C}^q)^*\right) \oplus \left((\mathbb{C}^p)^* \otimes \mathbb{C}^q\right).$$

1.1. $(+2^{\mathbf{r}}, +1^{\mathbf{p-r}}, -1^{\mathbf{q-r}}), r \ge 1.$

$$e = \sum_{i=1}^{r} e_i \otimes \varphi'_{q-r+i}, \qquad f = \sum_{i=1}^{r} \varphi_i \otimes e'_{q-r+i},$$
$$h(e_i) = \begin{cases} e_i & \text{if } 1 \leqslant i \leqslant r \\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} -e'_i & \text{if } q-r+1 \leqslant i \leqslant q \\ 0 & \text{otherwise} \end{cases}$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ S(GL(r) × GL(p - r) × GL(q - r) × GL(r)).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong S(GL(r) \times GL(p-r) \times GL(q-r))$, the GL(r) factor of L_e is embedded diagonally, $A \mapsto (A, A)$, into the $GL(r) \times GL(r)$ factor of L. For r = p = q, the connected component of L_e is isomorphic to SL(r).

 $1.2. \ (-\mathbf{2^r},+\mathbf{1^{p-r}},-\mathbf{1^{q-r}}), \ r \geqslant 1.$

$$e = \sum_{i=1}^{r} \varphi_{p-r+i} \otimes e'_i, \qquad f = \sum_{i=1}^{r} e_{p-r+i} \otimes \varphi'_i,$$
$$h(e_i) = \begin{cases} -e_i & \text{if } p-r+1 \leqslant i \leqslant p \\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} e'_i & \text{if } 1 \leqslant i \leqslant r \\ 0 & \text{otherwise} \end{cases}$$

The centralizers of h and e are the same as in the previous case up to switching the two factors of K as well as the role of p and q.

1.3.
$$(+2^{\mathbf{r}}, -2^{\mathbf{s}}, +1^{\mathbf{p-r-s}}, -1^{\mathbf{q-r-s}}), r, s \ge 1$$

$$e = \sum_{i=1}^{r} e_i \otimes \varphi'_{q-r+i} + \sum_{i=1}^{s} \varphi_{p-s+i} \otimes e'_i,$$

$$h(e_i) = \begin{cases} e_i & \text{if } 1 \leqslant i \leqslant r \\ -e_i & \text{if } p-s+1 \leqslant i \leqslant p \\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} e'_i & \text{if } 1 \leqslant i \leqslant s \\ -e'_i & \text{if } q-r+1 \leqslant i \leqslant q \\ 0 & \text{otherwise} \end{cases},$$

$$f = \sum_{i=1}^{r} \varphi_i \otimes e'_{q-r+i} + \sum_{i=1}^{s} e_{p-s+i} \otimes \varphi'_i.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ S(GL(r) × GL(p - r - s) × GL(s) × GL(s) × GL(q - r - s) × GL(r)).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong S(GL(r) \times GL(p-r-s) \times GL(s) \times GL(q-r-s))$, the GL(r) and GL(s) factors of L_e are embedded diagonally, respectively, into the $GL(r) \times GL(r)$ and $GL(s) \times GL(s)$ factors of L.

1.4.
$$(+3^2, +1^{p-4}), q = 2$$
.

$$e = e_1 \otimes \varphi'_1 + e_2 \otimes \varphi'_2 + \varphi_{p-1} \otimes e'_1 + \varphi_p \otimes e'_2,$$

$$h(e_i) = \begin{cases} 2e_i & \text{if } 1 \leqslant i \leqslant 2\\ -2e_i & \text{if } p-1 \leqslant i \leqslant p \\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = 0 \ \forall \ i,$$

$$f = 2(\varphi_1 \otimes e'_1 + \varphi_2 \otimes e'_2 + e_{p-1} \otimes \varphi'_1 + e_p \otimes \varphi'_2).$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ S(GL(2) × GL(p - 4) × GL(2) × GL(2)).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong S(GL(2) \times GL(p-4))$, the GL(2) factor of L_e is embedded diagonally, $A \mapsto (A, A, A)$, into the GL(2) × GL(2) × GL(2) factor of L. For p = 4, the connected component of L_e is isomorphic to SL(2).

1.5.
$$(-3^2, -1^{q-4}), p = 2.$$

$$e = e_1 \otimes \varphi'_{q-1} + e_2 \otimes \varphi'_q + \varphi_1 \otimes e'_1 + \varphi_2 \otimes e'_2,$$

$$h(e_i) = 0 \ \forall \ i, \quad h(e'_i) = \begin{cases} 2e'_i & \text{if } 1 \leqslant i \leqslant 2\\ -2e'_i & \text{if } q-1 \leqslant i \leqslant q\\ 0 & \text{otherwise} \end{cases},$$

$$f = 2(\varphi_1 \otimes e'_{q-1} + \varphi_2 \otimes e'_q + e_1 \otimes \varphi'_1 + e_2 \otimes \varphi'_2).$$

The centralizers of h and e are the same as in the previous case up to switching the two factors of K as well as the role of p and q.

$$1.6. \ (+3, +2^{\mathbf{r}}, -2^{\mathbf{s}}, +1^{\mathbf{p-r-s-2}}, -1^{\mathbf{q-r-s-1}}).$$

$$e = e_{1} \otimes \varphi'_{q-r} + \sum_{i=1}^{r} e_{i+1} \otimes \varphi'_{q-r+i} + \sum_{i=1}^{s} \varphi_{p-s+i-1} \otimes e'_{i} + \varphi_{p} \otimes e'_{q-r},$$

$$h(e_{i}) = \begin{cases} 2e_{i} & \text{if } i = 1 \\ e_{i} & \text{if } 2 \leq i \leq r+1 \\ -e_{i} & \text{if } p-s \leq i \leq p-1 \\ -2e_{i} & \text{if } i = p \\ 0 & \text{otherwise} \end{cases},$$

$$f = 2\varphi_{1} \otimes e'_{q-r} + \sum_{i=1}^{r} \varphi_{i+1} \otimes e'_{q-r+i} + \sum_{i=1}^{s} e_{p-s+i-1} \otimes \varphi'_{i} + 2e_{p} \otimes \varphi'_{q-r}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ S(GL(1)×GL(r)×GL(p-r-s-2)×GL(s)×GL(1)×GL(s)×GL(q-r-s)×GL(r)). The centralizer of e is $K_e = L_e K_e^u$ where $L_e \cong S(GL(1) \times GL(r) \times GL(p - r - s - 2) \times GL(s) \times GL(q - r - s - 1))$, the $GL(1) \times GL(q - r - s - 1)$ factor of L_e is embedded as

$$(z, A) \mapsto (z, z, (A, z))$$

into $\operatorname{GL}(1) \times \operatorname{GL}(1) \times (\operatorname{GL}(q-r-s-1) \times \operatorname{GL}(1))$ and $\operatorname{GL}(q-r-s-1) \times \operatorname{GL}(1)$ is included in the $\operatorname{GL}(q-r-s)$ factor of L, the $\operatorname{GL}(r)$ and $\operatorname{GL}(s)$ factors of L_e are embedded diagonally, respectively, into the $\operatorname{GL}(r) \times \operatorname{GL}(r)$ and $\operatorname{GL}(s) \times \operatorname{GL}(s)$ factors of L. The quotient $\operatorname{Lie} Q^{\mathrm{u}}/\operatorname{Lie} K_e^{\mathrm{u}}$ is the sum of two simple L_e -modules of dimension r and s, respectively, as follows. In $\mathfrak{k}(1)$ there are exactly two simple L_e -submodules, $W_{0,1}, W_{1,1}$, of highest weight ω_{r-1} w.r.t. the semisimple part of the $\operatorname{GL}(r)$ factor, isomorphic as L_e -modules but lying in two distinct isotypical L-components. Similarly, in $\mathfrak{k}(1)$ there are exactly two simple L_e -submodules, $W_{0,2}, W_{1,2}$, of highest weight ω_1 w.r.t. the semisimple part of the $\operatorname{GL}(s)$ factor, isomorphic as L_e -modules but lying in two distinct isotypical L-components. Let V be the L_e -complement of $W_{0,1} \oplus W_{1,1} \oplus W_{0,2} \oplus W_{1,2}$ in $\operatorname{Lie} Q^{\mathrm{u}}$. As L_e -module, $\operatorname{Lie} K_e^{\mathrm{u}}$ is the direct sum of V, of a simple L_e -submodule of $W_{0,1} \oplus W_{1,1}$ which projects non-trivially on both summands $W_{0,1}$ and $W_{1,1}$, and of a simple L_e -submodule of $W_{0,2} \oplus W_{1,2}$ which projects non-trivially on both summands $W_{0,2}$ and $W_{1,2}$.

$$1.7. \ (-3, +2^{\mathbf{r}}, -2^{\mathbf{s}}, +1^{\mathbf{p-r-s-1}}, -1^{\mathbf{q-r-s-2}}).$$

$$e = \sum_{i=1}^{r} e_i \otimes \varphi'_{q-r+i-1} + e_{p-s} \otimes \varphi'_q + \varphi_{p-s} \otimes e'_1 + \sum_{i=1}^{s} \varphi_{p-s+i} \otimes e'_{i+1},$$

$$h(e_i) = \begin{cases} e_i & \text{if } 1 \leqslant i \leqslant r \\ -e_i & \text{if } p-s+1 \leqslant i \leqslant p \\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} 2e'_i & \text{if } i = 1 \\ e'_i & \text{if } 2 \leqslant i \leqslant s+1 \\ -e'_i & \text{if } q-r \leqslant i \leqslant q-1 \\ -2e'_i & \text{if } i = q \\ 0 & \text{otherwise} \end{cases}$$

$$f = \sum_{i=1}^{r} \varphi_i \otimes e'_{q-r+i-1} + 2\varphi_{p-s} \otimes e'_q + 2e_{p-s} \otimes \varphi'_1 + \sum_{i=1}^{s} e_{p-s+i} \otimes \varphi'_{i+1}.$$

The centralizers of h and e are the same as in the previous case up to switching the two factors of K as well as the role of p and q, and the role of r and s, respectively.

2. $\operatorname{SO}(2n+1)/\operatorname{SO}(2n-1) \times \operatorname{SO}(2)$. $K = \operatorname{SO}(2n-1) \times \operatorname{SO}(2), n > 2, \mathfrak{p} = V(\omega_1) \oplus V(\omega_1)$ as K^{ss} -module.

Let us fix a basis $e_1, \ldots, e_{n-1}, e_0, e_{-n+1}, \ldots, e_{-1}$ of \mathbb{C}^{2n-1} , a symmetric bilinear form β such that $\beta(e_i, e_j) = \delta_{i,-j}$ for all i, j. Similarly, let us fix a basis e'_1, e'_{-1} of \mathbb{C}^2 and a symmetric bilinear form β' such that $\beta'(e'_i, e'_j) = \delta_{i,-j}$ for all i, j. For convenience, let us denote by $\varphi'_1, \varphi'_{-1}$ the dual basis of $(\mathbb{C}^2)^*$. Then K = $\mathrm{SO}(\mathbb{C}^{2n-1}, \beta) \times \mathrm{SO}(\mathbb{C}^2, \beta')$ and

$$\mathfrak{p} = \mathbb{C}^{2n-1} \otimes (\mathbb{C}^2)^*.$$

2.1. $(+2^2, +1^{2n-3})$, I and II.

Case (I)

$$e = e_1 \otimes \varphi'_{-1}, \qquad f = -e_{-1} \otimes \varphi'_1,$$

$$h(e_i) = \begin{cases} e_i & \text{if } i = 1\\ -e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} e'_i & \text{if } i = 1\\ -e'_i & \text{if } i = -1 \end{cases}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(1) × SO(2n - 3) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong \operatorname{GL}(1) \times \operatorname{SO}(2n-3)$, the $\operatorname{GL}(1)$ factor of L_e is embedded skew-diagonally, $z \mapsto (z, z^{-1})$, into the $\operatorname{GL}(1) \times \operatorname{GL}(1)$ factor of L.

Case (II)

$$e = e_1 \otimes \varphi'_1, \qquad f = -e_{-1} \otimes \varphi'_{-1},$$

$$h(e_i) = \begin{cases} e_i & \text{if } i = 1\\ -e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} -e'_i & \text{if } i = 1\\ e'_i & \text{if } i = -1 \end{cases}$$

The centralizer of h is the same as in case (I).

The centralizer of e is also the same, except that the GL(1) factor of L_e is embedded diagonally, $z \mapsto (z, z)$, into the $GL(1) \times GL(1)$ factor of L.

2.2. $(+3, +1^{2n-3}, -1)$.

$$e = e_1 \otimes (\varphi'_1 - \varphi'_{-1}), \qquad f = e_{-1} \otimes (\varphi'_1 - \varphi'_{-1}),$$
$$h(e_i) = \begin{cases} 2e_i & \text{if } i = 1\\ -2e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = 0 \ \forall \ i.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(1) × SO(2n - 3) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong O(1) \times SO(2n-3)$, the O(1) factor of L_e is embedded diagonally into the $GL(1) \times GL(1)$ factor of L.

2.3. $(-3, +1^{2n-2})$, I and II.

Case (I)

$$e = e_0 \otimes \varphi'_{-1}, \qquad f = -2e_0 \otimes \varphi'_1,$$

$$h(e_i) = 0 \ \forall \ i, \quad h(e'_i) = \begin{cases} 2e'_i & \text{if } i = 1\\ -2e'_i & \text{if } i = -1 \end{cases}.$$

Here the centralizer of h is $K_h = K \cong SO(2n-1) \times GL(1)$. The centralizer of e is $K_e \cong S(O(2n-2) \times O(1))$ embedded as

$$(A,z) \mapsto ((A,z), z^{-1})$$

into $S(O(2n-2) \times O(1)) \times GL(1)$, where $S(O(2n-2) \times O(1))$ is included in the SO(2n-1) factor of K.

Case (II)

$$e = e_0 \otimes \varphi'_1, \qquad f = -2e_0 \otimes \varphi'_{-1},$$

28

$$h(e_i) = 0 \ \forall \ i, \quad h(e'_i) = \begin{cases} -2e'_i & \text{if } i = 1\\ 2e'_i & \text{if } i = -1 \end{cases}$$

The centralizers of h and e are the same as in case (I).

2.4. $(+3^2, +1^{2n-5})$.

$$e = e_1 \otimes \varphi'_{-1} - e_2 \otimes \varphi'_1, \qquad f = 2(e_{-2} \otimes \varphi'_{-1} - e_{-1} \otimes \varphi'_1)$$
$$h(e_i) = \begin{cases} 2e_i & \text{if } 1 \leqslant i \leqslant 2\\ -2e_i & \text{if } -2 \leqslant i \leqslant -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = 0 \ \forall \ i.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(2) × SO(2n - 5) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong SO(2n - 5) \times GL(1)$, the GL(1) factor of L_e is embedded as

$$z \mapsto ((z, z^{-1}), z^{-1})$$

into $(GL(1) \times GL(1)) \times GL(1)$ included into the $GL(2) \times GL(1)$ factor of L.

3. Sp(2n)/GL(n).

 $K = \operatorname{GL}(n), n \ge 2, \mathfrak{p} = V(2\omega_1) \oplus V(2\omega_{n-1})$ as K^{ss} -module.

Let us fix a basis e_1, \ldots, e_n of \mathbb{C}^n and denote by $\varphi_1, \ldots, \varphi_n$ the dual basis of $(\mathbb{C}^n)^*$. Then $K = \operatorname{GL}(\mathbb{C}^n)$ and

$$\mathfrak{p} = \mathsf{S}^2(\mathbb{C}^n) \oplus \mathsf{S}^2(\mathbb{C}^n)^*.$$

3.1. $(+2^{r}, +1^{2n-2r}).$

$$e = \sum_{i=1}^{r} e_i e_{r-i+1}, \qquad h(e_i) = \begin{cases} e_i & \text{if } 1 \leq i \leq r \\ 0 & \text{otherwise} \end{cases}, \qquad f = \sum_{i=1}^{r} \varphi_i \varphi_{r-i+1}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL $(r) \times$ GL(n - r).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong O(r) \times GL(n-r)$.

3.2. $(-2^{r}, +1^{2n-2r}).$

$$e = \sum_{i=1}^{r} \varphi_{n-r+i} \varphi_{n-i+1}, \qquad f = \sum_{i=1}^{r} e_{n-r+i} e_{n-i+1},$$
$$h(e_i) = \begin{cases} -e_i & \text{if } n-r+1 \leqslant i \leqslant n \\ 0 & \text{otherwise} \end{cases}.$$

The centralizers of h and e are the same as in the previous case up to an external automorphism of K.

29

3.3.
$$(+2^{r}, -2^{s}, +1^{2n-2r-2s}).$$

$$e = \sum_{i=1}^{r} e_i e_{r-i+1} + \sum_{i=1}^{s} \varphi_{n-s+i} \varphi_{n-i+1}, \qquad f = \sum_{i=1}^{r} \varphi_i \varphi_{r-i+1} + \sum_{i=1}^{s} e_{n-s+i} e_{n-i+1},$$
$$h(e_i) = \begin{cases} e_i & \text{if } 1 \leqslant i \leqslant r \\ -e_i & \text{if } n-s+1 \leqslant i \leqslant n \\ 0 & \text{otherwise} \end{cases}$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ $\operatorname{GL}(r) \times \operatorname{GL}(n - r - s) \times \operatorname{GL}(s)$.

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong O(r) \times GL(n-r-s) \times O(s)$.

4. $\operatorname{SO}(2n)/\operatorname{SO}(2n-2) \times \operatorname{SO}(2)$. $K = \operatorname{SO}(2n-2) \times \operatorname{SO}(2), n > 4, \mathfrak{p} = V(\omega_1) \oplus V(\omega_1)$ as K^{ss} -module.

Let us fix a basis $e_1, \ldots, e_{n-1}, e_{-n+1}, \ldots, e_{-1}$ of \mathbb{C}^{2n-2} , a symmetric bilinear form β such that $\beta(e_i, e_j) = \delta_{i,-j}$ for all i, j. Similarly, let us fix a basis e'_1, e'_{-1} of \mathbb{C}^2 and a symmetric bilinear form β' such that $\beta'(e'_i, e'_j) = \delta_{i,-j}$ for all i, j. For convenience, let us denote by $\varphi'_1, \varphi'_{-1}$ the dual basis of $(\mathbb{C}^2)^*$. Then $K = \mathrm{SO}(\mathbb{C}^{2n-2}, \beta) \times \mathrm{SO}(\mathbb{C}^2, \beta')$ and

$$\mathfrak{p} = \mathbb{C}^{2n-2} \otimes (\mathbb{C}^2)^*.$$

4.1. $(+2^2, +1^{2n-4})$, I and II. Case (I)

$$e = e_1 \otimes \varphi'_{-1}, \qquad f = -e_{-1} \otimes \varphi'_1,$$

$$h(e_i) = \begin{cases} e_i & \text{if } i = 1\\ -e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} e'_i & \text{if } i = 1\\ -e'_i & \text{if } i = -1 \end{cases}$$

.

•

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(1) × SO(2n - 4) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong \operatorname{GL}(1) \times \operatorname{SO}(2n-4)$, the $\operatorname{GL}(1)$ factor of L_e is embedded skew-diagonally, $z \mapsto (z, z^{-1})$, into the $\operatorname{GL}(1) \times \operatorname{GL}(1)$ factor of L.

Case (II)

$$e = e_1 \otimes \varphi'_1, \qquad f = -e_{-1} \otimes \varphi'_{-1},$$
$$h(e_i) = \begin{cases} e_i & \text{if } i = 1\\ -e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = \begin{cases} -e'_i & \text{if } i = 1\\ e'_i & \text{if } i = -1 \end{cases}$$

The centralizer of h is the same as in case (I).

The centralizer of e is also the same, except that the GL(1) factor of L_e is embedded diagonally, $z \mapsto (z, z)$, into the $GL(1) \times GL(1)$ factor of L.

 $4.2. (+3, +1^{2n-4}, -1).$

$$e = e_1 \otimes (\varphi'_1 - \varphi'_{-1}), \qquad f = e_{-1} \otimes (\varphi'_1 - \varphi'_{-1}),$$
$$h(e_i) = \begin{cases} 2e_i & \text{if } i = 1\\ -2e_i & \text{if } i = -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = 0 \ \forall \ i.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(1) × SO(2n - 4) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong O(1) \times SO(2n-4)$, the O(1) factor of L_e is embedded diagonally into the $GL(1) \times GL(1)$ factor of L.

4.3. $(-3, +1^{2n-3})$, I and II.

Case (I)

$$e = (e_{n-1} - e_{-n+1}) \otimes \varphi'_{-1}, \qquad f = (e_{n-1} - e_{-n+1}) \otimes \varphi'_{1},$$
$$h(e_{i}) = 0 \ \forall \ i, \quad h(e'_{i}) = \begin{cases} 2e'_{i} & \text{if } i = 1\\ -2e'_{i} & \text{if } i = -1 \end{cases}.$$

Here the centralizer of h is $K_h = K \cong SO(2n-2) \times GL(1)$. The centralizer of e is $K_e \cong SO(2n-3) \times O(1)$, embedded as

Tanzer of e is $\Lambda_e = 50(2\pi - 3) \times O(1)$, embedded

$$(A,z)\mapsto ((A,z),z^{-1})$$

into $(SO(2n - 3) \times O(1)) \times GL(1)$, where $SO(2n - 3) \times O(1)$ is included in the SO(2n - 2) factor of K.

Case (II)

$$e = (e_{n-1} - e_{-n+1}) \otimes \varphi'_1, \qquad f = (e_{n-1} - e_{-n+1}) \otimes \varphi'_{-1},$$
$$h(e_i) = 0 \ \forall \ i, \quad h(e'_i) = \begin{cases} -2e'_i & \text{if } i = 1\\ 2e'_i & \text{if } i = -1 \end{cases}.$$

The centralizers of h and e are the same as in case (I).

4.4. $(+3^2, +1^{2n-6})$.

$$e = e_1 \otimes \varphi'_{-1} - e_2 \otimes \varphi'_1, \qquad f = 2(e_{-2} \otimes \varphi'_{-1} - e_{-1} \otimes \varphi'_1),$$
$$h(e_i) = \begin{cases} 2e_i & \text{if } 1 \leqslant i \leqslant 2\\ -2e_i & \text{if } -2 \leqslant i \leqslant -1\\ 0 & \text{otherwise} \end{cases}, \quad h(e'_i) = 0 \ \forall \ i.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(2) × SO(2n - 6) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong SO(2n-6) \times GL(1)$, the GL(1) factor of L_e is embedded as

$$z \mapsto ((z, z^{-1}), z^{-1})$$

into $(GL(1) \times GL(1)) \times GL(1)$ included into the $GL(2) \times GL(1)$ factor of L.

5. SO(2n)/GL(n).

 $K = \operatorname{GL}(n), n \ge 4, \mathfrak{p} = V(\omega_2) \oplus V(\omega_{n-2})$ as K^{ss} -module.

Let us fix a basis e_1, \ldots, e_n of \mathbb{C}^n and denote by $\varphi_1, \ldots, \varphi_n$ the dual basis of $(\mathbb{C}^n)^*$. Then $K = \operatorname{GL}(\mathbb{C}^n)$ and

$$\mathfrak{p} = \Lambda^2(\mathbb{C}^n) \oplus \Lambda^2(\mathbb{C}^n)^*.$$

5.1. $(+2^{r}, +1^{n-2r})$.

$$e = \sum_{i=1}^{r} e_i \wedge e_{2r-i+1}, \qquad h(e_i) = \begin{cases} e_i & \text{if } 1 \leq i \leq 2r \\ 0 & \text{otherwise} \end{cases}, \qquad f = \sum_{i=1}^{r} \varphi_i \wedge \varphi_{2r-i+1}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL $(2r) \times$ GL(n - 2r).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong \operatorname{Sp}(2r) \times \operatorname{GL}(n-2r)$.

5.2.
$$(-2^{\mathbf{r}}, +1^{\mathbf{n-2r}}).$$

$$e = \sum_{i=1}^{r} \varphi_{n-2r+i-1} \wedge \varphi_{n-i+1}, \qquad f = \sum_{i=1}^{r} e_{n-2r+i-1} \wedge e_{n-i+1},$$
$$h(e_i) = \begin{cases} -e_i & \text{if } n-2r+1 \leqslant i \leqslant n \\ 0 & \text{otherwise} \end{cases}.$$

The centralizers of h and e are the same as in the previous case up to an external automorphism of K.

5.3.
$$(+2^{r}, -2^{s}, +1^{n-2r-2s}).$$

$$e = \sum_{i=1}^{r} e_i \wedge e_{2r-i+1} + \sum_{i=1}^{s} \varphi_{n-2s+i-1} \wedge \varphi_{n-i+1},$$
$$h(e_i) = \begin{cases} e_i & \text{if } 1 \leqslant i \leqslant 2r \\ -e_i & \text{if } n-2s+1 \leqslant i \leqslant n \\ 0 & \text{otherwise} \end{cases},$$

$$f = \sum_{i=1}^{r} \varphi_i \wedge \varphi_{2r-i+1} + \sum_{i=1}^{s} e_{n-2s+i-1} \wedge e_{n-i+1}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL $(2r) \times$ GL $(n - 2r - 2s) \times$ GL(2s).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong \operatorname{Sp}(2r) \times \operatorname{GL}(n - 2r - 2s) \times \operatorname{Sp}(2s)$.

32

5.4. (+3, +1ⁿ⁻³).

$$e = e_1 \wedge e_2 + \varphi_2 \wedge \varphi_n, \qquad f = 2(\varphi_1 \wedge \varphi_2 + e_2 \otimes e_n)$$
$$h(e_i) = \begin{cases} 2e_i & \text{if } i = 1\\ -2e_i & \text{if } i = n\\ 0 & \text{otherwise} \end{cases}.$$

Let $Q = L Q^{u}$ be the corresponding parabolic subgroup of K, so that $L = K_{h} \cong$ GL(1) × GL(n - 2) × GL(1).

The centralizer of e is $K_e = L_e Q^u$ where $L_e \cong \operatorname{GL}(1) \times \operatorname{GL}(n-3)$, L_e is embedded as

$$(z,A) \mapsto (z,(z^{-1},A),z)$$

into $\operatorname{GL}(1) \times (\operatorname{GL}(1) \times \operatorname{GL}(n-3)) \times \operatorname{GL}(1)$, and $(\operatorname{GL}(1) \times \operatorname{GL}(n-3))$ is included into the $\operatorname{GL}(n-2)$ factor of L.

Appendix B. Tables of spherical nilpotent K-orbits in \mathfrak{p} in the classical Hermitian cases

In Tables 1–6, for every spherical nilpotent orbit $Ke \subset \mathfrak{p}$, we report its signed partition (column 2), the Kostant-Dynkin diagram and the height of Ge (columns 3 and 4), the Kostant-Dynkin diagram and the \mathfrak{p} -height of Ke (columns 5 and 6), the codimension of $\overline{Ke} \setminus Ke$ in \overline{Ke} (column 7) and the weight semigroup of \overline{Ke} (column 8).

The generators of the weight semigroups given in the tables are expressed in terms of the fundamental weights of K^{ss} , the semisimple part of K, plus a multiple of χ , where χ denotes the character of the 1-dimensional center of K on \mathfrak{p}_1 , as defined in Section 4.

Recall that the fundamental weights of an irreducible root system of rank n are denoted by $\omega_1, \ldots, \omega_n$ (and enumerated as in Bourbaki). For notational convenience, we set $\omega_i = 0$ if $i \leq 0$ or i > n.

In Tables 7–11, for every spherical nilpotent orbit Ke in \mathfrak{p} , we report the Luna diagram and the set of spherical roots of the spherical system of $K\pi(e)$, where the definition of $\pi(e)$ is as follows. Recall that in the Hermitian case $\mathfrak{p} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$, so that $e = e_1 + e_2$ with $e_1 \in \mathfrak{p}_1$ and $e_2 \in \mathfrak{p}_2$. Therefore, we set $\pi(e) = [e_1] \in \mathbb{P}(\mathfrak{p}_1)$ if $e_2 = 0$, $\pi(e) = [e_2] \in \mathbb{P}(\mathfrak{p}_2)$ if $e_1 = 0$, and $\pi(e) = ([e_1], [e_2]) \in \mathbb{P}(\mathfrak{p}_1) \times \mathbb{P}(\mathfrak{p}_2)$ otherwise.

	Signed partition	Diagram of Ge	ht(e)	Diagram of Ke	$ht_{\mathfrak{p}}(e)$	$\operatorname{codim}(\overline{Ke}\smallsetminus Ke)$	Generators of $\Gamma(\overline{Ke})$
1.1	$(+2^r, +1^{p-r}, -1^{q-r})$ $r \ge 1$	$\underbrace{(\underbrace{0\dots0}_{r-1}10\dots01\underbrace{0\dots0}_{r-1})}_{p-1} \text{if } 2r < p+q$ $\underbrace{(\underbrace{0\dots0}_{p-1}20\dots0)}_{p-1} \text{if } r=p=q$	2	$\underbrace{(\underbrace{0\dots0}_{r-1}10\dots0,\ 0\dots01\underbrace{0\dots0}_{r-1};\ 0)}_{q-1} \text{if } r < p,q$ $\underbrace{(\underbrace{0\dots0}_{r-1}10\dots0,\ 0\dots01\underbrace{0\dots0}_{r-1};\ 1)}_{q-1} \text{if } r = p < q$ $\underbrace{(\underbrace{0\dots0}_{q-1}10\dots0,\ 0\dots0;\ 1)}_{q-1} \text{if } r = q < p$ $\underbrace{(\underbrace{0\dots0}_{q-1}0\dots0,\ 0\dots0;\ 2)}_{r-1} \text{if } r = p = q$	2	p+q-2r+1	$\omega_{p-i} + \omega'_i - i\chi \ (i = 1, \dots, r)$
1.2	$(-2^r, +1^{p-r}, -1^{q-r})$ $r \ge 1$	$\underbrace{(\underbrace{0\dots0}_{r-1} 10\dots01\underbrace{0\dots0}_{r-1})}_{p-1} \text{if } 2r < p+q$ $\underbrace{(\underbrace{0\dots0}_{p-1} 20\dots0)}_{p-1} \text{if } r=p=q$	2	$(0 \dots 01 \underbrace{0 \dots 0}_{r-1}, \underbrace{0 \dots 0}_{r-1} 10 \dots 0; -2) \text{if } r < p, q$ $(0 \dots 0, \underbrace{0 \dots 0}_{p-1} 10 \dots 0; -2) \text{if } r = p < q$ $(0 \dots 01 \underbrace{0 \dots 0}_{q-1}, 0 \dots 0; -2) \text{if } r = q < p$ $(0 \dots 0, 0 \dots 0; -2) \text{if } r = p = q$	2	p + q - 2r + 1	$\omega_i + \omega'_{q-i} + i\chi \ (i=1,\ldots,r)$
1.3	$(+2^r, -2^s, +1^{p-r-s}, -1^{q-r-s})$ $r, s \ge 1$	$\underbrace{(\underbrace{0\dots0}_{r+s-1}10\dots01\underbrace{0\dots0}_{r+s-1})}_{p-1} \text{if } 2(r+s) < p+q$ $\underbrace{(\underbrace{0\dots0}_{p-1}20\dots0)}_{p-1} \text{if } r+s=p=q$	2	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2	p+q-2r-2s+1	$\omega_j + \omega'_{q-j} + j\chi \ (j = 1, \dots, s)$ $\omega_{p-i} + \omega'_i - i\chi \ (i = 1, \dots, r)$
1.4	$(+3^2, +1^{p-4}), q=2$	(020020)	4	$\begin{array}{ll} (020\dots 020,\ 0;\ -2) & \text{if } p>4\\ (040,\ 0;\ -2) & \text{if } p=4 \end{array}$	2	p-3	
1.5	$(-3^2, -1^{q-4}), p=2$	(020020)	4	$\begin{array}{ll} (0, \ 020 \dots 020; \ -2) & \text{if } q > 4 \\ (0, \ 040; \ -2) & \text{if } q = 4 \end{array}$	2	q-3	$ \begin{array}{c} \omega + \omega_1' - \chi, \ \omega + \omega_{q-1}' + \chi, \\ \omega_1' + \omega_{q-1}', \ \omega_2' - 2\chi, \ \omega_{q-2}' + 2\chi \end{array} $

Table 1: $G = \mathsf{A}_{p+q-1}, K^{ss} = \mathsf{A}_{p-1} \times \mathsf{A}_{q-1} \ (p,q \ge 1)$

	Signed partition	Diagram of Ge	ht(e)	Diagram of Ke	$\operatorname{ht}_{\mathfrak{p}}(e)$	$\boxed{\operatorname{codim}(\overline{Ke}\smallsetminus Ke)}$	Generators of $\Gamma(\overline{Ke})$
$ \begin{array}{c} 1.6 \\ (r = s = 0, \ q = 1) \end{array} $	$(+3,+1^{p-2})$	(2002)	4	$\begin{array}{ll} (4; \ -2) & \text{if } p = 2\\ (20 \dots 02; \ -2) & \text{if } p > 2 \end{array}$	2	p - 1	$\omega_1 + \chi, \ \omega_{p-1} - \chi$
$ \begin{array}{c} 1.6 \\ (r = s = 0, \ q > 1) \end{array} $	$(+3,+1^{p-2},-1^{q-1})$	(2002)	4	$\begin{array}{ll} (4, \ 0 \dots 0; \ -2) & \text{if } p = 2 \\ (20 \dots 02, \ 0 \dots 0; \ -2) & \text{if } p > 2 \end{array}$	2	1	$\omega_1 + \omega'_{q-1} + \chi, \ \omega_{p-1} + \omega'_1 - \chi, \ \omega_1 + \omega_{p-1}$
$\begin{array}{c} 1.6\\ (0 < r + s \leqslant q - 1) \end{array}$	$(+3, +2^r, -2^s, +1^{p-r-s-2}, -1^{q-r-s-1})$	$(1\underbrace{00}_{r+s-1}1001\underbrace{00}_{r+s-1}1)$	4	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3	p+q-2(r+s+1)	$ \begin{split} & \omega_{j} + \omega_{q-j}' + j\chi \ (j = 1, \dots, s+1) \\ & \omega_{p-i} + \omega_{i}' - i\chi \ (i = 1, \dots, r+1) \\ & \omega_{j} + \omega_{p-i} + \omega_{i-1}' + \\ & \omega_{q-j+1}' + (j-i)\chi \end{split} \begin{pmatrix} 1 \leq i \leq r+1 \\ 1 \leq j \leq s+1 \\ i+j < q+1 \end{pmatrix} $
$ \begin{array}{c} 1.7 \\ (r = s = 0, \ p = 1) \end{array} $	$(-3, -1^{q-2})$	(2002)	4	(2002; -2) if $q > 2$	2	q-1	$\omega_1'-\chi,\ \omega_{q-1}'+\chi$
$ \begin{array}{c} 1.7 \\ (r = s = 0, \ p > 1) \end{array} $	$(-3,+1^{p-1},-1^{q-2})$	(2002)	4	$\begin{array}{ll} (0\dots 0,\ 4;\ -2) & \text{if } q=2\\ (0\dots 0,\ 20\dots 02;\ -2) & \text{if } q>2 \end{array}$	2	1	$\omega_1 + \omega'_{q-1} + \chi, \ \omega_{p-1} + \omega'_1 - \chi, \ \omega'_1 + \omega'_{q-1}$
$ \begin{array}{c} 1.7\\ (0 < r + s \leq p - 1) \end{array} $	$(-3, +2^r, -2^s, +1^{p-r-s-1}, -1^{q-r-s-2})$	$(1 \underbrace{0 \dots 0}_{r+s-1} 10 \dots 01 \underbrace{0 \dots 0}_{r+s-1} 1)$	4	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	3	p+q-2(r+s+1)	$ \begin{split} \omega_{j} + \omega_{q-j}' + j\chi & (j = 1, \dots, s+1) \\ \omega_{p-i} + \omega_{i}' - i\chi & (i = 1, \dots, r+1) \\ \omega_{j-1} + \omega_{p-i+1} + \\ \omega_{i}' + \omega_{q-j}' + (j-i)\chi & \begin{pmatrix} 1 \leq i \leq r+1 \\ 1 \leq j \leq s+1 \\ i+j < p+1 \end{pmatrix} \end{split} $

Table 2: $G = \mathsf{A}_{p+q-1}, K^{ss} = \mathsf{A}_{p-1} \times \mathsf{A}_{q-1} \ (p,q \ge 1) \ (\text{continued})$

	Signed partition	Diagram of Ge	$\operatorname{ht}(e)$	Diagram of Ke	$\operatorname{ht}_{\mathfrak{p}}(e)$	$\operatorname{codim}(\overline{Ke}\smallsetminus Ke)$	Generators of $\Gamma(\overline{Ke})$
2.1	(+2 ² , +1 ²ⁿ⁻³) (I) or (II)	(0100)	2	$\begin{array}{ccc} (I) & (10\dots 0; \ 0) \\ (II) & (10\dots 0; \ -2) \end{array}$	2	2(n-1)	$ \begin{array}{cc} (\mathrm{I}) & \omega_1 - \chi \\ (\mathrm{II}) & \omega_1 + \chi \end{array} $
2.2	$(+3,+1^{2n-3},-1),$	(200)	2	(200; -2)	2	1	$\omega_1 + \chi, \ \omega_1 - \chi$
2.3	$(-3, +1^{2n-2})$ (I) or (II)	(200)	2	$\begin{array}{ccc} (I) & (0 \dots 0; \ 2) \\ (II) & (0 \dots 0; \ -2) \end{array}$	2	1	(I) $\omega_1 - \chi, -2\chi$ (II) $\omega_1 + \chi, 2\chi$
2.4	$(+3^2,+1^{2n-5})$	(0200)	4	(0200; -2)	2	2(n-2)	$\omega_1 + \chi, \ \omega_1 - \chi, \ \omega_2$

Table 3: $G = B_n, K^{ss} = B_{n-1} (n > 2)$

	Signed partition	Diagram of Ge	ht(e)	Diagram of Ke	$ht_{\mathfrak{p}}(e)$	$codim(\overline{Ke} \smallsetminus Ke)$	Ge
3.1	$(+2^r,+1^{2n-2r}), r \ge 1$	$(\underbrace{00}_{r-1} 100) \text{if } r < n$ $(002) \text{if } r = n$	2	$\underbrace{(\underbrace{00}_{r-1} 100; 0) \text{if } r < n}_{(00; 2) \text{if } r = n}$	2	n-r+1	$2\omega_{n-1}$
3.2	$(-2^r, +1^{2n-2r}), r \ge 1$	$\underbrace{(\underbrace{00}_{r-1} 100)}_{(002)} \text{if } r < n$	2	$(001\underbrace{00}_{r-1}; -2) \text{if } r < n$ $(00; -2) \text{if } r = n$	2	n-r+1	$2\omega_i$
3.3	$(+2^r, -2^s, +1^{2n-2r-2s}), r, s \ge 1$	$(\underbrace{0\dots 0}_{r+s-1}10\dots 0) \text{if } r+s < n$ $(0\dots 02) \text{if } r+s = n$	2	$\underbrace{(\underbrace{0\dots0}_{r-1}10\dots01\underbrace{0\dots0}_{s-1}; -2)}_{r-1} \text{if } r+s < n$	2	n - r - s + 1	$\frac{2\omega_j}{2\omega_{n-1}}$

Table 4: $G = \mathsf{C}_n, K^{ss} = \mathsf{A}_{n-1} \ (n \ge 2)$

	Signed partition	Diagram of Ge	ht(e)	Diagram of Ke	$ht_{\mathfrak{p}}(e)$	$\operatorname{codim}(\overline{Ke}\smallsetminus Ke)$	Generators of $\Gamma(\overline{Ke})$
4.1	(+2 ² , +1 ²ⁿ⁻⁴) (I) or (II)	(0100)	2	$\begin{array}{ccc} (I) & (10\dots 0; \ 0) \\ (II) & (10\dots 0; \ -2) \end{array}$	2	2n - 3	$ \begin{array}{cc} (\mathrm{I}) & \omega_1 - \chi \\ (\mathrm{II}) & \omega_1 + \chi \end{array} $
4.2	$(+3,+1^{2n-4},-1),$	(200)	2	(200; -2)	2	1	$\omega_1 + \chi, \ \omega_1 - \chi$
4.3	$(-3, +1^{2n-3})$ (I) or (II)	$(20\ldots 0)$	2	$\begin{array}{ccc} (I) & (0 \dots 0; \ 2) \\ (II) & (0 \dots 0; \ -2) \end{array}$	2	1	(I) $\omega_1 - \chi, -2\chi$ (II) $\omega_1 + \chi, 2\chi$
4.4	$(+3^2,+1^{2n-6})$	(0200)	4	(0200; -2)	2	2n - 5	$\omega_1 + \chi, \ \omega_1 - \chi, \ \omega_2$

Table 5: $G = \mathsf{D}_n, K^{ss} = \mathsf{D}_{n-1} (n > 4)$

	Signed partition	Diagram of Ge	ht(e)	Diagram of Ke	$ht_{\mathfrak{p}}(e)$	$\operatorname{codim}(\overline{Ke}\smallsetminus Ke)$	Generators of $\Gamma(\overline{Ke})$
5.1	$(+2^r,+1^{n-2r}), r \ge 1$	$\begin{array}{ccc} \underbrace{(0\ldots 0}_{2r-1} 10\ldots 0) & \text{if } 2r < n-1 \\ (0\ldots 011) & \text{if } 2r = n-1 \\ (0\ldots 02) & \text{if } 2r = n \end{array}$	2	$\underbrace{(\underbrace{0\dots0}_{2r-1} 10\dots0; 0)}_{(0\dots01; 1)} \text{if } 2r < n-1$ $\underbrace{(0\dots01; 1)}_{(0\dots0; 2)} \text{if } 2r = n-1$ $\underbrace{(0\dots0; 2)}_{1} \text{if } 2r = n$	2	2n - 4r + 3	$\omega_{n-2i} - i\chi \ (i = 1, \dots, r)$
5.2	$(-2^r, +1^{n-2r}), r \ge 1$	$\begin{array}{c} \underbrace{(0\ldots 0}_{2r-1} 10\ldots 0) & \text{if } 2r < n-1 \\ (0\ldots 011) & \text{if } 2r = n-1 \\ (0\ldots 02) & \text{if } 2r = n \end{array}$	2	$(0 \dots 01 \underbrace{0 \dots 0}_{2r-1}; -2) \text{if } 2r < n$ $(0 \dots 0; -2) \text{if } 2r = n$	2	2n - 4r + 3	$\omega_{2i} + i\chi \ (i = 1, \dots, r)$
5.3	$(+2^r, -2^s, +1^{n-2r-2s}), r, s \ge 1$	$\begin{array}{ccc} (\underbrace{0\dots 0}_{2r+2s-1} & 10\dots 0) & \text{if } 2(r+s) < n-1 \\ (0\dots 011) & \text{if } 2(r+s) = n-1 \\ (0\dots 02) & \text{if } 2(r+s) = n \end{array}$	2	$\underbrace{(\underbrace{0\dots0}_{2r-1}10\dots01\underbrace{0\dots0}_{2s-1}; -2)}_{2r-1} \text{if } 2(r+s) < n$ $\underbrace{(\underbrace{0\dots0}_{2r-1}20\dots0; -2)}_{2r-1} \text{if } 2(r+s) = n$	2	2n - 4r - 4s + 3	$\omega_{2j} + j\chi \ (j = 1, \dots, s)$ $\omega_{n-2i} - i\chi \ (i = 1, \dots, r)$
5.4	$(+3,+1^{n-3})$	(0200)	4	(2002; -2)	2	3	$\omega_1 + \omega_{n-1}, \ \omega_2 + \chi, \ \omega_{n-2} - \chi$

Table 6: $G = \mathsf{D}_n, K^{ss} = \mathsf{A}_{n-1} \ (n \ge 4)$

Generators of
$$\Gamma(\overline{Ke})$$

 $\omega_{n-i} - i\chi \ (i = 1, ..., r)$
 $\omega_i + i\chi \ (i = 1, ..., r)$
 $\omega_j + j\chi \ (j = 1, ..., s)$
 $\omega_{n-i} - i\chi \ (i = 1, ..., r)$

	Signed partition	Diagram of $K\pi(e)$	Spherical roots
1.1	$(+2^r, +1^{p-r}, -1^{q-r}), r \ge 1$		$\alpha_{p-r+1} + \alpha'_{r-1}, \dots, \alpha_{p-1} + \alpha'_1$
1.2	$(-2^r, +1^{p-r}, -1^{q-r}), r \ge 1$		$\alpha_1 + \alpha'_{q-1}, \dots, \alpha_{r-1} + \alpha'_{q-r+1}$
1.3	$(+2^r, -2^s, +1^{p-r-s}, -1^{q-r-s}), r, s \ge 1$		$\alpha_1 + \alpha'_{q-1}, \dots, \alpha_{s-1} + \alpha'_{q-s+1}, \ \alpha_{p-r+1} + \alpha'_{r-1}, \dots, \alpha_{p-1} + \alpha'_1$
1.4	$(+3^2, +1^{p-4}), q=2$		$\alpha_1, \alpha_{p-1}, \alpha'$
1.5	$(-3^2, -1^{q-4}), p = 2$		$\alpha, \alpha'_1, \alpha'_{q-1}$
$ \begin{array}{c} 1.6\\(r+s < q-1) \end{array} $	$(+3, +2^r, -2^s, +1^{p-r-s-2}, -1^{q-r-s-1})$		$\alpha_1, \ldots, \alpha_s, \ \alpha_{p-r}, \ldots, \alpha_{p-1}, \ \alpha'_1, \ldots, \alpha'_r, \ \alpha'_{r+1} + \ldots + \alpha'_{q-s-1}, \ \alpha'_{q-s}, \ldots, \alpha'_{q-1}$
$ \begin{array}{c} 1.6\\(r+s=q-1)\end{array} $	$(+3,+2^r,-2^s,+1^{p-q-1})$		$\alpha_1,\ldots,\alpha_s, \ \alpha_{p-r},\ldots,\alpha_{p-1}, \ \alpha'_1,\ldots,\alpha'_{q-1}$
$ \begin{array}{c} 1.7\\(r+s < p-1) \end{array} $	$(-3, +2^r, -2^s, +1^{p-r-s-1}, -1^{q-r-s-2})$		$\alpha_1, \ldots, \alpha_s, \ \alpha_{s+1} + \ldots + \alpha_{p-r-1}, \ \alpha_{p-r}, \ldots, \alpha_{p-1}, \ \alpha'_1, \ldots, \alpha'_r, \ \alpha'_{q-s}, \ldots, \alpha'_{q-1}$
$ \begin{array}{c} 1.7\\(r+s=p-1)\end{array} $	$(-3, +2^r, -2^s, -1^{q-p-1})$		$\alpha_1, \ldots, \alpha_{p-1}, \ \alpha'_1, \ldots, \alpha'_r, \ \alpha'_{q-s}, \ldots, \alpha'_{q-1}$

Table 7: $G = \mathsf{A}_{p+q-1}, K^{ss} = \mathsf{A}_{p-1} \times \mathsf{A}_{q-1} \ (p,q \ge 1)$

	Signed partition	Diagram of $K\pi(e)$	Spherical roots
2.1	(+2 ² , +1 ²ⁿ⁻³) (I) or (II)	₀	none
2.2	$(+3,+1^{2n-3},-1)$	•••••	none
2.3	$(-3, +1^{2n-2})$ (I) or (II)	2 	$2(\alpha_1 + \ldots + \alpha_{n-1})$
2.4	$(+3^2,+1^{2n-5})$		α1

Table 8: $G = B_n, K^{ss} = B_{n-1} (n > 2)$

	Signed partition	Diagram of $K\pi(e)$	Spherical roots
3.1	$(+2^r,+1^{2n-2r})$		$2\alpha_{n-r+1},\ldots,2\alpha_{n-1}$
3.2	$(-2^r, +1^{2n-2r})$	• • • • • • • • • • • • • • • • • • •	$2\alpha_1,\ldots,2\alpha_{r-1}$
3.3	$(+2^r, -2^s, +1^{2n-2r-2s})$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$2\alpha_1,\ldots,2\alpha_{s-1},\ 2\alpha_{n-r+1},\ldots,2\alpha_{n-1}$

Table 9: $G = \mathsf{C}_n, K^{\mathrm{ss}} = \mathsf{A}_{n-1} \ (n \ge 2)$

	Signed partition	Diagram of $K\pi(e)$	Spherical roots
4.1	(+2 ² , +1 ²ⁿ⁻⁴) (I) or (II)	·····	none
4.2	$(+3,+1^{2n-4},-1)$	·····	none
4.3	(-3, $+1^{2n-3}$) (I) or (II)	•<	$2(\alpha_1 + \ldots + \alpha_{n-3}) + \alpha_{n-2} + \alpha_{n-1}$
4.4	$(+3^2,+1^{2n-6})$	$ \stackrel{\circ}{\underset{\circ}{\overset{\circ}{}} \bullet} \bullet $	α_1

Table 10: $G = \mathsf{D}_n, K^{ss} = \mathsf{D}_{n-1} (n > 4)$

	Signed partition	Diagram of $K\pi(e)$	Spherical roots
5.1	$(+2^r, +1^{n-2r})$	·•	$\alpha_{n-2i-1} + \alpha_{n-2i} + \alpha_{n-2i+1} \ (i = 1, \dots, r-1)$
5.2	$(-2^r,+1^{n-2r}),$	· · · · · · · · · · · · · · · · · · ·	$\alpha_{2i-1} + \alpha_{2i} + \alpha_{2i+1} \ (i = 1, \dots, r-1)$
5.3	$(+2^r, -2^s, +1^{n-2r-2s})$	$\cdots \circledast \cdots \circledast \circledast \cdots \cdots \circledast \circledast \cdots \cdots \circledast \cdots \circledast \cdots \circledast \circledast \cdots \cdots \circledast \circledast \cdots \cdots \circledast $	$\begin{array}{c} \alpha_{2j-1} + \alpha_{2j} + \alpha_{2j+1} & (j = 1, \dots, s-1) \\ \alpha_{n-2i-1} + \alpha_{n-2i} + \alpha_{n-2i+1} & (i = 1, \dots, r-1) \end{array}$
5.4	$(+3,+1^{n-3})$	$\odot \vdots \vdots \vdots \vdots \vdots \vdots \vdots \vdots \vdots $	$\alpha_2 + \ldots + \alpha_{n-2}$

Table 11: $G = \mathsf{D}_n, K^{\mathrm{ss}} = \mathsf{A}_{n-1} \ (n \ge 4)$

References

- D. N. Akhiezer, Equivariant completions of homogeneous algebraic varieties by homogeneous divisors, Ann. Global Anal. Geom. 1 (1983) 49–78.
- [2] R. Avdeev, Normalizers of Solvable Spherical Subgroups, Math. Notes 94 (2013), 20-31.
- [3] B. Binegar, On a class of multiplicity-free nilpotent K_C-orbits, J. Math. Kyoto Univ. 47 (2007), 735–766.
- [4] P. Bravi, R. Chirivì and J. Gandini, Regular functions on spherical nilpotent orbits in complex symmetric pairs: classical non-Hermitian cases, Kyoto J. Math. 57 (2017), to appear. doi:10.1215/21562261-2017-0013.
- [5] P. Bravi, R. Chirivì, J. Gandini and A. Maffei, Standard monomial theory for wonderful varieties, Algebr. Represent. Theory 19 (2016), 489–509.
- [6] P. Bravi, J. Gandini and A. Maffei, Projective normality of model varieties and related results, Represent. Theory 20 (2016), 39–93.
- [7] P. Bravi and G. Pezzini, The spherical systems of the wonderful reductive subgroups, J. Lie Theory 25 (2015), 105–123.
- [8] P. Bravi and G. Pezzini, Primitive wonderful varieties, Math. Z. 282 (2016), 1067–1096.
- [9] R. Chirivì, C. De Concini and A. Maffei, On normality of cones over symmetric varieties, Tohoku Math. J. (2) 58 (2006) 599–616.
- [10] R. Chirivì and A. Maffei, Projective normality of complete symmetric varieties, Duke Math. J. 122 (2004), 93–123.
- [11] D.H. Collingwood and W.M. McGovern, Nilpotent Orbits in Semisimple Lie Algebras, Van Nostrand Reinhold Co., New York, 1993.
- [12] C. De Concini, Normality and non normality of certain semigroups and orbit closures, in: Algebraic transformation groups and algebraic varieties, Encyclopaedia Math. Sci. 132, Springer, Berlin, 2004, pp. 15–35.
- [13] D.R. King, Classification of spherical nilpotent orbits in complex symmetric space, J. Lie Theory 14 (2004), 339–370.
- [14] D.R. King, Small spherical nilpotent orbits and K-types of Harish Chandra modules, arXiv:0701034 [math.RT].
- [15] F. Knop, The Luna-Vust Theory of spherical embeddings, in Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), 225–249, Manoj Prakashan, Madras, 1991.
- [16] F. Knop, Automorphisms, root systems, and compactifications of homogeneous varieties, J. Amer. Math. Soc. 9 (1996) 153–174.
- [17] H. Kraft and C. Procesi, On the geometry of conjugacy classes in classical groups, Comment. Math. Helv. 57 (1982), no. 4, 539–602.
- [18] K. Nishiyama, Classification of spherical nilpotent orbits for U(p, p), J. Math. Kyoto Univ. 44 (2004), 203–215.
- [19] R. Richardson, G. Röhrle and R. Steinberg, Parabolic subgroups with abelian unipotent radical, Invent. Math. 110 (1992), 649–671.
- [20] D.A. Timashev, Homogeneous spaces and equivariant embeddings, Encyclopaedia Math. Sci. 138, Springer, Heidelberg, 2011.

 $E\text{-}mail\ address: \texttt{braviQmat.uniroma1.it}$

 $Current \ address:$ Dipartimento di Matematica, Sapienza Università di Roma, Piazzale Aldo Moro 5, 00185 Roma, Italy

$E\text{-}mail\ address: \texttt{jacopo.gandini@sns.it}$

Current address: Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy