











# Publisher's Note: "Laser-driven strong shocks with infrared lasers at intensity of $10^{16}$ W/cm<sup>2</sup>" [Phys. Plasmas 26, 112708 (2019)]

Cite as: Phys. Plasmas 27, 039903 (2020); <https://doi.org/10.1063/5.0005221>

Submitted: 26 November 2019 . Published Online: 02 March 2020

 L. Antonelli, J. Trela,  F. Barbato, G. Boutoux, Ph. Nicolaï, D. Batani,  V. Tikhonchuk,  D. Mancelli, A. Tentori,  S. Atzeni,  A. Schiavi, F. Baffigi,  G. Cristoforetti,  S. Viciani,  L. A. Gizzi, M. Smid,  O. Renner,  J. Dostal,  R. Dudzak, L. Juha, and M. Krus



View Online



Export Citation



CrossMark

## ARTICLES YOU MAY BE INTERESTED IN


[Laser-driven strong shocks with infrared lasers at intensity of  \$10^{16}\$  W/cm<sup>2</sup>](#)  
Phys. Plasmas 26, 112708 (2019); <https://doi.org/10.1063/1.5119697>

[The experimental investigation of the hohlraum energetics of two-entrance holes spherical hohlraum at the 100 kJ level laser facility](#)  
Phys. Plasmas 27, 032702 (2020); <https://doi.org/10.1063/1.5136274>

[Mixing in ICF implosions on the National Ignition Facility caused by the fill-tube](#)  
Phys. Plasmas 27, 032703 (2020); <https://doi.org/10.1063/1.5125599>

AIP Advances  
Fluids and Plasmas Collection

READ NOW



# Publisher's Note: "Laser-driven strong shocks with infrared lasers at intensity of $10^{16}$ W/cm<sup>2</sup>" [Phys. Plasmas 26, 112708 (2019)]

Cite as: Phys. Plasmas **27**, 039903 (2020); doi: [10.1063/5.0005221](https://doi.org/10.1063/5.0005221)

Submitted: 26 November 2019 · Published Online: 2 March 2020















View Online



Export Citation



CrossMark

L. Antonelli,<sup>1</sup>  J. Trela,<sup>2</sup> F. Barbato,<sup>2</sup>  C. Boutoux,<sup>2</sup> Ph. Nicolai,<sup>2,a)</sup> D. Batani,<sup>2</sup> V. Tikhonchuk,<sup>2,3</sup>  D. Mancelli,<sup>2,4</sup>  A. Tentori,<sup>2</sup> S. Atzeni,<sup>5</sup>  A. Schiavi,<sup>5</sup>  F. Baffigi,<sup>6</sup> G. Cristoforetti,<sup>6</sup>  S. Viciani,<sup>6</sup>  L. A. Gizzi,<sup>6</sup>  M. Smid,<sup>3,7</sup> O. Renner,<sup>3,8</sup>  J. Dostal,<sup>3,8</sup>  R. Dudzak,<sup>3</sup>  L. Juha,<sup>3,8</sup> and M. Krus<sup>3</sup>

## AFFILIATIONS

<sup>1</sup>York Plasma Institute, Department of Physics, University of York, York YO10 5DQ, United Kingdom

<sup>2</sup>Université de Bordeaux, CNRS, CEA, CELIA (Centre Lasers Intenses et Applications), UMR 5107, F-33405 Talence, France

<sup>3</sup>Institute of Physics and ELI Beamlines, Czech Academy of Sciences, Czech Republic

<sup>4</sup>Donostia International Physics Center (DIPC), Donostia/San Sebastian, Basque Country, Spain

<sup>5</sup>Dipartimento SBAI, Università degli Studi di Roma "La Sapienza," Via Antonio Scarpa 14, 00161 Roma, Italy

<sup>6</sup>Intense Laser Irradiation Laboratory, INO CNR (National Council of Research), Pisa, Italy

<sup>7</sup>Helmholtz-Zentrum Dresden-Rosendorf, Dresden, Germany

<sup>8</sup>Institute of Plasma Physics, Czech Academy of Sciences, Prague, Czech Republic

<sup>a)</sup>Author to whom correspondence should be addressed: [philippe.nicolai@u-bordeaux.fr](mailto:philippe.nicolai@u-bordeaux.fr)

<https://doi.org/10.1063/5.0005221>

This article was originally published online on 19 November 2019 with an error in the affiliations. The affiliation "Institute of Physics and ELI Beamlines, Czech Academy of Sciences, Czech Republic" was missing from the originally published version, causing

the remaining affiliations to be misnumbered. The correct affiliations appear above. AIP Publishing apologizes for this error. All online versions of the article were corrected on 26 November 2019; the article is correct as it appears in the printed version of the journal.