

Fast cubature of high dimensional biharmonic potential based on Approximate Approximations

Flavia Lanzara* · Vladimir Maz'ya · Gunther Schmidt

Received: date / Accepted: date

Abstract We derive new formulas for the high dimensional biharmonic potential acting on Gaussians or Gaussians times special polynomials. These formulas can be used to construct accurate cubature formulas of an arbitrary high order which are fast and effective also in very high dimensions. Numerical tests show that the formulas are accurate and provide the predicted approximation rate $O(h^8)$ up to the dimension 10^7 .

Keywords Biharmonic potential · higher dimensions · separated representations · multidimensional convolution.

Mathematics Subject Classification (2010) 65D32 · 41A30 · 41A63

1 Introduction

The present paper is devoted to the approximation of the high dimensional biharmonic potential

$$\begin{aligned}\mathcal{B}_n f(\mathbf{x}) &= \frac{\Gamma(n/2)}{4\pi^{n/2}(n-2)(n-4)} \int_{\mathbb{R}^n} \frac{f(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|^{n-4}} d\mathbf{y}, \quad n \geq 3, n \neq 4, \\ \mathcal{B}_4 f(\mathbf{x}) &= -\frac{1}{8\pi^2} \int_{\mathbb{R}^4} \ln|\mathbf{x}-\mathbf{y}| f(\mathbf{y}) d\mathbf{y}\end{aligned}\tag{1.1}$$

(cf. [26, p.235] or [28, p.100]) for integrable f , by using approximate approximations (cf.[25] and the references therein). The potential (1.1) satisfies the equation $\mathcal{B}_n \Delta^2 f = \Delta^2 \mathcal{B}_n f = f$ for any smooth and rapidly decaying function f on \mathbb{R}^n ([29, p.118]). Equations involving

F. Lanzara
Mathematics Department, Sapienza University, P.le Aldo Moro 2, 00185 Rome, Italy
E-mail: lanzara@mat.uniroma1.it
ORCID <http://orcid.org/0000-0002-2052-4202>

V. Maz'ya
Department of Mathematics, University of Linköping, 581 83 Linköping, Sweden
RUDN University, 6 Miklukho-Maklay St, Moscow, 117198, Russia

G. Schmidt
Lichtenberger Str. 12, 10178 Berlin, Germany

the biharmonic operator in high dimension appear, for example, in conformal geometry (cf., e.g., [8] for $n = 3, 4$ and [23] for $n \geq 4$) or in the fourth order analogue of the Gel'fand problem (cf. [9]). Approximate approximations allow to construct efficient high order cubature formulas for convolution integral operators even with singular kernel functions (cf. [24]). Due to the operation number proportional to h^{-n} , where h denotes size of a uniform grid on the support of the density, these methods are practical only for small n . By combining approximate approximations with separated representations (also called tensor structured approximations) introduced in [3,4], we derive a method for approximating volume potentials which is accurate and fast in high dimensions.

In the last years modern methods based on tensor structured approximations have been applied successfully to some class of multidimensional integral operators (e.g. [5,6,7,10,11,12,13,14,15]). In this paper, for the first time to our knowledge, the fast cubature of the high dimensional biharmonic potential of an arbitrary high order is considered. We derive new formulas for the biharmonic potential acting on Gaussians or special polynomial times Gaussians. These formulas can be used to construct fast and high-order accurate cubature formulas in high dimensions. We report on numerical experiments which show approximation order $O(h^8)$ up to dimension $n = 10^7$.

First results on the fast cubature of high dimensional harmonic potential in the framework of approximate approximations have been obtained in [16,17]. The procedure has been applied in [18,19] to advection-diffusion potentials and in [20] to parabolic problems of second order. In [22,21] our approach has been extended to the Schrödinger equation and to the computation of the diffraction potential, where standard cubature methods are very expensive due to the fast oscillations of the kernel. Here we consider the problem of constructing fast cubature formulas for higher order problems.

We construct an approximation of $\mathcal{B}_n f$ if we replace f by functions with analytically known biharmonic potential. Specifically, we approximate the density $f \in C_0^N(\mathbb{R}^n)$ with the approximate quasi-interpolant

$$f_{h,\mathcal{D}}(\mathbf{x}) = \mathcal{D}^{-n/2} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) \eta \left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}} \right), \quad (1.2)$$

where h and \mathcal{D} are positive and η is a smooth and rapidly decaying function of the Schwarz space $\mathcal{S}(\mathbb{R}^n)$. The generating function is chosen so that $\mathcal{B}_n \eta$ can be computed analytically or efficiently numerically. If the generating function η satisfies the moment condition of order N

$$\int_{\mathbb{R}^n} \mathbf{x}^\alpha \eta(\mathbf{x}) d\mathbf{x} = \delta_{0,\alpha}, \quad 0 \leq |\alpha| < N, \quad (1.3)$$

then ([25, p.21])

$$|f(\mathbf{x}) - f_{h,\mathcal{D}}(\mathbf{x})| \leq c(\sqrt{\mathcal{D}}h)^N \|\nabla_N f\|_{L_\infty} + \sum_{k=0}^{N-1} \varepsilon_k (h\sqrt{\mathcal{D}})^k |\nabla_k f(\mathbf{x})|$$

with

$$0 < \varepsilon_k \leq \sum_{\mathbf{m} \in \mathbb{Z}^n \setminus \{0\}} |\nabla_k \mathcal{F} \eta(\sqrt{\mathcal{D}}\mathbf{m})|; \quad \lim_{\mathcal{D} \rightarrow \infty} \sum_{\mathbf{m} \in \mathbb{Z}^n \setminus \{0\}} |\nabla_k \mathcal{F} \eta(\sqrt{\mathcal{D}}\mathbf{m})| = 0$$

and

$$|\nabla_k f(\mathbf{x})| = \sum_{|\alpha|=k} \frac{|\partial^\alpha f(\mathbf{x})|}{\alpha!}.$$

Here $\mathcal{F}\eta$ denotes the Fourier transform of η

$$\mathcal{F}\eta(\mathbf{y}) = \int_{\mathbb{R}^n} \eta(\mathbf{x}) e^{-2i\pi(\mathbf{x}, \mathbf{y})} d\mathbf{x}$$

and we use the multi-index notation. Hence, for any *saturation error* $\varepsilon > 0$, one can fix the parameter $\mathcal{D} > 0$ so that

$$|f(\mathbf{x}) - f_{h, \mathcal{D}}(\mathbf{x})| = \mathcal{O}((\sqrt{\mathcal{D}}h)^N + \varepsilon) \|f\|_{W_\infty^N},$$

where $W_\infty^N = W_\infty^N(\mathbb{R}^n)$ denotes the Sobolev space of L_∞ -functions whose generalized derivatives up to the order N also belong to L_∞ . Then the linear combination

$$\mathcal{B}_n f_{h, \mathcal{D}}(\mathbf{x}) = \frac{h^4}{\mathcal{D}^{n/2-2}} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) \mathcal{B}_n \eta \left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}} \right) \quad (1.4)$$

gives rise to a new class of semi-analytic cubature formulas with the property that, for any prescribed accuracy $\varepsilon > 0$, one can fix the parameter \mathcal{D} so that (1.4) differs in some uniform or L_p -norm from the integral (1.1) by

$$\mathcal{O}((\sqrt{\mathcal{D}}h)^N + (\sqrt{\mathcal{D}}h)^4 \varepsilon) \quad \text{as} \quad h \rightarrow 0,$$

where N is determined by (1.3). Estimates of the cubature error for general generating functions are proved in Section 2.

Therefore, to construct cubature formulas for (1.1) it remains to compute the integral $\mathcal{B}_n \eta$. This can be taken analytically or transformed to a simple one-dimensional integral. We choose the generating functions

$$\eta(\mathbf{x}) = \pi^{-n/2} L_{M-1}^{(n/2)}(|\mathbf{x}|^2) e^{-|\mathbf{x}|^2}$$

with the generalized Laguerre polynomials

$$L_k^{(\gamma)}(y) = \frac{e^y y^{-\gamma}}{k!} \frac{d^k}{dy^k} (e^{-y} y^{k+\gamma}), \quad \gamma > -1,$$

orthogonal in the space $L_2(0, \infty)$ with weight $y^\gamma e^{-y}$. Since the function η satisfies the moment conditions (1.3) with $N = 2M$ ([25, p.55]), formula (1.4) gives rise to semi-analytic cubature formulas for $\mathcal{B}_n f$ of order h^{2M} modulo the saturation error. In Section 3 we describe these formulas when $M = 1$ that is for the exponential $\eta(\mathbf{x}) = \pi^{-n/2} e^{-|\mathbf{x}|^2}$ and, in Section 4, when $M > 1$.

If the generating function is the tensor product of one-dimensional functions of the form

$$\eta(\mathbf{x}) = \prod_{j=1}^n \pi^{-1/2} L_{M-1}^{(1/2)}(x_j^2) e^{-x_j^2},$$

each of them satisfying the moment conditions (1.3) of order $2M$, then $\mathcal{B}_n \eta$ is transformed to a one-dimensional integral with a separable integrand, i.e., a product of functions depending only on one of the variables. This is considered in Section 5 where we obtain, for example, the integral representation

$$\mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) = \frac{1}{16} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{n/2}} t dt, \quad n \geq 5.$$

These one-dimensional integrals with separable integrand in combination with a quadrature rule lead to accurate separated representations of the potential acting on the generating function. In Section 6 for functions f with separated representations, i.e., within a given accuracy they can be represented as a sum of products of univariate functions, we derive formulas which reduce the n -dimensional convolution (1.1) to one-dimensional discrete convolutions. Thus for the computation of (1.1) only one-dimensional operations are used. We derive formulas of an arbitrary order fast and accurate in high dimensions. We provide results of numerical experiments which show that even for very high space dimensions the approximations preserve the predicted convergence order 2, 4, 6 and 8 of the cubature.

2 Cubature error

The estimate of the cubature error

$$\mathcal{B}_n f_{h,\mathcal{D}}(\mathbf{x}) - \mathcal{B}_n f(\mathbf{x}) = \mathcal{B}_n(f_{h,\mathcal{D}} - f)(\mathbf{x})$$

for the biharmonic potential (1.1) is a consequence of the structure of the quasi-interpolation error, which is proved in general form in [25, Thm 2.28]. Suppose that f has generalized derivatives of order N . Using Taylor expansions of f for the nodes $h\mathbf{m}$, $\mathbf{m} \in \mathbb{Z}^n$, and Poisson's summation formula the quasi-interpolant can be written as

$$f_{h,\mathcal{D}}(\mathbf{x}) = (-h\sqrt{\mathcal{D}})^N f_N(\mathbf{x}) + \sum_{|\alpha|=0}^{N-1} \frac{(h\sqrt{\mathcal{D}})^{|\alpha|}}{\alpha!(2\pi i)^{|\alpha|}} \partial^\alpha f(\mathbf{x}) \sigma_\alpha(\mathbf{x}, \eta, \mathcal{D}) \quad (2.1)$$

with the function

$$f_N(\mathbf{x}) = \frac{1}{\mathcal{D}^{n/2}} \sum_{|\alpha|=N} \frac{N}{\alpha!} \sum_{\mathbf{m} \in \mathbb{Z}^n} \left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}} \right)^\alpha \eta \left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}} \right) \int_0^1 s^{N-1} \partial^\alpha f(s\mathbf{x} + (1-s)h\mathbf{m}) ds,$$

containing the remainder of the Taylor expansions, and the fast oscillating functions

$$\sigma_\alpha(\mathbf{x}, \eta, \mathcal{D}) = \sum_{\mathbf{v} \in \mathbb{Z}^n} \partial^\alpha \mathcal{F} \eta(\sqrt{\mathcal{D}}\mathbf{v}) e^{\frac{2\pi i}{h}(\mathbf{x}, \mathbf{v})}. \quad (2.2)$$

It follows from (2.2) that due to the moment condition (1.3) the second sum in (2.1) transforms to

$$f(\mathbf{x}) + \sum_{|\alpha|=0}^{N-1} \frac{(h\sqrt{\mathcal{D}})^{|\alpha|}}{\alpha!(2\pi i)^{|\alpha|}} \partial^\alpha f(\mathbf{x}) \varepsilon_\alpha(\mathbf{x}, \eta, \mathcal{D}),$$

where we denote

$$\varepsilon_\alpha(\mathbf{x}, \eta, \mathcal{D}) = \sum_{\mathbf{v} \in \mathbb{Z}^n \setminus \{0\}} \partial^\alpha \mathcal{F} \eta(\sqrt{\mathcal{D}}\mathbf{v}) e^{\frac{2\pi i}{h}(\mathbf{x}, \mathbf{v})} = \sigma_\alpha(\mathbf{x}, \eta, \mathcal{D}) - \delta_{0|\alpha|}.$$

We denote by $W_p^N = W_p^N(\mathbb{R}^n)$ the Sobolev space of $L_p = L_p(\mathbb{R}^n)$ functions whose generalized derivatives up to order N belong to L_p , with the norm

$$\|f\|_{W_p^N} = \sum_{l=0}^N |f|_{W_p^l}, \quad |f|_{W_p^l} = \sum_{|\alpha|=l} \|\partial^\alpha f\|_{L_p}.$$

If $f \in W_p^N$ with $N > n/p$, $1 \leq p \leq \infty$, then f_N can be estimated by

$$\|f_N\|_{L_p} \leq C_N |f|_{W_p^N}, \quad |f|_{W_p^N} = \sum_{|\alpha|=N} \|\partial^\alpha f\|_{L_p},$$

with a constant C_N not depending on f, h and \mathcal{D} . Hence (2.1) leads to the representation of the quasi-interpolation error

$$f_{h,\mathcal{D}}(\mathbf{x}) - f(\mathbf{x}) = (-h\sqrt{\mathcal{D}})^N f_N(\mathbf{x}) + \sum_{|\alpha|=0}^{N-1} \frac{(h\sqrt{\mathcal{D}})^{|\alpha|}}{\alpha!(2\pi i)^{|\alpha|}} \partial^\alpha f(\mathbf{x}) \varepsilon_\alpha(\mathbf{x}, \eta, \mathcal{D}), \quad (2.3)$$

which implies in particular the error estimate in L_p

$$\|f - f_{h,\mathcal{D}}\|_{L_p} \leq C_N (h\sqrt{\mathcal{D}})^N |f|_{W_p^N} + \sum_{k=0}^{N-1} \frac{(h\sqrt{\mathcal{D}})^k}{(2\pi)^k} \sum_{|\alpha|=k} \frac{\|\varepsilon_\alpha(\cdot, \eta, \mathcal{D})\|_{L_\infty} \|\partial^\alpha f\|_{L_p}}{\alpha!}. \quad (2.4)$$

Thus the quasi-interpolation error consists of a term ensuring $\mathcal{O}(h^N)$ -convergence and of the so-called saturation error, which, in general, does not converge to zero as $h \rightarrow 0$. However, due to the fast decay of $\partial^\alpha \mathcal{F}\eta$, one can choose \mathcal{D} large enough to ensure that

$$\|\varepsilon_\alpha(\cdot, \mathcal{D}, \eta)\|_{L_\infty} \leq \sum_{\mathbf{v} \in \mathbb{Z}^n \setminus \{0\}} |\partial^\alpha \mathcal{F}\eta(\sqrt{\mathcal{D}}\mathbf{v})| < \varepsilon \quad (2.5)$$

for given small $\varepsilon > 0$.

From Sobolev's theorem we have that for $n \geq 5$, $1 < p < n/4$, and $q = np/(n-4p)$ the integral (1.1) converges absolutely for almost every \mathbf{x} and the operator \mathcal{B}_n is a bounded mapping from L_p into L_q (cf. [29, p. 119]). Hence

$$\|\mathcal{B}_n f_{h,\mathcal{D}} - \mathcal{B}_n f\|_{L_q} \leq A_{p,q} \|f_{h,\mathcal{D}} - f\|_{L_p}, \quad (2.6)$$

where $A_{p,q}$ denotes the norm of $\mathcal{B}_n : L_p \rightarrow L_q$. Then, from (2.4) and (2.5),

Theorem 1 *Let $n \geq 5$, $1 < p < n/4$, $q = np/(n-4p)$ and $f \in W_p^N$ with $N > n/p$. Then, for any $\varepsilon > 0$ there exists $\mathcal{D} > 0$ such that*

$$\|\mathcal{B}_n f_{h,\mathcal{D}} - \mathcal{B}_n f\|_{L_q} \leq A_{p,q} \left(C_N (h\sqrt{\mathcal{D}})^N |f|_{W_p^N} + \varepsilon \sum_{k=0}^{N-1} \frac{(h\sqrt{\mathcal{D}})^k}{(2\pi)^k} \|\nabla_k f\|_{L_p} \right). \quad (2.7)$$

We used the notation

$$\|\nabla_k f\|_{L_p} = \sum_{|\alpha|=k} \frac{\|\partial^\alpha f\|_{L_p}}{\alpha!}.$$

It turns out, that under the conditions of Theorem 1 the cubature formula $\mathcal{B}_n f_{h,\mathcal{D}}$ converges to $\mathcal{B}_n f$. Since the biharmonic potential is a smoothing operator and by (2.3) the saturation error of the quasi-interpolant is a small, fast oscillating function, estimate (2.7) can be sharpened to the form that $\mathcal{B}_n f_{h,\mathcal{D}}$ approximates $\mathcal{B}_n f$ with the error $\mathcal{O}(h^N + \varepsilon h^4)$.

We denote by $H_p^s = H_p^s(\mathbb{R}^n)$ the Bessel potential space defined as the closure of compactly supported smooth functions with respect to the norm

$$\|u\|_{H_p^s} = \|\mathcal{F}^{-1}((1+4\pi^2|\cdot|^2)^{s/2} \mathcal{F}u)\|_{L_p} = \|(1-\Delta)^{s/2} u\|_{L_p}.$$

We shall use the error estimate for the quasi-interpolant (1.2) in the spaces H_p^s obtained in [25, p.83] which yields, in the case $s = -4$, the following result.

Theorem 2 [25, p.83] Suppose that $\eta \in \mathcal{S}(\mathbb{R}^n)$ satisfies the moment conditions (1.3) of order N . Then, for any $f \in W_p^L$, $1 < p < \infty$, $L \geq N \geq 4$ with $L > n/p$, there exist constants c_η and c_p , not depending on f and h such that $f_{h,\mathcal{D}}$ defined in (1.2) satisfies

$$\|f - f_{h,\mathcal{D}}\|_{H_p^{-4}} \leq c_\eta (h\sqrt{\mathcal{D}})^N \|f\|_{W_p^L} + c_p h^4 \sum_{k=0}^{N-5} \frac{(h\sqrt{\mathcal{D}})^k}{(2\pi)^{k+4}} \varepsilon_k(\mathcal{D}) \sum_{|\alpha|=k} \|\partial^\alpha f\|_{W_p^4} \quad (2.8)$$

with the numbers

$$\varepsilon_k(\mathcal{D}) = \max_{|\alpha|=k} \sum_{\mathbf{v} \in \mathbb{Z}^n \setminus \{0\}} \left| \partial^\alpha \mathcal{F} \eta(\sqrt{\mathcal{D}} \mathbf{v}) \right|.$$

Theorem 2 leads to the following error estimate for the quasi-interpolation procedure.

Theorem 3 Suppose that $\eta \in \mathcal{S}(\mathbb{R}^n)$ satisfies the moment conditions (1.3) of order N . Let $n \geq 5$, $1 < p < n/4$, $q = np/(n-4p)$, $r = np/(n-2p)$ and $f \in W_p^L$ with $L \geq N \geq 4$ and $L > n/p$. Then there exist constants c_η , c_p , c_q and c_r , not depending on f, h, \mathcal{D} , such that

$$\begin{aligned} \|\mathcal{B}_n f - \mathcal{B}_n f_{h,\mathcal{D}}\|_{L_q} &\leq c_\eta (h\sqrt{\mathcal{D}})^N \|f\|_{W_p^L} + \\ &h^4 \sum_{k=0}^{N-5} (h\sqrt{\mathcal{D}})^k \frac{\varepsilon_k(\mathcal{D})}{(2\pi)^{k+2}} \sum_{l=0}^4 \left(A_{p,q} c_p |f|_{W_p^{l+k}} + 2B_{r,q} c_r |f|_{W_r^{l+k}} + c_q |f|_{W_q^{l+k}} \right). \end{aligned}$$

Proof Let \mathcal{L}_n is the inverse of the Laplace operator. It is known that, for $n \geq 3$, $1 < r < n/2$ and $q = nr/(n-2r)$,

$$\|\mathcal{L}_n(f_{h,\mathcal{D}} - f)\|_{L_q} \leq B_{r,q} \|f_{h,\mathcal{D}} - f\|_{L_r} \quad (2.9)$$

where $B_{r,q}$ denotes the norm $\mathcal{L}_n : L_r \rightarrow L_q$. Hence, keeping in mind (2.6) and (2.9), we obtain

$$\begin{aligned} \|\mathcal{B}_n(f - f_{h,\mathcal{D}})\|_{L_q} &= \|\mathcal{B}_n(I - \Delta)^2(I - \Delta)^{-2}(f - f_{h,\mathcal{D}})\|_{L_q} \\ &\leq A_{p,q} \|(I - \Delta)^{-2}(f - f_{h,\mathcal{D}})\|_{L_p} + 2B_{r,q} \|(I - \Delta)^{-2}(f - f_{h,\mathcal{D}})\|_{L_r} + \|(I - \Delta)^{-2}(f - f_{h,\mathcal{D}})\|_{L_q} \\ &= A_{p,q} \|f - f_{h,\mathcal{D}}\|_{H_p^{-4}} + 2B_{r,q} \|f - f_{h,\mathcal{D}}\|_{H_r^{-4}} + \|f - f_{h,\mathcal{D}}\|_{H_q^{-4}}. \end{aligned}$$

Using the estimate (2.8) and the continuous embeddings $W_p^L \subset W_r^{L-2} \subset W_q^{L-4}$ ([29, p.124]) the assertion follows immediately. \square

3 Action on Gaussians

Consider the generating function $\eta_2(\mathbf{x}) = \pi^{-n/2} e^{-|\mathbf{x}|^2}$. The moment conditions (1.3) are fulfilled with $N = 2$. If we replace f in (1.1) by

$$f_{h,\mathcal{D}}(\mathbf{x}) = (\pi\mathcal{D})^{-n/2} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) e^{-|\mathbf{x}-h\mathbf{m}|^2/(h^2\mathcal{D})},$$

then we obtain a cubature for (1.1)

$$\mathcal{B}_n f_{h,\mathcal{D}}(\mathbf{x}) = \frac{(h\sqrt{\mathcal{D}})^4}{(\pi\mathcal{D})^{n/2}} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) \Phi_2 \left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}} \right) \quad (3.1)$$

with

$$\Phi_2(\mathbf{x}) := \pi^{n/2} \mathcal{B}_n \eta_2(\mathbf{x}) = \mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}).$$

Theorem 4 Let $n \geq 3, n \neq 4$. The biharmonic potential acting on the Gaussian allows the following representation

$$\mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) = \frac{1}{4(n-2)(n-4)} {}_1F_1\left(\frac{n-4}{2}, \frac{n}{2}, -|\mathbf{x}|^2\right), \quad (3.2)$$

where ${}_1F_1$ denotes the Kummer or confluent hypergeometric function.

Proof The cubature of the 3-dimensional biharmonic potential $\mathcal{B}_3 f$ is considered in [25, p.119]. To determine the action of the biharmonic potential on the Gaussian the general formula

$$(\mathcal{Q} * e^{-|\cdot|^2})(\mathbf{x}) = \frac{2\pi^{n/2} e^{-|\mathbf{x}|^2}}{|\mathbf{x}|^{n/2-1}} \int_0^\infty Q(r) e^{-r^2} I_{n/2-1}(2r|\mathbf{x}|) r^{n/2} dr \quad (3.3)$$

with the modified Bessel functions of the first kind I_n ([1, p.374]) and $Q(r) = -r/(8\pi)$ is used. Then (3.3) gives

$$\mathcal{B}_3(e^{-|\cdot|^2})(\mathbf{x}) = -\frac{e^{-|\mathbf{x}|^2}}{8} - \frac{\sqrt{\pi} \operatorname{erf}(|\mathbf{x}|)}{16|\mathbf{x}|} (2|\mathbf{x}|^2 + 1). \quad (3.4)$$

(3.4) can also be expressed by means of the confluent hypergeometric functions as

$$\mathcal{B}_3(e^{-|\cdot|^2})(\mathbf{x}) = -\frac{1}{4} {}_1F_1\left(-\frac{1}{2}, \frac{3}{2}, -|\mathbf{x}|^2\right).$$

Let $n \geq 5$. The convolution of two radial functions can be transformed to a one-dimensional integral by using the Fourier transforms of the radial functions. Indeed (cf. [25, (2.15) p.22])

$$\int_{\mathbb{R}^n} Q(|\mathbf{x}-\mathbf{y}|) f(|\mathbf{y}|) d\mathbf{y} = \frac{2\pi}{|\mathbf{x}|^{n/2-1}} \int_0^\infty \mathcal{F}Q(r) \mathcal{F}f(r) J_{n/2-1}(2\pi r|\mathbf{x}|) r^{n/2} dr. \quad (3.5)$$

Since $\mathcal{F}(e^{-|\cdot|^2})(\mathbf{x}) = \pi^{n/2} e^{-\pi^2|\mathbf{x}|^2}$ and $\mathcal{F}(|\cdot|^{4-n})(\mathbf{x}) = \frac{\pi^{n/2-4}}{\Gamma(\frac{n-4}{2})} |\mathbf{x}|^{-4}$ (cf. [27, p.156]) we have from (3.5) that

$$\mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) = \frac{\pi^{n/2-3}}{8|\mathbf{x}|^{n/2-1}} \int_0^\infty e^{-\pi^2 r^2} J_{n/2-1}(2\pi r|\mathbf{x}|) r^{n/2-4} dr,$$

where we used the relation

$$\Gamma\left(\frac{n}{2}\right) = \frac{(n-2)(n-4)}{4} \Gamma\left(\frac{n-4}{2}\right).$$

This integral can be expressed by means of the Kummer or confluent hypergeometric function ${}_1F_1(a, c, z)$ (cf. [2, (8.6.14)]). The representation formula (3.2) follows. \square

In particular, for $n = 5$, (3.2) gives

$$\mathcal{B}_5(e^{-|\cdot|^2})(\mathbf{x}) = \frac{1}{16} \left(\frac{e^{-|\mathbf{x}|^2}}{|\mathbf{x}|^2} + \sqrt{\pi} \frac{\operatorname{erf}(|\mathbf{x}|)}{2|\mathbf{x}|^3} (2|\mathbf{x}|^2 - 1) \right)$$

and, for $n = 6$, (cf. [1, 13.6]).

$$\mathcal{B}_6(e^{-|\cdot|^2})(\mathbf{x}) = \frac{e^{-|\mathbf{x}|^2} - 1 + |\mathbf{x}|^2}{16|\mathbf{x}|^4}.$$

In dimension $n = 4$ the following representation formula holds.

Theorem 5 *The biharmonic potential acting on the Gaussian function admits the representation*

$$\mathcal{B}_4(e^{-|\cdot|^2})(\mathbf{x}) = \frac{1}{16} \left(\frac{e^{-|\mathbf{x}|^2} - 1}{|\mathbf{x}|^2} - 2 \log |\mathbf{x}| - E_1(|\mathbf{x}|^2) \right) \quad (3.6)$$

where $E_1(r)$ is the exponential integral

$$E_1(r) = \int_r^\infty \frac{e^{-t}}{t} dt.$$

Proof We use that the radial function $g(r) = \Phi_2(\mathbf{x})$, $r = |\mathbf{x}|$, is solution of the differential equation

$$\frac{1}{r^3} \frac{d}{dr} \left(r^3 \frac{d}{dr} \left(\frac{1}{r^3} \frac{d}{dr} \left(r^3 \frac{d}{dr} \right) \right) \right) g(r) = e^{-r^2}, \quad r > 0,$$

satisfying the conditions

$$\begin{aligned} g(r) &\approx -\frac{1}{8} \log r \quad \text{as } r \rightarrow \infty, \\ g(0) &= -\frac{1}{4} \int_0^\infty r^3 \log(r) e^{-r^2} dr = \frac{\gamma-1}{16}, \quad g'(0) = 0 \end{aligned}$$

with the Euler constant γ . Denote by \mathcal{L}_n the inverse of the Laplace operator, the harmonic potential

$$(\mathcal{L}_n f)(\mathbf{x}) = -\frac{\Gamma(n/2-1)}{4\pi^{n/2}} \int_{\mathbb{R}^n} \frac{f(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|^{n-2}} d\mathbf{y},$$

which provides the unique solution of the Poisson equation

$$\Delta u = f \quad \text{in } \mathbb{R}^n, \quad |u(\mathbf{x})| \leq c|\mathbf{x}|^{n-2} \quad |\mathbf{x}| \rightarrow \infty.$$

Hence we have

$$\frac{1}{r^3} \frac{d}{dr} \left(r^3 \frac{d}{dr} \right) g = \mathcal{L}_4(e^{-|\cdot|^2})(\mathbf{x}).$$

From the relation (cf. [25, p.75])

$$\mathcal{L}_4(e^{-|\cdot|^2})(\mathbf{x}) = \frac{e^{-|\mathbf{x}|^2} - 1}{4|\mathbf{x}|^2}$$

we deduce that g solves

$$g''(r) + 3\frac{g'(r)}{r} = \frac{e^{-r^2} - 1}{4r^2}, \quad g(0) = \frac{\gamma-1}{16}, \quad g'(0) = 0.$$

We obtain

$$g'(r) = \frac{1}{8} \left(\frac{1}{r^3} - \frac{e^{-r^2}}{r^3} - \frac{1}{r} \right)$$

and, finally

$$\begin{aligned} g(r) &= \frac{1}{8} \int_0^r \left(\frac{1}{s^3} - \frac{e^{-s^2}}{s^3} - \frac{1}{s} \right) ds + g(0) = \frac{1}{16} \int_0^{r^2} \left(\frac{1}{t} - \frac{e^{-t}}{t} - 1 \right) \frac{dt}{t} \\ &+ g(0) = \frac{1}{16} \left(\frac{e^{-r^2} - 1}{r^2} - E_1(r^2) - 2 \log r \right) \end{aligned}$$

with the exponential integral E_1 ([1, 5.1.1]). \square

(3.1), together with (3.2) and (3.6), gives rise to second order semi-analytic cubature formulas for the biharmonic operator in any dimension $n \geq 3$.

4 Action on higher-order radial basis functions

In this Section we consider as generating functions the radial functions

$$\eta_{2M}(\mathbf{x}) = \pi^{-n/2} L_{M-1}^{(n/2)}(|\mathbf{x}|^2) e^{-|\mathbf{x}|^2}, \quad (4.1)$$

which satisfy the moment conditions of order $2M$ ([25, p.56]) and give rise to approximation formulas (1.2) of order $2M$ modulo the saturation error.

If we give an analytic formula for the biharmonic potential applied to η_{2M} , that is

$$\Phi_{2M}(\mathbf{x}) := \pi^{n/2} \mathcal{B}_n \eta_{2M}(\mathbf{x}) = \mathcal{B}_n(L_{M-1}^{(n/2)}(|\cdot|^2) e^{-|\cdot|^2})(\mathbf{x}),$$

then we obtain the following semi-analytic cubature for (1.1)

$$\mathcal{B}_n f_{h,\mathcal{D}}(\mathbf{x}) = \frac{(h\sqrt{\mathcal{D}})^4}{(\pi\mathcal{D})^{n/2}} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) \Phi_{2M}\left(\frac{\mathbf{x} - h\mathbf{m}}{h\sqrt{\mathcal{D}}}\right). \quad (4.2)$$

Theorem 6 For $M > 1$ we have

$$\Phi_{2M}(\mathbf{x}) = \mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) + \frac{1}{16|\mathbf{x}|^{n-2}} \gamma\left(\frac{n}{2} - 1, |\mathbf{x}|^2\right) + \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(n/2-1)}(|\mathbf{x}|^2)}{(j+1)(j+2)}$$

with the lower incomplete Gamma function

$$\gamma(a, x) = \int_0^x t^{a-1} e^{-t} dt.$$

Proof We use the relation [25, (3.18)]

$$L_{M-1}^{(n/2)}(|\mathbf{x}|^2) e^{-|\mathbf{x}|^2} = \sum_{j=0}^{M-1} \frac{(-1)^j}{j!4^j} \Delta^j e^{-|\mathbf{x}|^2} = e^{-|\mathbf{x}|^2} - \frac{1}{4} \Delta e^{-|\mathbf{x}|^2} + \sum_{j=2}^{M-1} \frac{(-1)^j}{j!4^j} \Delta^j e^{-|\mathbf{x}|^2}.$$

Let \mathcal{L}_n be the inverse of the Laplace operator, that is $\mathcal{L}_n \Delta = I$. Then $\mathcal{B}_n \Delta = \mathcal{L}_n$ and we have

$$\mathcal{B}_n(L_{M-1}^{(n/2)}(|\cdot|^2) e^{-|\cdot|^2})(\mathbf{x}) = \mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) - \frac{1}{4} \mathcal{L}_n(e^{-|\cdot|^2})(\mathbf{x}) + \sum_{j=2}^{M-1} \frac{(-1)^j}{j!4^j} \Delta^{j-2} e^{-|\mathbf{x}|^2}.$$

From the relations ([25, p.75])

$$\mathcal{L}_n(e^{-|\cdot|^2})(\mathbf{x}) = -\frac{1}{4|\mathbf{x}|^{n-2}} \gamma\left(\frac{n}{2} - 1, |\mathbf{x}|^2\right), \quad n \geq 3,$$

with the lower incomplete Gamma function $\gamma(a, \mathbf{x})$ and ([25, (4.24)])

$$\sum_{j=2}^{M-1} \frac{(-1)^j}{j!4^j} \Delta^{j-2} e^{-|\mathbf{x}|^2} = \frac{1}{16} \sum_{s=0}^{M-3} \frac{(-1)^s}{4^s (s+2)!} \Delta^s e^{-|\mathbf{x}|^2} = \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(n/2-1)}(|\mathbf{x}|^2)}{(j+1)(j+2)}$$

we conclude the proof. \square

From the relation ([1, 6.5.16])

$$\gamma\left(\frac{1}{2}, x^2\right) = \sqrt{\pi} \operatorname{erf}(x)$$

and the recurrence relation ([1, 6.5.22])

$$\gamma(a+1, x^2) = a\gamma(a, x) - e^{-x} x^2$$

we see that in the case of odd space dimension the biharmonic potential of the Gaussian is expressed using the error function erf. In the case of even space dimension the biharmonic potential of the Gaussian is expressed by elementary functions since ([1, 6.5.13])

$$\gamma(k, x) = (k-1)! \left(1 - e^{-x} \sum_{j=0}^{k-1} \frac{x^j}{j!} \right), \quad k \in \mathbf{N}.$$

In particular we have

$$\Phi_{2M}(\mathbf{x}) = -\frac{e^{-|\mathbf{x}|^2}}{8} - \frac{\sqrt{\pi}|\mathbf{x}|}{8} \operatorname{erf}(|\mathbf{x}|) + \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(1/2)}(|\mathbf{x}|^2)}{(j+1)(j+2)} \quad \text{for } n=3,$$

$$\Phi_{2M}(\mathbf{x}) = -\frac{\log|\mathbf{x}|}{8} - \frac{E_1(|\mathbf{x}|^2)}{16} + \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(1)}(|\mathbf{x}|^2)}{(j+1)(j+2)} \quad \text{for } n=4,$$

$$\Phi_{2M}(\mathbf{x}) = \frac{\sqrt{\pi} \operatorname{erf}(|\mathbf{x}|)}{16|\mathbf{x}|} + \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(3/2)}(|\mathbf{x}|^2)}{(j+1)(j+2)} \quad \text{for } n=5,$$

$$\Phi_{2M}(\mathbf{x}) = \frac{1 - e^{-|\mathbf{x}|^2}}{16|\mathbf{x}|^2} + \frac{e^{-|\mathbf{x}|^2}}{16} \sum_{j=0}^{M-3} \frac{L_j^{(2)}(|\mathbf{x}|^2)}{(j+1)(j+2)} \quad \text{for } n=6.$$

Theorem 6 shows that Φ_{2M+2} can be obtained from Φ_{2M} by adding some rapidly decaying terms. We conclude that the approximation of the density f by the quasi-interpolant (1.2) with the basis functions (4.1) leads to the semi-analytic approximation of the biharmonic potential (4.2) and the corresponding analytic expression for Φ_{2M} has to be used.

5 Action on tensor product basis functions

In this section we take the tensor product generating function

$$\eta_{2M}(\mathbf{x}) = \prod_{j=1}^n \tilde{\eta}_{2M}(x_j); \quad \tilde{\eta}_{2M}(x_j) = \frac{(-1)^{M-1}}{2^{2M-1} \sqrt{\pi} (M-1)!} \frac{H_{2M-1}(x_j) e^{-x_j^2}}{x_j}, \quad (5.1)$$

which satisfies the moment conditions of order $2M$, where H_k are the Hermite polynomials

$$H_k(x) = (-1)^k e^{x^2} \left(\frac{d}{dx} \right)^k e^{-x^2}.$$

The n -dimensional potential \mathcal{B}_n applied to the basis function η_{2M} can be transformed to a one-dimensional integral with separable integrand, i.e., a product of functions depending only on one of the variables. In Section 6 we will show how these one-dimensional integrals, in combination with a quadrature rule, lead to accurate separated representations of

the potential acting on the generating function. Hence, for functions f with separated representations, we derive fast formulas which reduces the n -dimensional convolution (1.1) by one-dimensional discrete convolutions.

We start with second order approximations, i.e. $M = 1$.

Theorem 7 *The biharmonic potential $\mathcal{B}_n(e^{-|\cdot|^2})$ admits the following one-dimensional integral representation*

$$\mathcal{B}_3(e^{-|\cdot|^2})(\mathbf{x}) = -\frac{1}{8} \int_0^\infty e^{-|\mathbf{x}|^2/(1+t)} \left(\frac{1}{(1+t)^{3/2}} + \frac{t|\mathbf{x}|^2}{(1+t)^{5/2}} \right) dt, \quad (5.2)$$

$$\mathcal{B}_n(e^{-|\cdot|^2})(\mathbf{x}) = \frac{1}{16} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{n/2}} t dt, \quad n \geq 5. \quad (5.3)$$

Proof We use the integral formula [1, 13.2.1]

$${}_1F_1(a, c, z) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 e^{z\tau} \tau^{a-1} (1-\tau)^{-a+c-1} d\tau, \quad \operatorname{Re}(c) > \operatorname{Re}(a) > 0. \quad (5.4)$$

Let $n \geq 5$. With the substitution $\tau = 1/(1+t)$ we get

$$\begin{aligned} {}_1F_1\left(\frac{n-4}{2}, \frac{n}{2}, -|\mathbf{x}|^2\right) &= \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n}{2}-2)} \int_0^\infty e^{-|\mathbf{x}|^2/(1+t)} \frac{1}{(1+t)^{n/2-3}} \frac{t}{1+t} \frac{dt}{(1+t)^2} \\ &= \frac{(n-2)(n-4)}{4} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{n/2}} t dt. \end{aligned}$$

From (3.2) in Theorem 4 we get (5.3). From the recurrence relation (cf. [1, 13.4.4]) we have

$${}_1F_1\left(-\frac{1}{2}, \frac{3}{2}, -|\mathbf{x}|^2\right) = {}_1F_1\left(\frac{1}{2}, \frac{3}{2}, -|\mathbf{x}|^2\right) + \frac{2}{3} |\mathbf{x}|^2 {}_1F_1\left(\frac{1}{2}, \frac{5}{2}, -|\mathbf{x}|^2\right).$$

The integral formula (5.4) gives

$${}_1F_1\left(\frac{1}{2}, \frac{3}{2}, -|\mathbf{x}|^2\right) = \frac{1}{2} \int_0^1 \frac{e^{-|\mathbf{x}|^2\tau}}{\tau^{1/2}} d\tau = \frac{1}{2} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{3/2}} dt,$$

$${}_1F_1\left(\frac{1}{2}, \frac{5}{2}, -|\mathbf{x}|^2\right) = \frac{3}{4} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{5/2}} t dt.$$

The integral representation (5.2) follows from (3.2). \square

In the next theorem we derive one-dimensional integral representations for the potential \mathcal{B}_n acting on the basis functions (5.1).

Theorem 8 For $M > 1$ we have

$$\begin{aligned} \mathcal{B}_3(\eta_{2M})(\mathbf{x}) = & -\frac{1}{8\pi^{3/2}} \left(\int_0^\infty \prod_{j=1}^3 \frac{e^{-x_j^2/(1+t)}}{\sqrt{1+t}} Q_M(x_j, t) dt \right. \\ & \left. + \int_0^\infty \left(\sum_{i=1}^3 \frac{e^{-x_i^2/(1+t)}}{\sqrt{1+t}} R_M(x_i, t) \prod_{\substack{j=1 \\ j \neq i}}^3 \frac{e^{-x_j^2/(1+t)}}{\sqrt{1+t}} Q_M(x_j, t) \right) t dt \right) \end{aligned} \quad (5.5)$$

$$\mathcal{B}_n(\eta_{2M})(\mathbf{x}) = \frac{1}{16\pi^{n/2}} \int_0^\infty \left(\prod_{j=1}^n \frac{e^{-x_j^2/(1+t)}}{\sqrt{1+t}} Q_M(x_j, t) \right) t dt, \quad n \geq 5 \quad (5.6)$$

with η_{2M} in (5.1) and

$$\begin{aligned} Q_M(x, t) &= \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{1}{(1+t)^k} H_{2k} \left(\frac{x}{\sqrt{1+t}} \right); \\ R_M(x, t) &= \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{1}{(1+t)^k} \mathcal{S}_{2k} \left(\frac{x}{\sqrt{1+t}} \right); \end{aligned} \quad (5.7)$$

$$\mathcal{S}_k(y) = y^2 H_k(y) - 2ky H_{k-1}(y) + k(k-1) H_{k-2}(y). \quad (5.8)$$

$Q_M(x, t)$ and $R_M(x, t)$ are polynomials in x whose coefficients depend on t .

Proof To get a one-dimensional integral representation of $\mathcal{B}_n(\prod_{j=1}^n \tilde{\eta}_{2M})$ we use the relation ([25, p.55])

$$\tilde{\eta}_{2M}(y) = \frac{1}{\sqrt{\pi}} \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dy^{2k}} e^{-y^2}.$$

Let $n \geq 5$. The solution of the equation

$$\Delta \Delta u = \prod_{j=1}^n \tilde{\eta}_{2M}(x_j)$$

is given by the integral

$$\begin{aligned} u(\mathbf{x}) &= \mathcal{B}_n \prod_{j=1}^n \tilde{\eta}_{2M}(x_j) = \frac{1}{16} \prod_{j=1}^n \frac{1}{\sqrt{\pi}} \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} \int_0^\infty \frac{e^{-x_j^2/(1+t)}}{(1+t)^{n/2}} t dt \\ &= \frac{1}{16} \int_0^\infty \left(\prod_{j=1}^n \frac{1}{\sqrt{\pi}} \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} e^{-x_j^2/(1+t)} \right) \frac{t dt}{(1+t)^{n/2}} \\ &= \frac{1}{16} \int_0^\infty \left(\prod_{j=1}^n \frac{1}{\sqrt{\pi}} \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{e^{-x_j^2/(1+t)}}{(1+t)^{k+1/2}} H_{2k} \left(\frac{x_j}{\sqrt{1+t}} \right) \right) t dt, \end{aligned}$$

that is (5.6).

Let $n = 3$. Keeping in mind (5.2), we get

$$\begin{aligned} \mathcal{B}_3(\eta_{2M})(\mathbf{x}) &= -\frac{1}{8\pi^{3/2}} \prod_{j=1}^3 \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} \int_0^\infty \frac{e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{3/2}} dt \\ &\quad - \frac{1}{8\pi^{3/2}} \prod_{j=1}^3 \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} \int_0^\infty \frac{|\mathbf{x}|^2 e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{5/2}} t dt. \end{aligned}$$

The first term in the r.h.s is similar to that considered in the case $n \geq 5$.

Concerning the second term, we have

$$\frac{d^{2k}}{dx^{2k}} \left(\frac{x^2}{1+t} e^{-x^2/(1+t)} \right) = \frac{1}{(1+t)^k} \frac{d^{2k}}{dy^{2k}} \left(y^2 e^{-y^2} \right)_{y=x/\sqrt{1+t}} = \frac{e^{-x^2/(1+t)}}{(1+t)^k} \mathcal{S}_{2k} \left(\frac{x}{\sqrt{1+t}} \right)$$

with $\mathcal{S}_k(y)$ in (5.8). Then

$$\sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx^{2k}} \left(\frac{x^2}{1+t} e^{-x^2/(1+t)} \right) = R_M(x, t) e^{-x^2/(1+t)}$$

with $R_M(x, t)$ defined in (5.7). It follows that the second integral can be written as

$$\begin{aligned} &\prod_{j=1}^3 \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} \int_0^\infty \frac{|\mathbf{x}|^2 e^{-|\mathbf{x}|^2/(1+t)}}{(1+t)^{5/2}} t dt \\ &= \int_0^\infty \prod_{j=1}^3 \sum_{k=0}^{M-1} \frac{(-1)^k}{k!4^k} \frac{d^{2k}}{dx_j^{2k}} |\mathbf{x}|^2 e^{-|\mathbf{x}|^2/(1+t)} \frac{t dt}{(1+t)^{5/2}} \\ &= \int_0^\infty \sum_{i=1}^3 R_M(x_i, t) \prod_{\substack{j=1 \\ j \neq i}}^3 Q_M(x_j, t) \frac{e^{-|\mathbf{x}|^2/(1+t)} t dt}{(1+t)^{3/2}}, \end{aligned}$$

which leads to (5.5). \square

The polynomials $Q_M(x, t)$ and $R_M(x, t)$ for $M = 1, 2, 3, 4$ are given by

$$\begin{aligned} Q_1(x, t) &= 1, & Q_2(x, t) &= -\frac{x^2}{(1+t)^2} + \frac{1}{2(1+t)} + 1, \\ Q_3(x, t) &= Q_2(x, t) + \frac{x^4}{2(1+t)^4} - \frac{3x^2}{2(1+t)^3} + \frac{3}{8(1+t)^2}, \\ Q_4(x, t) &= Q_3(x, t) - \frac{x^6}{6(1+t)^6} + \frac{5x^4}{4(1+t)^5} - \frac{15x^2}{8(1+t)^4} + \frac{5}{16(1+t)^3}, \\ R_1(x, t) &= \frac{x^2}{1+t}, & R_2(x, t) &= -\frac{x^4}{(1+t)^3} + \frac{x^2}{1+t} + \frac{5x^2}{2(1+t)^2} - \frac{1}{2(1+t)}, \\ R_3(x, t) &= R_2(x, t) + \frac{x^6}{2(1+t)^5} - \frac{7x^4}{2(1+t)^4} + \frac{39x^2}{8(1+t)^3} - \frac{3}{4(1+t)^2}, \\ R_4(x, t) &= R_3(x, t) - \frac{x^8}{6(1+t)^7} + \frac{9x^6}{4(1+t)^6} - \frac{65x^4}{8(1+t)^5} + \frac{125x^2}{16(1+t)^4} - \frac{15}{16(1+t)^3}. \end{aligned}$$

6 Separated representation of the biharmonic potential and numerical results

In this section we consider the fast computation of the biharmonic potential based on (5.1). From (4.2) and (5.6) we derive the cubature formula, valid for $n \geq 5$

$$\mathcal{B}_{M,h}^{(n)} f(\mathbf{x}) = \frac{(h\sqrt{\mathcal{D}})^4}{16(\pi\mathcal{D})^{n/2}} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) \int_0^\infty \left(\prod_{j=1}^n \frac{e^{-(x_j - hm_j)^2 / (h^2 \mathcal{D}(1+t))}}{\sqrt{1+t}} Q_M \left(\frac{x_j - hm_j}{h\sqrt{\mathcal{D}}}, t \right) \right) t dt.$$

At the grid points $h\mathbf{k} = (hk_1, \dots, hk_n)$ we obtain

$$\mathcal{B}_{M,h}^{(n)} f(h\mathbf{k}) = \frac{(h\sqrt{\mathcal{D}})^4}{16} \sum_{\mathbf{m} \in \mathbb{Z}^n} f(h\mathbf{m}) a_{\mathbf{k}-\mathbf{m}}^{(M)} \quad (6.1)$$

where

$$a_{\mathbf{k}}^{(M)} = \frac{1}{(\pi\mathcal{D})^{n/2}} \int_0^\infty \left(\prod_{j=1}^n \frac{e^{-k_j^2 / (\mathcal{D}(1+t))}}{\sqrt{1+t}} Q_M \left(\frac{k_j}{\sqrt{\mathcal{D}}}, t \right) \right) t dt. \quad (6.2)$$

The product structure of the integrand leads to new cubature formulas if the density f admits the so-called separated representation. The idea is the following. If f is given as product of univariate functions

$$f(\mathbf{x}) = \prod_{j=1}^n f_j(x_j)$$

then the values on the grid $h\mathbf{k}$ of the cubature formula can be written as

$$\mathcal{B}_{M,h}^{(n)} f(h\mathbf{k}) = \frac{(h\sqrt{\mathcal{D}})^4}{16(\pi\mathcal{D})^{n/2}} \int_0^\infty \left(\prod_{j=1}^n \sum_{m_j \in \mathbb{Z}} f_j(hm_j) \frac{e^{-(k_j - m_j)^2 / (\mathcal{D}(1+t))}}{\sqrt{1+t}} Q_M \left(\frac{k_j - m_j}{\sqrt{\mathcal{D}}}, t \right) \right) t dt.$$

A suitable quadrature of this integral with nodes τ_s and quadrature weights ω_s leads to

$$\mathcal{B}_{M,h}^{(n)} f(h\mathbf{k}) \approx \frac{(h\sqrt{\mathcal{D}})^4}{16(\pi\mathcal{D})^{n/2}} \sum_s \omega_s \tau_s \prod_{j=1}^n \sigma_j(k_j, \tau_s)$$

with

$$\sigma_j(k, \tau) = \sum_{m \in \mathbb{Z}} f_j(hm) \frac{e^{-(k-m)^2 / (\mathcal{D}(1+\tau))}}{\sqrt{1+\tau}} Q_M \left(\frac{k-m}{\sqrt{\mathcal{D}}}, \tau \right).$$

Then the value of the integral operator on the grid $h\mathbf{k}$ can be obtained by computing one-dimensional sums, and therefore the computational complexity of the algorithm scales linearly in the physical dimension.

We use an efficient quadrature based on the classical trapezoidal rule, which is exponentially converging for rapidly decaying smooth functions on the real line. We make the substitutions

$$t = e^\xi, \quad \xi = a(\sigma + e^\sigma), \quad \sigma = b(u - e^{-u})$$

with positive constants a and b proposed in [30] (see also [16, 17]). Then the integrals (6.2) are transformed to integrals over \mathbb{R} with integrands decaying doubly exponentially in u . After the substitution we have

$$a_{\mathbf{k}}^{(M)} = \frac{1}{(\pi\mathcal{D})^{n/2}} \int_{-\infty}^\infty \left(\prod_{j=1}^n \frac{e^{-k_j^2 / (\mathcal{D}(1+\Phi(u)))}}{\sqrt{1+\Phi(u)}} Q_M \left(\frac{k_j}{\sqrt{\mathcal{D}}}, \Phi(u) \right) \right) \Phi(u) \Phi'(u) du$$

with the functions

$$\begin{aligned}\Phi(u) &= \exp(ab(u - e^{-u}) + a \exp(b(u - e^{-u}))), \\ \Phi'(u) &= \Phi(u)ab(1 + e^{-u})(1 + \exp(b(u - e^{-u}))).\end{aligned}$$

Thus the trapezoidal rule of step τ can provide very accurate approximations of the integral for a relatively small number of nodes τs

$$a_{\mathbf{k}}^{(M)} \approx \frac{\tau}{(\pi \mathcal{D})^{n/2}} \sum_s \left(\prod_{j=1}^n \frac{e^{-k_j^2 / (\mathcal{D}(1 + \Phi(\tau s)))}}{\sqrt{1 + \Phi(\tau s)}} Q_M \left(\frac{k_j}{\sqrt{\mathcal{D}}}, \Phi(\tau s) \right) \right) \Phi(\tau s) \Phi'(\tau s).$$

Assume that f , within a prescribed accuracy, can be represented as sum of products of one-dimensional functions

$$f(\mathbf{x}) = \sum_{p=1}^P \beta_p \prod_{j=1}^n f_j^{(p)}(x_j) + \mathcal{O}(\varepsilon) \quad (6.3)$$

with suitable functions $f_j^{(p)}$ chosen such that the separation rank P is small. We derive the approximation of the convolutional sum (6.1) using one-dimensional operations

$$\begin{aligned}\mathcal{B}_{M,h}^{(n)} f(h\mathbf{k}) &\approx \frac{(h\sqrt{\mathcal{D}})^4}{16(\pi \mathcal{D})^{n/2}} \tau \sum_{p=1}^P \beta_p \sum_s \Phi(\tau s) \Phi'(\tau s) \\ &\times \prod_{j=1}^n \sum_{m_j} f_j^{(p)}(hm_j) \left(\frac{e^{-(k_j - m_j)^2 / (\mathcal{D}(1 + \Phi(\tau s)))}}{\sqrt{1 + \Phi(\tau s)}} Q_M \left(\frac{k_j - m_j}{\sqrt{\mathcal{D}}}, \Phi(\tau s) \right) \right).\end{aligned}$$

We provide results of some experiments which show the accuracy and numerical order of the method. We compute the biharmonic potential of the density

$$f(\mathbf{x}) = 4e^{-|\mathbf{x}|^2} (n(n+2) - 4(n+2)|\mathbf{x}|^2 + 4|\mathbf{x}|^4), \quad (6.4)$$

which has exact values $\mathcal{B}_n f(\mathbf{x}) = e^{-|\mathbf{x}|^2}$. In Table 1 we compare the exact values of $\mathcal{B}_n f$ and the approximate values $\mathcal{B}_{4,0.025}^{(n)} f$ at some grid points $(x_1, 0, \dots, 0) \in \mathbb{R}^n$ for space dimensions $n = 5, 10, 10^2, \dots, 10^8$.

In Table 2 we report on the absolute errors and approximation rates for the biharmonic potential $\mathcal{B}_n f(1, 0, \dots, 0)$ in the space dimensions $n = 5 \times 10^k$, $k = 0, \dots, 4$. The approximate values are computed by the cubature formulas $\mathcal{B}_{M,h}^{(n)}$ for $M = 1, 2, 3, 4$. We use uniform grids of size $h = 0.1 \times 2^{-k}$, $k = 1, \dots, 5$. For high dimensional cases the second order formula fails whereas the eighth order formula $\mathcal{B}_{4,h}^{(n)}$ approximates with the predicted approximation rates. Table 3 shows that the cubature method is effective also for much higher space dimensions and the approximation rate is reached. For all calculations the same quadrature rule is used for computing the one-dimensional integral, the parameters are $\mathcal{D} = 5$, $a = 6$ and $b = 5$, $\tau = 0.003$ and 300 summands in the quadrature sum.

In the remainder of this section we compute the 3-dimensional biharmonic potential by means of the approximating formula (5.5). For functions of the form (6.3) we obtain that, at

Table 1 Exact value of $\mathcal{B}_n f(x_1, 0, \dots, 0)$, absolute error and relative error using $\mathcal{D}_{4,0.025}^{(n)}$

n		5		10		100	
x_1	exact	abs. error	rel. error	abs. error	rel. error	abs. error	rel. error
0	0.100E+01	0.129E-09	0.129E-09	0.258E-09	0.258E-09	0.258E-08	0.258E-08
1	0.368E+00	0.286E-10	0.777E-10	0.760E-10	0.207E-09	0.930E-09	0.253E-08
2	0.183E-01	0.171E-11	0.933E-10	0.404E-11	0.220E-09	0.465E-10	0.254E-08
3	0.123E-03	0.112E-12	0.910E-09	0.943E-13	0.764E-09	0.381E-12	0.309E-08
4	0.113E-06	0.435E-13	0.386E-06	0.948E-14	0.843E-07	0.973E-14	0.864E-07
n		1000		10000		100000	
x_1	exact	abs. error	rel. error	abs. error	rel. error	abs. error	rel. error
0	0.100E+01	0.258E-07	0.258E-07	0.258E-06	0.258E-06	0.258E-05	0.258E-05
1	0.368E+00	0.947E-08	0.257E-07	0.948E-07	0.258E-06	0.949E-06	0.258E-05
2	0.183E-01	0.472E-09	0.257E-07	0.472E-08	0.258E-06	0.472E-07	0.258E-05
3	0.123E-03	0.324E-11	0.263E-07	0.319E-10	0.258E-06	0.318E-09	0.258E-05
4	0.113E-06	0.123E-13	0.110E-06	0.385E-13	0.342E-06	0.300E-12	0.266E-05
n		1000000		10000000		100000000	
x_1	exact	abs. error	rel. error	abs. error	rel. error	abs. error	rel. error
0	0.100E+01	0.258E-04	0.258E-04	0.258E-03	0.258E-03	0.258E-02	0.258E-02
1	0.368E+00	0.949E-05	0.258E-04	0.948E-04	0.258E-03	0.947E-03	0.258E-02
2	0.183E-01	0.472E-06	0.258E-04	0.472E-05	0.258E-03	0.472E-04	0.258E-02
3	0.123E-03	0.318E-08	0.258E-04	0.318E-07	0.258E-03	0.318E-06	0.258E-02
4	0.113E-06	0.291E-11	0.259E-04	0.290E-10	0.258E-03	0.290E-09	0.258E-02

the points of the uniform grid $\{h\mathbf{k}\}$, the 3-dimensional integral $\mathcal{B}_3 f$ is approximated by

$$\begin{aligned}
\mathcal{B}_3 f(h\mathbf{k}) \approx & -\frac{h^4 \mathcal{D}^{5/2}}{8\pi^{3/2}} \tau \sum_{p=1}^P \beta_p \sum_s \Phi'(\tau s) \\
& \times \left(\prod_{i=1}^3 \sum_{m_i} f_j^{(p)}(hm_i) \frac{e^{-(k_i-m_i)^2/(\mathcal{D}(1+\Phi(\tau s)))}}{\sqrt{1+\Phi(\tau s)}} Q_M\left(\frac{k_i-m_i}{\sqrt{\mathcal{D}}}, \Phi(\tau s)\right) \right. \\
& + \Phi(\tau s) \sum_{i=1}^3 \sum_{m_i} \frac{e^{-(k_i-m_i)^2/(\mathcal{D}(1+\Phi(\tau s)))}}{\sqrt{1+\Phi(\tau s)}} R_M\left(\frac{k_i-m_i}{\sqrt{\mathcal{D}}}, \Phi(\tau s)\right) f_i^{(p)}(hm_i) \\
& \left. \times \prod_{\substack{j=1 \\ j \neq i}}^3 \sum_{m_j} \frac{e^{-(k_j-m_j)^2/(\mathcal{D}(1+\Phi(\tau s)))}}{\sqrt{1+\Phi(\tau s)}} Q_M\left(\frac{k_j-m_j}{\sqrt{\mathcal{D}}}, \Phi(\tau s)\right) f_j^{(p)}(hm_j) \right).
\end{aligned}$$

In Table 4 we report on the relative and absolute errors, and the approximation rate for the 3-dimensional biharmonic potential $\mathcal{B}_3 f$ at the point $(1, 1, 1)$ of the density (6.4), which has exact value e^{-3} . The numerical results confirm the h^{2M} convergence of the cubature formula when $M = 1, 2, 3, 4$.

Acknowledgement

The second author was supported by the RUDN University Program 5-100.

Table 2 Absolute errors and approximation rates for $\mathcal{B}_n f(1, 0, \dots, 0)$ using $\mathcal{B}_{M,h}^{(n)}$

$M = 4$										
n	5		50		500		5000		50000	
h^{-1}	error	rate	error	rate	error	rate	error	rate	error	rate
10	0.15E-05		0.25E-04		0.26E-03		0.26E-02		0.25E-01	
20	0.70E-08	7.77	0.11E-06	7.81	0.12E-05	7.81	0.12E-04	7.81	0.12E-03	7.76
40	0.29E-10	7.94	0.46E-09	7.95	0.47E-08	7.95	0.47E-07	7.95	0.47E-06	7.95
80	0.15E-12	7.55	0.18E-11	7.99	0.19E-10	7.99	0.19E-09	7.99	0.19E-08	7.99
160	0.38E-13	2.02	0.10E-13	7.44	0.86E-13	7.75	0.84E-12	7.80	0.61E-11	8.26
$M = 3$										
10	0.30E-04		0.60E-03		0.62E-02		0.58E-01		0.30E+00	
20	0.53E-06	5.83	0.10E-04	5.86	0.11E-03	5.85	0.11E-02	5.74	0.11E-01	4.82
40	0.86E-08	5.96	0.17E-06	5.96	0.17E-05	5.96	0.17E-04	5.96	0.17E-03	5.94
80	0.13E-09	5.99	0.26E-08	5.99	0.27E-07	5.99	0.27E-06	5.99	0.27E-05	5.99
160	0.21E-11	5.97	0.41E-10	6.00	0.43E-09	6.00	0.43E-08	6.00	0.43E-07	6.00
$M = 2$										
10	0.74E-03		0.15E-01		0.13E+00		0.36E+00		0.37E+00	
20	0.49E-04	3.91	0.10E-02	3.89	0.10E-01	3.63	0.92E-01	1.98	0.35E+00	0.08
40	0.31E-05	3.98	0.63E-04	3.98	0.67E-03	3.96	0.66E-02	3.79	0.61E-01	2.50
80	0.20E-06	3.99	0.40E-05	3.99	0.42E-04	3.99	0.42E-03	3.98	0.42E-02	3.87
160	0.12E-07	4.00	0.25E-06	4.00	0.26E-05	4.00	0.26E-04	4.00	0.26E-03	3.99
$M = 1$										
10	0.26E-01		0.37E+00		0.37E+00		0.37E+00		0.37E+00	
20	0.68E-02	1.95	0.35E+00	0.07	0.35E+00	0.07	0.37E+00	0.00	0.37E+00	0.00
40	0.17E-02	1.99	0.20E+00	0.82	0.20E+00	0.82	0.37E+00	0.00	0.37E+00	0.00
80	0.43E-03	2.00	0.65E-01	1.61	0.65E-01	1.61	0.32E+00	0.22	0.37E+00	0.00
160	0.11E-03	2.00	0.17E-01	1.90	0.17E-01	1.90	0.14E+00	1.15	0.37E+00	0.01

Table 3 Absolute errors and approximation rates for $\mathcal{B}_n f(1, 0, \dots, 0)$ using $\mathcal{B}_{M,h}^{(n)}$

$M = 4$						
n	100000		1000000		10000000	
h^{-1}	error	rate	error	rate	error	rate
10	0.49E-01		0.28E+00		0.37E+00	
20	0.23E-03	7.71	0.23E-02	6.90	0.23E-01	4.02
40	0.95E-06	7.95	0.95E-05	7.95	0.95E-04	7.91
80	0.37E-08	7.99	0.37E-07	7.99	0.37E-06	7.99
160	0.13E-10	8.20	0.20E-09	7.58	0.11E-08	8.41
$M = 3$						
10	0.36E+00		0.37E+00		0.37E+00	
20	0.21E-01	4.08	0.16E+00	1.16	0.37E+00	0.00
40	0.35E-03	5.92	0.35E-02	5.57	0.33E-01	3.47
80	0.55E-05	5.99	0.55E-04	5.98	0.55E-03	5.92
160	0.86E-07	6.00	0.86E-06	6.00	0.86E-05	6.00

References

1. M. Abramowitz, I.A. Stegun, Handbook of Mathematical Functions, Dover Publ., New York (1968)
2. H. Bateman, A. Erdélyi, Tables of Integral transforms, Vol. 2, Mc Graw Hill (1954)
3. G. Beylkin, M.J. Mohlenkamp, Numerical-operator calculus in higher dimensions. Proc. Nat. Acad. Sci. USA, 99, 10246-10251 (2002)
4. G. Beylkin, M.J. Mohlenkamp, Algorithms for numerical analysis in high dimensions, SIAM J. Sci. Comput., 26(6), 2133–2159 (2005)

Table 4 Relative errors, absolute errors and approximation rates for $\mathcal{B}_3 f(1, 1, 1)$ using $\mathcal{B}_{M,h}^{(3)}$.

h^{-1}	$M = 4$			$M = 3$		
	absolute error	relative error	rate	absolute error	relative error	rate
10	0.236E-06	0.474E-05		0.822E-05	0.165E-03	
20	0.965E-09	0.194E-07	7.93	0.137E-06	0.275E-05	5.91
40	0.381E-11	0.765E-10	7.99	0.217E-08	0.436E-07	5.98
80	0.150E-13	0.301E-12	7.99	0.341E-10	0.685E-09	5.99
160	0.438E-14	0.879E-13	1.77	0.538E-12	0.108E-10	5.99
h^{-1}	$M = 2$			$M = 1$		
	absolute error	relative error	rate	absolute error	relative error	rate
10	0.217E-03	0.435E-02		0.359E-02	0.722E-01	
20	0.143E-04	0.287E-03	3.92	0.925E-03	0.186E-01	1.96
40	0.907E-06	0.182E-04	3.98	0.233E-03	0.468E-02	1.99
80	0.569E-07	0.114E-05	3.99	0.583E-04	0.117E-02	2.00
160	0.356E-08	0.715E-07	4.00	0.146E-04	0.293E-03	2.00

5. G. Beylkin, R. Cramer, G. Fann, R. J. Harrison, Multiresolution separated representations of singular and weakly singular operators, *Appl. Comput. Harmon. Anal.* 23 (2), 235–253 (2007)
6. G. Beylkin, G. Fann, R. J. Harrison, C. Kurcz, L. Monzón, Multiresolution representation of operators with boundary conditions on simple domains, *Appl. Comput. Harmon. Anal.* 33 (1), 109–139 (2012)
7. G. Beylkin, C. Kurcz, L. Monzón, Fast convolution with the free space Helmholtz Green's function, *J. Comput. Phys.* 228 (8), 2770–2791 (2009)
8. S.-Y. A. Chang, P. C. Yang, Fourth order equations in conformal geometry, *Séminaires & Congrès*, 4, 155–165 (2000)
9. L. Dupaigne, M. Ghergu, O. Goubet, G. Warnault, The Gel'fand Problem for the Biharmonic Operator, *Arch. Rational Mech. Anal.*, 208, 725–752 (2013)
10. I. P. Gavriljuk, W. Hackbusch, B. N. Khoromskij, Hierarchical tensor-product approximation to the inverse and related operators for high-dimensional elliptic problems, *Computing* 74 (2), 131–157 (2005)
11. W. Hackbusch, Efficient convolution with the Newton potential in d dimensions, *Numer. Math.* 110 (4), 449–489 (2008)
12. W. Hackbusch, B. N. Khoromskij, Tensor-product approximation to operators and functions in high dimensions, *J. Complexity* 23 (4-6), 697–714 (2007)
13. W. Hackbusch, B. N. Khoromskij, Tensor-product approximation to multidimensional integral operators and Green's functions, *SIAM J. Matrix Anal. Appl.* 30 (3), 1233–1253 (2008)
14. B. N. Khoromskij, Fast and accurate tensor approximation of a multivariate convolution with linear scaling in dimension, *J. Comput. Appl. Math.* 234 (11), 3122–3139 (2010)
15. B. N. Khoromskij, Range-separated tensor format for many-particle modeling, *Siam J. Sci. Comput.*, 40, A1034–A1062 (2018)
16. F. Lanzara, V. Maz'ya, G. Schmidt, On the fast computation of high dimensional volume potentials, *Math. Comput.*, 80, 887–904 (2011)
17. F. Lanzara, V. Maz'ya, G. Schmidt, Accuracy cubature of volume potentials over high-dimensional half-spaces, *J. Math. Sciences*, 173, 683–700 (2011)
18. F. Lanzara, V. Maz'ya, G. Schmidt, Fast cubature of volume potentials over rectangular domains by approximate approximations, *Appl. Comput. Harmon. Anal.* 36, 167–182 (2014)
19. F. Lanzara, G. Schmidt, On the computation of high-dimensional potentials of advection-diffusion operators, *Mathematika* 61, 309–327 (2015)
20. F. Lanzara, V. Maz'ya, G. Schmidt, Approximation of solutions to multidimensional parabolic equations by approximate approximations, *Appl. Comput. Harmon. Anal.* 41, 749–767 (2016)
21. F. Lanzara, V. Maz'ya, G. Schmidt, Accurate Computation of the high dimensional diffraction potential over hyper-rectangles, *Bulletin of TICMI* 22, 91–102 (2018)
22. F. Lanzara, V. Maz'ya, G. Schmidt, A fast solution method for time dependent multidimensional Schrödinger equations. *Applicable Analysis* 98, 408–429 (2019)
23. C. S. Lin, A classification of solutions of a conformally invariant fourth order equation in \mathbb{R}^n , *Comment. Math. Helv.* 73, 206–231 (1998)
24. V. Maz'ya, G. Schmidt, "Approximate Approximations" and the cubature of potentials. *Rend. Mat. Acc. Lincei* 6, 161–184 (1995)
25. V. Maz'ya, G. Schmidt, *Approximate Approximations*, AMS (2007)

-
26. D. Mitrea, *Distributions, Partial Differential Equations, and Harmonic Analysis*, Springer (2013)
 27. U. Neri, *Singular Integrals*, Lecture Notes in Mathematics, Springer (1971)
 28. O. Norbert, P. Wagner, *Fundamental solutions of linear Partial Differential Operators*, Springer (2015)
 29. E.M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press (1970)
 30. H. Takahasi and M. Mori, Doubly exponential formulas for numerical integration, *Publ. RIMS, Kyoto Univ.* 9, 721-741(1974)