

A C^1 REGULARITY RESULT FOR THE INHOMOGENEOUS NORMALIZED INFINITY LAPLACIAN

GRAZIANO CRASTA, ILARIA FRAGALÀ

ABSTRACT. We prove that the unique solution to the Dirichlet problem with constant source term for the inhomogeneous normalized infinity Laplacian on a convex domain of \mathbb{R}^N is of class C^1 . The result is obtained by showing as an intermediate step the power-concavity (of exponent $1/2$) of the solution.

1. INTRODUCTION

This paper is focused on the regularity of the unique solution to the Dirichlet problem

$$(1) \quad \begin{cases} -\Delta_{\infty}^N u = 1 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is an open bounded convex subset of \mathbb{R}^N and $\Delta_{\infty}^N u$ is the normalized infinity Laplacian. The symbolic definition of $\Delta_{\infty}^N \varphi$ for a smooth function φ is

$$(2) \quad \Delta_{\infty}^N \varphi(x) := \begin{cases} \frac{1}{|\nabla \varphi(x)|^2} \langle \nabla^2 \varphi(x) \nabla \varphi(x), \nabla \varphi(x) \rangle & \text{if } \nabla \varphi(x) \neq 0 \\ [\lambda_{\min}(\nabla^2 \varphi(x)), \lambda_{\max}(\nabla^2 \varphi(x))] & \text{if } \nabla \varphi(x) = 0, \end{cases}$$

being $\lambda_{\min}(\nabla^2 \varphi(x))$ and $\lambda_{\max}(\nabla^2 \varphi(x))$ respectively the minimum and the maximum eigenvalue of the Hessian matrix $\nabla^2 \varphi(x)$; for the detailed interpretation of the pde $-\Delta_{\infty}^N u = 1$ in the viscosity sense, we refer to Section 2 below.

The normalized infinity Laplace operator has recently attracted an increasing interest for its applications and connections with different areas, such as mass transportation [8], shape metamorphism [4], and especially differential games [13, 17]. In fact, according to Kohn and Serfaty [13], and Peres et al. [17], the equation $-\Delta_{\infty}^N u = 1$ is satisfied by the continuum value of a differential game called ‘‘tug of war’’ (a description can be found for instance in [2]).

The development of the existence, uniqueness, and regularity theory for boundary value problems involving the normalized infinity Laplace operator is still at its early stages.

The well-posedness of the Dirichlet problem (1) (possibly with a more general source term f) has been proved independently in [17] with probabilistic methods, in [15] with pde methods (see also [16]), and in [2] using a finite difference approach. Let us mention that the existence and uniqueness questions have been attacked also in the parabolic framework, see the interesting paper [12] on the evolution governed by $\Delta_{\infty}^N u$.

About regularity theory, so far no result seems to be available. As it is well-known, the fundamental contributions on regularity for pde’s involving the infinity Laplace operator are the celebrated works by Evans-Smart [10] and Evans-Savin [9], which concern infinity harmonic function and establish they are differentiable in any space dimension and $C^{1,\alpha}$

Date: June 17, 2015.

2010 Mathematics Subject Classification. Primary 49K20, Secondary 35J57, 35J70, 49N60.

Key words and phrases. viscosity solutions, infinity laplacian, tug of war, power-concavity.

in dimension two. Recently, some of these results have been extended to a class of inhomogeneous Dirichlet problems for the not-normalized infinity Laplacian in any dimension: the everywhere differentiability property has been obtained by Lindgren [14], and the C^1 regularity in case of a constant source term on a convex domain has been proved in our previous paper [6].

In case of the normalized operator, its definition via a dichotomy involving the maximum and minimum directions of the Hessian necessarily augments the difficulty of enforcing regularity techniques, and in fact no result beyond Lipschitz regularity is currently known. Aim of this paper is to present two new regularity results for the unique solution to problem (1) on convex domains.

The first result establishes that the solution is power-concave, precisely, $1/2$ -concave, see Theorem 6. To prove such result, we apply the convex envelope method by Alvarez-Lasry-Lions we already exploited in our previous paper [6], but in the current case this requires a more delicate procedure which is outlined for the benefit of the reader at the beginning of Section 3. In particular, this procedure exploits as a crucial tool a comparison principle proved in [2].

The second result is obtained as a consequence of the first one, and states that the solution is of class $C^1(\Omega)$, see Theorem 16. The proof relies on the local semiconcavity of the solution, combined with an estimate for semiconcave functions near singular points proved in [6].

The paper is organized as follows. In Section 2 we provide and discuss the definitions and properties of viscosity solution to problem (1) and to more general second order equations that we need to consider in the proofs. In Sections 3 and 4 we state and prove respectively the power-concavity and the C^1 regularity of the solution.

2. PRELIMINARIES

In this section we review the definition of normalized infinity Laplace operator, and that of viscosity sub- and super-solutions of (1), as well as of more general second order equations (that we shall need to use). Afterwards, we give some remarks to enlighten some basic features of solutions.

For a C^2 function φ defined in a neighborhood of $x \in \mathbb{R}^n$, we define the (*not normalized*) *infinity Laplace operator*

$$\Delta_\infty \varphi(x) := \langle \nabla^2 \varphi(x) \nabla \varphi(x), \nabla \varphi(x) \rangle$$

and the operators

$$\Delta_\infty^+ \varphi(x) := \begin{cases} |\nabla \varphi(x)|^{-2} \Delta_\infty \varphi(x), & \text{if } \nabla \varphi(x) \neq 0, \\ \lambda_{\max}(\nabla^2 \varphi(x)), & \text{if } \nabla \varphi(x) = 0, \end{cases}$$

$$\Delta_\infty^- \varphi(x) := \begin{cases} |\nabla \varphi(x)|^{-2} \Delta_\infty \varphi(x), & \text{if } \nabla \varphi(x) \neq 0, \\ \lambda_{\min}(\nabla^2 \varphi(x)), & \text{if } \nabla \varphi(x) = 0, \end{cases}$$

where, for a symmetric matrix $A \in \mathbb{R}_{\text{sym}}^{n \times n}(n)$, $\lambda_{\min}(A)$ and $\lambda_{\max}(A)$ denote respectively the minimum and the maximum eigenvalue of A .

In the following, if $u, v: \Omega \rightarrow \mathbb{R}$ are two functions and $x \in \Omega$, by

$$u \prec_x v$$

we mean that $u(x) = v(x)$ and $u(y) \leq v(y)$ for every $y \in \Omega$.

Moreover we recall that *second order sub-jet* (resp. *super-jet*), $J_{\Omega}^{2,-}u(x)$ (resp. $J_{\Omega}^{2,+}u(x)$), of a function $u \in C(\overline{\Omega})$ at a point $x \in \overline{\Omega}$, is by definition the set of pairs $(p, A) \in \mathbb{R}^n \times \mathbb{R}_{\text{sym}}^{n \times n}$ such that, as $y \rightarrow x$, $y \in \overline{\Omega}$, it holds

$$(3) \quad u(y) \geq (\leq) u(x) + \langle p, y - x \rangle + \frac{1}{2} \langle A(y - x_0), y - x_0 \rangle + o(|y - x|^2).$$

Definition 1. Let $f: \Omega \rightarrow \mathbb{R}$ be a continuous function and consider the normalized infinity Laplace equation

$$(4) \quad -\Delta_{\infty}^N u = f(x) \quad \text{in } \Omega.$$

(i) An upper semicontinuous function $u: \Omega \rightarrow \mathbb{R}$ is a *viscosity sub-solution* of (4) if, for every $x \in \Omega$,

$$-\Delta_{\infty}^+ \varphi(x) \leq f(x) \quad \forall \varphi \in C^2(\Omega) \text{ s.t. } u \prec_x \varphi.$$

The explicit formulation reads

$$\begin{cases} -\Delta_{\infty} \varphi(x) \leq f(x) |\nabla \varphi(x)|^2 & \forall \varphi \in C^2(\Omega) \text{ s.t. } u \prec_x \varphi, \text{ if } \nabla \varphi(x) \neq 0 \\ -\lambda_{\max}(\nabla^2 \varphi(x)) \leq f(x) & \forall \varphi \in C^2(\Omega) \text{ s.t. } u \prec_x \varphi, \text{ if } \nabla \varphi(x) = 0, \end{cases}$$

or in terms of super-jets

$$\begin{cases} -\langle Xp, p \rangle \leq f(x) |\nabla \varphi(x)|^2 & \forall (p, X) \in J_{\Omega}^{2,+}u(x), \text{ if } p \neq 0 \\ -\lambda_{\max}(X) \leq f(x) & \forall (p, X) \in J_{\Omega}^{2,+}u(x), \text{ if } p = 0. \end{cases}$$

(ii) A lower semicontinuous function $u: \Omega \rightarrow \mathbb{R}$ is a *viscosity super-solution* of (4) if, for every $x \in \Omega$,

$$-\Delta_{\infty}^- \varphi(x) \geq f(x) \quad \forall \varphi \in C^2(\Omega) \text{ s.t. } \varphi \prec_x u.$$

The explicit formulation reads

$$\begin{cases} -\Delta_{\infty} \varphi(x) \geq f(x) |\nabla \varphi(x)|^2 & \forall \varphi \in C^2(\Omega) \text{ s.t. } \varphi \prec_x u, \text{ if } \nabla \varphi(x) \neq 0 \\ -\lambda_{\min}(\nabla^2 \varphi(x)) \geq f(x) & \forall \varphi \in C^2(\Omega) \text{ s.t. } \varphi \prec_x u, \text{ if } \nabla \varphi(x) = 0, \end{cases}$$

or in terms of super-jets

$$\begin{cases} -\langle Xp, p \rangle \geq f(x) |\nabla \varphi(x)|^2 & \forall (p, X) \in J_{\Omega}^{2,-}u(x), \text{ if } p \neq 0 \\ -\lambda_{\min}(X) \geq f(x) & \forall (p, X) \in J_{\Omega}^{2,-}u(x), \text{ if } p = 0. \end{cases}$$

(iii) A function $u \in C(\Omega)$ is a *viscosity solution* of (4) if u is both a viscosity sub-solution and a viscosity super-solution of (4).

(iv) A function $u \in C(\overline{\Omega})$ is a *viscosity solution* of (1) if $u = 0$ on $\partial\Omega$ and u is a viscosity solution of (4) in Ω .

Definition 2. Let $I \subset \mathbb{R}$ be an open interval, let $H: \Omega \times I \times \mathbb{R}^n \times \mathbb{R}_{\text{sym}}^{n \times n}(n) \rightarrow \mathbb{R}$, and consider the equation

$$(5) \quad H(x, u, \nabla u, \nabla^2 u) = 0 \quad \text{in } \Omega.$$

(i) An upper semicontinuous function $u: \Omega \rightarrow I$ is a *viscosity sub-solution* of (5) if, for every $x \in \Omega$,

$$H_*(x, u(x), \nabla \varphi(x), \nabla^2 \varphi(x)) \leq 0 \quad \forall \varphi \in C^2(\Omega) \text{ s.t. } u \prec_x \varphi,$$

where H_* is the lower semicontinuous envelope of H .

(ii) A lower semicontinuous function $u: \Omega \rightarrow I$ is a *viscosity super-solution* of (5) if, for every $x \in \Omega$,

$$H^*(x, u(x), \nabla\varphi(x), \nabla^2\varphi(x)) \geq 0 \quad \forall \varphi \in C^2(\Omega) \text{ s.t. } u \prec_x \varphi,$$

where H^* is the upper semicontinuous envelope of H .

(iii) A function $u \in C(\Omega, I)$ is a *viscosity solution* of (5) if u is both a viscosity sub-solution and a viscosity super-solution of (4).

(iv) A function $u \in C(\bar{\Omega}, I)$ is a *viscosity solution of the homogeneous Dirichlet problem for equation (5)* if $u = 0$ on $\partial\Omega$ and u is a viscosity solution of (5) in Ω .

Remark 3. By taking the function

$$H(p, A) := -\left\langle A \frac{p}{|p|} \frac{p}{|p|} \right\rangle, \quad \forall (p, A) \in (\mathbb{R}^n \setminus \{0\}) \times \mathbb{R}_{\text{sym}}^{n \times n}(n),$$

and computing its upper and lower semicontinuous envelopes, we see that Definition 2 gives back Definition 1. (Note that $H_* = -(-H)^*$ and $H^* = -(-H)_*$). This argument justifies the apparently strange notions of the operators Δ_∞^+ and Δ_∞^- .

Remark 4. It is clear that the viscosity solution of the Dirichlet problem (1) is also a viscosity solution of

$$(6) \quad \begin{cases} -\Delta_\infty u = |\nabla u|^2 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

On the other hand, the converse is not true, and in fact the Dirichlet problem (6) has not, in general, a unique solution. To shed some light on this feature, it is enough to look at the one-dimensional case. If Ω is the interval $(-R, R)$, problems (1) and (6) read respectively

$$(7) \quad \begin{cases} -u'' = 1 & \text{in } (-R, R), \\ u(\pm R) = 0. \end{cases} \quad \begin{cases} -u''(u')^2 = (u')^2 & \text{in } (-R, R), \\ u(\pm R) = 0. \end{cases}$$

It is immediate to check that the function

$$u_r(x) := \begin{cases} \frac{R^2 - r^2}{2} & \text{if } |x| \leq r \\ \frac{R^2 - x^2}{2} & \text{if } r \leq |x| \leq R \end{cases}$$

is a solution to the second problem in (7) for every $r \in [0, R]$, whereas it is a solution to the first problem in (7) only for $r = 0$. Indeed, if $r \in (0, R]$, for every x with $|x| < r$ there exist smooth functions φ such that $\varphi \prec_x u$ but violate the condition $-\varphi''(x) \geq 1$, so that u is not a super-solution to $-u'' = 1$.

Remark 5. The viscosity solution u to problem (1) is strictly positive in Ω . Indeed, it is nonnegative by the comparison result proved in [2, Thm. 2.18]. Assume by contradiction that $u(x_0) = 0$ at some point $x_0 \in \Omega$. Then the function $\varphi \equiv 0$ touches u from below at x_0 , and hence u cannot be a viscosity supersolution to the equation $-\Delta_\infty^N u = 1$ at x_0 .

3. POWER-CONCAVITY OF SOLUTIONS ON CONVEX DOMAINS

In this section we prove:

Theorem 6. *Assume that Ω is an open bounded convex subset of \mathbb{R}^n , and let u be the solution to problem (1). Then $u^{1/2}$ is concave in Ω .*

Our proof strategy is the following.

Step 1. We show that the map $u \mapsto w := -u^{1/2}$ establishes a one-to-one correspondence between positive viscosity sub- and super-solution of $-\Delta_\infty^N u = 1$ in Ω and a “restricted class” of, respectively, negative viscosity super- and sub-solution of the equation

$$(8) \quad F(w, \nabla w, \nabla^2 w) = 0 \quad \text{in } \Omega,$$

where the function $F: (-\infty, 0) \times \mathbb{R}^n \times \mathbb{R}_{\text{sym}}^{n \times n}(n) \rightarrow \mathbb{R}$ is defined by

$$(9) \quad F(w, p, A) := -\langle Ap, p \rangle - \frac{1}{w} \left(|p|^4 + \frac{1}{2}|p|^2 \right).$$

Step 2. Under the additional assumption that

$$(10) \quad \text{the convex set } \Omega \text{ satisfies an interior sphere condition,}$$

we can adapt the convex envelope method of Alvarez, Lasry and Lions (see [1]), proving that the convex envelope w_{**} of a (restricted) super-solution w of (8) is still a (restricted) super-solution.

Step 3. Using the comparison principle proved in [2, Thm. 2.18], we conclude that, if (10) is fulfilled, then w is convex, namely $u^{1/2}$ is concave.

Step 4. By approximating Ω with outer parallel sets, we finally show that the assumption (10) can be removed.

Remark 7. The assumption (10) is used in a crucial way to prove Lemma 11 below, so that we do not need to impose *state constraints boundary conditions* on $\partial\Omega$ (see [1, Definition 2]). In this respect, we mention that the power concavity of solutions to (8) has been discussed by Juutinen in [11]; nevertheless, as kindly pointed out by the Author himself, his proof is flawed precisely in the argument used to show the validity of these state constraints boundary conditions. (See Lemma 4.1 in [11] where, at boundary points, the emptiness of $J_\Omega^{2,-}$ instead of $\bar{J}_\Omega^{2,-}$ is proved.)

Remark 8. The comparison principle proved in [2, Thm. 2.18], that we use as a crucial tool in Step 3, holds true for solutions to problem (1). In view of the one-to one correspondence mentioned in Step 1, it is therefore irremissible to deal with “restricted” solutions to the equation (8), according to Definition 9 below.

Step 1. We set the following

Definition 9. We say that w is a *restricted viscosity super-solution* of (8), if it satisfies

$$(11) \quad \forall x \in \Omega, \forall \psi \prec_x w \implies \begin{cases} F(w(x), \nabla \psi(x), \nabla^2 \psi(x)) \geq 0, \\ \lambda_{\min}(\nabla^2 \psi(x)) \leq -\frac{1}{2w(x)} \quad \text{if } \nabla \psi(x) = 0. \end{cases}$$

We say that w is a *restricted viscosity sub-solution* of (8), if it satisfies

$$(12) \quad \forall x \in \Omega, \forall w \prec_x \psi \implies \begin{cases} F(w(x), \nabla \psi(x), \nabla^2 \psi(x)) \leq 0, \\ \lambda_{\max}(\nabla^2 \psi(x)) \geq -\frac{1}{2w(x)} \quad \text{if } \nabla \psi(x) = 0. \end{cases}$$

Lemma 10. *A positive upper semicontinuous function $u: \Omega \rightarrow \mathbb{R}^+$ is a viscosity sub-solution of $-\Delta_\infty^N u = 1$ in Ω if and only if the function $w = -u^{1/2}$ is a restricted viscosity super-solution of (8). Likewise, a positive lower semicontinuous function $u: \Omega \rightarrow \mathbb{R}^+$ is a viscosity super-solution of $-\Delta_\infty^N u = 1$ in Ω if and only if the function $w = -u^{1/2}$ is a restricted viscosity sub-solution of (8).*

Proof. We are going to prove only the first part of the statement. To that aim it is enough to observe that

$$u \prec_x \varphi \iff \psi := -\varphi^{1/2} \prec_x w,$$

and the test functions φ and ψ satisfy

$$\nabla\psi(x) = -\frac{\nabla\varphi(x)}{2\varphi(x)^{1/2}}, \quad \nabla^2\psi(x) = \frac{1}{4\varphi(x)^{3/2}} \nabla\varphi(x) \otimes \nabla\varphi(x) - \frac{1}{2\varphi(x)^{1/2}} \nabla^2\varphi(x).$$

In particular, $\nabla\varphi(x) = 0$ if and only if $\nabla\psi(x) = 0$ and, if this is the case, $\lambda_{\max}(\nabla^2\varphi(x)) \geq -1$ if and only if $\lambda_{\min}(\nabla^2\psi(x)) \leq -1/(2w(x))$. \square

Step 2. Let us show that, if (10) is satisfied and w is a restricted viscosity solution to

$$(13) \quad \begin{cases} -\Delta_\infty w - \frac{1}{w} (|\nabla w|^4 + \frac{1}{2}|\nabla w|^2) = 0 & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega, \end{cases}$$

then w is convex. We denote by w_{**} the largest convex function below w . We first establish that, under the assumption (10), for every $x \in \Omega$, in the characterization

$$w_{**}(x) = \inf \left\{ \sum_{i=1}^k \lambda_i w(x_i) : x = \sum_{i=1}^k \lambda_i x_i, x_i \in \bar{\Omega}, \lambda_i > 0, \sum_{i=1}^k \lambda_i = 1, k \leq n+1 \right\}$$

the infimum can be attained only at interior points $x_i \in \Omega$:

Lemma 11. *Assume (10), and let u be the solution to problem (1). Set $w := -u^{1/2}$. For a fixed $x \in \Omega$, let $x_1, \dots, x_k \in \bar{\Omega}$, $\lambda_1, \dots, \lambda_k > 0$, with $\sum_{i=1}^k \lambda_i = 1$, be such that*

$$x = \sum_{i=1}^k \lambda_i x_i, \quad w_{**}(x) = \sum_{i=1}^k \lambda_i w(x_i).$$

Then $x_1, \dots, x_k \in \Omega$.

Proof. Assume by contradiction that at least one of the x_i 's, say x_1 , belongs to $\partial\Omega$. Let $B_R(y) \subset \Omega$ be a ball such that $\partial B_R(y) \cap \partial\Omega = \{x_1\}$. Since $-\Delta_\infty u = |\nabla u|^2$, by Lemma 2.2 in [5] the function $\tilde{u} := -u$ enjoys the property of comparison with cones from above according to Definition 2.3 in the same paper. Then, by Lemma 2.4 in [5], the function

$$r \mapsto \max_{x \in \partial B_r(y)} \frac{\tilde{u}(x) - \tilde{u}(y)}{r} = - \min_{x \in \partial B_r(y)} \frac{u(x) - u(y)}{r}$$

is monotone nondecreasing on the interval $(0, R)$. Namely, for all $r \in (0, R)$, there holds

$$(14) \quad \min_{x \in \partial B_r(y)} \frac{u(x) - u(y)}{|x - y|} \geq \min_{x \in \partial B_R(y)} \frac{u(x) - u(y)}{|x - y|} = -\frac{u(y)}{R},$$

where the last equality comes from the fact that u is non-negative in Ω (cf. Remark 5). By (14), we have

$$u(x) \geq u(y) \left(1 - \frac{|x - y|}{R} \right) \quad \forall x \in B_R(y),$$

and hence

$$(15) \quad w(x) \leq w(y) \left(1 - \frac{|x - y|}{R} \right)^{1/2} \quad \forall x \in B_R(y).$$

Let us define the unit vector $\zeta := (x - x_1)/|x - x_1|$ and let $\nu = (y - x_1)/|y - x_1|$ denote the inner normal of $\partial\Omega$ at x_1 . Since Ω is a convex set and $x \in \Omega$, we have that $\langle \zeta, \nu \rangle > 0$ and $x_1 + t\zeta \in B_R(y)$ for $t > 0$ small enough. Moreover, w_{**} is affine on $[x_1, x]$: indeed, since

the epigraph of w_{**} is the convex envelope of the epigraph of w , it is readily seen that w_{**} is affine on the whole set of convex combinations of the points $\{x_1, \dots, x_k\}$. Taking into account that $w_{**}(x_1) = w(x_1) = 0$, we infer that there exists $\mu > 0$ such that

$$w(x_1 + t\zeta) \geq w_{**}(x_1 + t\zeta) = -\mu t \quad \forall t \in [0, 1].$$

From (15) we obtain

$$-\mu t \leq w(y) \left(1 - \frac{|t\zeta - R\nu|}{R}\right)^{1/2} = w(y) \left(\langle \zeta, \nu \rangle \frac{t}{R} + o(t)\right)^{1/2}, \quad t \rightarrow 0^+,$$

and, recalling that $w(y) < 0$,

$$\mu t^{1/2} \geq K + o(1), \quad t \rightarrow 0^+$$

with $K > 0$, a contradiction. \square

On the basis of the lemma just proved, we obtain:

Lemma 12. *Assume (10), and let w be a restricted viscosity super-solution to (13). Then also w_{**} is a restricted viscosity super-solution to the same problem.*

Proof. Let w be a restricted viscosity super-solution to (13). In order to show that w_{**} is still a restricted viscosity super-solution to the same problem, we begin by observing that that w_{**} agrees with w on $\partial\Omega$, namely $w_{**} = 0$ on $\partial\Omega$ (since [1, Lemma 4.1] applies). Now let us check that w_{**} satisfies (11). In terms of sub-jets, such property can be rephrased as

$$(16) \quad \forall x \in \Omega, \forall (p, A) \in J_{\Omega}^{2,-} w_{**}(x) \implies \begin{cases} F(w_{**}(x), p, A) \geq 0 & \text{if } p \neq 0, \\ \lambda_{\min}(A) \leq -\frac{1}{2w_{**}(x)} & \text{if } p = 0. \end{cases}$$

Let $x \in \Omega$ and consider first the case when $(p, A) \in J_{\Omega}^{2,-} w_{**}(x)$, with $p \neq 0$.

For every $\epsilon > 0$ small enough, applying Proposition 1 in [1] and Lemma 11, we obtain points $x_1, \dots, x_k \in \Omega$, positive numbers $\lambda_1, \dots, \lambda_k$ satisfying $\sum_{i=1}^k \lambda_i = 1$, and elements $(p, A_i) \in \bar{J}_{\Omega}^{2,-} w(x_i)$, with A_i positive semidefinite, such that

$$\sum_{i=1}^k \lambda_i x_i = x, \quad \sum_{i=1}^k \lambda_i w(x_i) = w_{**}(x), \quad A - \epsilon A^2 \leq \left(\sum_{i=1}^k \lambda_i A_i^{-1} \right)^{-1} =: B.$$

We recall that, here and in the sequel, it is not restrictive to assume that the matrices A, A_1, \dots, A_k are positive definite, since the case of degenerate matrices can be handled as in [1], p. 273. Moreover we recall that the ‘‘closure’’ $J_{\Omega}^{2,-} v(x_0)$ is the set of $(p, A) \in \mathbb{R}^n \times \mathbb{R}_{\text{sym}}^{n \times n}$ for which there is a sequence $(p_j, A_j) \in J_{\Omega}^{2,-} v(x_j)$ (according to (3)) such that $(x_j, v(x_j), p_j, A_j) \rightarrow (x_0, v(x_0), p, A)$.

Then, since by assumption w is a super-solution to (13), we have $F(w(x_i), p, A_i) \geq 0$, i.e.

$$-w(x_i) \leq \frac{1}{\langle A_i p, p \rangle} \left(|p|^4 + \frac{1}{2} |p|^2 \right),$$

so that

$$-\frac{1}{\sum_{i=1}^k \lambda_i w(x_i)} \left(|p|^4 + \frac{1}{2} |p|^2 \right) \geq \left(\sum_{i=1}^k \lambda_i \frac{1}{\langle A_i p, p \rangle} \right)^{-1}.$$

Then, using the degenerate ellipticity of F and the concavity of the map $Q \mapsto 1/\text{tr}((p \otimes p)Q^{-1})$ (see [1], p. 286), we obtain

$$\begin{aligned} F(w_{**}(x), p, A - \epsilon A^2) &\geq -\langle Bp, p \rangle - \frac{1}{\sum_{i=1}^k \lambda_i w(x_i)} \left(|p|^4 + \frac{1}{2}|p|^2 \right) \\ &\geq -\langle Bp, p \rangle + \left(\sum_{i=1}^k \lambda_i \frac{1}{\langle A_i p, p \rangle} \right)^{-1} \\ &\geq 0. \end{aligned}$$

It remains to consider the case when $(0, A) \in J_{\Omega}^{2,-} w_{**}(x)$. We have to show that

$$\lambda_{\min}(A) \leq -\frac{1}{2w_{**}(x)}.$$

In terms of test functions, this is equivalent to prove that

$$\psi \prec_x w_{**}, \nabla \psi(x) = 0 \implies \lambda_{\min}(\nabla^2 \psi(x)) \leq -\frac{1}{2w_{**}(x)}.$$

Since w_{**} is a convex function, the conditions $\psi \prec_x w_{**}$ and $\nabla \psi(x) = 0$ imply that x is a minimum point of w_{**} . In particular, we must have $w(x_1) = \dots = w(x_k) = w_{**}(x)$.

If $k = 1$, then $w_{**}(x) = w(x)$, $B = A_1$ and $\lambda_{\min}(A - \epsilon A^2) \leq \lambda_{\min}(B) = \lambda_{\min}(A_1) \leq -1/(2w(x))$, so that the required inequality follows.

Assume now that $k > 1$, so that x is not a strict minimum point of w_{**} . Since x belongs to the relative interior of the convex polyhedron with vertices x_1, \dots, x_k , if we choose $q := (x_1 - x)/|x_1 - x|$ we get that $w_{**}(x + tq)$ is constant for $|t|$ small enough, so that $\psi(x + tq) \leq w_{**}(x) = \psi(x)$ for $|t|$ small. Hence

$$\lambda_{\min}(\nabla^2 \psi(x)) \leq \langle \nabla^2 \psi(x) q, q \rangle \leq 0 < -\frac{1}{2w_{**}(x)}$$

completing the proof. \square

Step 3. Let us prove that, under the additional assumption (10), the unique solution u to (1) is $1/2$ -power concave.

Let $w = -u^{1/2}$. By Lemma 10, w is a restricted super-solution to (13) hence, by Lemma 12, also w_{**} is a restricted super-solution to (13). Invoking again Lemma 10, the function $v := (w_{**})^2$ is a viscosity sub-solution to (1). By the comparison principle proved in [2, Thm. 2.18] we deduce that $v \leq u$, i.e. $(w_{**})^2 \leq w^2$. On the other hand, since $w_{**} \leq w$ (by definition of convex envelope) and $w \leq 0$, we have that $(w_{**})^2 \geq w^2$, so that $w = w_{**}$, namely w is a convex function.

Step 4. Let us finally show that the conclusions of Step 3 (i.e. the power concavity of u) remains true if Ω is any bounded convex domain.

For $\epsilon \in (0, 1]$ let Ω_ϵ denote the outer parallel body of Ω defined by

$$\Omega_\epsilon := \{x \in \mathbb{R}^n : \text{dist}(x, \Omega) < \epsilon\},$$

and let u_ϵ denote the solution to

$$\begin{cases} -\Delta_\infty^N u_\epsilon = 1 & \text{in } \Omega_\epsilon \\ u_\epsilon = 0 & \text{on } \partial\Omega_\epsilon. \end{cases}$$

Since Ω_ε satisfies an interior sphere condition (of radius ε), by Step 2 the function $u_\varepsilon^{1/2}$ is concave in Ω_ε . Therefore, to show that $u^{1/2}$ is concave in $\bar{\Omega}$, it is enough to show that, as $\varepsilon \rightarrow 0$, $u_\varepsilon \rightarrow u$ uniformly in $\bar{\Omega}$. In turn, by Theorem 5.3 in [15], this convergence holds true provided that $u_\varepsilon|_{\partial\Omega}$ converges uniformly to 0.

Let $y \in \partial\Omega$, and let $x_\varepsilon \in \partial\Omega_\varepsilon$ be such that $|x_\varepsilon - y| = \varepsilon$. Let us consider the polar quadratic polynomial

$$\eta(x) := \frac{1}{2} \text{diam}(\Omega_\varepsilon) |x - x_\varepsilon| - \frac{1}{2} |x - x_\varepsilon|^2.$$

Since $u_\varepsilon \leq \eta$ on $\partial\Omega_\varepsilon$, and u_ε enjoys the comparison with quadratic cones (see [15, Theorem 2.2] or [2, Lemma 5.1]), we have $u_\varepsilon \leq \eta$ on Ω_ε , and, in particular,

$$u_\varepsilon(y) \leq \frac{\varepsilon}{2} (\text{diam}(\Omega) + 1) - \frac{1}{2} \varepsilon^2.$$

Hence $u_\varepsilon|_{\partial\Omega}$ converges uniformly to 0 in $\partial\Omega_\varepsilon$.

4. LOCAL SEMICONCAVITY AND C^1 -REGULARITY OF SOLUTIONS

In this section we show the local semiconcavity and the C^1 -regularity of the unique solution to problem (1).

We recall that $u: \Omega \rightarrow \mathbb{R}$ is called *semiconcave (with constant C) in Ω* if

$$u(\lambda x + (1-\lambda)y) \geq \lambda u(x) + (1-\lambda)u(y) - C \frac{\lambda(1-\lambda)}{2} |x-y|^2 \quad \forall [x, y] \subset \Omega \text{ and } \forall \lambda \in [0, 1].$$

We say that u is *locally semiconcave in Ω* if it is semiconcave on compact subsets of Ω .

Proposition 13. *Assume that Ω is an open bounded convex subset of \mathbb{R}^n , and let u be the solution to problem (1). Then u is locally semiconcave in Ω .*

Proof. Let K be a compact convex subset of Ω , and let M be the Lipschitz constant of $v := u^{1/2}$ in K . We claim that u is semiconcave in K with semiconcavity constant $C = 2M^2$. Namely, given $x, y \in K$ and $\lambda \in [0, 1]$ and using the concavity of v established in Theorem 6, we get

$$\begin{aligned} u(\lambda x + (1-\lambda)y) - \lambda u(x) - (1-\lambda)u(y) &+ \frac{C}{2} \lambda(1-\lambda) |x-y|^2 \\ &\geq [\lambda v(x) + (1-\lambda)v(y)]^2 - \lambda v(x)^2 - (1-\lambda)v(y)^2 + M^2 \lambda(1-\lambda) |x-y|^2 \\ &= \lambda(1-\lambda) [M^2 |x-y|^2 - |v(x) - v(y)|^2] \geq 0. \end{aligned} \quad \square$$

Remark 14. In the above result the semiconcavity property of u is stated just *locally* in Ω . The reason can be understood by inspection of the proof, and analyzing the behaviour of the constant M appearing therein as the compact set $K \uparrow \Omega$. Indeed, recalling that the function $w = -v = -u^{1/2}$ satisfies (15), choosing $x = x_1 + \lambda\nu$, and taking into account that $w(x_1) = 0$, it is readily seen that

$$\lim_{\lambda \rightarrow 0^+} \frac{w(x_1 + \lambda\nu) - w(x_1)}{\lambda} \leq \lim_{\lambda \rightarrow 0^+} \frac{w(y)}{\lambda} \left(\frac{\lambda}{R} \right)^{1/2} = -\infty.$$

This means that the normal derivative of w with respect to the external normal is $+\infty$ at every boundary point of Ω (so that $M \rightarrow +\infty$ as $K \uparrow \Omega$).

Next, let us quote an estimate for locally semiconcave functions near singular points that we are going to exploit in order to arrive at the C^1 -regularity; such result was proved in our previous paper [6, Thm. 8] (see also [7, Thm. 5]).

Given a function $u \in C(\Omega)$, we denote by $\Sigma(u)$ the singular set of u , namely the set of points where u is not differentiable. For every $x_0 \in \Sigma(u)$, the *super-differential* of u at x_0 , which is defined by

$$D^+u(x_0) := \left\{ p \in \mathbb{R}^n : \limsup_{x \rightarrow x_0} \frac{u(x) - u(x_0) - \langle p, x - x_0 \rangle}{|x - x_0|} \leq 0 \right\},$$

turns out to be a nonempty compact convex set different from a singleton. In particular, $D^+u(x_0) \setminus \text{extr } D^+u(x_0)$ is not empty and contains non-zero elements.

Theorem 15. ([6, Theorem 8]) *Let $u: \Omega \rightarrow \mathbb{R}$ be a locally semiconcave function, let $x_0 \in \Sigma(u)$, and let $p \in D^+u(x_0) \setminus \text{extr } D^+u(x_0)$. Let $R > 0$ be such that $\overline{B}_R(x_0) \subset \Omega$, and let C denote the semiconcavity constant of u on $\overline{B}_R(x_0)$. Then there exist a constant $K > 0$ and a unit vector $\zeta \in \mathbb{R}^n$ satisfying the following property:*

$$(17) \quad u(x) \leq u(x_0) + \langle p, x - x_0 \rangle - K |\langle \zeta, x - x_0 \rangle| + \frac{C}{2} |x - x_0|^2 \quad \forall x \in \overline{B}_R(x_0).$$

In particular, for every $c > 0$, setting $\delta := \min\{K/c, R\}$, it holds

$$(18) \quad u(x) \leq u(x_0) + \langle p, x - x_0 \rangle - c \langle \zeta, x - x_0 \rangle^2 + \frac{C}{2} |x - x_0|^2 \quad \forall x \in \overline{B}_\delta(x_0).$$

Furthermore, if $p \neq 0$ then the vector ζ can be chosen so that $\langle \zeta, p \rangle \neq 0$.

We are now ready to give our C^1 -regularity result:

Theorem 16. *Let $u \in C(\Omega)$ be a viscosity solution to $-\Delta_\infty u = f(x, u)$ in Ω . If u is locally semiconcave in Ω , then u is everywhere differentiable (hence of class C^1) in Ω . In particular, if Ω is an open bounded convex subset of \mathbb{R}^n , and u is the unique solution to problem (1), then $u \in C^1(\Omega)$.*

Proof. Let $u \in C(\Omega)$ be a locally semiconcave viscosity solution to $-\Delta_\infty u = f(x, u)$ in Ω . Assume by contradiction that $\Sigma(u) \neq \emptyset$. Without loss of generality we can assume that $0 \in \Sigma(u)$. Let $p \in D^+u(0) \setminus \text{extr } D^+u(0)$, $p \neq 0$. By Theorem 15, there exists a unit vector $\zeta \in \mathbb{R}^n$, with $\langle \zeta, p \rangle \neq 0$, such that, for every $c > 0$, the inequality

$$u(x) \leq u(0) + \langle p, x \rangle - c \langle \zeta, x \rangle^2 + \frac{C}{2} |x|^2$$

holds true for all $x \in B_\delta(0)$, with δ depending on c . Thus, setting $\varphi(x) := u(0) + \langle p, x \rangle - c \langle \zeta, x \rangle^2 + \frac{C}{2} |x|^2$, it holds $u \prec_0 \varphi$. Since $\nabla \varphi(0) = p \neq 0$, we have

$$\Delta_\infty^+ \varphi(0) = -2c \langle \zeta, p \rangle^2 + C|p|^2,$$

choosing $c > 0$ large enough we get $-\Delta_\infty^+ \varphi(0) > f(0, u(0))$, a contradiction.

Since u is differentiable everywhere in Ω , by [3, Prop. 3.3.4] we conclude that $u \in C^1(\Omega)$. Finally, if Ω is an open bounded convex subset of \mathbb{R}^n , and u is the unique solution to problem (1), the first part of the statement just proved applies (and hence $u \in C^1(\Omega)$) because we know from Proposition 13 that u is locally semiconcave. \square

REFERENCES

- [1] O. Alvarez, J.-M. Lasry, and P.-L. Lions, *Convex viscosity solutions and state constraints*, J. Math. Pures Appl. (9) **76** (1997), no. 3, 265–288. MR 1441987 (98k:35045)
- [2] S.N. Armstrong and C.K. Smart, *A finite difference approach to the infinity Laplace equation and tug-of-war games*, Trans. Amer. Math. Soc. **364** (2012), no. 2, 595–636. MR 2846345
- [3] P. Cannarsa and C. Sinestrari, *Semiconcave functions, Hamilton-Jacobi equations and optimal control*, Progress in Nonlinear Differential Equations and their Applications, vol. 58, Birkhäuser, Boston, 2004.
- [4] G. Cong, M. Esser, B Parvin, and G. Bebis, *Shape metamorphism using p -Laplace equation*, Proceedings of the 17th International Conference on Pattern Recognition **4** (2004), 15–18.
- [5] M.G. Crandall, L.C. Evans, and R.F. Gariepy, *Optimal Lipschitz extensions and the infinity Laplacian*, Calc. Var. Partial Differential Equations **13** (2001), no. 2, 123–139.
- [6] G. Crasta and I. Fragalà, *On the Dirichlet and Serrin problems for the inhomogeneous infinity Laplacian in convex domains: Regularity and geometric results*, to appear in Arch. Rational Mech. Anal., 2015.
- [7] G. Crasta and I. Fragalà, *A symmetry problem for the infinity Laplacian*, Int. Mat. Res. Not. IMRN (2014), doi: 10.1093/imrn/rnu204
- [8] L.C. Evans and W. Gangbo, *Differential equations methods for the Monge-Kantorovich mass transfer problem*, Mem. Amer. Math. Soc. **137** (1999), no. 653.
- [9] L.C. Evans and O. Savin, *$C^{1,\alpha}$ regularity for infinity harmonic functions in two dimensions*, Calc. Var. Partial Differential Equations **32** (2008), 325–347.
- [10] L.C. Evans and C.K. Smart, *Everywhere differentiability of infinity harmonic functions*, Calc. Var. Partial Differential Equations **42** (2011), 289–299.
- [11] P. Juutinen, *Concavity maximum principle for viscosity solutions of singular equations*, NoDEA Nonlinear Differential Equations Appl. **17** (2010), 601–618.
- [12] P. Juutinen and B. Kawohl, *On the evolution governed by the infinity Laplacian*, Math. Ann. **335** (2006), no. 4, 819–851.
- [13] R. V. Kohn and S. Serfaty, *A deterministic-control-based approach to motion by curvature*, Comm. Pure Appl. Math. **59** (2006), no. 3, 344–407.
- [14] E. Lindgren, *On the regularity of solutions of the inhomogeneous infinity Laplace equation*, Proc. Amer. Math. Soc. **142** (2014), no. 1, 277–288. MR 3119202
- [15] G. Lu and P. Wang, *A PDE perspective of the normalized infinity Laplacian*, Comm. Partial Differential Equations **33** (2008), no. 10-12, 1788–1817. MR 2475319 (2009m:35150)
- [16] G. Lu and P. Wang, *Infinity Laplace equation with non-trivial right-hand side*, Electron. J. Differential Equations (2010), No. 77, 12.
- [17] Y. Peres, O. Schramm, S. Sheffield, and D. B. Wilson, *Tug-of-war and the infinity Laplacian*, J. Amer. Math. Soc. **22** (2009), no. 1, 167–210.

(Graziano Crasta) DIPARTIMENTO DI MATEMATICA “G. CASTELNUOVO”, UNIV. DI ROMA I, P.LE A. MORO 2 – 00185 ROMA (ITALY)

E-mail address: `crasta@mat.uniroma1.it`

(Ilaria Fragalà) DIPARTIMENTO DI MATEMATICA, POLITECNICO, PIAZZA LEONARDO DA VINCI, 32 – 20133 MILANO (ITALY)

E-mail address: `ilaria.fragala@polimi.it`