



Recent Stressful Life Events in Euthymic Major Depressive Disorder Patients: Sociodemographic and Clinical Characteristics

Gianluca Serafini^{1,2*}, Xenia Gonda^{3,4,5}, Giovanna Canepa^{1,2}, Pierre A. Geoffroy^{6,7}, Maurizio Pompili⁸ and Mario Amore^{1,2}

¹ Section of Psychiatry, Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genoa, Genoa, Italy, ² IRCCS Ospedale Policlinico San Martino, Genoa, Italy, ³ Department of Psychiatry and Psychotherapy, St. Rokus Clinical Center, Semmelweis University, Budapest, Hungary, ⁴ MTA-SE Neuropsychopharmacology Research Group, Hungarian Academy of Sciences and Semmelweis University, Budapest, Hungary, ⁵ NAP-2-SE New Antidepressant Target Research Group, Hungarian Brain Research Programme, Semmelweis University, Budapest, Hungary, ⁶ Department of Psychiatry and Addiction Medicine, AP-HP, Hôpital Bichat-Claude Bernard, Paris, France, ⁷ NeuroDiderot, Inserm, Paris University, Paris, France, ⁸ Department of Neurosciences, Suicide Prevention Center, Sant'Andrea Hospital, University of Rome, Rome, Italy

OPEN ACCESS

Edited by:

Manit Srisurapanont,
Chiang Mai University, Thailand

Reviewed by:

George P. Chrousos,
National and Kapodistrian University of
Athens, Greece
Liliana Dell'Osso,
University of Pisa, Italy

*Correspondence:

Gianluca Serafini
gianluca.serafini@unige.it

Specialty section:

This article was submitted to
Mood and Anxiety Disorders,
a section of the journal
Frontiers in Psychiatry

Received: 26 May 2020

Accepted: 17 August 2020

Published: 11 September 2020

Citation:

Serafini G, Gonda X, Canepa G,
Geoffroy PA, Pompili M and Amore M
(2020) Recent Stressful Life Events in
Euthymic Major Depressive Disorder
Patients: Sociodemographic and
Clinical Characteristics.
Front. Psychiatry 11:566017.
doi: 10.3389/fpsy.2020.566017

Background: Stressful life events (SLE) may influence the illness course and outcome. This study aimed to characterize socio-demographic and clinical features of euthymic major depressive disorder (MDD) outpatients with SLE compared with those without.

Methods: The present sample included 628 (mean age=55.1 ± 16.1) currently euthymic MDD outpatients of whom 250 (39.8%) reported SLE and 378 (60.2%) did not.

Results: After univariate analyses, outpatients with SLE were most frequently widowed and lived predominantly with friends/others. Moreover, relative to outpatients without SLE, those with SLE were more likely to have a family history of suicidal behavior, manifested melancholic features, report a higher Coping Orientation to the Problems Experienced (COPE) positive reinterpretation/growth and less likely to have a comorbid panic disorder, residual interepisodic symptoms, use previous psychiatric medications, and currently use of antidepressants. Having a family history of suicide (OR=9.697; $p \leq .05$), history of psychotropic medications use (OR=2.888; $p \leq .05$), and reduced use of antidepressants (OR=.321; $p = .001$) were significantly associated with SLE after regression analyses. Mediation analyses showed that the association between current use of antidepressants and SLE was mediated by previous psychiatric medications.

Conclusion: Having a family history of suicide, history of psychotropic medications use, and reduced use of antidepressants is linked to a specific “at risk” profile characterized by the enhanced vulnerability to experience SLE.

Keywords: negative distressing/stressful life events, family history of suicide, major depressive disorder, previous psychiatric medications, antidepressant medications

BACKGROUND

Stressful life events (SLE) may be defined as: “environmental events or chronic conditions that objectively threaten the physical and/or psychological health or well-being of individuals of a particular age in a particular society” (1). Life events may be described as social and environmental occurrences leading to psychophysiological modification in the general population during the time course (2). Life events are considered positive or negative according to the individual’s subjective experience. Negative life events, which may have a prominent role in the development and clinical course of psychiatric disorders may be identified in various social/environmental contexts (2) and may be commonly defined as: 1) maltreatment and violence; 2) loss events; 3) intrafamilial problems; 4) school and interpersonal problems (3, 4). These type of life events are usually associated with undesirable affective experiences exerting deleterious effects on the individual mental health and social/environmental adaptation (4). Generally, SLE such as perceived loss, humiliation, entrapment, and perceived danger seem to provide interesting information about the course and outcome of psychiatric conditions (5). Notably, certain types of SLE such as involuntary occupation loss and marital separation may enhance suicide risk (6, 7), and influence the time to remission of major depressive disorder (MDD) (8) as well as substance use (9).

Based on stress vulnerability models, having experienced SLE is generally linked to poor mental health (10) with the development of major psychiatric conditions generally associated with the exposure to SLE (11). Having experienced more proximal life personal and situational factors may also enhance suicide risk (12) with the cumulative effect of multiple life events playing a more relevant effect than that of specific single life events (13).

While depression has a moderate heritability, environmental events account for approximately 60% of the variation in depression (14) on a general population level depending on the severity of the illness, with 80% of depressive episodes preceded by a major stress exposure (15) and etiologically relevant distal and proximal stressors occurring quite frequently, about at least once every 3–4 years.

SLE are able to strongly predict both the onset (16) and recurrence (17) of depressive episodes, including suicidal behavior. Evidence also suggested that negative life adversities are important risk factors particularly for depressed subjects with a negative cognitive style (18). This hypothesis postulated that exposure to SLE may determine sensitization or kindling and may enhance the individual vulnerability to manifest depressive episodes, although the patient is not directly experiencing a relevant psychosocial stressor. Socio-demographic and clinical variables such as gender potentially mediate the relation between SLE and depression. For instance, when compared to males, females seem to be more vulnerable to the adverse effects of psychosocial stressors (19). Studies (19, 20) stressed the link between the personality trait “neuroticism,” which increases the likelihood of maladaptive responses and coping in the face of negative life events and increased sensitivity to stress and severity

of MDD. Importantly, Kendler and colleagues (20) found that, in a sample of more than 7500 twins, higher neuroticism enhanced the likelihood of depressive episodes in response to SLE.

The above indicates that SLE should be taken into consideration in affective disorder patients due to their impact on illness course and outcomes including suicide. Thus, the present study aimed to compare socio-demographic/clinical characteristics (including marital and living level, family history of psychiatric conditions and/or suicidal behavior, melancholic characteristics in the past, residual interepisodic symptoms, psychiatric medications in the past, current antidepressant medications use, and specific psychometric measures) of euthymic MDD outpatients experiencing significant SLE in the previous 6 months and those who did not, focusing on specific features that may explain the differential burden of disease directly linked to having or not experienced recent SLE. Our aim was also to test whether the effect of current antidepressants use on SLE was mediated by specific variables (i.e., previous psychiatric medications, family history of suicide). Here, we mainly hypothesize that previous psychiatric medications have a significant direct/indirect effect on SLE.

METHODS

Sample

The present sample is composed of 628 currently euthymic MDD outpatients with age ranging from 18 to 85 years (mean age = 55.1 ± 16.1). Clinically, euthymia was defined using a Montgomery–Asberg Depression Rating Scale (MADRS) (21) score of <10 . Our sample included predominantly single MDD outpatients (52.9%) and, in a relatively smaller proportion, recurrent MDD outpatients (47.1%). Participants are all consecutive euthymic unipolar outpatients receiving only maintenance treatment at the time of evaluation. Overall, 488 subjects (77.7%) of our sample were treated with antidepressant medications and 140 individuals (22.3%) received only psychotherapy as maintenance. In particular, psychopharmacological treatments and psychopathological conditions were stable for at least 6 months before assessment. All euthymic MDD outpatients were recruited from our catchment area and have been followed/treated by our university outpatient service for at least 12 months. In addition, they were regularly followed by our local psychiatric services.

Measures and Study Design

An investigator (GC) who was adequately trained to improve the interrater reliability administered to participants the following psychometric instruments: 1) Clinical Global Impression (CGI) (22); 2) Beck Hopelessness Scale (BHS) (23); 3) Hamilton Anxiety Rating Scale (HARS) (24); 4) Global Assessment of Functioning Scale (GAF) (American Psychiatric Association, 1987) (25); 5) Childhood Trauma Questionnaire (CTQ) (26); 6) Coping Orientation to the Problems Experienced (COPE) (27); 7) Toronto Alexithymia Scale (TAS-20) (28) to collect other relevant clinical information.

All clinical information were retraced, as specified, by using clinical files and lifetime computerized medical records. All outpatients accepted voluntarily to participate in the present study and provided their informed consent to participate. The local Ethical Review Board regularly approved the specified study design.

Ascertainment of Recent Stressful Life Events

Significant stressful/distressing life-events in the previous 6 months were collected based on self-report and described as follows: 1) maltreatment and violence (sexual/physical/emotional abuse, emotional/physical neglect, witnessing home/community aggression); 2) loss events (separations, death of a parent or close friend); 3) intrafamilial problems (parental divorce, family instability, social or economic problems); 4) school and interpersonal problems (failure of grade in school or in exam, breaking up with a close friend, and poor social relationships) (for more details, see 3 and 4).

Procedure and Data Collection

All MDD outpatients were recruited at the Department of Neuroscience (DINOEMI), University of Genoa, outpatient service, between July 2014 and February 2020. We used the following inclusion criteria: 1) a diagnosis of single/recurrent MDD; 2) euthymia; 3) a current age of >18 years. Depressive symptoms related to MDD were classified based on the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5) (29). First, an investigator (GC) examined clinical records to retrace subjective histories and clinical variables; then, two senior authors (GS, MA) systematically and independently verified data by using the Mini International Neuropsychiatric Interview (MINI) (30) updated to map to DSM-5 (29). Direct interviews with patients/family members as well as investigation of existing medical records allow the detailed collection of clinical information regarding affective episodes prior the patients' recruitment in our outpatient service. Specifically, we performed a comprehensive data collection using the following variables: 1) socio-demographic information; 2) individual data (e.g., personal autonomy, lifetime substance abuse/dependence, use of psychiatric medications and psychotherapy in the past); 3) positive history of clinical conditions/negative outcome in family (e.g., psychiatric disorders and suicidal behavior); 4) comorbid conditions (e.g., medical/psychiatric disorders); 5) illness course features (e.g., manifested melancholic features and/or current atypical depressive characteristics, seasonality, psychotic symptoms at first episode, single/recurrent episode, residual interepisodic symptoms, illness duration and untreated illness, current illness episode, number of previous affective episodes, age of illness onset, age at first treatment and first hospitalization, and lifetime suicide attempts).

Exclusion criteria were as follows: 1) all clinical conditions that may influence the ability to complete the evaluations including conditions such as Alzheimer's disease, delirium, or severe neurological disorders like intellectual disability; 3) denial of informed consent, 4) a history of active abuse/dependence

(throughout the previous 6 months). Lifetime substance use emerging during mental examination, was not considered a specific exclusion criterion. Intellectual disability was investigated using the Wechsler Adult Intelligence Scale (31).

Statistical Analyses

Subjects were categorized based on the presence/absence of SLE in the previous 6 months and divided into two groups similarly to existing published studies (32) outpatients with recent SLE and 2) outpatients without recent SLE. Data for categorical variables were analyzed using Student's t-tests and Pearson chi-square/Fisher's exact test in contingency tables (χ^2). The Kolmogorov-Smirnov test was performed to confirm whether all the investigated variables in the sample followed the normal distribution. Significance was set at $P \leq 0.05$ (two-tailed).

A binary logistic regression analysis considering specific socio-demographic characteristics (marital status that was dichotomized in married/unmarried status and living status that was dichotomized in living alone or not alone) together with other variables such as family history of suicidal behavior, comorbid panic disorder, previous psychiatric medications, manifested melancholic features in the past, interepisodic residual symptoms, current antidepressants use, and COPE positive reinterpretation and growth was carried out to investigate the contribution of these characteristics in predicting presence/absence of SLE.

Multicollinearity and extreme cases were excluded, and the normality of residuals (histograms and P-P plots) was appropriately checked.


To assess whether the effect of current antidepressants use on SLE was mediated by specific variables (i.e., previous psychiatric medications, family history of suicide), single-mediator models were tested according to the strategy recommended by Preacher and Hayes (33). In a single-mediator model, an independent variable (X = current antidepressants use) is hypothesized to act on the outcome variable (Y = SLE) in two ways: X change a mediator (e.g., M_i = previous psychiatric medications or family history of suicide; path A_i) that, in turn, changes an outcome variable (Y ; path B_i), or X changes Y directly (path C') (for more details, see **Figure 1**).

The Statistical Package for Social Sciences (SPSS) for Windows 21.0 was used to carry out all the statistical analyses.

RESULTS

Socio-Demographic and Clinical Variables of Outpatients With Significant Stressful Life Events in the Previous 6 Months Compared With Those Without

The sample includes 628 euthymic MDD outpatients who were consecutively and voluntarily recruited at the outpatient service of Department of Neuroscience (DINOEMI), University of Genoa from the catchment area. Socio-demographic and clinical features of outpatients are summarized in **Table 1**.



Frequency of Recent Stressful Life Events and Associated Socio-Demographic and Clinical Characteristics

Among subjects who were recruited, 250 (39.8%) reported significant SLE in the previous 6 months and 378 (60.2%) did not.

Those reporting recent SLE [87 males (36.7%) and 163 females (41.7%)] differed from individuals who did not report recent SLE. Specifically, when compared with those who did not report recent SLE subjects with recent SLE differed in terms of both marital and living status ($\chi^2_{(3)}=12.963$, $p=.005$; $\chi^2_{(3)}=9.729$, $p \leq .05$), respectively.

Relative to outpatients without recent stressful life events, those who have experienced these events were more likely to have a family history of suicidal behavior (77.8 vs. 22.2%) ($\chi^2_{(1)}=5.818$, $p \leq .05$), less likely to have a comorbid panic disorder (23.3 vs. 76.7%) ($\chi^2_{(1)}=3.595$, $p \leq .05$) and less likely to have used psychiatric medications (42.2 vs. 57.8%) ($\chi^2_{(1)}=6.000$, $p \leq .01$).

When compared to subjects without SLE in the previous 6 months, subjects with recent stressful events were also more likely to have exhibited melancholic characteristics in the past (61.3 vs. 38.7%) ($\chi^2_{(1)}=6.538$, $p=.01$) and less likely to manifest residual interepisodic symptoms (43.4 vs. 56.6%) ($\chi^2_{(1)}=3.024$, $p \leq .05$) and use current antidepressant medications (36.7 vs. 63.3%) ($\chi^2_{(1)}=9.273$, $p \leq .005$).

The Kolmogorov–Smirnov test did not find any significant abnormality concerning the continuous data. Moreover, subjects with recent SLE were more likely to have a higher COPE positive reinterpretation and growth (11.2 ± 5.9 vs. 10.3 ± 2.5 , $t_{457} = 9.422$, $p \leq .05$) when compared to those without SLE in the previous 6 months (for more details, see **Tables 1** and **2**).

Multivariate Regression Analyses Including Stressful Life Events as Dependent Variable in the Total Sample

All clinical variables which were significant at the univariate analyses were entered into a binary logistic regression analysis

with SLE as dependent variable in order to identify independent factors which resulted associated with. Specific socio-demographic characteristics such as marital and living status as well as clinical variables (e.g., family history of suicidal behavior, comorbid panic disorder, previous psychiatric medications, manifested melancholic features in the past, interepisodic residual symptoms, current antidepressants use, and COPE positive reinterpretation and growth) were introduced into the regression model. The amount of variation in the first dependent variable (presence/absence of recent SLE) that was accounted for all predictors (R^2 -value) was 18.3% ($p \leq .001$) (see **Table 3**).

Having a family history of suicide with an OR of 9.697 ($p \leq .05$), history of psychotropic medications use with an OR of 2.888 ($p \leq .01$), and reduced use of antidepressant drugs with an OR of .321 ($p=.001$) were all significantly associated with recent SLE.

Mediation Analyses

In a regression model with current antidepressants use and family history of suicide as predictors and SLE as criterion, the regression models were both significant ($R^2 = .020$, $B = -.590$, $SE=.195$, $Wald = 9.145$, $p = .002$, $OR = .554$) ($R^2 = .020$, $B = 1.738$, $SE=.809$, $Wald = 4.620$, $p = .032$, $OR = 5.687$), respectively. When considering the effect of current antidepressants use on SLE in a single mediator model (considering first family history of suicide and later previous psychiatric medications as possible mediators), the results were as follows.

Family history of suicide had a significant direct effect on SLE (family history of suicide: omnibus test of direct effect of M_i on Y : $R^2 = .020$, $B = 1.738$, $SE=.809$, $Wald = 4.620$, $p = .032$, $OR = 5.687$) but a nonsignificant indirect effect (family history of suicide: omnibus test of direct effect of X on M_i : $R^2 = .000$, $B = -.012$, $SE=.811$, $Wald = .000$, $p = .988$, $OR = .988$). Previous psychiatric medications had a significant direct effect on SLE

TABLE 1 | Socio-demographic and clinical variables (categorical and quantitative) in patients with significant distressing life-events in the last 6 months (N=250) compared with those without (N=378).

Variables	Subjects with significant distressing life events in the previous 6 months (N=250)		Subjects without significant distressing life events in the previous 6 months (N=378)		Statistic (χ^2)	*p
	N	%	N	%		
Gender						
Male	87	36.7	150	63.3	$\chi^2_{(1)}=1.527$.125*
Female	163	41.7	228	58.3		
Marital status						
Single	62	33.7	122	66.3	$\chi^2_{(3)}=12.963$.005
Married	116	38.5	185	61.5		
Divorced	35	43.8	45	56.3		
Widowed	34	59.6	23	40.4		
Living status						
Alone	66	42.9	88	57.1	$\chi^2_{(3)}=9.729$.021
With family	163	37.0	278	63.0		
With friends	9	64.3	5	35.7		
With others	6	75.0	2	25.0		
Working status						
Employed	113	40.6	165	59.4	$\chi^2_{(3)}=3.635$.304
Unemployed	59	45.7	70	54.3		
Retired	65	36.5	113	63.5		
Student	8	30.8	18	69.2		
Socio-economic level						
Below average	76	39.6	116	60.4	$\chi^2_{(2)}=.659$.719
Average	154	40.6	225	59.4		
Above average	14	34.1	27	65.9		
Positive history of psychiatric conditions in family						
No	163	38.3	263	61.7	$\chi^2_{(1)}=.367$.302*
Yes	78	40.8	113	59.2		
Family history of suicidal behavior						
No	144	38.1	234	61.9	$\chi^2_{(1)}=5.818$.020*
Yes	7	77.8	2	22.2		
Psychiatric comorbidity						
No	197	40.3	292	59.7	$\chi^2_{(1)}=.077$.431*
Yes	53	39.0	83	61.0		
Comorbid anxiety						
No	234	39.5	359	60.5	$\chi^2_{(1)}=.747$.240*
Yes	16	47.1	18	52.9		
Comorbid panic disorder						
No	243	40.7	354	59.3	$\chi^2_{(1)}=3.595$.041*
Yes	7	23.3	23	76.7		
Comorbid phobias						
No	247	39.6	376	60.4	$\chi^2_{(1)}=2.072$.177*
Yes	3	75	1	25		
Comorbid obsessive compulsive disorder						
No	250	40.2	372	59.8	$\chi^2_{(1)}=3.342$.078*
Yes	0	0.0	5	100		
Comorbid post-traumatic stress disorder						
No	250	39.9	376	60.1	$\chi^2_{(1)}=.664$.601*
Yes	0	0.0	1	100.0		
Comorbid personality disorder, cluster A						
No	250	40	375	60.0	$\chi^2_{(1)}=1.331$.361*
Yes	0	0.0	2	100.0		
Comorbid personality disorder, cluster B						
No	241	39.7	366	60.3	$\chi^2_{(1)}=.227$.399*
Yes	9	45.0	11	55.0		
Comorbid personality disorder, cluster C						
No	247	39.7	375	60.3	$\chi^2_{(1)}=.852$.315*
Yes	3	60.0	2	40.0		

(Continued)

TABLE 1 | Continued

Variables	Subjects with significant distressing life events in the previous 6 months (N=250)		Subjects without significant distressing life events in the previous 6 months (N=378)		Statistic (χ^2)	♦p
	N	%	N	%		
Psychiatric medications in the past						
No	36	30.0	84	70.0	$\chi^2_{(1)}=6.000$.009*
Yes	213	42.2	292	57.8		
Non psychiatric medications in the past						
No	121	41.0	174	59.0	$\chi^2_{(1)}=.379$.297*
Yes	127	38.6	202	61.4		
Lifetime substance abuse/dependence						
No	237	39.8	359	60.2	$\chi^2_{(1)}=.003$.563*
Yes	11	39.3	17	60.7		
Psychotherapy in the past						
No	203	39.3	313	60.7	$\chi^2_{(1)}=.002$.523*
Yes	40	39.6	61	60.4		
Single/recurrent episode						
Single	129	41.5	182	58.5	$\chi^2_{(1)}=.842$.203*
Recurrent	93	37.7	154	62.3		
First depressive episode						
Depressive	9	23.1	30	72.9	$\chi^2_{(1)}=.237$.414*
Nondepressive	9	28.1	23	71.9		
Current catatonic characteristics						
No	159	39.7	242	60.3	$\chi^2_{(1)}=1.962$.222*
Yes	0	0.0	3	100.0		
Current melancholic characteristics						
No	141	37.9	231	62.1	$\chi^2_{(1)}=6.538$.010*
Yes	19	61.3	12	38.7		
Current atypical depressive characteristics						
No	153	40.3	227	59.7	$\chi^2_{(1)}=.875$.239*
Yes	7	30.4	16	69.6		
Psychotic symptoms at first episode						
No	140	40.2	208	59.8	$\chi^2_{(1)}=1.573$.176*
Yes	6	60.0	4	40		
BMI						
<18	10	50.0	10	50.0	$\chi^2_{(3)}=4.977$.173
18–25	191	41.7	267	58.3		
25–30	39	32.0	83	68.0		
>30	9	34.6	17	65.4		
Residual interepisodic symptoms						
No	116	36.4	203	63.6	$\chi^2_{(1)}=3.024$.049*
Yes	118	43.4	154	56.6		
Lifetime suicide attempts						
None	240	40.3	183	92.9	$\chi^2_{(2)}=2.292$.318
0–3	7	25.9	13	6.6		
>3	1	33.3	0	0.0		
Seasonality						
No	144	39.0	225	61.0	$\chi^2_{(1)}=.649$.340
Yes	2	25.0	6	75.0		
Adherence to treatment						
Bad	11	36.7	19	63.3	$\chi^2_{(1)}=.43$.500*
Good	132	38.6	210	61.4		
Current anxyolytic medications						
None	96	41.6	135	58.4	$\chi^2_{(3)}=.641$.887
Short half-life	75	38.9	118	61.1		
Middle half-life	57	37.7	94	62.3		
High half-life	19	40.4	28	59.6		
Current antidepressant medications						
No	70	51.1	67	48.9	$\chi^2_{(1)}=9.273$.002*
Yes	179	36.7	309	63.3		

(Continued)

TABLE 1 | Continued

Variables	Subjects with significant distressing life events in the previous 6 months (N=250)		Subjects without significant distressing life events in the previous 6 months (N=378)		Statistic (χ^2)	♦p
	N	%	N	%		
Personal autonomy						
Bad	18	52.9	140	38.3	$\chi^2_{(1)}=2.809$.069*
Good	16	47.1	226	61.7		
	Subjects with significant distressing life-events in the previous 6 months (N=250)		Subjects without significant distressing life-events in the previous 6 months (N=378)		Statistic (Student's t-test)	p
	Mean	SD	Mean	SD		
Current age	55.1	15.7	54.8	16.4	$T_{626}=-.605$.841
Educational level (years)	11.3	3.5	11.6	3.5	$T_{583}=.426$.332
Number of illness episodes	1.8	1.4	1.9	1.5	$T_{499} = 1.202$.549
Age at first treatment	47.8	16.7	45.2	16.6	$T_{550}=.055$.084
Duration of untreated illness (years)	3.7	26.4	1.7	3.3	$T_{468} = 4.582$.223
Duration of illness (years)	9.6	12.9	11.8	14.2	$T_{528} = 2.455$.070
Duration of substance abuse (years)	0.4	2.5	0.4	2.4	$T_{609}=-.017$.962
MADRS total score	6.5	3.3	6.2	3.5	$T_{203} = 1.686$.530
CGI severity (item 1)	2.1	4.1	1.4	2.5	$T_{174} = 5.844$.139
BHS total score	10.7	6.2	11.03	5.8	$T_{429}=.619$.545
HARS total score	7.9	8.4	8.2	7.5	$T_{174}=.864$.791
GAF Mental Composite Score	11.1	2.5	11.6	4.2	$T_{463}=.560$.161
GAF Physical Composite Score	14.9	1.9	14.8	1.8	$T_{450} = 1.732$.790
CTQ emotional abuse	3.2	5.03	3.3	4.7	$T_{422}=.149$.842
CTQ physical abuse	1.5	3.2	1.6	4.01	$T_{440}=.303$.800
CTQ sexual abuse	1.7	4.1	1.2	3.5	$T_{431} = 7.205$.177
CTQ emotional neglect	13.1	7.1	14.04	6.7	$T_{425}=.775$.188
CTQ physical neglect	5.3	3.4	5.2	3.5	$T_{119}=.311$.917
COPE active	11.2	5.9	10.3	2.5	$T_{457}=.9422$.036
COPE planning	10.6	4.6	10.5	3.6	$T_{445}=.908$.877
COPE suppression	10.1	5.1	9.9	3.4	$T_{453} = 2.102$.643
COPE behavioral disengagement	8.8	3.6	9.1	4.2	$T_{449}=.101$.477
COPE denial	7.7	4.4	7.3	2.4	$T_{450} = 3.111$.147
COPE mental disengagement	8.9	2.6	9.1	4.2	$T_{446}=.082$.553
COPE substance abuse	5.02	2.3	5.3	3.1	$T_{460} = 2.211$.286
COPE emotional social support	4.4	72.6	9.7	3.9	$T_{454} = 2.983$.234
COPE instrumental social support	10.5	4.6	10.3	3.6	$T_{466}=.699$.727
COPE venting emotions	10.7	4.7	10.2	3.3	$T_{443} = 1.384$.138
COPE positive reinterpretations	10.1	4.3	10.2	3.8	$T_{468}=.603$.904
COPE restraint	9.7	5.1	10.3	4.8	$T_{445}=.066$.188
COPE acceptance	9.9	3.6	10.1	4.8	$T_{447}=.048$.652
COPE turning to religion	8.8	5.3	8.6	4.5	$T_{460}=.142$.670
COPE humor	6.7	2.9	6.7	3.02	$T_{433}=.045$.879
TAS total score	63.2	13.6	63.8	18.6	$T_{370} = 1.237$.710
TAS difficulties in identifying feelings	17.4	7.1	17.5	8.9	$T_{419}=.090$.874
TAS difficulties in communicating feelings to others	15.3	4.4	15.4	5.7	$T_{425}=.020$.833
TAS thoughts oriented to external context	27.7	5.4	28.6	9.2	$T_{431} = 3.091$.246
DAI total score	12.9	4.3	14.03	4.8	$T_{277}=.897$.066

BHS, Beck Hopelessness Scale; COPE, Coping Orientation to the Problems Experienced; CTQ, Childhood Trauma Questionnaire; DAI, Drug Attitude Inventory; HARS, Hamilton Anxiety Rating Scale; MADRS, Montgomery-Asberg Depression Rating Scale; TAS, Toronto Alexithymia Scale; *Fisher's exact test (significance two-tailed); percentages were calculated per column.

(previous psychiatric medications: omnibus test of direct effect of Mi on Y: $R^2 = .013$, $B = .532$, $SE=.219$, $Wald = 5.917$, $p = .015$, $OR = 1.702$) and a significant indirect effect (previous psychiatric medications: omnibus test of direct effect of X on Mi: $R^2 = .121$, $B = 1.561$, $SE=.219$, $Wald = 50.777$, $p = <.001$, $OR = 4.765$).

DISCUSSION

Our study investigating the association of recent SLE in euthymic depressive patients indicated that the experience of recent stressful events was associated with several clinical, course, and sociodemographic characteristics, some of which may be clinically

TABLE 2 | Medical comorbidity in patients with significant distressing life-events in the previous 6 months (N=250) compared with those without (N=378).

	Subjects with significant distressing life-events in the previous 6 months (N=250)		Subjects without significant distressing life-events in the previous 6 months (N=378)		Statistic (χ^2)	p
	N	%	N	%		
Medical comorbidities						
No	143	65.0	77	35.0	$\chi^2_{(1)}=3.193$.064*
Yes	233	57.7	171	42.3		
Neurological comorbid disorders						
No	358	60.6	233	39.4	$\chi^2_{(1)}=.475$.304*
Yes	18	54.5	15	45.5		
Cardiological comorbid disorders						
No	175	89.3	170	95.0	$\chi^2_{(1)}=.001$.555*
Yes	21	10.7	9	5.0		
Endocrinological comorbid disorders						
No	341	59.5	232	40.5	$\chi^2_{(1)}=1.625$.130*
Yes	35	68.6	16	31.4		
Inflammatory/immunological comorbid disorders						
No	342	59.8	230	40.2	$\chi^2_{(1)}=.623$.262*
Yes	34	65.4	18	34.6		
Metabolic disorders						
No	366	60.4	240	39.6	$\chi^2_{(1)}=.171$.427*
Yes	10	55.6	8	44.4		
Mild cognitive impairment (amnesic)						
No	196	100.0	178	99.4	$\chi^2_{(1)}=1.098$.477*
Yes	0	0.0	1	0.6		
Comorbid polmonary diseases						
No	194	99.0	179	100.0	$\chi^2_{(1)}=1.836$.273*
Yes	2	1.0	0	0.0		
Comorbid cancer						
No	189	96.4	169	94.4	$\chi^2_{(1)}=.878$.246*
Yes	7	3.6	10	5.6		
Comorbid chronic renal failure						
No	196	100.0	178	99.4	$\chi^2_{(1)}=1.098$.477*
Yes	0	0.0	1	0.6		
Comorbid HIV						
No	194	99.0	179	100.0	$\chi^2_{(1)}=1.836$.273*
Yes	1	1.0	1	0.0		

*HIV, human immunodeficiency virus; Fisher's exact test (significance two-tailed); percentages were calculated per column.

TABLE 3 | Multiple regression model of significant distressing life-events in the previous 6 months in the total sample (N=628).

Variable	B	SE	Wald	P	OR	95% CI OR	
						Lower	Upper
Married status	.021	.276	.006	.940	1.021	.595	1.754
Living status	-.010	.010	1.136	.286	.990	.971	1.009
Family history of suicide	2.272	1.146	3.931	.047	9.697	1.026	91.629
Comorbid panic disorders	-1.022	.681	2.248	.134	.360	.095	1.369
Previous psychiatric medications	1.061	.406	6.808	.009	2.888	1.302	6.407
Melancholic features in the past	.864	.592	2.129	.145	2.373	.743	7.576
Interepisodic residual symptoms	.391	.290	1.819	.177	1.479	.838	2.610
Current antidepressants use	-1.136	.349	10.627	.001	.321	.162	.636
COPE active	.065	.037	3.054	.081	1.067	9.92	1.147
Constant	-2.814	.910	9.572	.002	.060		

Dependent variable: SLE. All predictors were entered in one block (hierarchical method). Model summary: R=.113, R²=.153; significance: P=≤.001. Bold values denote statistical significance.

utilizable in understanding and managing the culminating effect of life stressors in influencing illness course and outcome.

Socio-Demographic Characteristics of Outpatients With Significant Stressful Life Events in the Previous 6 Months Compared With Those Without

Euthymic outpatients who experienced SLE differed from those who did not experience SLE in terms of both marital and living status.

Existing evidence suggested that being widowed is generally perceived as one of the most SLE (34) linked to relevant impairments in the general well-being of the bereaved spouse. According to previously published studies (35), widowers and widows should be considered at higher risk to experience lower subjective well-being even several years after the death of the spouse. In line with these findings, Carnelley and colleagues (36) found that widowers/widows experienced a significant distress and reduced subjective wellbeing even decades after the death. Similarly, we found a higher ratio of those widowed among those experiencing recent distressing events.

On the other hand, conversely to existing evidence suggesting that subjects who have more contact with friends and family are happier than those who have less contact (37), we found that outpatients who experienced SLE lived predominantly with friends or others. However, living with family rather than living with friends may be generally interpreted as a protective factor for MDD patients. This discrepancy could be related, in our opinion, to the fact that, although being recruited in stable psychopathological conditions, participants of the present study are predominantly affected by MDD (single episode) which is a condition for which many of them might have searched a significant social contact or asked support due to their illness.

Clinical Characteristics of Euthymic Outpatients Experiencing Significant Stressful Life Events in the Previous 6 Months Compared With Those Without

As for clinical variables, outpatients experiencing significant SLE were more likely to have a family history of suicidal behavior, manifested melancholic features in the past, report a higher COPE positive reinterpretation and growth, and less likely to have a comorbid panic disorder, residual interepisodic symptoms, past psychiatric medication use, and current antidepressant medication use.

In line with previous evidence (38) suggesting that the occurrence of SLE was higher and the associated perceived experience was more deleterious in suicidal patients, we found an association between SLE in the previous six months and family history of suicidal behavior. Specifically, this could be explained by the assumption that the interpersonal pattern linked to family history of suicide may consistently influence the ability to maintain stable and significant relations with others as supported by Rajalin and colleagues (39).

Moreover, we found that subjects with recent SLE were more likely to have manifested melancholic features in the past. Existing studies analyzed the association between depression subtypes and

the likelihood to experience life adversities. For instance, Kohn and colleagues (40) reported that, in a sample of 115 MDD patients, subjects with nonmelancholic depression or first depressive episode more frequently experienced ≥ 3 life adversities when compared to severe, melancholic, or recurrent depression. However, they investigated only a subset of MDD patients which is more prone to suffer from a cluster of negative life events. In addition, while the study of Kohn and colleagues (40) analyzed a sample of patients with mild-moderate depression, we investigated a larger (N=628) sample of MDD euthymic subjects. Thus, the main differences between our study and the study of Kohn and colleagues (40) may be related to the different study sample selection and different clinical characteristics.

Finally, we found that those who experienced SLE were more likely to report higher COPE positive reinterpretation and growth. This may be explained by the fact that the attitudes to cope with stress in our MDD patients, the psychopathological conditions of which were stable for at least 6 months before assessment, were more pronounced than those of other patient populations. Clinicians and health professionals should take appropriately into account these data in order to conduct psychotherapeutic interventions which are directed toward further coping with stress and create treatment strategies which may result especially useful in improving functional capacity in the identified stable phases of MDD.

Predictors of Stressful Life Events and Mediators of the Associations With Stressful Life Events

After binary regression analysis, MDD patients with recent SLE were more likely to have a family history of suicide (OR=9.68) and a history of psychotropic medications use (OR=2.89), but they were less likely to use currently antidepressants (OR=.32). In particular, outpatients who have experienced recent SLE were more likely to report a family history of suicide, more likely to have used previous psychiatric medications, and less likely to currently use antidepressants drugs than those without recent SLE in the previous 6 months.

Generally, having a family history of suicidal behavior is also associated with a greater propensity to SLE; thus, clinicians should be warned about carefully searching for suicidal thoughts in this patient subgroup as these individuals might exhibit a worse combination of biological, psychological, and environmental risk for suicide. In particular, in individuals with family history of suicidal behavior, the increased likelihood to experience stressful life events may enhance the perception of fewer reasons for living, thus elevating the risk of suicidal behavior, although in our study we did not observe significant differences in terms of suicide attempts between those with and without recent stressful life events. Consistent with this hypothesis, evidence documented that having a family history of suicide or attempted suicide is a significant risk factor for suicidal behavior and supposed the existence of a specific genetic component of suicidal behavior (41, 42). In line with these results, Rajalin and colleagues (39), recently found in a sample of 181 suicide attempters that having a family history of suicide was linked to factor 2 (nonassertive; exploitable; overly nurturant, and

intrusive), in particular with Intrusive scale scores of the Inventory of Interpersonal Problems (IIP). Subjects with higher scores in the Intrusive subscale of the IIP are described to be excessively controlling, not careful to respect individual and other's boundaries as well as more prone to take charge of other subjects' problems (39). This may imply that these individuals pervasively experience disappointment and presumably reject in social contexts appearing unable to identify themselves in positive arguments with others. According to these findings, subjects who have experienced recent SLE and report a positive history of suicide among family members seem to exhibit specific deficits in interpersonal functioning. Moreover, evidence (43) clearly suggested that having a family history of suicide may be considered not only as a biological but also a psychological risk factor for familial transmission of suicide. Unfortunately, in our study we were not able to investigate the impact of interpersonal difficulties in subjects with family history of suicidal behavior who have also experienced recent SLE. In addition to the identified at risk profile, the participants of our study were also more likely to be widowed and have used psychiatric medications in the past.

Surprisingly, after multivariate analysis, subjects with significant SLE were less likely to currently use antidepressant drugs compared to those without recent stressful life events. This finding may be presumably explained by the fact that the majority of our study participants suffer predominantly from reactive and situational depression, but it is also possible that individuals who experienced recent SLE in our study may rely on more resilient social groups in which they received social support even in the context of negative interpersonal events or may have benefited from a psychotherapeutic approach. Such support may have helped them to better cope with SLE, thus reducing the current use of antidepressant medications. However, most patients do not frequently consult psychiatrists or mental health professionals due to stigma and discrimination as well as fear of disability and chronicity related to psychiatric conditions. This may, at least partially, justify the reduced or absent psychiatric services attendance in the past.

Finally, according to mediation analyses the association between the reduced use of antidepressant drugs and SLE was mediated by the history of psychotropic medications use. Thus, with this result we confirm the initial hypothesis based on which previous psychiatric medications had a significant direct/indirect effect on SLE. Specifically, subjects with a reduced use of antidepressants may be more vulnerable to experience SLE, in particular whether they have been treated in the past with psychiatric medications (that may have been prematurely discontinued by patients). This result is in line with the questioned long-term partial benefits of psychiatric medications (44) and may reflect, in our opinion, the limited potential of these compounds, which are detrimentally interrupted by patients on their own initiative (45, 46), on protecting against long-term exposure to stressors.

Unfortunately, whether the use of pharmacological medications is not combined with other available treatment strategies (e.g., psychotherapy), depressed individuals who have been treated only with psychoactive agents may be insufficiently protected in the long-term period towards potential illness recurrences (47,

48) neither they seem to have developed specific coping strategies to successfully cope with psychosocial distress in daily life. To further confirm our findings, it has been reported that the current use of antidepressants may help subjects to face intense emotional stimuli and stressful cues (49), presumably by attenuating the brain expression and/or actions of pro-inflammatory mediators. Here, we add that having undergone a psychiatric treatment (that may have been prematurely discontinued) is associated with the more frequent occurrence of SLE, if MDD patients are not adequately treated with antidepressant medications.

LIMITATIONS OF THE STUDY

The present study should be considered in the light of the following limitations/shortcomings. First, SLE were not measured using specific psychometric instruments such as the Social Readjustment Rating Scale (SRRS) which assesses the frequency of 43 common stressful life events occurring over the previous 12 months (34), that would allow us to rate the impact of specific stressful life experiences on a specific severity scale. In addition, we were also unable to use specific psychometric tools like the Karolinska Self Harm History Interview (50), including specific questions regarding family history of suicidal behavior in order to investigate the family history of completed suicide in a more detailed manner. In addition, the cross-sectional nature of the present study potentially leading to type I errors needs to be considered as an additional caveat. Moreover, we were not able to collect information regarding positive life events in the present study and thus we mainly explored the impact of negative ones. Additionally, by setting a P value of $\leq .05$, some significant differences in our univariate analyses may lead to false-positive findings due to multiple comparisons. Unfortunately, we were not able to apply an adjustment for multiple comparisons (e.g., Bonferroni's adjustment)

Furthermore, we mainly analyzed a specific list of SLE: maltreatment and violence; loss events; intrafamilial problems; school and interpersonal problems based on our clinical experience as well as according to previously published studies (3, 4) without considering other types of SLE such as specific interpersonal stressors that may influence the onset/maintenance of depression in specific sociocultural environments (39).

Additionally, we did not take into account the possible link between the time from the occurrence of these events and recent SLE. For instance, one can inquire that having experienced the loss of a loved one by suicide even in the past may have influenced the likelihood to experience recent SLE. Although we did not find in our sample any recent loss event directly related to the loss of a loved one by suicide, the association between family history of suicide in the past and SLE may be affected by this caveat. Furthermore, we were not able in this study to investigate the effect of previous SLE, such as childhood traumatic experiences. Moreover, psychological factors which may further enhance the risk of negative outcome such as neuroticism, that were comprehensively examined in existing

studies (19), were not considered in our study. Also, the role of potential protective factors like social support (51) or coping skills (52) were not taken specifically into account. Finally, while previous studies (53) supported a significant dose-response effect of the number of life events on illness course and outcome, here we were not able to identify the exact number of SLE.

However, despite the mentioned shortcomings, the recruitment of a large homogeneous sample of outpatients with MDD who were clearly euthymic together with the investigation of the effects of a comprehensive set of potential socio-demographic and clinical variables on the outcome need to be considered as major strengths of this study.

CONCLUSION

Having a family history of suicide and history of psychotropic medications use but even the reduced use of antidepressant drugs, of which some (e.g., duloxetine) act as inhibitors of both norepinephrine and serotonin transporters (54) and others (e.g., agomelatine) are of particular interest due to their ability to enhance neuroplasticity mechanisms (55), may reflect a specific “at risk” clinical profile characterized by the enhanced vulnerability to experience SLE. This finding is in line with the expression of a complement of genes that correlates with the degree of adversity, based on the assumption that early adversities may predispose selective molecular systems and related gene expression profiles associated with impaired stress response and/or differences in the sensitivity of specific genes involved in the response to stress signals (56, 57). Further additional studies are needed in order to test these exploratory findings.

REFERENCES

- Grant KE, Compas BE, Stuhlmacher AF, Thurm AE, McMahon SD, Halpert JA. Stressors and child and adolescent psychopathology: Moving from markers to mechanisms of risk. *Psychol Bull* (2003) 129:447–66. doi: 10.1037/0033-2909.129.3.447
- Monroe SM, Harkness KL. Life stress, the ‘kindling’ hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychol Rev* (2005) 112:417–45. doi: 10.1037/0033-295X.112.2.417
- Serafini G, Muzio C, Piccinini G, Flouri E, Ferrigno G, Pompili M, et al. Life adversities and suicidal behavior in young individuals: a systematic review. *Eur Child Adolesc Psychiatry* (2015) 24:1423–46. doi: 10.1007/s00787-015-0760-y
- Zhou L, Chen J. Life events and hopelessness depression: The influence of affective experience. *PLoS One* (2017) 12:e0187898. doi: 10.1371/journal.pone.0187898
- Kendler KS, Hettema JM, Butera F, Gardner CO, Prescott CA. Life event dimensions of loss, humiliation, entrapment, and danger in the prediction of onsets of major depression and generalized anxiety. *Arch Gen Psychiatry* (2003) 60:789–96. doi: 10.1001/archpsyc.60.8.789
- Pompili M, Innamorati M, Lamis DA, Erbutto D, Venturini P, Ricci F, et al. The associations among childhood maltreatment, “male depression” and suicide risk in psychiatric patients. *Psychiatry Res* (2014) 220:571–8. doi: 10.1016/j.psychres.2014.07.056
- Kölves K, Ide N, De Leo D. Fluctuations of suicidality in the aftermath of a marital separation: 6-month follow-up observations. *J Affect Disord* (2012) 142:256–63. doi: 10.1016/j.jad.2012.04.036
- Spinhoven P, Elzinga BM, Hovens JG, Roelofs K, van Oppen P, Zitman FG, et al. Positive and negative life events and personality traits in predicting course of depression and anxiety. *Acta Psychiatr Scand* (2011) 124:462–73. doi: 10.1111/j.1600-0447.2011.01753.x
- Gallo WT, Bradley EH, Siegel M, Kasl SV. Health effects of involuntary job loss among older workers: Findings from the health and retirement survey. *J Gerontol B Psychol Sci Soc Sci* (2000) 55B:S131–40. doi: 10.1093/geronb/55.3.s131
- Turner RJ, Lloyd DA. Lifetime traumas and mental health: the significance of cumulative adversity. *J Health Soc Behav* (1995) 36:360–76. doi: 10.2307/2137325
- Mayo D, Corey S, Kelly LH, Yohannes S, Youngquist AL, Stuart BK, et al. The role of trauma and stressful life events among individuals at clinical high risk for psychosis: A review. *Front Psychiatry* (2017) 20:55. doi: 10.3389/fpsy.2017.00055
- Oquendo MA, Perez-Rodriguez MM, Poh E, Sullivan G, Burke AK, Sublette ME, et al. Life events: a complex role in the timing of suicidal behavior among depressed patients. *Mol Psychiatry* (2014) 19:902–9. doi: 10.1038/mp.2013.128
- Furniss T, Beyer T, Müller JM. Impact of life events on child mental health before school entry at age six. *Eur Child Adolesc Psychiatry* (2009) 18:717–24. doi: 10.1007/s00787-009-0013-z
- Rutter M, Silberg J. Gene-environment interplay in relation to emotional and behavioral disturbance. *Annu Rev Psychol* (2002) 53:463–90. doi: 10.1146/annurev.psych.53.100901.135223
- Maciejewski PK, Mazure CM. Stressful life events and depression. *Am J Psychiatry* (2000) 157:1344–5. doi: 10.1176/appi.ajp.157.8.1344-a
- Kendler KS, Karkowski LM, Prescott CA. Stressful life events and major depression: risk period, longterm contextual threat, and diagnostic specificity. *J Nerv Ment Dis* (1988) 186:661–9. doi: 10.1097/00005053-199811000-00001
- Bos EH, Bouhuys AL, Geerts E, van Os TW, Ormel J. Stressful life events as a link between problems in nonverbal communication and recurrence of depression. *J Affect Disord* (2007) 97:161–9. doi: 10.1016/j.jad.2006.06.011

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material; further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Local Ethical Review Board of the University of Genoa, Liguria, Italy. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

GS designed the study and wrote the manuscript. GC managed the literature searches and analyses. XG and PG contributed to write the first draft of the manuscript. MA provided the intellectual impetus and MP supervised the study design and search strategy. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

This work was developed within the framework of the DINOGMI Department of Excellence of MIUR 2018-2022 (law 232/2016). Xenia Gonda was supported by the Janos Bolyai Research Fellowship of the Hungarian Academy of Sciences.

18. Post RM. Transduction of psychosocial stress into the neurobiology of recurrent affective disorder. *Am J Psychiatry* (1992) 149:999–1010. doi: 10.1176/ajp.149.8.999
19. Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex, and stressful life events in the prediction of episodes of major depression. *Am J Psychiatry* (2004) 161:631–6. doi: 10.1176/appi.ajp.161.4.631
20. Kendler KS, Gardner CO. A longitudinal etiologic model for symptoms of anxiety and depression in women. *Psychol Med* (2011) 41:2035–45. doi: 10.1017/S0033291711000225
21. Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry* (1979) 134:382–9. doi: 10.1192/bjp.134.4.382
22. Guy W. *ECDEU Assessment Manual for Psychopharmacology*. Rockville, MD (US): Department of Health, Education, and Welfare (1976).
23. Beck AT. *Beck Hopelessness Scale*. The Psychological Corporation, San Antonio. (1988).
24. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol* (1959) 32:50–5. doi: 10.1111/j.2044-8341.1959.tb00467.x
25. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders. 3rd edn, revised*. DSM-III-R: Washington, DC (1987).
26. Bernstein DP, Ahluvalia T, Pogge D, Handelsman L. Validity of the Childhood Trauma Questionnaire in an adolescent psychiatric population. *J Am Acad Child Adolesc Psychiatry* (1997) 36:340–8. doi: 10.1097/00004583-199703000-00012
27. Carver CS, Scheier MF, Weintraub JK. Assessing coping strategies: A theoretically based approach. *J Pers Soc Psychol* (1989) 56:267. doi: 10.1037//0022-3514.56.2.267
28. Bagby RM, Parker JD, Taylor GJ. The twenty-item Toronto Alexithymia Scale–I. Item selection and cross-validation of the factor structure. *J Psychosom Res* (1994) 38:23–32. doi: 10.1016/0022-3999(94)90005-1
29. American Psychiatric Association. DSM-5 Task Force. In: *Diagnostic and statistical manual of mental disorders: DSM-5™, 5th ed*. Arlington, VA: American Psychiatric Publishing (2013).
30. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* (1998) 59(Suppl 20):22–33; quiz 34–57.
31. Wechsler D. *WISC-IV Wechsler Intelligence Scale for Children: Technical and interpretative manual*. Psychological Corporation: San Antonio (TX) (2003).
32. Mazure CM, Maciejewski PK, Jacobs SC, Bruce ML. Stressful life events interacting with cognitive/personality styles to predict late-onset major depression. *Am J Geriatr Psychiatry* (2002) 10:297–304. doi: 10.1097/00019442-200205000-00009
33. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* (2008) 40:879–91. doi: 10.3758/brm.40.3.879
34. Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res* (1967) 11:213–8. doi: 10.1016/0022-3999(67)90010-4
35. Yap SCY, Anusic I, Lucas RE. Does personality moderate reaction and adaptation to major life events? Evidence from the British Household Panel Study. *J Res Pers* (2012) 46:477–88. doi: 10.1016/j.jrp.2012.05.005
36. Carnelley KB, Wortman CB, Bolger N, Burke CT. The time course of grief reactions to spousal loss: Evidence from a national probability sample. *J Pers Soc Psychol* (2006) 91:476–92. doi: 10.1037/0022-3514.91.3.476
37. Lucas RE, Dyrenforth PS. Does the Existence of Social Relationships Matter for Subjective Well-being. In: Vohs KD, Finkel EJ, editors. *Self and relationships: Connecting intrapersonal and interpersonal processes*. New York: Guilford Press (2006).
38. Fennig S, Horesh N, Aloni D, Apter A, Weizman A, Fennig S. Life events and suicidality in adolescents with schizophrenia. *Eur Child Adolesc Psychiatry* (2005) 14:454–60. doi: 10.1007/s00787-005-0498-z
39. Rajalin M, Hirvikoski T, Salander Renberg E, Åsberg M, Jokinen J. Family history of suicide and interpersonal functioning in suicide attempters. *Psychiatry Res* (2017) 247:310–4. doi: 10.1016/j.psychres.2016.11.029
40. Kohn Y, Zislin J, Agid O, Hanin B, Troudart T, Shapira B, et al. Increased prevalence of negative life events in subtypes of major depressive disorder. *Compr Psychiatry* (2001) 42:57–63. doi: 10.1053/comp.2001.19753
41. Pompili M, Gibiino S, Innamorati M, Serafini G, Del Casale A, De Risio L, et al. Prolactin and thyroid hormone levels are associated with suicide attempts in psychiatric patients. *Psychiatry Res* (2012) 200:389–94. doi: 10.1016/j.psychres.2012.05.010
42. Roy A, Segal NL, Centerwall BS, Robinette CD. Suicide in twins. *Arch Gen Psychiatry* (1991) 48:29–33. doi: 10.1001/archpsyc.1991.01810250031003
43. Tidemalm D, Runeson B, Waern M, Frisell T, Carlström E, Lichtenstein P, et al. Familial clustering of suicide risk: a total population study of 11.4 million individuals. *Psychol Med* (2011) 41:2527–34. doi: 10.1017/S0033291711000833
44. Gøtzsche PC, Young AH, Crace J. Does long term use of psychiatric drugs cause more harm than good? *BMJ* (2015) 350:h2435. doi: 10.1136/bmj.h2435
45. McCann DJ, Petry NM, Bresell A, Isacson E, Wilson E, Alexander RC. Medication Nonadherence, “Professional Subjections,” and Apparent Placebo Responders: Overlapping Challenges for Medications Development. *J Clin Psychopharmacol* (2015) 35:566–73. doi: 10.1097/JCP.0000000000000372
46. Lanouette NM, Folsom DP, Sciolla A, Jeste DV. Psychotropic medication nonadherence among United States Latinos: a comprehensive literature review. *Psychiatr Serv* (2009) 60:157–74. doi: 10.1176/appi.ps.60.2.157
47. Jawad I, Watson S, Haddad PM, Talbot PS, McAllister-Williams RH. Medication nonadherence in bipolar disorder: a narrative review. *Ther Adv Psychopharmacol* (2018) 8:349–63. doi: 10.1177/2045125318804364
48. Swann AC. Approaches to preventing relapse in bipolar disorder: addressing nonadherence and prodromal symptoms. *J Clin Psychiatry* (2010) 71:e35. doi: 10.4088/JCP.9075tx4c
49. Castanon N, Leonard BE, Neveu PJ, Yirmiya R. Effects of antidepressants on cytokine production and actions. *Brain Behav Immun* (2002) 16:569–74. doi: 10.1016/s0889-1591(02)00008-9
50. Jokinen J, Forslund K, Ahnemark E, Gustavsson JP, Nordström P, Asberg M. Karolinska Interpersonal Violence Scale predicts suicide in suicide attempters. *J Clin Psychiatry* (2010) 71:1025–32. doi: 10.4088/JCP.09m05944blu
51. Dalgard OS, Dowrick C, Lehtinen V, Vazquez-Barquero JL, Casey P, Wilkinson G, et al. Negative life events, social support and gender difference in depression: A multinational community survey with data from the ODIN study. *Soc Psychiatry Psychiatr Epidemiol* (2006) 41:444–51. doi: 10.1007/s00127-006-0051-5
52. Wilhelm K, Parker G, Dewhurst J. Examining sex differences in the impact of anticipated and actual life events. *J Affect Disord* (1998) 48:37–45. doi: 10.1016/s0165-0327(97)00138-9
53. Liu X, Tein JY. Life events, psychopathology, and suicidal behavior in Chinese adolescents. *J Affect Disord* (2005) 86:195–203. doi: 10.1016/j.jad.2005.01.016
54. Girardi P, Pompili M, Innamorati M, Mancini M, Serafini G, Mazzarini L, et al. Duloxetine in acute major depression: review of comparisons to placebo and standard antidepressants using dissimilar methods. *Hum Psychopharmacol* (2009) 24:177–90. doi: 10.1002/hup.1005
55. Pompili M, Serafini G, Innamorati M, Venturini P, Fusar-Poli P, Sher L, et al. P. Agomelatine, a novel intriguing antidepressant option enhancing neuroplasticity: a critical review. *World J Biol Psychiatry* (2013) 14:412–31. doi: 10.3109/15622975.2013.765593
56. Miller GE, Chen E, Parker KJ. Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving towards a model of behavioral and biological mechanisms. *Psychol Bull* (2011) 137:959–97. doi: 10.1037/a0024768
57. Zhang TY, Bagot R, Parent C, Nesbitt C, Bredy TW, Caldji C, et al. Maternal programming of defensive responses through sustained effects on gene expression. *Biol Psychol* (2006) 73:72–89. doi: 10.1016/j.biopsycho.2006.01.009

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Serafini, Gonda, Canepa, Geoffroy, Pompili and Amore. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.