

Journal Pre-proof

Contrasting Epidemiology and Genetic Variation of *Plasmodium vivax* Infecting Duffy Negatives across Africa

Eugenia Lo, Gianluca Russo, Kareen Pestana, Daniel Kepple, Beka Raya Abargero, Ghyslaine Bruna Djeunang Dongho, Karthigayan Gunalan, Louis H Miller, Muzamil Mahdi Abdel Hamid, Delenasaw Yewhalaw, Giacomo Maria Paganotti



PII: S1201-9712(21)00410-0

DOI: <https://doi.org/10.1016/j.ijid.2021.05.009>

Reference: IJID 5393

To appear in: *International Journal of Infectious Diseases*

Received Date: 13 April 2021

Revised Date: 6 May 2021

Accepted Date: 7 May 2021

Please cite this article as: { doi: <https://doi.org/>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Contrasting Epidemiology and Genetic Variation of *Plasmodium vivax* Infecting Duffy Negatives across Africa

Eugenia Lo^{1*}, Gianluca Russo^{2*}, Kareen Pestana¹, Daniel Kepple¹, Beka Raya Abargero³, Ghyslaine Bruna Djeunang Dongho^{2,4}, Karthigayan Gunalan⁵, Louis H Miller⁵, Muzamil Mahdi Abdel Hamid⁶, Delenasaw Yewhalaw³, Giacomo Maria Paganotti^{7,8,9}

¹ Biological Sciences, University of North Carolina at Charlotte, USA

² Department of Public Health and Infectious diseases, Sapienza University, Rome, Italy

³ Tropical Infectious Disease Research Center, Jimma University, Jimma, Ethiopia

⁴ Evangelical University of Cameroon, Bandjoun, Cameroon

⁵ Laboratory of Malaria and Vector Research, NIAID/NIH, Bethesda, USA

⁶ Department of Parasitology and Medical Entomology, Institute of Endemic Diseases, University of Khartoum, Khartoum, Sudan

⁷ Botswana-University of Pennsylvania Partnership, Gaborone, Botswana

⁸ Division of Infectious Diseases, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

⁹ Department of Biomedical Sciences, Faculty of Medicine, University of Botswana, Gaborone, Botswana

* Correspondence authors: Eugenia Lo; email: eugenia.lo@uncc.edu; Gianluca Russo; email: gianluca.russo@uniroma1.it

Highlights

- **Vivax malaria is spreading across Africa causing symptomatic and asymptomatic cases**
- ***P. vivax* infections in Duffy-negative populations have low parasitemia**
- **Sensitive and standardized methods are needed to detect vivax malaria in Africa**
- ***P. vivax* in Duffy-negative Africans have evolved multiple times independently**
- **Public health significance of vivax malaria in Africa should no longer be neglected**

Abstract

Objectives. *Plasmodium vivax* malaria was thought to be rare in Africans who lack the Duffy blood group antigen expression. However, recent studies indicate that *P. vivax* can infect Duffy-negative individuals and has penetrated into areas of high Duffy-negativity across Africa. This study compares epidemiological and genetic features of *P. vivax* between African regions.

Methods. We utilized a standardized approach to identify and quantify *P. vivax* from Botswana, Ethiopia, and Sudan, where Duffy-positive and Duffy-negative individuals coexist. We sequenced Duffy Binding Protein (DBP) gene and inferred genetic relationships among all Africa *P. vivax*.

Results. Among 1,215 febrile patients, the proportions of Duffy negativity range from 20-36% in East Africa to 84% in Southern Africa. *P. vivax* prevalence among Duffy-

negative populations ranging from averaged 9.2% in Sudan to 86% in Botswana.

Parasite density in Duffy-negative is significantly lower than in Duffy-positive infections.

P. vivax in Duffy-negative populations were not monophyletic. Duffy-negative and Duffy-positive *P. vivax* shared similar DBP haplotypes and occurred in multiple well-supported clades.

Conclusions. Duffy-negative Africans are not resistant to *P. vivax* and the public health significance should not be neglected. This study highlights need for standardized approach and more resources/training to diagnosis of vivax malaria in Africa.

Keywords: Malaria, *Plasmodium vivax*, Duffy negatives, Sub-Saharan Africa, Genetic relationships, Molecular epidemiology

Introduction

Plasmodium vivax malaria was previously thought to be rare or absent in African populations who lack the Duffy blood group antigen expression (Miller et al., 1976, Howes et al., 1995). A point mutation (c.1-67T>C; rs2814778) in the GATA-1 transcription factor binding site of the Duffy antigen/receptor for chemokines (*DARC*) gene promoter alters erythroid expression, eliminating Duffy antigen expression on the surface of the red blood cells (Tournamille et al. 1995, King et al., 2011). However, recent studies reported several cases of *P. vivax* infection in Duffy-negative people in different parts of Africa (Zimmerman, 2017, Gunalan et al., 2018), including countries where Duffy-negatives are predominant (Brazeau et al., 2018, Mendes et al., 2011, Motshoge et al., 2016, Niangaly et al., 2017, Russo et al., 2017) (**Table 1**). In addition, 29 African countries including six previously undocumented endemic countries (Benin,

Comoros, Mozambique, Senegal, Zambia and Zimbabwe) have reported *P. vivax* clinical cases, infected vectors or asymptomatic parasitemia (Niang et al., 2018, Oboh et al., 2020, Poirier et al., 2016). These reports indicate that the endemic range of *P. vivax* has extended beyond East Africa and penetrated into areas of very high Duffy-negativity (Gunalan et al., 2018, Twohig et al., 2019). While *P. falciparum* is considered to be the deadliest malaria parasite with the most severe clinical outcomes, *P. vivax* is more widespread and often associated with high levels of morbidity. Compared to *P. falciparum*, *P. vivax* has a broader temperature tolerance, an earlier onset of gametocyte development, and can form dormant hypnozoites causing relapse (Livingstone, 1984), enabling *P. vivax* to spread through the diverse African climate and outcompete *P. falciparum* (Battle et al., 2019). Primaquine and 8-aminoquinoline are antimalarials effective in clearing hypnozoites and preventing relapses, but they may promote hemolysis in subjects with G6PD deficiency (Baird, 2019). These factors make *P. vivax* malaria difficult to control and eliminate, highlighting the concern of this 'new' *P. vivax* strains that infect Duffy-negative hosts to spread through much of Africa and result in substantial, negative public health and economic impacts.

There is a major knowledge gap in the *P. vivax* invasion mechanisms in Duffy-negative erythrocytes. In *P. falciparum*, erythrocyte invasion involves multiple interactions between parasite ligands and host receptors, some of which have overlapping and partially redundant roles (Cowman et al. 2017, Kumar and Tolia, 2019). Several established invasion ligands from Erythrocyte Binding Antigens such as EBA-175, EBA-181/JESEBL and EBA-140/BAEBL and Reticulocyte binding homolog proteins such as RH1, RH2a, RH2b, RH4 and RH5 are used by *P. falciparum* for

invasion (Gunalan et al., 2013, Kumar and Tolia, 2019). In *P. vivax*, only a single *P. vivax* ligand-receptor interaction has so far been studied in any detail, *P. vivax* Duffy Binding Protein (*PvDBP1*). Previous study has shown that mutations in *PvDBP1* region II unique to *P. vivax* in Duffy-negative people in Ethiopia did not lead to binding of Duffy-negative erythrocytes (Gunalan et al., 2016). Salvador (Sal) I *P. vivax* infects Squirrel monkeys without *PvDBP1* binding to Squirrel monkey erythrocytes (Gunalan et al., 2019). Further, EBP/DBP2 region II, a paralog of *PvDBP1*, was shown to bind to Duffy-positive and Duffy-null human erythrocytes at low frequency (Gunalan et al., 2016, Ntumngia et al., 2016), despite being deleted in Sal-I *P. vivax* (Hester et al., 2013). Recently, reticulocyte binding protein RBP2b of *P. vivax* was shown to bind to transferrin receptor in the reticulocytes (Gruszczyk et al., 2018). These findings suggested that there are other Duffy-independent pathways that enable erythrocyte invasion and explain the widespread phenomenon of *P. vivax* infections in Africa.

Despite the fact that several case reports from almost all countries across the African continent are emerging from various entomological and serological studies, community surveys, and clinical records (Gunalan et al., 2018, Twohig et al., 2019), the documentation of *P. vivax* infections across Africa is diverse, context-specific, and primarily driven by the specific objectives of isolated clinical or epidemiological activities. The varied diagnostic and methodological approaches used across studies have limited our ability to identify distinct epidemiological characteristics of *P. vivax* between regions (**Table 1**). This situation is concerning because there is no comprehensive genetic and epidemiological data of *P. vivax* in Africa available to National Malaria Programs or World Health Organization to assess impacts and confer control strategies. Therefore,

in this study, we utilized a standardized assay to examine the epidemiological attributes of *P. vivax* in three African countries where Duffy-positive and Duffy-negative individuals coexist. Specifically, we (1) compared the prevalence of Duffy negativity and *P. vivax* infections among countries; (2) compared *P. vivax* parasitemia between Duffy-negative and Duffy-positive infections collected from the same area; and (3) inferred the genetic relationships among the African *P. vivax* isolates. The epidemiological and genetic features of *P. vivax* from different parts of Africa will fill critical gap in understanding how widespread this phenomenon is impacting malaria control and the important effect of *P. vivax* as a cause of anemia.

Materials and Methods

Study sites and sample collection

A total of 1,215 febrile patients were collected from seven study sites in three countries including (1) Jimma and Bonga in Ethiopia; (2) Khartoum, River Nile, and New Halfa in Sudan; and (3) Tutume and Kweneng East in Botswana (**Figure 1**). Finger-prick blood samples were obtained from patients who visited the health facilities. Thick and thin blood smears were prepared for microscopic screening. Three to four blood spots were blotted on Whatman 3MM filter paper from each participant. Parasite DNA was extracted from dried blood spots by the Saponin/Chelex method (Bereczky et al., 2005). Eluted DNA was used for PCR diagnosis, quantification and genotyping of malaria parasites.

Molecular screening of *P. vivax*

Parasite gene copy number was estimated using the SYBR Green detection method (Lo et al., 2015) using *P. vivax*-specific primers that targeted the 18S rRNA genes (detail in **Supplementary File 1**). Each assay included positive controls of *P. vivax* Pakchong (MRA-342G) and Nicaragua (MRA-340G) isolates, in addition to negative controls. A standard curve was produced from a ten-fold dilution series of the *P. vivax* control plasmid to determine the amplification efficiency (E). Melting curve analyses were performed to confirm the specificity of gene amplifications. The mean threshold cycle (*C_t*) and standard error were calculated from three independent assays of each sample. The amount of parasite density in a sample was calculated using the follow equation: $\text{Parasite density}_{\text{sample}} = 2^{E \times (40 - C_{t\text{sample}})}$. The differences in the log-transformed parasite density between samples among the study sites were assessed for significance by one-tailed t-tests.

Duffy blood group genotyping

For all febrile patients, we first employed qPCR-based TaqMan assay to examine the point mutation (c.1-67T>C; rs2814778) of the *DARC* gene (**Supplementary File 1**). A no-template control was used in each assay. The *Fy* genotypes were determined by the allelic discrimination plot based on the fluorescent signal emitted from the allele-specific probes. For *P. vivax* positive samples, a 1,100-bp fragment of the *DARC* gene was further amplified using published primers (Menard et al., 2010). PCR products were sequenced to confirm the *Fy* genotypes.

Phylogenetic analyses of *P. vivax* from Duffy negative and Duffy positive samples

We amplified and obtained *PvDBP* sequences of 4 Duffy-positive and 4 Duffy-negative *P. vivax* samples from Botswana, 107 Duffy-positive and 9 Duffy-negative *P. vivax* samples from Ethiopia, and 53 Duffy-positive and 16 Duffy-negative *P. vivax* samples from Sudan (Genbank accession number: MZ062224-MZ062409). These sequences were aligned with 36 previously published *P. vivax* isolates from other parts of Africa including Uganda ($n=31$), Madagascar ($n=4$), and Mauritania ($n=1$; **Supplementary File 2**). Duffy status of the published sequences are unknown. The *DBP* sequence of Sal-1 (NC_009911.1) and *EBP* sequence of *P. cynomolgi* (Y11396.1) were used as outgroups. Phylogenetic trees were reconstructed using the maximum likelihood method implemented in RAxML v8.0 with 500 bootstrap replicates to assess clade support (details in **Supplementary File 1**). We further examined the nucleotide and haplotype diversity of *PvDBP* sequences in Duffy-negative and Duffy-positive samples using DnaSP v6.12.03.

Results and Discussion

Contrasting proportion of Duffy-negatives and *P. vivax* prevalence

Duffy genotyping shows different proportions of Duffy-negative among febrile patients in Botswana, Ethiopia, and Sudan (**Figure 1**). In Botswana, the proportion of Duffy-negative was 83.5% (147/176) among febrile patients (**Figure 1**). In Kweneng East, our qPCR analyses indicated that 3% (9 out of 301) of the febrile patients were *P. vivax* positive. Among them, eight were Duffy-negative (C/C) and one was Duffy-positive (T/C) (**Table 2; Supplementary File 3**). In Tutume, 6.8% (12/176) of the febrile patients were detected with *P. vivax* and 10 of them were Duffy-negative. Vivax malaria was first

reported in asymptomatic children in a survey during the 2012-2013 transmission season [10]. The average rate of asymptomatic *P. vivax* cases was 4.7%, but with large variation among districts. Compared to other parts of Botswana, Tutume and Kweneng East accounted for most of the *P. vivax* cases, with previously reported rates of 16.9% (54/320) and 13.6% (93/686), respectively (Motshoge et al., 2016).

In Ethiopia, the proportion of Duffy-negative was 35.9% (235/655) among febrile patients (**Figure 1**), similar to our earlier finding in Asendabo indicating that 35.1% (137/390) of the general population was Duffy-negative (Lo et al., 2015). Among the 358 febrile patient samples collected in Jimma, 36% (129/358) were Duffy-negatives (**Figure 1**) and 37.4% (134/358) were detected with *P. vivax* (**Table 2**). About 11.9% (16/134) of the confirmed *P. vivax* infections were from Duffy-negatives. Likewise, in Bonga, 30.3% (125/413) of the febrile patients were detected with *P. vivax* and 3.2% (4/125) were from Duffy-negatives (**Table 2**). For these 20 Duffy-negatives *P. vivax* infections, microscopy, nested and quantitative PCRs indicated that 16 were single infections and four were mixed with *P. falciparum*. Vivax malaria is a significant problem in Ethiopia (Lo et al., 2015, Woldearegai et al., 2013). Our previous study has shown that the asymptomatic prevalence of *P. vivax* is 5.9% (23/390) in Asendabo and Duffy-negatives accounted for 8.7% (2/23) of the *P. vivax* infections (Lo et al., 2015). A lower proportion of Duffy-negativity in febrile patients and the general population in Ethiopia, as compared to Botswana, is consistent with the ethnic diversity and complex admixture history in East Africa (Hollfelder et al., 2017, Pickrell et al., 2014).

In Sudan, the proportion of Duffy-negative was 20% (77/384) among febrile patients (**Figure 1**). Over a 6-month collection period between 2018 and 2019, 101 out

of 831 febrile patients were confirmed as *P. vivax* positive by qPCR assays (**Table 2**). Further testing revealed that 4 of the 101 *P. vivax* samples were mixed with *P. falciparum*. The highest rate of *P. vivax* infection was observed in River Nile, of which 24.4% (52/213) of the febrile patients were confirmed with *P. vivax* and Duffy-negatives accounted for 3.8% (2/52) of these infections (**Table 2**). In Khartoum, 8% (42/525) of the febrile patients were *P. vivax*-positive and Duffy-negatives accounted for 9.5% (4/42) of these infections. In New Halfa, despite a smaller sample size, 7.5% (7/93) of the febrile patients was *P. vivax*-positive and Duffy-negatives accounted for 14.3% (1/7) of these infections (**Table 2**). Across the country, there has been an increase in *P. vivax* detection and reports in recent years (Albsheer et al., 2019) and our findings indicated that the infection rate in Duffy-negative individuals varies among study sites.

Historical movement and genetic admixture explain distribution of Duffy-negative people in Africa

Historical human movement and human genetics are highly relevant to the distribution of Duffy-negative people and *P. vivax* in Africa. Recent genome-wide studies of African populations have refined earlier models of the continent's history and its impact on genetic diversity of its inhabitants (Choudhury et al., 2020). Our data showing a Duffy negative rate of 83.5% among febrile patients in Botswana (**Figure 1**) is consistent with the Bantu expansion and admixture theories (Choudhury et al., 2020, Grollemunda et al., 2015). The Bantu expansion and population admixture are two main historical events that shape the present distribution and genetic make-up of ethnic groups across Africa. The Bantu and Khoisan are two major ethnic groups in West-Central and

Southern Africa, with the Bantu heartland in the region between southern Nigeria and Cameroon where malaria transmission was and still is endemic (Grollemunda et al., 2015). A component of Bantu ancestry (likely Duffy-negative) was found in the Southern African Khoisan, which were originally and mostly Duffy-positive ancestors (Hamblin et al., 2002, Petersen et al., 2013). The Duffy-negative allele from Bantu of West-Central Africa may have reached south of the continent within the last 750 years and mixed with the indigenous Khoisan, resulting in a variable Khoisan ancestry (Busby et al., 2016, Schuster et al., 2010).

While the direction of the Bantu expansion is still in debate, there is evidence showing that the Bantu migrated towards East Africa where other ethnic groups such as the Cushitic and Nilotic dominated, potentially around 2,000 years ago (Pickrell et al., 2014). Our data showing a Duffy negative rate of 20-36% in Southwestern Ethiopia and East Sudan (**Figure 1**) is consistent with the complex admixture history. The Ethiopian and Sudanese population, with an admixture of several Eurasian ancestries and some Nilotic and Semitic-Cushitic components, migrated south after the Bantu expansion 2-5 thousand years (Hollfelder et al., 2017, Pickrell et al., 2014). Many population groups in Sudan are dominated by Nilotic and Eurasian admixtures with minimal West African component. One such exception is the Afro-Asiatic speaking Hausa population in the Middle Eastern Sudan, which have migrated from West Africa within the past 300 years (Hollfelder et al., 2017). These migrations could have spread *P. vivax* from West-Central to other parts of Africa.

Low parasitemia in symptomatic Duffy-negative *P. vivax* infections and implications on invasion mechanism

In Botswana, Ethiopia, and Sudan, Duffy-positive and Duffy-negative individuals coexist. *P. vivax* parasite density in Duffy-negative infected individuals is significantly lower than the Duffy-positive infected individuals, regardless of geographical differences (**Figure 2**). Duffy-positive individuals with heterozygous C/T and homozygous T/T were not significantly different in parasitemia. Both genotypes showed significantly higher parasitemia than Duffy negative C/C. The Duffy-negative *P. vivax* samples in Ethiopia and Sudan showed a greater range of parasitemia variation than those in Botswana. This may be due to differences in sample size (**Figure 2; Supplementary File 3**). In very few cases the asexual parasites were detected by microscopy in Duffy-negative individuals. For example, among the 20 *P. vivax* infections identified in Duffy-negative patients from Ethiopia, only four were microscopic-positive and they all showed a relatively higher parasitemia compared to the submicroscopic infections. The Duffy-negative individuals who were infected with *P. vivax* were mostly submicroscopic and exhibited fever at the time of sample collection. Without highly sensitive diagnostic tools and vigorous on-site training and screening of *P. vivax* in different parts of Africa, the public health burden, economic impact, and severity associated with vivax malaria could have been vastly underestimated. The clinical spectrum of *P. vivax* malaria ranges from asymptomatic parasitemia and uncomplicated febrile illness to severe and fatal malaria (Naing et al., 2014). Moreover, *P. vivax* can cause anemia during chronic undetected infections (Niangaly et al., 2017). Other severe clinical manifestations include multiorgan dysfunction associated with anemia and thrombocytopenia, spontaneous

abortions, premature and low birth weight in pregnant women (Naing et al., 2014).

These clinical features have mostly been described for Duffy-positive populations. It is unclear if the spectrum of clinical symptoms is different in Duffy-negative patients in Africa.

Low parasitemia observed in Duffy-negative individuals might suggest a low invasion capability of *P. vivax* in Duffy-negative individuals. Recent study has shown that mutations in *PvDBP1* region II unique to *P. vivax* in Duffy-negative people in Ethiopia did not lead to binding of Duffy-negative erythrocytes (Gunalana et al., 2016). Also, Sal-I *P. vivax* infects Squirrel monkeys without *PvDBP1* binding to Squirrel monkey erythrocytes (Gunalana et al., 2019). These findings suggested that there are other Duffy-independent pathways that enable erythrocyte invasion. For example, EBP/DBP2 region II has shown to bind to Duffy-positive and Duffy-negative human erythrocytes at low frequency (Gunalana et al., 2016, Ntumngia et al., 2016). CD71 (Transferrin Receptor 1, TfR1) has been shown to bind readily to the reticulocyte binding proteins (*PvRBP2b*) based on *in vitro* experiments (Chan et al., 2020, Gruszczyk et al., 2018). Given reticulocytes constitute only a small fraction of all red blood cells, invasion via this RBP2b-TfR1 pathway may result in only a small number of infected erythrocytes and this may explain the considerably low parasitemia observed in Duffy-negative *P. vivax* infections (**Figure 2**). Further, recent transcriptomic study has also indicated that genes belonging to tryptophan-rich antigen and merozoite surface protein families were highly expressed in the *Saimiri*-infected *P. vivax*, of which erythrocytes did not bind to DBP1 from the Belem isolate of *P. vivax* (Gunalana et al., 2019). There is growing evidence that members of the tryptophan-rich antigen gene

family are involved in erythrocyte invasion (Zeeshan et al., 2015). Various other invasion ligands may also mediate the recognition and invasion to reticulocytes, providing a potential mechanism for variations in reticulocyte preference (Baquero et al., 2017, Moreno-Pérez et al., 2017). Successful schizont development has been shown to be associated with increased younger reticulocytes in the Indian *P. vivax* isolates (Lim et al., 2016). The low prevalence of schizonts in peripheral blood has led to the hypothesis that *P. vivax* could be sequestering in reticulocyte-rich zones such as the bone marrow (Mayor and Alano, 2015), resulting in lower detectable parasitemia. Future studies should clarify the expression and role of various *P. vivax* ligand proteins and their respective receptors in Duffy-negative erythrocyte invasion.

Genetic relationships and origin hypotheses of *P. vivax* in Duffy-negative Africans

Maximum likelihood analyses of the African *P. vivax* isolates based on *PvDBP* indicated that *P. vivax* from Duffy-negative individuals were not monophyletic but found in multiple well-supported clades (clades I-III in **Figure 3**). These clades did not show clear geographical boundary but a mixture of *P. vivax* from different African countries. For instance, Duffy-negative *P. vivax* from Botswana, Ethiopia, and Sudan were closely related to Duffy-positive *P. vivax* from the same area, as well as to *P. vivax* from neighboring Uganda (clade II; bootstrap 91%). The Duffy-negative *P. vivax* were clustered together with the Duffy-positive ones without genetic distinction. The present data may imply that Duffy-negative and Duffy-positive individuals shared similar *P. vivax* strains possibly by the same ancestral origin or through recent transmission. The evolution of *PvDBP* region II could be also driven by functional selection rather than by

geographical isolation. Interestingly, Duffy-negative *P. vivax* samples from Ethiopia and Sudan showed a higher nucleotide and haplotype diversity than the Duffy-positive ones, despite a smaller sample size (**Table 3**). Among all geographical isolates, *P. vivax* from Uganda and Madagascar had the highest level of genetic variation, though Duffy status of these samples are unclear (**Table 3**). These findings offered a hypothesis on the origin of Duffy-negative *P. vivax*, but *PvDBP* could be biased by selection or has limited resolution. Extensive phylogenetic analyses using whole genome sequences of Duffy-negative *P. vivax* from West-Central, Southern, and East Africa, together with the existing data of the *P. vivax*-like isolates in African apes are needed to adjudicate these origin hypotheses.

Previous studies indicated that *P. vivax* in Southeast Asia and South America evolved in a clade of parasites that infect African monkeys (Loy et al., 2018). *Plasmodium vivax* in African apes might present a substantial parasite reservoir from which Duffy-positive and Duffy-negative human infections arose from. There are two hypotheses concerning the origin of *P. vivax* in Duffy-negative Africans (**Figure 4**). The first hypothesis posits that the ancestral *P. vivax* infected all African primates including apes and Duffy-positive humans (Liu et al., 2014) (**Figure 4A**). One of these ancestral lineages evolved to a Duffy-independent pathway and subsequently spread to different parts of Africa via human migration (Choudhury et al., 2020, Grollemunda et al., 2015). The geographical overlap between apes and humans, e. g, in Cameroon and the Democratic Republic of Congo suggest a West-Central African origin of *P. vivax* in Duffy-negatives (Liu et al., 2014). The second hypothesis posits that the ancestral *P. vivax* infected only non-human primates in Africa until some of the lineages crossed the

species barrier and gave rise to the parasite population currently infecting Duffy-positive humans (Prugnolle et al., 2013). It is possible that Duffy-negative *P. vivax* observed today across Africa represent separate lineages that were derived multiple times independently from Duffy-positive individuals (**Figure 4B**). Previous phylogenies based on nuclear genes and partial mitochondrial genomes revealed incongruent genetic relationships (Liu et al., 2014, Prugnolle et al., 2013), possibly due to incomplete lineage sorting or lack of phylogenetic signal (Maddison and Knowles, 2006). Moreover, no African *P. vivax* isolates from Duffy-positive and Duffy-negative individuals were included. Future studies should employ genome-based phylogenetic approach and molecular dating analyses to clarify the origin of *P. vivax* in Africa.

Conclusions

With the increasing number of *P. vivax* cases reported in Duffy-negative individuals as well as across the continent, vivax malaria is no longer a rare but a growing and possibly widespread phenomenon in Africa. To the best of our knowledge, this paper is the first using a standardized approach to characterize and compare the epidemiological and genetic features of Duffy-negative *P. vivax* from different parts of Africa. The generally low parasitemia observed in the Duffy-negative infections may suggest a less efficient but continuously evolving invasion mechanism that allows a greater negative public health impact in Africa in coming years. The genetic relatedness based on *PvDBP* sequences suggested similar strains shared between Duffy-negative and Duffy-positive populations, though the transmission capability of *P. vivax* in Duffy-negative individuals is still unclear. Further investigations are needed to unveil the

invasion and transmission mechanisms of these infections. These data would help predict the scale of disease spread and improve existing malaria control measures, beyond *P. falciparum* in Africa. On the public health front-end, there should be more resources and training allocated to diagnosis and treatment of vivax malaria, given its unique ability in causing relapse and other longer-term health problems such as anemia in asymptomatic infections. Duffy-negative Africans are not resistant to *P. vivax* infection and the public health significance of vivax malaria in Africa should no longer be neglected.

Authors' contributions

EL, GR, MMAH, DY and GMP conceived and designed the study; EL, GR, BR, GBDD, MMAH, DY and GMP collected the samples; EL, KP, DK, KG, and GMP collected and analyzed the data; EL, GR, KG, LHM and GMP wrote the paper. All authors read and approved the final manuscript.

Funding

This research was funded by National Institutes of Health (R15 AI138002). GMP was supported also by Penn Center for AIDS Research (grant # P30 AI045008).

Ethical approval

Scientific and ethical clearance was given by the institutional scientific and ethical review boards of Jimma University (Ethiopia), the Ethics Committee of the Institute of Endemic Diseases, University of Khartoum (Reference number: 9/2016), the Health

Research and Development Division of the Botswana Ministry of Health and Wellness (Reference number: HPDME: 13/18/1), and University of North Carolina at Charlotte (USA). Written informed consent/assent was obtained from all consenting heads of households, parents/guardians (for minors under 18 years old), and each individual who participated in this study.

Conflict of interest

The authors declare no conflict of interest.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank the students and staffs from Jimma University, University of Khartoum, University of Botswana, and University of North Carolina at Charlotte for their technical assistance in sample collection and processing; the communities and hospitals for their support and willingness to participate in this research.

References

1. Albsheer M, Pestana K, Ahmed S, Elfaki M, Gamil E, Ahmed SM, et al. Distribution of Duffy Phenotypes among *Plasmodium vivax* Infections in Sudan. *Genes* 2019;10:437.
2. Baird JK. 8-Aminoquinoline therapy for latent malaria. *Clin Microbiol Rev* 2019;32:e00011-19.

3. Baquero LA, Moreno-Pérez DA, Garzón-Ospina D, Forero-Rodríguez J, Ortiz-Suárez HD, Patarroyo MA. PvGAMA reticulocyte binding activity: predicting conserved functional regions by natural selection analysis. *Parasit Vectors* 2017;10: 251.
4. Battle KE, Lucas TCD, Nguyen M, Howes RE, Nandi AK, Twohig KA, et al. Mapping the global endemicity and clinical burden of *Plasmodium vivax*, 2000–17: a spatial and temporal modelling study. *Lancet* 2019;394:P332-343.
5. Berczky S, Martensson A, Gil JP, Farnert A. Short report: Rapid DNA extraction from archive blood spots on filter paper for genotyping of *Plasmodium falciparum*. *Am J Trop Med Hyg* 2005;72:249–51.
6. Brazeau NF, Whitesell AN, Doctor SM, Keeler C, Mwandagalirwa MK, Tshetu AK, et al. *Plasmodium vivax* Infections in Duffy-negative individuals in the Democratic Republic of the Congo. *Am J Trop Med Hyg* 2018;99:1128-1133.
7. Busby GBJ, Band G, Le QS, Jallow M, Bougama E, Mangano VD, et al. Malaria genomic epidemiology network. Admixture into and within sub-Saharan Africa. *eLife* 2016;5:e15266.
8. Chan LJ, Dietrich MH, Nguitragool W, Tham WH. *Plasmodium vivax* Reticulocyte Binding Proteins for invasion into reticulocytes. *Cell Microbiol* 2020;22:e13110.
9. Choudhury A, Aron S, Botigué LR, Sengupta D, Botha G, et al. High-depth African genomes inform human migration and health. *Nature* 2020;586:741–748.
10. Cowman A, Tonkin CJ, Tham WH, Duraisingh MT. The molecular basis of erythrocyte invasion by malaria parasites. *Cell Host Microbe* 2017;22:232-245.

11. Grollemunda R, Branforda S, Bostoenb K, Meadea A, Vendittia C, Page M. Bantu expansion shows that habitat alters the route and pace of human dispersals. *Proc Natl Acad Sci USA* 2015;112:13296-301.
12. Gruszczyk J, Kanjee U, Chan LJ, Menant S, Malleret B, *et al.* Transferrin receptor 1 is a reticulocyte-specific receptor for *Plasmodium vivax*. *Science* 2018;359:48-55.
13. Gunalan K, Gao X, Yap SS, Huang X, Preiser PR. The role of the reticulocyte-binding-like protein homologues of *Plasmodium* in erythrocyte sensing and invasion. *Cell Microbiol* 2013;15:35-44.
14. Gunalana K, Lo E, Hostetler JB, Yewhalaw D, Mu J, Neafsey DE, Yan G, Miller LH. Role of *Plasmodium vivax* Duffy-binding protein 1 in invasion of Duffy-null Africans. *Proc Natl Acad Sci USA* 2016;113:6271.
15. Gunalan K, Niangaly A, Thera MA, Doumbo OK, Miller LH. *Plasmodium vivax* Infections of Duffy-negative erythrocytes: Historically undetected or a recent adaptation? *Trends Parasitol* 2018;34:420-429.
16. Gunalan K, Sá JM, Barros RRM, Anzick SL, Caleon RL, Mershon JP, *et al.* Transcriptome profiling of *Plasmodium vivax* in *Samira* monkeys identifies potential ligands for invasion. *Proc Natl Acad Sci USA* 2019;116: 7053–61.
17. Hamblin MT, Thompson EE, Di Rienzo A. Complex signatures of natural selection at the Duffy blood group locus. *Am J Hum Genet* 2002;70:369-383.
18. Hollfelder N, Schlebusch CM, Günther T, Babiker H, Hassan HY, Jakobsson M. Northeast African genomic variation shaped by the continuity of indigenous groups and Eurasian migrations. *PLoS Genet* 2017;13:e1006976.

19. Hester J, Chan ER, Menard D, Mercereau-Puijalon O, Barnwell J, Zimmerman PA, Serre D. De novo assembly of a field isolate genome reveals novel *Plasmodium vivax* erythrocyte invasion genes. PLoS Negl Trop Dis 2013;7:e2569.
20. Howes RRE, Patil APAAP, Piel FBF, Nyangiri OAOOA, Kabaria CWCCW, Gething PW, et al. The global distribution of the Duffy blood group. Nat Commun 2011;2:266.
21. King CL, Adams JH, Xianli J, Grimberg BT, McHenry AM, Greenberg LJ, Siddiqui A, Howes RE, da Silva-Nunes M, Ferreira MU, Zimmerman PA. Fy(a)/Fy(b) antigen polymorphism in human erythrocyte Duffy antigen affects susceptibility to *Plasmodium vivax* malaria. Proc Natl Acad Sci USA 2011;108:20113-8.
22. Kumar H, Tolia NH. Getting in: The structural biology of malaria invasion. PLoS Pathog 2019; 15: e1007943.
23. Lim C, Pereira L, Saliba KS, Mascarenhas A, Maki JN, Chery L, Gomes E, et al. Reticulocyte preference and stage development of *Plasmodium vivax* isolates. J Infect Dis 2016;214:1081–1084.
24. Liu W, Li Y, Shaw KS, Learn GH, Plenderleith LJ, Malenke JA, Sundararaman SA, et al. African origin of the malaria parasite *Plasmodium vivax*. Nat Commun 2014;5:3346.
25. Livingstone FB. The Duffy blood groups, vivax malaria, and malaria selection in human populations: a review. Hum Biol 1984;56:413–25.
26. Lo EYY, Delenasaw Y, Zhong D, Zemene E, Degefa T, Tushune K, et al. Molecular epidemiology of *Plasmodium vivax* and *Plasmodium falciparum* malaria among Duffy-positive and Duffy-negative populations in Ethiopia. Malar J 2015;14:84-94.

27. Loy DE, Plenderleith LJ, Sundararaman SA, Liu W, Gruszczyk J, Chen YJ, et al. Evolutionary history of human *Plasmodium vivax* revealed by genome-wide analyses of related ape parasites. Proc Natl Acad Sci USA 2018;115:E8450-8459.
28. Maddison WP, Knowles LL. Inferring phylogeny despite incomplete lineage sorting. Syst Biol 2006;55:21-30.
29. Mayor A, Alano P. Bone marrow reticulocytes: a *Plasmodium vivax* affair? Blood 2015;125:1203-1205.
30. Menard D, Barnadas C, Bouchier C, Henry-Halldin C, Gray LR, Ratsimbaoa A, et al. *Plasmodium vivax* clinical malaria is commonly observed in Duffy-negative Malagasy people. Proc Natl Acad Sci USA 2010;107:5967-5971.
31. Mendes C, Dias F, Figueiredo J, Mora VG, Cano J, de Sousa B, et al. Duffy negative antigen is no longer a barrier to *Plasmodium vivax*--molecular evidences from the African West Coast (Angola and Equatorial Guinea). PLoS Negl Trop Dis 2011;5:e1192.
32. Miller LH, Mason SJ, Clyde DF, McGinniss MH. The resistance factor to *Plasmodium vivax* in blacks. The Duffy-blood-group genotype, FyFy. N Engl J Med 1976;295:302-304.
33. Moreno-Pérez DA, Baquero LA, Chitiva-Ardila DM, Patarroyo MA. Characterising PvRBSA: an exclusive protein from *Plasmodium* species infecting reticulocytes. Parasit Vectors 2017;10:243.
34. Motshoge T, Ababio GK, Aleksenko L, Read J, Peloewetse E, Loeto M, et al. Molecular evidence of high rates of asymptomatic *P. vivax* infection and very low *P. falciparum* malaria in Botswana. BMC Infect Dis 2016;16:520.

35. Naing C, Whittaker MA, Nyunt Wai V, Mak JW. Is *Plasmodium vivax* Malaria a Severe Malaria? A Systematic Review and Meta-Analysis. *PLoS Negl Trop Dis* 2014;8:e3071.
36. Niang M, Sane R, Sow A, Sadio BD, Chy S, Legrand E, Faye O, et al. Asymptomatic *Plasmodium vivax* infections among Duffy-negative population in Kedougou, Senegal. *Trop Med Health* 2018;46:45.
37. Niangaly A, Gunalan K, Ouattara A, Coulibaly D, MS'a J, Adams M, Travassos MA, et al. *Plasmodium vivax* infections over 3 years in Duffy blood group negative Malians in Bandiagara, Mali. *Am J Trop Med Hyg* 2017;97:744-752.
38. Ntumngia FB, Thomson-Luque R, Torres Lde M, Gunalan K, Carvalho LH, Adams JH. A novel erythrocyte binding protein of *Plasmodium vivax* suggests an alternate invasion pathway into Duffy-positive reticulocytes. *MBio* 2016;7:e01261-01216.
39. Oboh MA, Singh US, Ndiaye D, Badiane AS, Ali NA, Bharti PK, Das A. Presence of additional *Plasmodium vivax* malaria in Duffy negative individuals from Southwestern Nigeria. *Malar J* 2020;19:229.
40. Petersen DC, Libiger O, Tindall EA, Hardie R-A, Hannick LI, Glashoff RH, et al. Complex patterns of genomic admixture within Southern Africa. *PLoS Genet* 2013;9:e1003309.
41. Pickrell JK, Patterson N, Loh PR, Lipson M, Berger B, Stoneking M, Pakendorf B, Reich D. Ancient west Eurasian ancestry in southern and eastern Africa. *Proc Natl Acad Sci USA* 2014;111:2632-7.

42. Poirier P, Doderer-Lang C, Atchade PS, Lemoine JP, de l'Isle MC, Abou-Bacar A, et al. The hide and seek of *Plasmodium vivax* in West Africa: report from a large-scale study in Beninese asymptomatic subjects. *Malar J* 2016;15:570.
43. Prugnolle F, Rougeron V, Becquarta P, Berryd A, Makangab B, Raholaa N, et al. Diversity, host switching and evolution of *Plasmodium vivax* infecting African great apes. *Proc Natl Acad Sci USA* 2013;110:8123-8128.
44. Russo G, Faggiono G, Paganoti GM, Djeunang Dongho GB, Pomponi A, De Santis R, et al. Molecular evidence of *Plasmodium vivax* infection in Duffy negative symptomatic individuals from Dschang, West Cameroon. *Malar J* 2017;16:74.
45. Schuster SC, Miller W, Ratan A, Tomsho LP, et al. Complete Khoisan and Bantu genomes from southern Africa. *Nature* 2010;463:943–947.
46. Tournamille C, Colin Y, Cartron JP, Le Van Kim C. Disruption of a GATA motif in the Duffy gene promoter abolishes erythroid gene expression in Duffy-negative individuals. *Nat Genet* 1995;10:224-228.
47. Twohig KA, Pfeffer DA, Baird JK, Price RN, Zimmerman PA, Hay SI, et al. Growing evidence of *Plasmodium vivax* across malaria endemic Africa. *PLoS Negl Trop Dis* 2019;13: e0007140.
48. Woldearegai TG, Kremsner PG, Kun JF, Mordmuller B. *Plasmodium vivax* malaria in Duffy-negative individuals from Ethiopia. *Trans R Soc Trop Med Hyg* 2013;107:328-31.
49. Zeeshan M, Tyagi RK, Tuagi K, Alam MS, Sharma YD. Host-parasite interaction: selective Pv-fam-a family proteins of *Plasmodium vivax* bind to a restricted number of human erythrocyte receptors. *J Infect Dis* 2015;211:1111-20.

50. Zimmerman PA. *Plasmodium vivax* Infection in Duffy-Negative People in Africa. Am J Trop Med Hyg 2017;97:636-638.

Journal Pre-proof

Tables and Figures

Figure 1. Map showing the distribution of study sites and the Duffy status of febrile patients included in the present study.




Figure 2. Comparison of *P. vivax* parasitemia based on quantitative PCR assays between Duffy negative and Duffy positive symptomatic infections among different geographical regions in Africa. Variations in parasitemia among samples were presented as boxplots showing the median and interquartile range values.




Figure 3. Phylogeny based on *PvDBP* sequences showing multiple source/origin of Duffy negative *P. vivax* in Africa. The reference *P. vivax* strain PVP01 isolated from an Indonesian patient was used as an outgroup. The size of the symbol indicates sample size of each *PvDBP* haplotype. No clear differentiation was observed between the Duffy negative and Duffy positive *P. vivax* but nested within one another, suggestive of similar DBP haplotypes.




Figure 4. Hypothetical models illustrating the genetic origin of *P. vivax* in Duffy-negative Africans in a phylogenetic context. (A) The null hypothesis posits that the ancestral *P. vivax* infected all African primates including apes and Duffy-positive humans forming a monophyletic clade. One of these ancestral lineages evolved to a Duffy-independent

pathway (dotted line) and subsequently spread to different parts of Africa via human migration. (B) An alternative hypothesis posits that the ancestral *P. vivax* infected only non-human primates in Africa until some of the lineages crossed the species barrier and gave rise to the parasite population currently infecting Duffy-positive humans. It is possible that Duffy-negative *P. vivax* observed today across Africa represent separate lineages that were derived multiple times independently from Duffy-positive individuals (dotted line) forming separate monophyletic clades.




Table 1. Summary of *P. vivax* infections with available Duffy blood group information in African countries from the literature.

Table 2. Comparison of *P. vivax* infection rate in Duffy-negative populations across different study sites in Botswana, Ethiopia, and Sudan.

Table 3. Nucleotide and haplotype diversity of *PvDBP* gene sequences between Duffy-negative and Duffy-positive samples from different African countries.

Journal Pre-proof

Table 1. Summary of *P. vivax* infections with available Duffy blood group information in African countries from the literature. nPCR: Nested PCR of *P. vivax* 18S rRNA gene; qPCR: Quantitative real-time PCR of *P. vivax* 18S rRNA gene.

Country	Sample collection year	Symptoms	Sample size	Duffy negative, <i>n</i> (%)	Malaria diagnostic method	<i>Plasmodium spp</i> pos (%)	<i>P. vivax</i> pos (% of <i>P. spp</i> +))	<i>Pv</i> + in Duffy neg (% of total <i>Pv</i> +))	Reference
East-Southern Africa									
Angola	2006-07	No	898 ^	*	nPCR	245/898 (28.9%)	7/245 (2.8%) ¹	7/7 (100%)*	Mendes C, et al. PLoS NTDs, 2011;5(6): e1192.
Ethiopia	2009	Yes	1,931	41/205 (20%) ²	nPCR	205/1,931 (10.6%)	111/205 (54.1%)	3/111 (2.7%)	Woldearegai TG, et al. Trans R Soc Trop Med Hyg 2013;107:328-31.
Ethiopia	2013-14	Yes	416	94/416 (29.7%)	qPCR	331/416 (79.5%)	197/331 (59.5%) ³	2/197 (1%) ⁴	Lo E, et al. Malaria J, 2015;14:84.
Ethiopia	2013-14	No	390	139/390 (35.6%)	qPCR	73/390 (18.7%)	24/73 (32.9%) ⁵	4/24 (16.6%)	Lo E, et al. Malaria J, 2015;14:84.
Kenya	1999-2000	Yes	31 ^ ⁶	31/31 ⁸ (100%)	microscopy	31/31 ⁶ (100%)	11/31 (35.4%)	9/11 (81.8%)	Ryan JR, et al. Am J Trop Med Hyg, 2006;75:575-81.
Madagascar	2006-07	Yes	183 ⁷	*	nPCR	183/183 (100%) ⁷	183/183 (100%) ⁷	17/183 (9.3%)*	Ménard D, et al. PNAS, 2010;107(13):5697-71.
Madagascar	2006-07	No	661 ^	476/661 (72%)	nPCR §	251/661 (38%)	86/251 (34.3%) ⁸	42/86 (48.8%)	Ménard D, et al. PNAS, 2010;107(13):5697-71.
Madagascar	2014	No	2,063	914/1,878 (48.7%)	nPCR	285/2,063 (13.8%) ⁹	137/285 (48.1%) ¹⁰	44/914 (4.8%)	Howes RE, et al. Am. J. Trop. Med. Hyg., 2018;99(4):995-1002.
West-Central Africa									
Benin	2009-10	No	84 ^^	*	nPCR	25/84 (29.8%)	13/25 (52%) ¹¹	13/13 (100%)*	Poirier P, et al., Malar J, 2016;15:570.
Botswana	2012	No	3,624 ²²	N.A.	nPCR	179/3,624 (5%)	169/179 (94.4%)	N.A.	Matshoge T, et al. BMC Inf Dis 2016;16:520.
Cameroon	N.A.	Yes	485	*	nPCR	201/485 (41.4%)	8/201 (4%) ¹²	8/8 (100%)*	Ngassa Mbenda HG & Das A. PLoS ONE 2014;9(8):e103262.
Cameroon	2008-09	No	269	*	nPCR	87/267 (32.3%)	13/87 (14.9%) ¹³	6/13 (46.1%)*	Fru-Cho J, et al. Malaria J, 2014;13:170.
Cameroon	2012-13	Yes	484	224/228 (98.3%) ¹⁵	nPCR	70/484 (14.4%)	27/70 (38.6%) ¹⁵	27/27 (100%)	Russo G et al. Malaria J, 2017;16(1):74.

Democratic Republic of Congo	2013-14	No	292 [^]	*	nPCR	194/292 (66.4%)	14/194 (7.2%)	14/14 (100%)*	Brazeau NF, et al. Am. J. Trop. Med. Hyg., 2018;99(5):1128–33.
Equatorial Guinea	2005	No	97	*	nPCR	84/97 (86.6%)	8/84 (9.5%) ¹⁶	8/8 (100%)*	Mendes C, et al. PLoS NTDs, 2011;5(6):e1192.
Mali	2009-11	No	300 [^]	*	qPCR	135/300 (45%)	25/135 (18.5%)	25/25 (100%)*	Niangaly A, et al. Am J Trop Med Hyg 2017;97(3):744-52.
Mauritania	2007-09	Yes	277	52/258 (20.1%)	qPCR	110/277 (39.7%)	110/110 (100%)	1/110 (0.9%)	Wurtz N et al. Malaria J, 2011;10:336.
Nigeria	2016-17	Yes	436	*	nPCR	256/436 (58.7%)	5/256 (1.9%) ¹⁷	5/5 (100%)*	Oboh MA, et al., Malar J, 2018;17:439 and 2020;19:229.
Senegal	2009-2013	Yes	263	N.A.	nPCR	164/263 (62.3%)	4/164 (2.4%) ²⁴	N.A.	Niang et al., Malar J, 2015; 14: 281.
Senegal	2010-2011	No	48(x4) ^{^18}	48/48 (100%)	nPCR	74/192 (38.5%)	15/74 (20.3%)	5/5 (100%)	Niang M, et al. Trop Med & Hyg 2018;46:45.
Sudan	2009	Yes	126	*	nPCR	N.A.	48/126 (38.1%)	4/48 (8.3%)*	Abdelraheem MH, et al. Trans R Soc Trop Med Hyg 2016;110:258-60
Sudan	2016	Yes	992 ¹⁹	*	microscopy	992/992 (100%) ¹⁹	190/992 (19.1%) ²⁰	34/190 (17.9%)*	Albsheer MMA, et al. Genes, 2019;10:437.
Uganda	2016	Yes	499 ²¹	N.A.	nPCR	499/499 (100%)	4/499 (0.8%) ²³	N.A.	Asua, V. et al. Am J Trop Med Hyg. 2017;97:753-57.

* Duffy-Ag assessed only among *Pv* pos patients; [^] only children; ^{^^} only blood-donors; N.A. not available.

§ Conventional PCR of genes *PvCOI* and *PvDBP*.

¹ 2 *Pf-Pv* co-infections; ² Duffy-Ag available only among *Plasmodium spp* pos; ³ 33 *Pf-Pv* co-infections; ⁴ 2 *Pf-Pv* co-infections; ⁵ 1 *Pf-Pv* co-infection; ⁶ 31 children Duffy neg affected by malaria enrolled in a precedent study (anemia study); ⁷ only *Pv* pos analysed (153 *Pv* mono-infections and 30 *Pf-Pv* co-infections); ⁸ 34 *Plasmodium* mixed-infections (species not specified); ⁹ 42 co-infections (25 *Pf-Pv*, 5 *Pf-Pm*, 9 *Pv-Pm*, 1 *Pv-Po*, 1 *Pf-Pv-Pm*, 1 *Pf-Pv-Pm-Po*); ¹⁰ 37 co-infections (25 *Pf-Pv*, 9 *Pv-Pm*, 1 *Pv-Po*, 1 *Pf-Pv-Pm*, 1 *Pf-Pv-Pm-Po*); ¹¹ 9 *Pf-Pv* and 1 *Pf-Pv-Pm* co-infections; ¹² 2 *Pf-Pv* co-infections; ¹³ 3 *Pf-Pv* and 1 *Pf-Pv-Pm* co-infections; ¹⁴ Duffy-Ag assessed among 228 participants (including all those infected); ¹⁵ 2 *Pf-Pv* co-infections; ¹⁶ 4 *Pf-Pv* co-infections; ¹⁷ 4 *Pf-Pv* co-infections; ¹⁸ 48 school children followed during 2 years (4 samples for each children, 192 total samples analyzed) ¹⁹ only *Plasmodium spp* pos sample analysed; ²⁰ 4 *Pf-Pv* co-infections; ²¹ all *Plasmodium* pos children aged 2 months-10 years; ²² all children aged 2-12 years; ²³ all 4 were co-infections: 3 *Pf-Pv* and 1 *Pv-Pm*; ²⁴ 2 *Pf-Pv* co-infections

Table 2 Comparison of *P. vivax* infection rate in Duffy-negative populations across different study sites in Botswana, Ethiopia, and Sudan based on febrile patient samples collected in this study.

Region	Country	Study site	Collection period	Type of collection	Total samples	Infection rate of <i>P. vivax</i>	Duffy-negative among <i>P. vivax</i> infections
Southern Africa							
	Botswana	Tutume	2017 - 2018	Symptomatic	176	12 (6.8%)	10 (83.3%)
		Kweneng East	2017 - 2018	Symptomatic	301	9 (3%)	8 (88.9%)
East Africa							
	Ethiopia	Jimma	April - October 2017	Symptomatic	358	134 (37.4%)	16 (11.9%)
		Bonga	October – November 2019	Symptomatic	297	76 (25.6%)	8 (10.5%)
	Sudan	River Nile	August 2018 – February 2019	Symptomatic	213	52 (24.4%)	2 (3.8%)
		Khartoum	August 2018 – February 2019	Symptomatic	525	42 (8%)	4 (9.5%)
		New Halfa	August 2018 – February 2019	Symptomatic	93	7 (7.5%)	1 (14.3%)

Table 3 Nucleotide and haplotype diversity of *PvDBP* gene sequences between Duffy-negative and Duffy-positive samples from different African countries.

Region	Country	Duffy	Total samples	Number of polymorphic sites	Nucleotide diversity (SD)	Number of haplotypes	Haplotype diversity (SD)
Central Africa							
	Botswana	Duffy-positive	4	0	0	1	0
		Duffy-negative	4	0	0	1	0
East Africa							
	Ethiopia	Duffy-positive	107	9	1.51×10^{-3} (1.5×10^{-4})	11	0.762 (0.031)
		Duffy-negative	9	9	4.18×10^{-3} (1.0×10^{-4})	4	0.694 (0.147)
	Sudan	Duffy-positive	53	4	3.03×10^{-3} (2.5×10^{-4})	6	0.720 (0.039)
		Duffy-negative	16	17	5.59×10^{-3} (8.3×10^{-4})	8	0.758 (0.110)
	Uganda	-	31	28	6.51×10^{-3} (7.7×10^{-4})	17	0.933 (0.027)
	Madagascar	-	4	6	7.08×10^{-3} (2.0×10^{-4})	3	0.833 (0.222)