

Early IgG / IgA response in hospitalized COVID-19 patients is associated with a less severe disease.

Giorgio Fedele , Gianluca Russo , Ilaria Schiavoni ,
Pasqualina Leone , Eleonora Olivetta , Valentina Perri ,
Maria Antonella Zingaropoli , Maria Rosa Ciardi , Patrizia Pasculli ,
Claudio Maria Mastroianni , Paola Stefanelli

PII: S0732-8893(21)00231-5
DOI: <https://doi.org/10.1016/j.diagmicrobio.2021.115539>
Reference: DMB 115539



To appear in: *Diagnostic Microbiology & Infectious Disease*

Received date: 23 February 2021
Revised date: 24 August 2021
Accepted date: 6 September 2021

Please cite this article as: Giorgio Fedele , Gianluca Russo , Ilaria Schiavoni , Pasqualina Leone , Eleonora Olivetta , Valentina Perri , Maria Antonella Zingaropoli , Maria Rosa Ciardi , Patrizia Pasculli , Claudio Maria Mastroianni , Paola Stefanelli , Early IgG / IgA response in hospitalized COVID-19 patients is associated with a less severe disease., *Diagnostic Microbiology & Infectious Disease* (2021), doi: <https://doi.org/10.1016/j.diagmicrobio.2021.115539>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Early IgG / IgA response in hospitalized COVID-19 patients is associated with a less severe disease.

Giorgio Fedele^{1*}, Gianluca Russo^{3*}, Ilaria Schiavoni¹, Pasqualina Leone¹, Eleonora Olivetta²,
Valentina Perri³, Maria Antonella Zingaropoli³, Maria Rosa Ciardi³, Patrizia Pasculli³, Claudio
Maria Mastroianni³, Paola Stefanelli¹.

*Equal contributors

¹Department Infectious Diseases, Istituto Superiore di Sanità, Italy

²National Centre for Global Health, Istituto Superiore di Sanità, Italy

³Department of Public Health and Infectious Diseases, Sapienza University of Rome, Italy

Corresponding author: Giorgio Fedele, Phone:+39-064990-2890, Fax:+39-064990-3168

giorgio.fedele@iss.it;

Running title: Antibody response and Covid-19 severity

Abstract: 127 words

Main text: 1363 words

Abstract

We determined the kinetics of anti-SARS-CoV-2 antibody response in fifteen hospitalized COVID-19 patients. Patients were divided into mild/moderate (mild, n=1; moderate, n=4) or severe (n=10) and virus-specific anti-Nucleocapsid IgM, anti-Spike IgG and anti-Spike IgA were measured in serial serum samples collected 0-15 days after hospital admission. Surrogate neutralization assays were performed by testing inhibition of ACE-2 binding to Spike. In three patients (2 severe and 1 moderate case), serum antibodies and T-cell memory were monitored six months after baseline. Although IgM response tended to appear first, patients affected by less severe disease were more prone to an early IgG/IgA response. Neutralization of Spike binding to ACE2 correlated with anti-Spike IgG and IgA. IgG and IgA antibody response persisted at the six months follow-up. A recall T-cell response to the Spike antigen was observed in two out of three patients, not related to disease severity.

Keywords: SARS-CoV-2; COVID-19; Serology; Disease Severity; T-cell immunity.

1. Introduction

The immune response against SARS-CoV-2 remains unclear and many caveats in its knowledge are still present. A correlation of clinical severity with high titres of SARS-CoV-2 antibodies, mainly immunoglobulin (Ig)G, was shown (Zhao et al, 2020; Long et al, 2020; Ma et al, 2020) and the importance of neutralizing humoral immunity on disease progression has been shown (Garcia-Beltran et al, 2021). Fewer studies assessed serum SARS-CoV-2 IgA; this Ig subclass becomes detectable 4-6 days after symptom onset with a higher and more persistent serum level than IgM (Padoan et al, 2020; Yu et al, 2020). IgA levels seem significantly higher in severe than in mild/moderate SARS-CoV-2 infections (Yu et al, 2020; Bartsch et al, 2021).

Early antibody responses appear to be correlated with disease outcome (Atyeo et al, 2020). Timing of IgM and IgG seroconversion is controversial, with serum IgM appearing before IgG or synchronously (Zhao et al, 2019; Xiang et al, 2020), or even later (To KKW et al, 2020; Zhang et al, 2020). There is growing evidence of pre-existing humoral and T-cell immunity caused by exposure to common cold coronaviruses (Ng KW et al, 2020; Sette and Crotty, 2020). Immunity to SARS-CoV-2 is also related to T-cell responses (McMahan et al, 2021). Patients who resolve infection have been shown to generate virus-specific T-cell responses (Sekine et al, 2020; Weiskopf et al, 2020). It has been proposed that SARS-CoV-2 induces multifunctional memory T cells which play a role in preventing recurrent episodes of severe SARS-CoV-2 infections (Sekine et al, 2020; Grifoni et al, 2020).

The objective of the study was the characterization of the serological response to SARS-CoV-2 in patients hospitalized with confirmed SARS-CoV-2 infection. We aimed at comparing the kinetics of IgM, IgG, IgA, and virus neutralization activity in a group of patients affected by

mild/moderate or severe SARS-CoV-2 infection. Six months after baseline, SARS-CoV-2 specific antibodies and T-cell responses were measured in a smaller number of individuals.

2. Material and methods

2.1 Clinical sample

Fifteen patients hospitalized with confirmed COVID-19 at the Infectious Diseases division of Policlinico Umberto 1st Hospital, Rome, Italy, were included in the study (Table1). The study was approved by local Ethic Committee and all participants have signed an informed consent. At the hospital admission all patients were affected by COVID-19 related pneumonia and, according to the PaO₂/FiO₂ ratio (partial oxygen pressure/oxygen flow), 10 patients were classified as having a severe disease (PaO₂/FiO₂ < 250), 4 having a moderate disease (PaO₂/FiO₂ > 250) in need of oxygen supply, and one mild disease only (PaO₂/FiO₂ > 250, no need of oxygen supply). Patients were re-tested for SARS-CoV-2 RNA 6 days after hospital admission and then every two days, until a negative NP swab result was obtained.

2.2 Serology

Serum samples were available at three different time-points: at hospital admission (T0), 5-10 days (T1), and 11-15 days (T2) after hospital admission. Immediately after collection, sera were shipped to the Department of Infectious Diseases, Istituto Superiore di Sanità (ISS); 0.5 ml aliquots were prepared and stored at -80 °C. Serology testing was performed at ISS by personnel blinded to specimen classification within one month from serum collection. ELISA assays were used to measure Anti-Nucleocapsid IgM (Dia-Pro diagnostics Bioprobes, Sesto San Giovanni (MI), Italy), anti-Spike IgG and anti-Spike IgA (both from Euroimmun, Lübeck, Germany), according to manufacturers' instructions. A surrogate neutralization assay (GenScript, Leiden, The Netherlands) was used to detect the virus neutralizing ability

of patients' sera. The assay detects and measures neutralizing antibodies blocking the Spike RBD-ACE2 interaction *in vitro* and was used according to manufacturer's instructions.

2.3 T-cell response

Additional blood samples from patients #4, #11 and #15 were collected six months after hospital discharge and humoral and T-cell immune responses to SARS-CoV-2 assessed. Since the second wave of Covid-19 cases hit the Policlinico Umberio 1st hospital at the end of September 2020, it was not possible to include in the 6-month follow-up study other patients, due to safety concerns. Virus-specific T-cell responses were measured by stimulating patients' peripheral blood mononuclear cells (PBMCs) with overlapping peptides covering the immunodominant sequence domains of the Spike protein (Miltenyi, Bergisch Gladbach, Germany). After overnight stimulation, cells were incubated with Live/Dead fixable violet dead cell stain kit used to exclude dead cells from the analyses (Thermo Fisher Scientific, Waltham, MA, USA). Cells were then fixed in paraformaldehyde 2%, permeabilized using a solution containing saponine 0.5% and stained with a predetermined optimal concentration of fluorochrome-conjugated Abs: anti-CD3-Horizon-BV510, anti-IL-2-FITC, anti-TNF α PE-Cy7 (all from BD Biosciences, Franklin Lakes, NJ, USA), anti-IL-4-APC, anti-IFN- γ -PerCP-Cy5.5 (both from Biolegend, San Diego, CA, USA), anti-CD8-PE (eBiosciences, Thermo Fisher Scientific). Cells were then acquired by flow cytometry in a Gallios Flow Cytometer (Beckman Coulter, Indianapolis, IN, USA) and compensation matrices were calculated with Kaluza Analysis software (Beckman Coulter).

3. Results

3.1 Clinical sample

Fifteen patients were included in the study (Table 1). Patients with mild/moderate SARS-CoV-2 infection had a lower mean age (59.0 vs 66.8; $p=0.27$) and higher median Ct values of PCR from nasopharyngeal (NP) samples (gene E: 32.10 vs 25.35, $p=0.05$; gene S: 31.00 vs 24.65, $p=0.058$) than those with severe infection. As expected, severely ill patients required more days for a negative result of naso-pharyngeal swab (median 16.0 vs 13.0; $p=0.29$).

3.2 Anti-SARS-CoV-2 IgM, IgG, IgA antibody responses

Serum samples were collected at hospital admission (T0), 5-10 days (T1), and 11-15 days (T2) after hospital admission. At T0, IgM specific for the viral nucleocapsid protein were above the positivity threshold in 9 patients (60.0 %). At T1, a positive IgM response was measured in 12 patients (80.0 %); at T2, 13 patients (86.7%) had a positive IgM response. The IgM response showed a slightly higher median value among severely affected patients at T1 and lower at T2 (Figure 1).

Anti-Spike IgG at T0 were above the positivity threshold in 5 patients (13.3 %). At T1 a positive IgG response was measured in 13 patients (86.7 %); at T2 14 patients (93.3%) had a positive IgG response. When patients were classified according to disease severity, we found that at T0 the median IgG response was higher in mild/moderate patients, although this difference was not statistically significant. IgG levels were higher among severely affected patients at T1 and T2 (Figure 1).

Anti-Spike IgA at T0 were above the positivity threshold in 11 out of 15 patients (73.3 %). At T1 and T2 a positive IgA response was measured in all the patients. Higher IgA levels were found at T0 in patients with a mild/moderate disease (Figure 1).

3.3 SARS-CoV-2 neutralization correlates with IgG and IgA serum levels in laboratory-confirmed COVID-19 patients

At hospital admission Spike RBD-ACE2 neutralization activity was measured in 13 patients (86.7 %). At T1 and T2 a neutralizing response was measured in all the patients. Patients with mild/moderate disease presented higher neutralizing antibody levels at hospital admission as compared to severe patients, although this was not statistically significant (Figure 1).

A statistically significant positive correlation of *in vitro* neutralization of Spike RBD-ACE2 binding with IgG and IgA serum levels at all the time-points was found, whereas no correlation was evident between anti-Nucleocapsid IgM and Spike RBD-ACE2 neutralizing activity (Figure 2).

3.4 Persistence of anti-SARS-CoV2 immunity six months after the infection

A 6-month follow-up analysis was performed on 3 out of 15 patients. We did not observe any decline in anti-Spike IgG levels (T2 mean \pm SD: 6.53 \pm 0.27; 6-month mean \pm SD: 6.21 \pm 0.9). Anti-Spike IgA levels declined but were still above the positivity threshold (T2 mean \pm SD: 11.70 \pm 0.56; 6-month mean \pm SD: 4.13 \pm 2.16). Sera retained surrogate neutralization activity (data not shown). Two patients (one moderate and one severe case) were able to mount a virus-specific T-cell response, marked by the production of TNF- α and IFN- γ by CD4 and CD8 T cells (Figure 3), while one patient was unresponsive. Profiles of T-cell response were rather different. A higher frequency of cytokine positive CD4⁺ vs. CD8⁺ T cells (0.91% vs 0.56%) was found in patient #10 (moderate), while in patient #15 (severe) this ratio was inverted (0.78% vs 0.96%). The proportion of CD4 and CD8 cells producing one, two or three cytokines was also different in the two responsive patients.

4. Discussion

The correlation of antibody responses to SARS-CoV-2 with COVID-19 disease severity is still unclear. Our data suggest that individuals with an earlier appearance of virus-specific IgG

and IgA potentially have a better outcome. In patients with a mild/moderate disease, IgG responses tended to occur simultaneously with IgM or even slightly earlier, while in patients with severe disease IgG and IgA were produced later during the course of the disease. Both IgG and IgA levels were correlated to serum neutralizing activity.

Our observations agree with a recent study by Carsetti and co-workers (2020) who found that IgA and IgG produced relatively late in the course of the infection characterize severe disease (Carsetti et al, 2020). Moreover, it has been shown that patients with more severe illness display higher antibody titers than those with milder disease (Long et al, 2020; Röltgen et al, 2020).

The presence of anti-SARS-CoV-2 IgG and IgA at T0 in mild/moderate patients suggests that some pre-existing immunity to the four endemic coronaviruses causing common cold in humans (HCoV-229E, -NL63, -OC43, and -HKU1) may play a role in counteracting the infection. These findings are supported by a recent study showing that COVID-19 disease appears less severe in patients with endemic coronavirus infection (Sagar et al, 2021).

5. Limitations

Before drawing conclusions, limitations of the study should be mentioned. The number of patients enrolled is small and probably not sufficient for making strong conclusions. Only 3 patients were used in the cellular immunity studies at 6 months. Although incomplete and preliminary, the 6-month follow-up showed the persistence of serological response in three patients. Data on T-cell response suggested the possibility of differential responses that could reflect asymptomatic exposure.

6. Conclusion

In conclusion, our data confirm the notion that an early IgG/IgA response focused on the Spike protein may help to constrain the viral infection, leading to a milder course of the disease. Further studies are required to investigate more in depth the duration and patterns of T-cell memory responses.

author statement

Giorgio Fedele: conceptualization, data curation, writing-original draft preparation and revised version; Gianluca Russo: conceptualization, data curation, writing-original draft preparation and revised version; Ilaria Schiavoni: investigation and Writing-Reviewing; Pasqualina Leone: investigation; Eleonora Olivetta: investigation; Valentina Perri, Maria Antonella Zingaropoli, Maria Rosa Ciardi and Patrizia Pasculli: resources; Claudio Maria Mastroianni: supervision; Paola Stefanelli: conceptualization and supervision.

Acknowledgements

The authors wish to thank the ISS working group on COVID-19: Stefano Fiore, Concetta Fabiani, Eleonora Benedetti, Angela Di Martino, Laura Calzoletti, Marzia Facchini, Giuseppina Di Mario, Giulietta Venturi, Claudia Fortuna, Antonello Amendola, Giulia Marsili.

Founding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors have nothing to disclose.

Sample CRediT author statement

Giorgio Fedele: conceptualization, data curation and writing-original draft preparation;
Gianluca Russo: conceptualization, data curation and writing-original draft preparation; Ilaria Schiavoni: investigation and Writing-Reviewing; Pasqualina Leone: investigation; Eleonora Olivetta: investigation; Valentina Perri, Maria Antonella Zingaropoli, Maria Rosa Ciardi and Patrizia Pasculli: resources; Claudio Maria Mastroianni: supervision; Paola Stefanelli: conceptualization and supervision.

References

1. Atyeo C, Fischinger S, Zohar T, Slein MD, Burke J, Loos C, et al. Distinct Early Serological Signatures Track with SARS-CoV-2 Survival. *Immunity*. 2020 Sep 15;53(3):524-532.e4. doi: 10.1016/j.immuni.2020.07.020.
2. Bartsch YC, Wang C, Zohar T, Fischinger S, Atyeo C, Burke JS, et al. Humoral signatures of protective and pathological SARS-CoV-2 infection in children. *Nat Med*. 2021 Feb 12. doi: 10.1038/s41591-021-01263-3.
3. Carsetti R, Zaffina S, Piano Mortari E, Terreri S, Corrente F, Capponi C, et al. Different Innate and Adaptive Immune Responses to SARS-CoV-2 Infection of Asymptomatic, Mild, and Severe Cases. *Front Immunol* 2020; 11:610300. [https://doi: 10.3389/fimmu.2020.610300](https://doi.org/10.3389/fimmu.2020.610300).
4. Garcia-Beltran WF, Lam EC, Astudillo MG, Yang D, Miller TE, Feldman J, et al. COVID-19-neutralizing antibodies predict disease severity and survival. *Cell*. 2021 Jan 21;184(2):476-488.e11. doi: 10.1016/j.cell.2020.12.015.

5. Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Rydyznski Moderbacher C, et al. Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed Individuals. *Cell* 2020; 181(7):1489-1501.e15. [https://doi: 10.1016/j.cell.2020.05.015](https://doi.org/10.1016/j.cell.2020.05.015).
6. Long QX, Tang XJ, Shi QL, Li Q, Deng HJ, Yuan J, et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med* 2020; 26(8):1200-1204. [https://doi: 10.1038/s41591-020-0965-6](https://doi.org/10.1038/s41591-020-0965-6).
7. Ma H, Zeng W, He H, Zhao D, Jiang D, Zhou P, et al. Serum IgA, IgM, and IgG responses in COVID-19. *Cellular & Molecular Immunology* 2020; 17(7):773-775. [https://doi: 10.1038/s41423-020-0474-z](https://doi.org/10.1038/s41423-020-0474-z).
8. McMahan K, Yu J, Mercado NB, Loos C, Tostanoski LH, Chandrashekar A, et al. Correlates of protection against SARS-CoV-2 in rhesus macaques. *Nature*. 2021 Feb;590(7847):630-634. doi: 10.1038/s41586-020-03041-6.
9. Ng KW, Faulkner N, Cornish GH, Rosa A, Harvey R, Hussain S, et al. Preexisting and de novo humoral immunity to SARS-CoV-2 in humans. *Science* 2020; 370(6522):1339-1343. [https://doi: 10.1126/science.abe1107](https://doi.org/10.1126/science.abe1107).
10. Padoan A, Sciacovelli L, Bassl D, Negrini D, Zuin S, Cosma C, et al. IgA-Ab response to spike glycoprotein of SARS-CoV-2 in patients with COVID-19: A longitudinal study. *Clin Chim Acta* 2020; 507:164-166. [https://doi: 10.1016/j.cca.2020.04.026](https://doi.org/10.1016/j.cca.2020.04.026).
11. Röltgen K, Powell AE, Wirz OF, Stevens BA, Hogan CA, Najeeb J, et al. Defining the features and duration of antibody responses to SARS-CoV-2 infection associated with disease severity and outcome. *Sci Immunol* 2020; 5(54):eabe0240. [https://doi: 10.1126/sciimmunol.abe0240](https://doi.org/10.1126/sciimmunol.abe0240).

12. Sagar M, Reifler K, Rossi M, Miller NS, Sinha P, White LF, et al. Recent endemic coronavirus infection is associated with less-severe COVID-19. *J Clin Invest* 2021; 131(1):e143380. <https://doi: 10.1172/JCI143380>.
13. Sekine T, Perez - Potti A, Rivera - Ballesteros O, Stralin K, Gorin JB, Olsson A, et al. Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID - 19. *Cell* 2020; 183(1): 158–168.e14. <https://doi: 10.1016/j.cell.2020.08.017>.
14. Sette A, Crotty S. Pre-existing immunity to SARS-CoV-2: the knowns and unknowns *Nat Rev Immunol* 2020; 20(8):457-458. <https://doi: 10.1038/s41577-020-0389-z>.
15. To KKW, Tsang OTY, Leung WS, Tam AR, Wu TC, Lung DC, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis.* 2020; 20(5):565-574. [https://doi: 10.1016/S1473-3099\(20\)30196-1](https://doi: 10.1016/S1473-3099(20)30196-1).
16. Weiskopf D, Schmitz KS, Raadsen MP, Grifoni A, Okba NMA, Endeman H, et al. Phenotype of SARS-CoV-2-specific T-cells in COVID-19 patients with acute respiratory distress syndrome. *Sci Immunol* 2020; 5(48):eabd2071. <https://doi: 10.1126/sciimmunol.abd2071>.
17. Xiang F, Wang X, He X, Peng Z, Yang B, Zhang J, et al. Antibody detection and dynamic characteristics in patients with COVID-19. *Clin Infect Dis* 2020; 71(8):1930-1934. <https://doi: 10.1093/cid/ciaa461>.
18. Yu HQ, Sun BQ, Fang ZF, Zhao JC, Liu XY, Li YM, et al. Distinct features of SARS-CoV-2-specific IgA response in COVID-19 patients. *Eur Respir J* 2020; 56(2):2001526. <https://doi:10.1183/13993003.01526-2020>.

19. Zhang W, Du RH, Li B, Zheng XS, Yang XL, Hu B, et al. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerging Microbes & Infections* 2020; 9(1): 386-89.
<https://doi:10.1080/22221751.2020.1729071>.
20. Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody response to SARS-CoV-2 in patients of novel coronavirus diseases 2019. *Clin Infect Dis* 2020; 71(16):2027-2034. [https://doi: 10.1093/cid/ciaa344](https://doi:10.1093/cid/ciaa344).

LEGEND TO THE FIGURES

Figure1. SARS-CoV-2 specific antibody response in mild/moderate and severe COVID-19 patients. Serological anti-Nucleocapsid IgM, anti-Spike IgG and anti-Spike IgA antibodies were tested by enzyme-linked immunosorbent assay. Sera neutralization activity was assessed by surrogate virus neutralization test based on antibody-mediated blockage of ACE2-Spike protein-protein interaction. Solid horizontal lines represent Mean. Dotted lines represent positivity thresholds. T0: hospital admission; T1: 5-10 days after hospital admission; T2: 11-15 days after hospital admission; S/Co: Signal to Cut-off (S/Co) ratio.




Figure 1. Fedele et al

Figure2. Serum IgG and IgA levels correlate with inhibition of SARS-CoV-2-ACE2 binding.

Linear regression correlating serum levels of anti-Nucleocapsid IgM (top), anti-Spike IgG (middle) and, anti-Spike IgA (bottom) antibodies with Spike-ACE2 binding neutralization; numerical values along horizontal axis represent Signal to Cut-off ratios for each serological assay. Dotted lines represent 95% confidence band. P values are shown.




Figure 2. Fedele et al

Figure 3. SARS-CoV-2 specific T-cell response six months after recovery. A) Dot plots show the percentage of CD4⁺ and CD8⁺ T cells producing IFN- γ , TNF- α and IL-2 in response to a pool of overlapping peptides covering the immunodominant sequence domains of Spike protein. B) Frequencies of mono- and multifunctional specific CD4⁺ and CD8⁺ T-cell responses.




Figure 3. Fedele et al

Table 1. Demographic and clinical characteristics of enrolled subjects

Patient Nr.	Sex	Age, years	Comorbidities	Days from symptoms onset to hospital admission	Clinical severity	Oxygen supply during hospital stay	CT value E gene	CT value S gene	Days to SARS-CoV-2 negative NP swab	Length of hospital stay, days
1	F	64	AI, DM	10	Mild	None	32.9	32.2	9	19
2	M	48	NHL	2	Moderate	VM	29.8	28.9	14	16
3	M	83	AI, IHD	7	Moderate	VM	28.2	27.1	17	35
4	M	42	DM	6	Moderate	VM	32.6	34.2	13	19
5	M	58	AI	13	Moderate	VM	32.1	31.0	12	19
6	M	62	None	6	Severe	HFNC/CPAP	22.2	21.2	NA	27
7	M	78	Lung cancer	1	Severe	HFNC/CPAP	14.9	14.8	NA	28
8	M	64	None	10	Severe	HFNC/CPAP	23.8	23.2	25	28
9	F	60	AI	4	Severe	HFNC/CPAP	20.3	19.3	17	21
10	M	69	None	3	Severe	HFNC/CPAP	31.0	32.0	12	14
11	M	52	None	3	Severe	HFNC/CPAP	23.6	22.9	25	30

12	M	78	AI	8	Severe	HFNC/CPAP	35.7	32.2	15	47
13	M	66	None	5	Severe	HFNC/CPAP	27.4	28.1	11	21
14	M	76	AI, COPD, HT	12	Severe	HFNC/CPAP	26.9	26.1	25	38
15	F	63	None	7	Severe	HFNC/CPAP	31.0	30.2	12	19

AI: Arterial hypertension; NHL: Non-Hodgkin Lymphoma; COPD: Chronic Obstructive Pulmonary Disease; IHD: Ischemic Heart Disease; DM: Diabetes Mellitus; HT: Hypothyroidism; HFNC: High Flux Nasal Cannula; CPAP: Continuous Positive Airway Pressure; VM: Venturi Mask; NA: not-applicable (patients died).

Journal Pre-proof