

Regeneration of Trabecular Meshwork in Primary Open Angle Glaucoma by Stem Cell Therapy

Enzo Maria Vingolo, Ayoub Chabib*

Department of Sense Organs, U.O.C. Ophthalmology, Sapienza University of Rome, Latina, Italy, Email: enzomaria.vingolo@uniroma1.it

ABSTRACT

Glaucoma is considered a neurodegenerative disease due to its pathophysiology and is one of the most common causes of vision loss worldwide. The treatment of glaucoma is essentially based on the reduction of IOP, through pharmacological therapies. The iPSCs enabling autologous transplantation and have the advantage of not generating immune-mediated rejection in the host, that iPSC can be induced to differentiate into a cell type that resembles to TM cells, The iPSC-TM. The data demonstrate that intraocular injection of iPSC-TM prevents the IOP elevation, the aqueous humor outflow reduction and results in preservation of Retinal Ganglion Cell (RGC) density in vivo mice models.

Key Words: Trabecular-meshwork; Glaucoma; POAG; IOP

Correspondence:

Chabib A, Department of Sense Organs,
U.O.C. Ophthalmology, Sapienza
University of Rome, Italy

E-mail: ayoub.chabib@gmail.com

INTRODUCTION

Glaucoma is considered a neurodegenerative disease due to its pathophysiology and is one of the most common causes of vision loss worldwide. The glaucoma disease causes death of Retinal Ganglion Cells (RGCs), as well as degeneration of the optic nerve head, the optic nerve, and the lateral geniculate nucleus, which leads to a gradual loss of vision. The most common type of glaucoma is the Primary Open Angle Glaucoma (POAG), which accounts for about 90% of all forms of glaucoma. The treatment of glaucoma is essentially based on the reduction of Intra Ocular Pressure (IOP), through pharmacological therapies, or in more severe cases through surgical procedures, which provide an aqueous humor outflow bypass [1].

STEM CELL OVERVIEW

The stem cells can be classified into Embryonic Stem (ES) cells, induced Pluripotent Stem Cells (iPSCs), and adult stem cells. One advantage of iPSCs is that these cells can be derived from readily available cell types of the intended recipient. The iPSCs enabling autologous transplantation and there by iPSCs are cells that have the advantage of not generating immune-mediated rejection in the host. The iPSCs also have the advantage circumventing the ethical and immunological disadvantages associated with ES cells therapy, since the iPSCs are autologous. However, there are safety concerns because the iPSCs may cause oncogenesis [2-5].

Structure, function and embryology of TM

The Trabecular Meshwork (TM) formation involves many genetic networks. Furthermore, the development of this complex tissue involves different molecular signals, among others PITX2, PITX3, PAX6, FOXC1, FOXE3, LMX1B and MAF. These specific gene regulatory networks are involved in tissue developments and include many transcription factors and molecular signals. The PAX6 is the most important eye development regulatory in a different organisms [6-8]. Numerous studies have noted that TM was abnormally formed in PITX2 and FOXC1 mice [9-12]. The LMX1B gene was shown to have an important role into the dysgenesis of the TM [13]. The heterozygous deficiency of BMP4 may cause an absent or hypo plastic TM or Schlemm's canal [14].

Functions and biological features of TM cells

TM cells constitute the proximal portion of the aqueous humor outflow pathway. To assure the effective outflow resistance regulation, the

TM tissue has an important function of biological filter self-cleaning, intercepting cellular debris and reducing the Reactive Oxygen Species (ROS). The TM cells have a macrophage-like activity to eliminate the cellular debris derived from the pigmented epithelia. The TM cells integrity is important for the regulation and the maintenance of homeostatic IOP. The TM cells dysfunction may cause an extra resistance that generates an IOP elevation [15].

After dexamethasone treatment the expression of myocilin by TM cells increases. The myocilin expression plays an important role in glucocorticoid-induced ocular hypertension [16-18]. Also the platelet-derived growth factor may increase cells division in TM. Besides, it enhances the phagocytic activity and promotes Extra-Cellular Matrix (ECM) secretion [19]. On the contrary, TM cells growth can be inhibited by the vascular endothelial cell growth factor [20].

TM cells loss and glaucoma

The TM cells decrease with age. This cells reduction is associated with glaucoma. Also the accumulation of ECM is associated with gap junction alteration and leading to TM cells death and IOP elevation. Moreover, the TM cells present damages probably caused by ROS [21,22]. This causes a diminished ability to drain humor aqueous.

Evidence of therapeutic implications for TM stem-cells

In 1982, unusual cells population located just beneath the Schwalbe's line in Rhesus monkeys was discovered by Raviola [23]. Afterwards, an increased TM cell division after Argon Laser Trabeculoplasty (ALT) in the anterior non-filtering portion of the TM [24].

Gonzalez, et al. noted that cultured TM cells can form the free-floating neuro spheres. This function is associated with neural stem cells. When the TM free-floating spheres were incubated, they evolved into cells morphologically indistinguishable from cultured TM cells. This discover,

This is an open access article distributed under the terms of the Creative Commons Attribution Noncommercial Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: invoice@jbclinpharm.org

Cite this article as: Chabib A, Vingolo EM. Regeneration of Trabecular Meshwork in Primary Open Angle Glaucoma by Stem Cell Therapy. J Basic Clin Phar. 2021;12:90-91.

indicating the possibility that they might differentiate spontaneously into TM cells [25].

The stem cells therapy can potentially restore the TM function and protect the optic nerve from the damage [26]. Replacing the damaged TM cells with healthy stem cells may restore the micro environment of filtering structures. This could trigger a reparative proliferation of stem cells and a restoration of the physiological aqueous outflow with a IOP reduction.

Du, et al. described the isolation of a stem cells population from human TM. This cells population in culture displaying homogeneous antigenic markers characterized for mesenchymal stem cells and expressing gene products associated with pluripotent stem cells. These cells are capable of differentiating into TM cells with TM markers [27,28].

Zhu, et al. demonstrate that iPSC can be induced to differentiate into a cell type that resembles to TM cells (iPSC-TM), The iPSC-TM responds to glucocorticoids exposure with myocilin secretion. For this purpose mouse iPSC-TM cells were induced from iPSC derived from fibroblasts isolated from transgenic animals. The data demonstrate that intraocular injection of iPSC-TM prevents the IOP elevation, the aqueous humor outflow reduction and results in preservation of RGC density in vivo mice models [29]. Moreover, Abu-Hassan, et al. observed that iPSC-TM cells can restore IOP in a human anterior segment ex vivo model [30].

CONCLUSION

The evidence reveals that there is a stem-like cells population located in the Schwalbe's ring. The increased TM cells division after ALT suggested the repopulation of the TM by stimulating cell division may restore the IOP. Recent progress in stem cell research provides an optimistic prospect on their use in regenerative medicine and tissue engineering.

CONFLICT OF INTEREST

Enzo M. Vingolo, Ayoub Chabib, declares that they have no conflict of interest.

REFERENCES

1. Fan BJ, Wiggs JL. Glaucoma: Genes, phenotypes, and new directions for therapy. *J Clin Invest.* 2010; 120(9):3064-3072.
2. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell J.* 2006; 126(4):663-676.
3. Stadtfeld M, Nagaya M, Utikal J, et al. Induced pluripotent stem cells generated without viral integration. *SCIEAS.* 2008; 322(5903):945-949.
4. Zhou H, Wu S, Joo JY, et al. Generation of induced pluripotent stem cells using recombinant proteins. *Cell stem cell.* 2009; 4(5):381-384.
5. Lin T, Ambasudhan R, Yuan X, et al. A chemical platform for improved induction of human iPSCs. *Nat methods.* 2009;6(11):805-808.
6. Cvekl A, Tamm ER. Anterior eye development and ocular mesenchyme: New insights from mouse models and human diseases. *Bioessays.* 2004;26(4):374-386.
7. Gehring WJ, Ikeo K. Pax 6: Mastering eye morphogenesis and eye evolution. *Trends Genet.* 1999;15(9):371-377.
8. Chow RL, Lang RA. Early eye development in vertebrates. *Annu Rev Cell Dev Biol.* 2001;17(1):255-296.
9. Kitamura K, Miura H, Miyagawa-Tomita S, et al. Mouse PITX2 deficiency leads to anomalies of the ventral body wall, heart, extra- and pericocular mesoderm and right pulmonary isomerism. *Development.* 1999;126(24):5749-5758.
10. Kidson SH, Kume T, Deng K, et al. The forkhead/winged-helix gene, Mf1, is necessary for the normal development of the cornea and formation of the anterior chamber in the mouse eye. *Devel Biol.* 1999;211(2):306-322.
11. Evans AL, Gage PJ. Expression of the homeobox gene PITX2 in neural crest is required for optic stalk and ocular anterior segment development. *Hum Mol Genet.* 2005; 14(22):3347-59.
12. Sowden JC. Molecular and developmental mechanisms of anterior segment dysgenesis. *Eye.* 2007;21(10):1310-1318.
13. Liu P, Johnson RL. LMX1B is required for murine trabecular meshwork formation and for maintenance of corneal transparency. *Dev Dyn.* 2010;239(8):2161-2171.
14. Chang B, Smith RS, Peters M, et al. Haploinsufficient BMP4 ocular phenotypes include anterior segment dysgenesis with elevated intraocular pressure. *BMC Genet.* 2001;2(1):1-2.
15. Lütjen-Drecoll E. Functional morphology of the trabecular meshwork in primate eyes. *Prog Retin Eye Res.* 1999;18(1):91-119.
16. Stamer WD, Seftor RE, Snyder RW, et al. Cultured human trabecular mesh work cells express aquaporin-1 water channels. *Cur eye res.* 1995;14(12):1095-1100.
17. Chang IL, Elner G, Yue YJ, et al. Expression of modified low-density lipoprotein receptors by trabecular meshwork cells. *Cur eye res.* 1991;10(12):1101-1112.
18. Gasiorowski JZ, Russell P. Biological properties of trabecular meshwork cells. *Exp Eye Res.* 2009;88(4):671-5.
19. Tamura M, Iwamoto Y. The effect of platelet-derived growth factor on phagocytosis of cultured human trabecular cells. *Exp Eye Res.* 1989;48(6):761-770.
20. Ando A, Kaneko SH, Matsumura M. The effect of growth factors on the proliferation of cultured porcine trabecular meshwork cells. *Nippon Ganka Gakkaï Zasshi.* 2004;108(9):549-553.
21. Chhunchha B, Singh P, Stamer WD, et al. PRDX6 retards senescence and restores trabecular meshwork cell health by regulating reactive oxygen species. *Cell Death Discov.* 2017;3(1):1-2.
22. Saccà SC, Izzotti A, Rossi P, et al. Glaucomatous outflow pathway and oxidative stress. *Exp Eye Res.* 2007;84(3):389-399.
23. Raviola G. Schwalbe line's cells: A new cell type in the trabecular meshwork of Macaca mulatta. *Invest ophthalmol vis sci.* 1982;22(1):45-56.
24. Acott TS, Samples JR, Bradley JM, et al. Trabecular repopulation by anterior trabecular meshwork cells after laser trabeculoplasty. *Am J Ophthalmol.* 1989;107(1):210-216.
25. Gonzalez P, Epstein DL, Luna C, et al. Characterization of free-floating spheres from Human Trabecular Meshwork (HTM) cell culture in vitro. *Exp Eye Res.* 2006;82(6):959-967.
26. Chamling X, Sluch VM, Zack DJ. The potential of human stem cells for the study and treatment of glaucoma. *Invest ophthalmol vis sci.* 2016;57(5):1-6.
27. Du Y, Roh DS, Mann MM, et al. Multipotent stem cells from trabecular meshwork become phagocytic TM cells. *Invest ophthalmol vis sci.* 2012;53(3):1566-1575.
28. Du Y, Yun H, Yang E, et al. Stem cells from trabecular meshwork home to TM tissue in vivo. *Invest ophthalmol vis sci.* 2013;54(2):1450-1459.
29. Zhu W, Gramlich OW, Laboissonniere L, et al. Transplantation of iPSC-derived TM cells rescues glaucoma phenotypes in vivo. *Proc Natl Acad Sci.* 2016;113(25):3492-3500.
30. Abu-Hassan DW, Li X, Ryan EI, et al. Induced pluripotent stem cells restore function in a human cell loss model of open-angle glaucoma. *Stem cells.* 2015;33(3):751-761.