

PAPER • OPEN ACCESS

## Results from the CUORE experiment

To cite this article: D Chiesa and for the CUORE Collaboration 2019 *J. Phys.: Conf. Ser.* **1137** 012052

View the [article online](#) for updates and enhancements.

You may also like

- [Theory of neutrinoless double-beta decay](#)  
J D Vergados, H Ejiri and F Šimković
- [Improved background rejection in neutrinoless double beta decay experiments using a magnetic field in a high pressure xenon TPC](#)  
J. Renner, A. Cervera, J.A. Hernando et al.
- [Progress Toward A 2 Measurement For The Majorana Demonstrator](#)  
T Gilliss, N Abgrall, S I Alvis et al.



The Electrochemical Society  
Advancing solid state & electrochemical science & technology

### 241st ECS Meeting

May 29 – June 2, 2022 Vancouver • BC • Canada

Extended abstract submission deadline: Dec 17, 2021

Connect. Engage. Champion. Empower. Accelerate.  
**Move science forward**



**Submit your abstract**



## Results from the CUORE experiment

D Chiesa<sup>1,2</sup>, D Q Adams<sup>1</sup>, C Alduino<sup>3</sup>, K Alfonso<sup>4</sup>,  
 F T Avignone III<sup>3</sup>, O Azzolini<sup>5</sup>, G Bari<sup>6</sup>, F Bellini<sup>7,8</sup>, G Benato<sup>9</sup>,  
 A Bersani<sup>10</sup>, M Biassoni<sup>2</sup>, A Branca<sup>11,12</sup>, C Brofferio<sup>1,2</sup>, C Bucci<sup>13</sup>,  
 A Caminata<sup>10</sup>, A Campani<sup>14,10</sup>, L Canonica<sup>15,13</sup>, X G Cao<sup>16</sup>,  
 S Capelli<sup>1,2</sup>, L Cappelli<sup>13,9,17</sup>, L Cardani<sup>8</sup>, P Carniti<sup>1,2</sup>, N Casali<sup>8</sup>,  
 L Cassina<sup>1,2</sup>, N Chott<sup>3</sup>, M Clemenza<sup>1,2</sup>, S Copello<sup>18,13</sup>, C Cosmelli<sup>7,8</sup>,  
 O Cremonesi<sup>2</sup>, R J Creswick<sup>3</sup>, J S Cushman<sup>19</sup>, A D'Addabbo<sup>13</sup>,  
 D D'Aguzzo<sup>13,20</sup>, I Dafinei<sup>8</sup>, C J Davis<sup>19</sup>, S Dell'Oro<sup>21</sup>,  
 M M Deninno<sup>6</sup>, S Di Domizio<sup>14,10</sup>, V Dompè<sup>13,18</sup>, A Drobizhev<sup>9,17</sup>,  
 D Q Fang<sup>16</sup>, M Faverzani<sup>1,2</sup>, E Ferri<sup>1,2</sup>, F Ferroni<sup>7,8</sup>, E Fiorini<sup>2,1</sup>,  
 M A Franceschi<sup>22</sup>, S J Freedman<sup>17,9,a</sup>, B K Fujikawa<sup>17</sup>, A Giachero<sup>1,2</sup>,  
 L Gironi<sup>1,2</sup>, A Giuliani<sup>23</sup>, L Gladstone<sup>15</sup>, P Gorla<sup>13</sup>, C Gotti<sup>1,2</sup>,  
 T D Gutierrez<sup>24</sup>, K Han<sup>25</sup>, K M Heeger<sup>19</sup>, R Hennings-Yeomans<sup>9,17</sup>,  
 R G Huang<sup>9</sup>, H Z Huang<sup>4</sup>, J Johnston<sup>15</sup>, G Keppel<sup>5</sup>,  
 Yu G Kolomensky<sup>9,17</sup>, A Leder<sup>15</sup>, C Ligi<sup>22</sup>, Y G Ma<sup>16</sup>, L Marini<sup>9,17</sup>,  
 M Martinez<sup>7,8,26</sup>, R H Maruyama<sup>19</sup>, Y Mei<sup>17</sup>, N Moggi<sup>27,6</sup>,  
 S Morganti<sup>8</sup>, S S Nagorny<sup>13,18</sup>, T Napolitano<sup>22</sup>, M Nastasi<sup>1,2</sup>,  
 C Nones<sup>28</sup>, E B Norman<sup>29,30</sup>, V Novati<sup>23</sup>, A Nucciotti<sup>1,2</sup>, I Nutini<sup>13,18</sup>,  
 T O'Donnell<sup>21</sup>, J L Ouellet<sup>15</sup>, C E Pagliarone<sup>13,20</sup>, M Pallavicini<sup>14,10</sup>,  
 V Palmieri<sup>5,a</sup>, L Pattavina<sup>13</sup>, M Pavan<sup>1,2</sup>, G Pessina<sup>2</sup>, C Pira<sup>5</sup>,  
 S Pirro<sup>13</sup>, S Pozzi<sup>1,2</sup>, E Previtali<sup>2</sup>, A Puiu<sup>1,2</sup>, F Reindl<sup>8</sup>, C Rosenfeld<sup>3</sup>,  
 C Rusconi<sup>3,13</sup>, M Sakai<sup>4</sup>, S Sangiorgio<sup>29</sup>, D Santone<sup>13,31</sup>, B Schmidt<sup>17</sup>,  
 N D Scielzo<sup>29</sup>, V Singh<sup>9</sup>, M Sisti<sup>1,2</sup>, D Speller<sup>19</sup>, L Taffarello<sup>11</sup>,  
 F Terranova<sup>1,2</sup>, C Tomei<sup>8</sup>, M Vignati<sup>8</sup>, S L Wagaarachchi<sup>9,17</sup>,  
 B S Wang<sup>29,30</sup>, H W Wang<sup>16</sup>, B Welliver<sup>17</sup>, J Wilson<sup>3</sup>, K Wilson<sup>3</sup>,  
 L A Winslow<sup>15</sup>, T Wise<sup>19,32</sup>, L Zanotti<sup>1,2</sup>, G Q Zhang<sup>16</sup>,  
 S Zimmermann<sup>33</sup>, and S Zucchelli<sup>27,6</sup>

<sup>1</sup> Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126, Italy

<sup>2</sup> INFN – Sezione di Milano Bicocca, Milano I-20126, Italy

<sup>3</sup> Department of Physics and Astronomy, University of South Carolina, Columbia, SC 29208, USA

<sup>4</sup> Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

<sup>5</sup> INFN – Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020, Italy

<sup>6</sup> INFN – Sezione di Bologna, Bologna I-40127, Italy

<sup>7</sup> Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185, Italy

<sup>8</sup> INFN – Sezione di Roma, Roma I-00185, Italy

<sup>9</sup> Department of Physics, University of California, Berkeley, CA 94720, USA

<sup>10</sup> INFN – Sezione di Genova, Genova I-16146, Italy

<sup>11</sup> INFN – Sezione di Padova, Padova I-35131, Italy

<sup>12</sup> Dipartimento di Fisica e Astronomia, Università di Padova, I-35131 Padova, Italy

<sup>13</sup> INFN – Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67100, Italy

<sup>14</sup> Dipartimento di Fisica, Università di Genova, Genova I-16146, Italy



<sup>15</sup> Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>16</sup> Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China

<sup>17</sup> Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>18</sup> INFN – Gran Sasso Science Institute, L'Aquila I-67100, Italy

<sup>19</sup> Wright Laboratory, Department of Physics, Yale University, New Haven, CT 06520, USA

<sup>20</sup> Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, Cassino I-03043, Italy

<sup>21</sup> Center for Neutrino Physics, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA

<sup>22</sup> INFN – Laboratori Nazionali di Frascati, Frascati (Roma) I-00044, Italy

<sup>23</sup> CSNSM, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France

<sup>24</sup> Physics Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA

<sup>25</sup> INPAC and School of Physics and Astronomy, Shanghai Jiao Tong University; Shanghai Laboratory for Particle Physics and Cosmology, Shanghai 200240, China

<sup>26</sup> Laboratorio de Física Nuclear y Astroparticulas, Universidad de Zaragoza, Zaragoza 50009, Spain

<sup>27</sup> Dipartimento di Fisica e Astronomia, Alma Mater Studiorum – Università di Bologna, Bologna I-40127, Italy

<sup>28</sup> Service de Physique des Particules, CEA / Saclay, 91191 Gif-sur-Yvette, France

<sup>29</sup> Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

<sup>30</sup> Department of Nuclear Engineering, University of California, Berkeley, CA 94720, USA

<sup>31</sup> Dipartimento di Scienze Fisiche e Chimiche, Università dell'Aquila, L'Aquila I-67100, Italy

<sup>32</sup> Department of Physics, University of Wisconsin, Madison, WI 53706, USA

<sup>33</sup> Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA


<sup>a</sup> Deceased

E-mail: [davide.chiesa@mib.infn.it](mailto:davide.chiesa@mib.infn.it)

**Abstract.** The Cryogenic Underground Observatory for Rare Events (CUORE) is the first bolometric experiment searching for neutrinoless double beta ( $0\nu\beta\beta$ ) decay that has been able to reach the 1-ton scale. The detector consists of an array of 988 TeO<sub>2</sub> crystals arranged in a cylindrical compact structure of 19 towers. The construction of the experiment and, in particular, the installation of all towers in the cryostat was completed in 2016 and data taking started in 2017. In this conference we present the  $0\nu\beta\beta$  decay results of CUORE from examining a total TeO<sub>2</sub> exposure of 86.3 kg yr, characterized by an effective energy resolution of 7.7 keV FWHM and a background in the region of interest of 0.014 counts/(keV kg yr). Based on these data, CUORE places a lower limit on the  $0\nu\beta\beta$  decay half-life of  $^{130}\text{Te}$   $T_{1/2}^{0\nu} > 1.5 \times 10^{25}$  yr (90% C.L.). We then discuss the latest updates in the analysis of background and in the evaluation of the half-life of  $2\nu\beta\beta$  decay of  $^{130}\text{Te}$ .

## 1. Introduction

The primary goal of the CUORE experiment is the search of the neutrinoless double beta ( $0\nu\beta\beta$ ) decay of  $^{130}\text{Te}$  [1]. The observation of this decay would imply that lepton number is not conserved and that neutrinos are massive Majorana particles [2]. The CUORE detector (Figure 1) is composed by 988 TeO<sub>2</sub> cubic crystals (5 cm side) operated as independent bolometers in a cryostat able to cool down about 1-ton detector at a temperature  $\sim 10$  mK. The crystals contain natural tellurium, which is about 34%  $^{130}\text{Te}$ , so they act both as sources and detectors of the  $0\nu\beta\beta$  decay. The total source-detector mass is 742 kg, corresponding to  $\sim 206$  kg of  $^{130}\text{Te}$ . Each crystal is equipped with a neutron transmutation doped (NTD) thermistor that reads out the energy depositions due to particle interactions. At cryogenic temperatures the heat capacity of a crystal is so small that a 1 MeV energy deposition causes a temperature variation of  $\sim 100$   $\mu\text{K}$ , that is converted into a measurable change in voltage by means of the NTD. When a  $\beta\beta$  decay



**Figure 1.** An illustration of the CUORE cryostat (left) and a picture of the CUORE detector (right).

occurs, the energy deposited in the bolometer by the electrons is detected. In the Standard Model,  $\beta\beta$  decay is an allowed transition if two (anti-)neutrinos are emitted to conserve lepton number. In this case, the energy carried by neutrinos is not detected and the  $2\nu\beta\beta$  decay produces a continuous spectrum. Conversely, the signature of the  $0\nu\beta\beta$  decay is a peak at the  $Q$ -value of the transition ( $Q_{\beta\beta}=2527.5$  keV for  $^{130}\text{Te}$ ), because all the energy is carried by the electrons.

The main challenge of CUORE is to maximize the experimental sensitivity, because  $0\nu\beta\beta$  is an extremely rare decay with  $T_{1/2}^{0\nu} > 10^{25}\text{yr}$ .


With non-zero background the sensitivity scales as:  $T_{1/2}^{0\nu} \propto \epsilon a \frac{Mt}{b \Delta E}$ , where  $\epsilon$  is the detection efficiency,  $a$  the isotopic abundance,  $M$  the active mass,  $t$  the livetime, and  $b$  and  $\Delta E$  the background rate and the energy resolution at  $Q_{\beta\beta}$ .

In the years prior to CUORE start-up, the collaboration endeavoured to reduce background and improve the energy resolution. The background in CUORE comes from natural radioactivity and to suppress it complementary strategies are put in place: (1) the CUORE detector is located at Laboratori Nazionali del Gran Sasso (Italy), where  $\sim 1400$  m of rock provide an effective shielding from cosmic rays; (2) the detector is surrounded by several layers of shielding to minimize the background due to environmental radioactivity inside the cavern; (3) all the materials used to build the detector and the cryostat were selected and cleaned to meet strict requirements about radio-purity; (4) analysis techniques based on pulse shape and time coincidence are used for event selection and background discrimination.


The CUORE-0 experiment [3], a single CUORE-like tower operated between 2013 and 2015, allowed to test the performance of the bolometers, proving that is possible to achieve a resolution of 5 keV FWHM [4, 5] and a background rate of  $10^{-2}$  counts/(keV kg yr) [6]. According to these design parameters, CUORE is expected to reach a sensitivity of  $9 \times 10^{25}\text{yr}$  after 5 years livetime [7].

## 2. Results

In 2017 CUORE collected the first physics data, corresponding to a  $\text{TeO}_2$  exposure of 86.3 kg yr. The CUORE cryostat is working very well and 92% of detector channels produced good quality data that were used for this first analysis. The physics data sets are bracketed by calibration measurements in which  $^{232}\text{Th}$  sources are temporarily deployed inside the detector region. The energy calibration curve is built using six  $\gamma$ -lines of the  $^{232}\text{Th}$  decay chain (from 239 keV to 2615 keV). Then, we build the physics spectrum applying selection criteria aimed at improving the experimental sensitivity. We also exclude events that simultaneously trigger more than one crystal, because most of the  $0\nu\beta\beta$  events ( $\sim 88\%$ ) are fully contained in a single crystal and, in this way, we reduce the background due to events depositing energy in multiple crystals. The overall detector efficiency is  $\sim 80\%$ .



**Figure 2.** The CUORE physics spectrum compared with the CUORE-0 one, showing the significant reduction of background rate in the energy region below the  $^{208}\text{Tl}$  line at 2615 keV. Figure coloured online.



**Figure 3.** Fit in the region of interest with 3 components: a constant background, a peak near 2506 keV attributed to  $^{60}\text{Co}$  and a posited peak at  $Q_{\beta\beta}$ .

In Figure 2 we show the first physics spectrum measured by CUORE. These data are characterized by an effective energy resolution of  $(7.7 \pm 0.5)$  keV FWHM at  $Q_{\beta\beta}$  and by a background rate of  $(0.014 \pm 0.002)$  counts/(keV kg yr) in the region of interest.

There is no evidence for  $0\nu\beta\beta$  decay of  $^{130}\text{Te}$  and we perform an Unbinned Extended Maximum Likelihood fit in the [2465–2575] keV range (Figure 3) to extract a new limit on its half-life:  $T_{1/2}^{0\nu} > 1.3 \times 10^{25}$  yr, 90% confidence level (C.L.).


This result is then combined with the ones from CUORE-0 and Cuoricino experiments, obtaining the best limit to date for the  $0\nu\beta\beta$  decay of  $^{130}\text{Te}$ :  $T_{1/2}^{0\nu} > 1.5 \times 10^{25}$  yr (90% C.L.) [8].

In order to analyze quantitatively the main sources of background, evaluate their impact in the  $0\nu\beta\beta$  ROI and measure the half-life of  $2\nu\beta\beta$  decay of  $^{130}\text{Te}$ , we develop a background model for CUORE, according to the methodology used for CUORE-0 and described with detail in [9]. The main background sources are expected to be contamination of  $\text{TeO}_2$  crystals and cryostat shields by natural radioisotopes ( $^{40}\text{K}$ ,  $^{232}\text{Th}$ ,  $^{238}\text{U}$  and their progeny). A small contribution from isotopes produced by cosmogenic activation or radioactive fallout and from environmental muons is also expected.


To identify the background sources, we extract a lot of information directly from CUORE data. We build different spectra by splitting the events depending on time coincidences and interaction position, and we analyze the  $\alpha$  and  $\gamma$  lines, which represent specific signatures of different contamination. In particular, we build the  $\mathcal{M}_1$  spectra with the events that triggered one bolometer, and the  $\mathcal{M}_2$  and  $\Sigma_2$  spectra with the events that simultaneously triggered two bolometers ( $\mathcal{M}_2$  comprises the two energies deposited in the crystals, while  $\Sigma_2$  includes the total energy). The  $\mathcal{M}_1$  events are further split into *Inner* and *Outer* layers, because the crystals in the *detector core* are more shielded from external radiation with respect to those closer to the cryostat shields. In this way, by exploiting the modularity of the CUORE detector, it is possible to better identify the position of the contaminants and distinguish among background sources that mainly produce  $\mathcal{M}_1$  events –like  $2\nu\beta\beta$  decay– with respect to other sources that produce a significant fraction of  $\mathcal{M}_2$  events.

We simulate the background sources using a Geant4-based Monte Carlo (MC) code that generates and propagates primary and secondary particles through the CUORE geometry. The MC outputs are then processed to include the detector features like resolution, energy thresholds and time coincidences, and to build the energy spectra produced in the detector by the different background sources. By comparing the features of MC and experimental data, we identify a list of background sources whose contributions, once summed, can fit the experimental spectra.





**Figure 4.** Top: The measured  $\mathcal{M}_1$ -Inner Layer spectrum and its reconstruction. Bottom: The ratio of the data to the reconstructed model with  $1\sigma$ ,  $2\sigma$  and  $3\sigma$  error bars. Figure coloured online.



**Figure 5.** Fit reconstruction of the  $^{130}\text{Te}$   $2\nu\beta\beta$  decay component in the  $\mathcal{M}_1$ -Inner Layer spectrum.

We reconstruct the CUORE background by fitting a linear combination of 60 source spectra to the experimental data. We perform a simultaneous Bayesian fit of the  $\mathcal{M}_1$ -Inner Layer,  $\mathcal{M}_1$ -Outer Layer,  $\mathcal{M}_2$  and  $\Sigma_2$  spectra. A variable binning of the spectra is adopted to avoid systematic uncertainties due to complicated line shapes. The fit is performed using the JAGS software package [10, 11], that samples the *joint posterior* probability distribution function of the fit parameters with a Markov Chain Monte Carlo (MCMC). The Bayesian approach allows to include the *a priori* information from previous experiments, radioassay measurements of materials and cosmogenic activation calculations.

As shown in Figure 4, we were able to reconstruct the major features of the experimental spectra. The  $2\nu\beta\beta$  decay of  $^{130}\text{Te}$  is one of the main components of the  $\mathcal{M}_1$  spectra (Figure 5) and, thanks to the model implemented, we were able to disentangle it from the contribution of the other background sources and to evaluate its half-life. The preliminary result, that is going to be published in a paper currently in preparation is:  $T_{1/2}^{2\nu} = (7.9 \pm 0.1_{\text{stat}} \pm 0.2_{\text{syst}}) \times 10^{20}$  yr. This result is consistent with previous measurements of this half-life [9, 12, 13] and represents the most precise and accurate measurement of  $2\nu\beta\beta$  decay of  $^{130}\text{Te}$  to date.

## Acknowledgments

The CUORE Collaboration thanks the directors and staff of the Laboratori Nazionali del Gran Sasso and the technical staff of our laboratories. This work was supported by the Istituto Nazionale di Fisica Nucleare (INFN); the National Science Foundation; the Alfred P. Sloan Foundation; the University of Wisconsin Foundation; and Yale University. This material is also based upon work supported by the US Department of Energy (DOE) Office of Science; and by the DOE Office of Science, Office of Nuclear Physics. This research used resources of the National Energy Research Scientific Computing Center (NERSC).

## References

- [1] Artusa D R *et al* (CUORE collaboration) 2015 *Advances in High Energy Physics* **2015** Article ID 879871
- [2] Majorana E 1937 *Il Nuovo Cimento* **14** 171
- [3] Alfonso K *et al* (CUORE collaboration) 2015 *Phys. Rev. Lett.* **115**(10) 102502
- [4] Alduino C *et al* (CUORE collaboration) 2016 *Journal of Instrumentation* **11** P07009
- [5] Alduino C *et al* (CUORE collaboration) 2016 *Phys. Rev. C* **93**(4) 045503
- [6] Alduino C *et al* (CUORE collaboration) 2017 *The European Physical Journal C* **77** 543
- [7] Alduino C *et al* (CUORE collaboration) 2017 *The European Physical Journal C* **77** 532
- [8] Alduino C *et al* (CUORE collaboration) 2018 *Phys. Rev. Lett.* **120**(13) 132501
- [9] Alduino C *et al* (CUORE collaboration) 2017 *The European Physical Journal C* **77** 13
- [10] Plummer M 2012 *JAGS Version 3.3.0 User Manual*
- [11] Chiesa D, Previtali E and Sisti M 2014 *Annals of Nuclear Energy* **70** 157–168
- [12] Arnaboldi C *et al* (MiDBD Collaboration) 2003 *Phys. Lett. B* **557** 167–175
- [13] Arnold R *et al* (NEMO-3 Collaboration) 2011 *Phys. Rev. Lett.* **107**(6) 062504