



Normality of Closures of Orthogonal Nilpotent Symmetric Orbits

M. Trevisiol¹

Received: 21 June 2021 / Accepted: 11 January 2022
© The Author(s) 2022

Abstract

We study closures of conjugacy classes in the symmetric matrices of the orthogonal group and we determine which one are normal varieties. In contrast to the result for the symplectic group where all classes have normal closure, there is only a relatively small portion of classes with normal closure. We perform a combinatorial computation on top of the same methods used by Kraft-Procesi and Ohta.

Introduction

In a fundamental paper of Kostant [4], the adjoint action on a reductive Lie algebra \mathfrak{g} defined over an algebraically closed field k of characteristic 0 is studied in detail. In the course of his analysis it arose the following problem. Let $A \in \mathfrak{g}$, let C_A be the conjugacy class of A and let $\overline{C_A}$ be the (Zarisky) closure of C_A . Is $\overline{C_A}$ always a normal variety?

In his paper Kostant showed that if A is a regular nilpotent element of \mathfrak{g} , so that $\overline{C_A}$ is the nilpotent cone of \mathfrak{g} , the normality is always the case.

In [3], the problem of normality of $\overline{C_A}$ is reduced to the case in which A is nilpotent, possibly changing the Lie algebra \mathfrak{g} .

Later, in [2, Theorem 4.1], Broer proved the normality of a handful of nilpotent orbits in \mathfrak{g} , including the regular and the subregular orbits (if \mathfrak{g} is simple).

In the fundamental paper [5] of Kraft and Procesi, the normality of $\overline{C_A}$ is proved for every nilpotent A in the case $\mathfrak{g} = \mathfrak{gl}_n$. Their method consists in constructing an auxiliary variety Z which is a normal complete intersection such that $\overline{C_A}$ is a *quotient* of Z .

Kraft and Procesi extended their method to the case where \mathfrak{g} is the orthogonal or symplectic Lie algebra in [7]. In that case not all nilpotent classes have normal

✉ M. Trevisiol
marco.trevisiol@sns.it

¹ Dipartimento di Matematica, Università di Roma “La Sapienza”, Piazzale Aldo Moro 5, 00185, Rome, Italy

closure. They obtained necessary and sufficient conditions on the partition of A in order to have the normality of $\overline{C_A}$, under suitable hypothesis on A .

In a subsequent paper by Sommers [13], the cases outside of this hypothesis (in [7]) are proved to always have normal closure. Sommers also solved the problem for the exceptional case E_6 in [12].

Kostant and Rallis generalized the study [4] of the adjoint action on \mathfrak{g} to the action on the symmetric space in [8]. We briefly describe their setting. Let θ be a Lie algebra automorphism of \mathfrak{g} of order 2. Then there exists a decomposition of \mathfrak{g} in θ -eigenspaces given by

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$$

where $\mathfrak{k} = \{A \in \mathfrak{g} : \theta(A) = A\}$ and $\mathfrak{p} = \{A \in \mathfrak{g} : \theta(A) = -A\}$. Let G be the adjoint group of \mathfrak{g} and $K \subseteq G$ be the subgroup of elements commuting with θ . We notice that \mathfrak{k} is a Lie subalgebra of \mathfrak{g} and $\text{Lie}K = \mathfrak{k}$. We remark that K is not necessarily connected. Moreover the adjoint action of G on \mathfrak{g} induces an action of K on \mathfrak{g} and both $\mathfrak{k}, \mathfrak{p}$ are K -stable. The pair $(\mathfrak{g}, \mathfrak{k})$ is called a symmetric pair and \mathfrak{p} is the symmetric space associate to it.

In their study, Kostant and Rallis focused on the action of K on \mathfrak{p} . Among other things, they already observed that the nilpotent cone is not irreducible nor normal in general.

In [14], Vinberg further generalized the study of the adjoint action to graded Lie algebras, also called θ -groups. Here θ is an automorphism of a Lie algebra \mathfrak{g} of finite order n , thus generalizing the case $n = 2$ of [8]. The graduation induced by θ , $\mathfrak{g} = \mathfrak{g}_0 \oplus \cdots \oplus \mathfrak{g}_{n-1}$, produces an action of the θ -fixed points of the adjoint group on \mathfrak{g}_1 .

After the work of Kostant, Rallis and Vinberg, the problem of the normality of $\overline{C_A}$ naturally generalizes to the case of symmetric spaces.

In [11], Sekiguchi studied deeply the geometry of the nilpotent cone in symmetric spaces. In particular he proved several results for the principal nilpotent elements (which are the analogue of the regular nilpotent elements) when the symmetric space is the orthogonal symmetric space or the symplectic symmetric space, that is, the symmetric spaces of the symmetric pairs $(\mathfrak{g}, \mathfrak{k}) = (\mathfrak{gl}(n), \mathfrak{o}(n))$ and $(\mathfrak{g}, \mathfrak{k}) = (\mathfrak{gl}(2m), \mathfrak{sp}(m))$ respectively.

The method of the auxiliary variety Z developed by Kraft and Procesi in [5–7] was adapted by Ohta in [10] to the study of the singularities of the orbits in the orthogonal and symplectic symmetric spaces. In particular he proved that $\overline{C_A}$ is always normal in the case of symplectic symmetric space. Moreover he found several orbits C_A in the case of the orthogonal symmetric space which have a non-normal closure.

It is well known that the conjugacy classes of the nilpotent cone of the orthogonal symmetric space are parametrized by the partitions of n (see section 1 for more details). The main purpose of this paper is to give a necessary and sufficient condition on the partition corresponding to the orbit of a nilpotent symmetric element A in order to have the normality of $\overline{C_A}$. The following theorem is the main result of the paper.

Theorem 1 *Let \mathfrak{p} be the symmetric space of the symmetric pair $(\mathfrak{gl}(n), \mathfrak{o}(n))$ and let $A \in \mathfrak{p}$ be a nilpotent element. Let $\lambda = (\lambda_1, \dots, \lambda_h)$ be the partition of A . Then $\overline{C_A}$ is normal if and only if*

$$\lambda_i - \lambda_{i+1} \leq 1 \quad \forall i = 1, \dots, h \tag{1}$$

with the convention that $\lambda_{h+1} = 0$.

The proof of Theorem 1 will be carried out in Section 8. The *only if* part was already proved in [10, 11] (see Section 8 for details).

We remark that, if n is even, C_A is not necessarily connected, as the group K is not connected. Therefore, sometimes, $\overline{C_A}$ is not normal just because it is not irreducible, being the union of two closure of $SO(n)$ -orbits (as an example, if one takes $\lambda = (2)$, one can show that $\overline{C_A}$ is the union of two lines which meet at 0). As the referee has kindly pointed out, it would be interesting to investigate the normality of the closure of the $SO(n)$ -orbits.

We summarize the content of the rest of the paper.

In Section 1 we recall some basic facts about symmetric nilpotent orbits for the orthogonal group. In Section 2 we recall from [5] the classification of nilpotent pairs via ab -diagrams and we describe the class of ab -diagrams which parametrize symmetric nilpotent pairs. In Section 3 we recall the construction of the variety Z and some of its properties which are needed for the proof. In Section 4 we define a condition on partitions which plays an important role in our investigation. In Section 5 we introduce some combinatorial description of pairs of partitions. In Section 7, resp. Section 8, we prove a complete intersection, resp. normality, condition for the variety Z using a combinatorial computation carried on in Section 6.

I am grateful to Claudio Procesi for proposing me this problem. I would like to thank Corrado De Concini, Andrea Maffei and Paolo Bravi for many useful comments. I am really grateful to the reviewers for their precise and useful comments. A special thank goes to my PhD advisor Giovanni Cerulli Irelli for many discussions about this problem and his precious help in editing this paper.

1 Symmetric Nilpotent Orbits

In this section we introduce settings and notations for the objects studied in the paper. We follow [7] and [8].

Let V be a vector space over \mathbb{C} together with a symmetric non-degenerate bilinear form $(-, -)$. Even if in this paper we will only be concerned with the orthogonal group $O(V)$, we will denote the isometry group with respect to $(-, -)$ as $G(V) := O(V)$ in order to keep the notation as close as possible with [7, 10]. Let $\mathfrak{gl}(V)$ be the space of linear endomorphisms of V . Clearly $G(V)$ acts on $\mathfrak{gl}(V)$ by conjugation. We denote by D^* the adjoint of an endomorphism D with respect to $(-, -)$. The endomorphism $\theta : D \mapsto -D^*$ of $\mathfrak{gl}(V)$ is involutive, therefore we have a decomposition into eigenspaces for θ as follows:

$$\mathfrak{gl}(V) = \mathfrak{k}(V) \oplus \mathfrak{p}(V),$$

where $\mathfrak{k}(V) = \{D : D = \theta(D) = -D^*\}$ is the Lie algebra of $G(V)$. Moreover the action of $G(V)$ leaves both $\mathfrak{k}(V)$, $\mathfrak{p}(V)$ stable.

In this paper we are concerned with the study of nilpotent orbits of $G(V)$ in $\mathfrak{p}(V)$. It is well known ([11, Sec. 3.1], [10, Sec. 0]) that the nilpotent orbits in $\mathfrak{p}(V)$ are completely determined by their Jordan form and every partition of $n := \dim V$ realizes a non-empty nilpotent orbit. We denote by $P(n)$ the set of partitions of n and by $C_\lambda \subseteq \mathfrak{p}(V)$ the nilpotent orbit corresponding to λ .

Let λ be a partition. We set $|\lambda| := n$ if $\lambda \in P(n)$ and we denote the dual partition of λ by $\widehat{\lambda}$. We frequently identify a partition with its Young diagram. This means that, if $\lambda = (\lambda_1, \dots, \lambda_h)$, $|\lambda| = \lambda_1 + \dots + \lambda_h$, λ_i are the rows of λ and if $\widehat{\lambda} = (\widehat{\lambda}_1, \dots, \widehat{\lambda}_t)$, $\widehat{\lambda}_j$ are the columns of λ .

We recall the dimension formula for the orbit C_λ from [10, Remark 8]:

$$\dim C_\lambda = \frac{1}{2} \left(n^2 - \sum_{i=1}^t \widehat{\lambda}_i^2 \right). \tag{2}$$


As the nilpotent cone of $\mathfrak{p}(V)$ is $G(V)$ -stable with only finitely many orbits, we have that orbit closure $\overline{C_\lambda}$ is $G(V)$ -stable, and the complement $\overline{C_\lambda} \setminus C_\lambda$ is a disjoint union of finitely many orbits. The relation $C_\mu \subseteq \overline{C_\lambda}$ produces a partial order on the partitions, called *dominance order* and denoted by $\mu \leq \lambda$, and it is given by

$$\sum_{i=1}^j \lambda_i \geq \sum_{i=1}^j \mu_i \quad \forall j \in \{1, \dots, h\}$$

or, equivalently,

$$\sum_{k>j} \widehat{\lambda}_k \geq \sum_{k>j} \widehat{\mu}_k \quad \forall j \in \{1, \dots, t\}.$$

We recall the notion of nilpotent pairs. We follow the same argument of [7]. Let V (resp. U) be a finite dimensional vector space over \mathbb{C} equipped with a non-degenerate symmetric bilinear form $(-, -)_V$ (resp. $(-, -)_U$). We denote $L(V, U) := \text{Hom}_{\mathbb{C}}(V, U)$, $L(V) := L(V, V)$ and we define $L_{V,U} := L(V, U) \times L(U, V)$. We can interpret $L_{V,U}$ as the representation variety of the quiver



with dimension vector $\underline{n} = (\dim V, \dim U)$. This means that

$$L_{V,U} = \left\{ V \begin{array}{c} \xrightarrow{A} \\ \xleftarrow{B} \end{array} U : A \in L(V,U), B \in L(U,V) \right\}.$$

The group $G(V) \times G(U)$ acts on $L_{V,U}$ by change of basis.

For every $A \in L(V, U)$ we define the adjoint map $A^* \in L(U, V)$ as the unique map such that

$$(Av, u)_U = (v, A^*u)_V \tag{3}$$

for all $v \in V, u \in U$.

Lemma 2 *Let $A \in L(V, U)$ be a linear map. Then*

$$rkA = rkA^*.$$

Proof By (3), we get that $\ker A \perp \text{Im } A^*$ and $\ker A^* \perp \text{Im } A$. □

A pair $(A, B) \in L_{V,U}$ is nilpotent if the endomorphism AB (or equivalently BA) is nilpotent. A pair $(A, B) \in L_{V,U}$ is symmetric if $B = A^*$. We define $N_{V,U}$ as the cone in $L_{V,U}$ of the symmetric nilpotent pairs.


As in [7], we have maps

$$\begin{array}{ccc} L_{V,U} & \xrightarrow{\pi} & L(U) \\ \rho \downarrow & & \\ L(V) & & \end{array}$$

defined by $\pi(A, B) = AB$, $\rho(A, B) = BA$. We can restrict π, ρ to the subspace of symmetric pairs, i. e. the image of $L(V, U) \rightarrow L_{V,U}$, $A \mapsto (A, A^*)$:

$$\begin{array}{ccc} L(V, U) & \xrightarrow{\pi} & \mathfrak{p}(U) \\ \rho \downarrow & & \\ \mathfrak{p}(V) & & \end{array}$$

The subvariety $N_{V,U} \subseteq L_{V,U}$ is $G(V) \times G(U)$ -stable. We call *nilpotent symmetric orbit* each of the $G(V) \times G(U)$ -orbits in $N_{V,U}$. We recall [5, Sec. 4.2] that an *ab*-diagram is a list of *ab*-strings, i.e., strings with letters *a* and *b* occurring on alternate positions. As shown in [5, Sec. 4.3], to each nilpotent pair $(A, B) \in L_{V,U}$ corresponds an *ab*-diagram which determines its $GL(V) \times GL(U)$ -orbit completely; i. e. the $GL(V) \times GL(U)$ -orbits correspond one to one to the *ab*-diagrams having $\dim V$ *a*'s and $\dim U$ *b*'s. The following picture illustrates how to associate an *ab*-diagram $\delta = \delta_{(A,B)}$ to a nilpotent pair (A, B) :



where $\{v_1, \dots, v_5\}$ is a suitable basis of V and $\{u_1, \dots, u_4\}$ is a suitable basis of U .

If δ is the *ab*-diagram of an element $p \in N_{V,U}$, we can retrieve the Young diagram of $\pi(p)$ (resp. $\rho(p)$) by suppressing the *a*'s (resp. the *b*'s) from δ . We denote by $\pi(\delta)$ (resp. $\rho(\delta)$) the Young diagram obtained in this manner.

Let $(A, B) \in L_{V,U}$ be a nilpotent pair with *ab*-diagram δ . For every $h \geq 1$, we want to infer the rank of the linear map $(AB)^{h-1}A : V \rightarrow V$ (resp. $(BA)^{h-1}B :$

$U \rightarrow U$) from δ . If $\gamma = c_1 \cdots c_\ell$ is a string, for any $1 \leq i \leq j \leq \ell$, we call $c_i \cdots c_j$ a *substring* of γ . A *substring* of δ is a substring of one of its *ab*-strings. For example, for $h = 1$, the rank of A equals the number of occurrences of the letter a which are not in the last position of a string; this is hence the number of occurrences of the substring $(ab) = (ab)^h$ in δ . With this in mind we prove the following Lemma.

Lemma 3 *Let (A, B) be a nilpotent pair with *ab*-diagram δ . For every $h \geq 1$, the $\text{rk}((AB)^{h-1}A)$ (resp. $\text{rk}((BA)^{h-1}B)$) equals the number of occurrences of the substring $(ab)^h$ (resp. $(ba)^h$) in δ .*

Proof The *ab*-diagram δ suggests the choice of basis of V and U for which the behavior of A and B is straightforward. For each *ab*-string $\delta_i = c_{i1}c_{i2} \dots c_{i\ell_i}$ of δ (where $c_{ij} \in \{a, b\}$), for each $j = 1, \dots, \ell_i$, we pick vectors $v_{ij} \in V$ (resp. $u_{ij} \in U$) if $c_{ij} = a$ (resp. $c_{ij} = b$), with the following properties: $\{v_{ij}\}_{i,j}$ (resp. $\{u_{ij}\}_{i,j}$) is a basis of V (resp. U) and $Av_{ij} = u_{i,j+1}$ (resp. $Bu_{ij} = v_{i,j+1}$) if $j + 1 \leq \ell_i$ or $Av_{i\ell_i} = 0$ (resp. $Bu_{i\ell_i} = 0$).

Notice that the map $(AB)^{h-1}A$ carries v_{ij} to $u_{i(j+2h-1)}$ if $j + 2h - 1 \leq \ell_i$ or to 0 otherwise. Therefore, $\text{rk}((AB)^{h-1}A)$ equals the number of pairs (i, j) such that $c_{ij} = a$ and $j + 2h - 1 \leq \ell_i$. Moreover, every such pair (i, j) corresponds one to one to the substring $c_{ij} \dots c_{i(j+2h-1)}$ of δ , which is the substring $(ab)^h$. A similar argument computes $\text{rk}((BA)^{h-1}B)$. □

2 Ortho-symmetric *ab*-Diagrams

In this section we give a combinatorial description of the *ab*-diagrams of the elements of $N_{V,U}$.

Let $X \in N_{V,U}$ be a nilpotent symmetric pair, let $G = G(V) \times G(U)$ and let $G.X$ be its orbit in $N_{V,U}$. We also consider $G' = \text{GL}(V) \times \text{GL}(U)$ acting on $L_{V,U}$, and the orbit $G'.X \subseteq L_{V,U}$.

The automorphism σ of G' defined by

$$(g_1, g_2)^\sigma = ((g_1^*)^{-1}, (g_2^*)^{-1})$$

has G as the set of fixed points. The automorphism (still denoted by σ) of $L_{V,U}$ defined by

$$\sigma(A, B) = (B^*, A^*)$$

has the symmetric pairs as the set of fixed points. It is straightforward to check that all hypothesis of [9, Prop. 2.1] hold. Thus we get that:

$$G'.X \cap N_{V,U} = G.X.$$

Let δ be an *ab*-diagram associated to an orbit $G'.Y$. We say that δ is *ortho-symmetric* if

$$G'.Y \cap N_{V,U} \neq \emptyset.$$

We are left with the following problem: which *ab*-diagrams are *ortho-symmetric*?

For any two pairs $(A, B) \in L_{V,U}$, $(A', B') \in L_{V',U'}$, the direct sum $(A, B) \oplus (A', B') = (A \oplus A', B \oplus B')$ is defined as an element in $L_{V \oplus V', U \oplus U'}$. If δ is the ab -diagram of (A, B) and δ' is the ab -diagram of (A', B') , the ab -diagram of the direct sum $(A, B) \oplus (A', B')$ is the disjoint union of δ and δ' .

We call a symmetric pair (A, A^*) indecomposable if it can not be written as a direct sum of two nontrivial symmetric pairs. We claim that the ab -diagrams of indecomposable nilpotent symmetric pairs are given in Table 1. This table closely follows table II in [7, 6.3].

Proposition 4 *An ab -diagram δ is ortho-symmetric if and only if it is a disjoint union of finitely many ab -diagrams from Table 1.*

Proof We say that an ab -diagram η has property \mathcal{P} if, for every $h \geq 1$, the number of occurrences of $(ab)^h$ in η equals the number of occurrences of $(ba)^h$ in η .

Every ab -diagrams $\alpha_k, \beta_k, \varepsilon_k$ in Table 1 has property \mathcal{P} . Indeed in the diagrams α_k and β_k , for every $h \geq 1$, there are exactly $k - h + 1$ occurrences of $(ab)^h$ and of $(ba)^h$. The diagram ε_k has property \mathcal{P} by symmetry of its two ab -strings.

Let $(A, A^*) \in N_{V,U}$ be a nilpotent symmetric pair with ab -diagram δ , an ortho-symmetric ab -diagram. For every positive integer h , we notice that the pair $((AA^*)^{h-1}A, (A^*A)^{h-1}A^*)$ is nilpotent symmetric. By Lemma 2, we have:

$$\text{rk}(AA^*)^{h-1}A = \text{rk}(A^*A)^{h-1}A^*,$$

which, by Lemma 3, implies that δ has property \mathcal{P} .

We claim that an ab -diagram δ with property \mathcal{P} is a disjoint union of finitely many ab -diagrams from Table 1. We proceed by contradiction: suppose that there exist ab -diagrams with property \mathcal{P} which are not disjoint union of finitely many ab -diagrams from Table 1 and take δ one with the least number of ab -strings. Let s be one of the ab -strings in δ with maximal length and let l be its length.

If $l = 2k + 1$ is odd, $s = \alpha_k$ or $s = \beta_k$. Let δ' be the ab -diagram such that $\delta = s \sqcup \delta'$ (we cannot have $\delta = s$, otherwise δ is already in Table 1).

If $l = 2k$ is even, without loss of generality, we assume that s starts with a . Then, in s there exists exactly one occurrence of $(ab)^k$ and no occurrence of $(ba)^k$. As δ satisfies property \mathcal{P} , by maximality of l , there must be an ab -string s' in δ of length $l = 2k$ starting with b . Thus $s \sqcup s' = \varepsilon_k$, so let δ' be the ab -diagram such that $\delta = s \sqcup s' \sqcup \delta'$ (again, it cannot be empty).

Table 1 Indecomposable ortho-symmetric ab -diagrams

type	α_k	β_k	ε_k
	$(ab)^k a = abab \cdots a$	$(ba)^k b = baba \cdots b$	$(ab)^k = abab \cdots b$ $(ba)^k = baba \cdots a$
#a	$k + 1$	k	$2k$
#b	k	$k + 1$	$2k$
	$k \geq 0$	$k \geq 0$	$k \geq 1$

In both cases, δ and $\delta \setminus \delta'$ have property \mathcal{P} , therefore also δ' has property \mathcal{P} . By minimality of δ , δ' is disjoint union of finitely many ab -diagrams from Table 1, thus so is δ . This contradicts the existence of such a δ and concludes the only if part of the proposition.

On the other hand, we show that every ab -diagram in Table 1 is ortho-symmetric. In order to show this, we just need to construct a symmetric pair (A, A^*) associated to each diagram $\alpha_k, \beta_k, \varepsilon_k$.

α_k : Let $D : \mathbb{C}^{k+1} \rightarrow \mathbb{C}^{k+1}$ be a symmetric nilpotent endomorphism of nilpotent order exactly $k + 1$. Let $D = I \circ X$ be the canonical decomposition of the map D through its image $D(\mathbb{C}^{k+1})$, so that $X : \mathbb{C}^{k+1} \rightarrow D(\mathbb{C}^{k+1})$ and $I : D(\mathbb{C}^{k+1}) \hookrightarrow \mathbb{C}^{k+1}$. Then D induces a non-degenerate bilinear symmetric form in $D(\mathbb{C}^{k+1})$, as shown in [7, 4], and $X = I^*$. As $\dim D(\mathbb{C}^{k+1}) = k$ and both X, I have maximal rank, we immediately get that the ab -diagram of the symmetric nilpotent pair (X, I) is α_k .

β_k : Let (A, A^*) be a pair with ab -diagram of type α_k . Then (A^*, A) is a pair with ab -diagram of type β_k .

ε_k : Let (A, B) be a nilpotent (not symmetric) pair between the spaces $(V, U) = (\mathbb{C}^k, \mathbb{C}^k)$ with ab -diagram $ab \cdots b$. Let (B^*, A^*) be the linear dual of (A, B) , so that it is a nilpotent pair between the dual spaces (V^*, U^*) with ab -diagram $ba \cdots a$. Then we can equip $V \oplus V^*$ and $U \oplus U^*$ with the non-degenerate symmetric bilinear form given by the duality pairing. Therefore $(A, B) \oplus (B^*, A^*)$ is a symmetric nilpotent pair and its ab -diagram has type ε_k .

□

3 The Variety Z

For each partition λ of n we are going to construct a variety $Z^{(\lambda)}$ (or just Z , if the partition λ is clear from the context) with the following property: there exists a quotient map $Z \rightarrow \overline{C}_\lambda$. In this way, we will be able to assert some properties of \overline{C}_λ by proving them for the variety Z . The construction of Z is analogous to the one carried out in [7] and it has already been described in [10]; here we just recall its definition.

Let $\lambda \in P(n)$ be a partition and let $t = \lambda_1$ be the number of columns of λ . As in [7], we define a variety Z using the representations of the quiver:

$$Q_t = 0 \begin{array}{c} \xrightarrow{x} \\ \xleftarrow{y} \end{array} 1 \begin{array}{c} \xrightarrow{x} \\ \xleftarrow{y} \end{array} \cdots \begin{array}{c} \xrightarrow{x} \\ \xleftarrow{y} \end{array} t.$$

Let

$$n_i = \sum_{j>i} \widehat{\lambda}_j$$

so that $n_0 = n$ and $n_t = 0$. We fix the dimension vector $\underline{n} = (n_0, \dots, n_t)$ and vector spaces V_i such that $\dim V_i = n_i$. We also fix a symmetric bilinear non-degenerate form on each V_i .

The variety Z consists of the representations of dimension vector \underline{n} of the quiver Q_t with relations:

- $xy = yx$;
- $y = x^*$ (the adjoint map as in (3)).

Therefore, each point $z \in Z$ is a sequence of maps

$$(A_1, B_1; A_2, B_2; \dots; A_t, B_t)$$

where $A_i \in L(V_{i-1}, V_i)$, $B_i \in L(V_i, V_{i-1})$, $A_i B_i = B_{i+1} A_{i+1}$ and $B_i = A_i^*$ for each i . As each B_i is adjoint to A_i , we get an inclusion

$$Z \hookrightarrow L(V_0, V_1) \times \dots \times L(V_{t-1}, V_t)$$

mapping

$$(A_1, B_1; A_2, B_2; \dots; A_t, B_t) \mapsto (A_1, \dots, A_t).$$

As in [7], we can think of Z as a schematic fiber in the following way. Let

$$M := L(V_0, V_1) \times \dots \times L(V_{t-1}, V_t),$$

$$N := \mathfrak{p}(V_1) \times \dots \times \mathfrak{p}(V_{t-1}),$$

$$\Phi : M \rightarrow N$$

$$(A_1, \dots, A_t) \mapsto (A_1 A_1^* - A_2^* A_2, \dots, A_{t-1} A_{t-1}^* - A_t^* A_t);$$

then we put $Z := \Phi^{-1}(0)$. As M is an affine space, this interpretation will be useful in Section 7 to show that Z is a complete intersection variety (at least for some λ).

As in [7], let $G = G(V_0) \times \dots \times G(V_t)$ be the group which acts on Z by change of basis and $H = G(V_1) \times \dots \times G(V_t) \subseteq G$. We have a map

$$\Theta : Z \rightarrow \overline{C_\lambda}$$

$$(A_1, B_1; \dots; A_t, B_t) \mapsto B_1 A_1.$$

By the same argument as in [7], we have the following:

Proposition 5 *The map Θ splits through the quotient by the group H as*

$$\Theta : Z \rightarrow Z/H \xrightarrow{\sim} \overline{C_\lambda};$$

moreover Θ is $G(V_0) = G/H$ -equivariant.

(In particular, $\Theta^{-1}(C_\mu)$ is stable under the action of $G(V_0)$ for each partition $\mu \leq \lambda$.)

By restricting the quiver Q_t to two consecutive vertices $i - 1, i$, we get a map $Z \rightarrow N_{V_{i-1}, V_i}$. As all the orbits in N_{V_{i-1}, V_i} are defined by their ortho-symmetric ab -diagram, we can map each point $z \in Z$ and $i = 1, \dots, t$ to its i th ab -diagram τ_i . Therefore, for each string of ab -diagrams

$$\tau = (\tau_1, \dots, \tau_t)$$

we define

$$Z_\tau = \{z \in Z : \forall i, \text{ the } i\text{th } ab\text{-diagram of } z \text{ is } \tau_i \}.$$

The following are necessary conditions on τ in order to have $Z_\tau \neq \emptyset$:

- (1) τ_i must have exactly n_{i-1} a 's and n_i b 's,
- (2) $\rho(\tau_i) = \pi(\tau_{i+1})$ for $i = 1, \dots, t - 1$,
- (3) each τ_i must be ortho-symmetric.

We will denote by $\Lambda^{(\lambda)}$ (or, shortly, by Λ) the set of all strings τ which verify the three conditions above.

As in [5, Lemma 5.4], [7, 8.2], $\Theta^{-1}(C_\lambda)$ is given by exactly one stratum Z_{τ^0} . The string of ab -diagrams τ^0 can be constructed in the following way: for all $i = 0, \dots, t - 1$, let λ_i be the partition of n_i given by the columns of λ with indices greater than i ; then let τ_{i+1} be the ab -diagram built by setting one a on each box in the Young diagram of λ_i and by setting one b between each pair of consecutive a 's.

Example 1 Let $\lambda = (3, 1)$ be a partition. Then the string of ab -diagrams τ^0 is the following:

$$\tau^0 = (\tau_1^0, \tau_2^0, \tau_3^0) = \left(\begin{array}{c|c|c|c} ababa & aba & a \\ \hline a & & \end{array} \right).$$

The stratum Z_{τ^0} is open in Z as it is the subvariety of the points

$$(A_1, B_1, \dots, A_t, B_t)$$

in Z where each A_i, B_i has maximal rank. We have the following key result after the manner of Kraft, Procesi and Ohta.

Proposition 6 ([7, 5.5] and [10, Prop. 8]) *The open set Z_{τ^0} is contained in the smooth locus of Z .*

We recall a dimension formula for the strata Z_τ due to Ohta [10, Prop. 6]. First, for each ab -diagram δ , we denote by a_i (resp. b_i) the number of rows of δ starting by a (resp. b) of length i . We define

$$\Delta(\delta) := \sum_{i \text{ odd}} a_i b_i$$

and

$$o(\delta) := \sum_{i \text{ odd}} (a_i + b_i).$$

Proposition 7 [10, Prop. 6] *Let $\tau = (\tau_1, \dots, \tau_t)$. We have*

$$\begin{aligned} \dim Z_\tau &= \frac{1}{2} \dim C_{\pi(\tau_1)} + \sum_{i=0}^{t-1} \left(\frac{1}{2} n_i n_{i+1} - \frac{1}{4} (n_i + n_{i+1}) \right) \\ &+ \sum_{i=1}^t \left(\frac{1}{4} o(\tau_i) - \frac{1}{2} \Delta(\tau_i) \right). \end{aligned} \tag{4}$$

In the following sections we focus on the combinatorics of the $\tau \in \Lambda$. We need tools to effectively manipulate formula (4), in particular we aim at proposition 10 and proposition 14 (see Section 6).

4 s-Step Condition

In this section we define a condition which plays a pivotal role in the subsequent study of the nilpotent symmetric orbits.

Definition 1 Given a partition $\lambda = (\lambda_1, \dots, \lambda_h)$, we say that λ satisfies the *s-step condition* if for every $i = 1, \dots, h$ we have

$$\lambda_i \leq \lambda_{i+1} + s$$

with the convention that $\lambda_{h+1} = 0$.

We want to highlight some simple properties of this definition. First of all, s_1 -step condition implies s_2 -step condition for each $s_2 \geq s_1$. Secondly, if λ satisfies the s -step condition, then if we remove the first row or the first column from λ , the resulting partition still satisfies the s -step condition.

Example 2 The single line partition (n) satisfies the n -step condition, but not the $(n - 1)$ -step condition. The triangular partition $(n, n - 1, \dots, 1)$ satisfies the 1-step condition. The 2-skew triangular partition $(2n, 2(n - 1), \dots, 2)$ satisfies the 2-step condition (but not the 1-step condition).

5 Differences Between Partitions


In this section we study some quantitative ways to address the difference between two partitions. Let $\lambda, \mu \in P(n)$ be two partitions of n such that $\lambda > \mu$.

Definition 2 If $\lambda = (\lambda_1, \dots, \lambda_k) \in P(n)$ and $\mu = (\mu_1, \dots, \mu_k) \in P(n)$, possibly allowing $\lambda_i = 0$ for some i , we define

$$q(\lambda, \mu) = \frac{1}{2} \sum_{i=1}^k |\lambda_i - \mu_i|.$$

Remark 1 By an easy computation modulo 2, it can be seen that $q(\lambda, \mu)$ is an integer.

Example 3 Let $\lambda = (7, 2, 2, 1)$ and $\mu = (5, 3, 1, 1, 1, 1)$. We can draw the Young diagram of both partitions and overlay one over the other, so that λ is given by white and gray boxes while μ is given by white and black boxes in the following diagram:



We can compute the q difference:

$$q(\lambda, \mu) = \frac{1}{2} ((7 - 5) + (3 - 2) + (2 - 1) + (1 - 1) + 1 + 1) = 3.$$

We can check that in this case $q(\lambda, \mu)$ is the number of gray boxes or the number of black boxes. This is an easy fact (it is not even needed that $\lambda > \mu$) that can be derived from the following Lemma.

Lemma 8 *Let $\lambda > \mu$ be two partitions of n . Then there exist partitions*

$$\lambda = \lambda^0 > \lambda^1 > \dots > \lambda^{q(\lambda, \mu)} = \mu$$

such that $q(\lambda^i, \lambda^{i+1}) = 1$ for each $i = 0, \dots, q(\lambda, \mu) - 1$.

Moreover, we can choose λ^i such that for each row r and for each column c the sequences of integers $\lambda_r^0, \dots, \lambda_r^{q(\lambda, \mu)}$ and $\widehat{\lambda}_c^0, \dots, \widehat{\lambda}_c^{q(\lambda, \mu)}$ are monotone.

Proof We proceed by induction on $q = q(\lambda, \mu)$. If $q = 1$ we trivially have $\lambda^1 = \mu$. If $q > 1$, we are going to build λ^1 such that $q(\lambda, \lambda^1) = 1$ and $q(\lambda^1, \mu) = q(\lambda, \mu) - 1$.

We build λ^1 starting from λ in this way. Let \tilde{i} be the first index such that $\lambda_{\tilde{i}} > \mu_{\tilde{i}}$ and let j be the first index such that $\lambda_j < \mu_j$. Clearly there must exist such indices because $\lambda \neq \mu$ and $|\lambda| = |\mu|$. Let $i \geq \tilde{i}$ be the last index such that $\lambda_i = \mu_i$ (possibly $i = \tilde{i}$). As $\lambda > \mu$, we must have $\tilde{i} < j$ and therefore $i < j$. We define

$$\lambda^1 = (\lambda_1, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, \dots, \lambda_{j-1}, \lambda_j + 1, \lambda_{j+1}, \dots, \lambda_k).$$

The sequence λ^1 is actually a partition of n as $|\lambda^1| = |\lambda| = n$, $\lambda_i - 1 \geq \lambda_{i+1}$ by the maximality of i and $\lambda_{j-1} \geq \lambda_j + 1$ by the minimality of j .

Moreover, both the following hold: $q(\lambda, \lambda^1) = 1$ and $q(\lambda^1, \mu) = q - 1$. We apply induction on the pair (λ^1, μ) and we conclude the proof of the first part of the Lemma.


To conclude the proof, we notice that if, for a row r , $\lambda_r > \mu_r$ (resp. $\lambda_r < \mu_r$), then we have $\lambda_r^i \geq \lambda_r^{i+1}$ (resp. $\lambda_r^i \leq \lambda_r^{i+1}$) by construction. The same holds true for the columns lengths. □

Example 4 Taking the partitions λ, μ defined in the previous example, the construction explained in the Lemma produces the following sequence of partitions:

$$(7, 2, 2, 1) > (6, 3, 2, 1) > (5, 3, 2, 1, 1) > (5, 3, 1, 1, 1).$$

Remark 2 In view of Lemma 8 we can interpret $q(\lambda, \mu)$ as the minimum number of boxes needed to lower in order to obtain μ from λ .

We consider two other ways of comparing partitions. Given a partition λ , for each box B in the Young diagram of λ we define its column number $c(B)$ (resp. row number $r(B)$) by counting the columns (resp. rows) starting from the leftmost column (resp. uppermost row). For example, given $\lambda = (7, 2, 2, 1)$ we have:



Remark 3 By summing up over the rows (resp. the columns) of λ , we have:

$$\sum_{B \in \lambda} c(B) = \sum_{i=1}^h \binom{\lambda_i + 1}{2} \quad \left(\text{resp. } \sum_{B \in \lambda} r(B) = \sum_{i=1}^t \binom{\widehat{\lambda}_i + 1}{2} \right). \tag{5}$$

Definition 3 Given $\lambda \geq \mu$ as before, we define:

$$c(\lambda, \mu) = \sum_{B \in \lambda} c(B) - \sum_{B \in \mu} c(B),$$

where B is selected among the boxes in the Young diagrams of λ and μ and, similarly,

$$r(\lambda, \mu) = \sum_{B \in \mu} r(B) - \sum_{B \in \lambda} r(B).$$

Example 5 Let $\lambda = (7, 2, 2, 1)$ and $\mu = (5, 3, 1, 1, 1, 1)$. We compute

$$\begin{aligned} c(\lambda, \mu) &= (28 + 3 + 3 + 1) - (15 + 6 + 1 + 1 + 1 + 1) = 10, \\ r(\lambda, \mu) &= (21 + 3 + 3 + 1 + 1) - (10 + 6 + 1 + 1 + 1 + 1 + 1) = 8. \end{aligned}$$

Remark 4 It is always guaranteed that $c(\lambda, \mu) \geq 0$ and $r(\lambda, \mu) \geq 0$ for every $\lambda \geq \mu$. We even have that

$$\lambda = \mu \iff c(\lambda, \mu) = 0 \iff r(\lambda, \mu) = 0. \tag{6}$$

In fact we have that $c(\lambda, \mu) \geq 0$ is equivalent to

$$\sum_i \binom{\lambda_i + 1}{2} \geq \sum_i \binom{\mu_i + 1}{2}. \tag{7}$$

As the function $n \mapsto \binom{n+1}{2}$ is convex, (7) is given by the Karamata’s inequality [1, Chapter 1, §28].

Remark 5 Given partitions $\lambda \geq \mu \geq \nu$, by definition of c and r we have:

$$\begin{aligned} c(\lambda, \nu) &= c(\lambda, \mu) + c(\mu, \nu), \tag{8} \\ r(\lambda, \nu) &= r(\lambda, \mu) + r(\mu, \nu). \tag{9} \end{aligned}$$

Remark 6 For every pair of partitions $\lambda \geq \mu$ we have that

$$c(\lambda, \mu) \geq q(\lambda, \mu). \tag{10}$$

In fact, when $q(\lambda, \mu) = 1$, by (6) we get (10). Otherwise, by Lemma 8, we have a sequence of partitions $\lambda^0, \dots, \lambda^{q(\lambda, \mu)}$. For each pair $(\lambda^i, \lambda^{i+1})$, we have $c(\lambda^i, \lambda^{i+1}) \geq 1$, therefore the conclusion follows by summing up each inequalities by (8).

Lemma 9 Let λ be a partition satisfying the s -step condition and $\mu < \lambda$ be another partition. Then

$$s \cdot r(\lambda, \mu) \geq c(\lambda, \mu) + q(\lambda, \mu). \tag{11}$$

Moreover, (11) holds strictly if there exists a column index c such that $\widehat{\mu}_c > \widehat{\lambda}_c + 1$ or a row index r such that $\mu_r > \lambda_r + 1$.

Proof Let $q = q(\lambda, \mu)$. By Lemma 8, we obtain a sequence

$$\lambda = \lambda^0 > \dots > \lambda^q = \mu,$$

with $q(\lambda^i, \lambda^{i+1}) = 1$ for $i = 0, \dots, q - 1$.

We want to prove

$$s \cdot r(\lambda^i, \lambda^{i+1}) \geq c(\lambda^i, \lambda^{i+1}) + 1 \tag{12}$$

for all $i = 0, \dots, q - 1$. Once (12) is proved, the conclusion will follow by summing up each of those inequalities by (9) and (8).


We start by noticing that the s -step condition implies that, for every pair of row indices $r_1 < r_2$,

$$\lambda_{r_1} - \lambda_{r_2} \leq s(r_2 - r_1).$$

Fix r_1 to be an index row such that $\lambda_{r_1} \geq \mu_{r_1}$ and fix r_2 to be an index row such that $\lambda_{r_2} \leq \mu_{r_2}$. Lemma 8 asserts that $\lambda_{r_j}^i$ is a monotone sequence for both $j = 1, 2$; therefore, for such r_1, r_2 , we get $\lambda_{r_1}^i - \lambda_{r_2}^i \leq \lambda_{r_1} - \lambda_{r_2}$, so we get

$$\lambda_{r_1}^i - \lambda_{r_2}^i \leq s(r_2 - r_1). \tag{13}$$

As $q(\lambda^i, \lambda^{i+1}) = 1$, we already observed that the Young diagrams of λ^i, λ^{i+1} have only one box in different positions. This means that there exist exactly two columns $c_{i,1} < c_{i,2}$ which are not equal between λ^i, λ^{i+1} . Similarly, there exist exactly two rows indices $r_{i,1} < r_{i,2}$.



We have $c(\lambda^i, \lambda^{i+1}) = c_{i,2} - c_{i,1}$ and $r(\lambda^i, \lambda^{i+1}) = r_{i,2} - r_{i,1}$. Moreover, by definition of λ^i and λ^{i+1} , we have $\lambda_{r_{i,1}}^i = c_{i,2}$ and $\lambda_{r_{i,2}}^i = c_{i,1} - 1$. Therefore, by (13), we get

$$s \cdot r(\lambda^i, \lambda^{i+1}) = s(r_{i,2} - r_{i,1}) \geq \lambda_{r_{i,1}}^i - \lambda_{r_{i,2}}^i = c_{i,2} - c_{i,1} + 1 = c(\lambda^i, \lambda^{i+1}) + 1.$$

Finally, we consider the additional hypothesis of the existence of a column index c such that $\widehat{\mu}_c > \widehat{\lambda}_c + 1$ or a row index r such that $\mu_r > \lambda_r + 1$. We are going to prove that there exists i such that (13) holds strictly.

Let

$$I_{\text{cols}}(c) = \left\{ i \in \{0, \dots, q - 1\} : \widehat{\lambda}_c^{i+1} > \widehat{\lambda}_c^i \right\}.$$

As $q(\lambda^i, \lambda^{i+1}) = 1$, we have $|\widehat{\lambda}_c^{i+1} - \widehat{\lambda}_c^i| \leq 1$ for each i . Therefore, if there exists c such that $\widehat{\mu}_c > \widehat{\lambda}_c + 1$, we have $|I_{\text{cols}}(c)| \geq 2$. Let j be the minimum in $I_{\text{cols}}(c)$.

We show that (13) holds strictly for the rows $r_{i,1}, r_{i,2}$ for every $i \in I_{\text{cols}}(c)$ such that $i \neq j$. Indeed, as $\lambda_{r_{i,2}}^i = c = \lambda_{r_{j,2}}^j$, we have

$$\begin{aligned} \lambda_{r_{i,1}}^i - \lambda_{r_{i,2}}^i &\leq \lambda_{r_{i,1}}^j - \lambda_{r_{j,2}}^j \leq \lambda_{r_{i,1}} - \lambda_{r_{j,2}} \\ &\leq s(r_{j,2} - r_{i,1}) \leq s(r_{i,2} - 1 - r_{i,1}) < s(r_{i,2} - r_{i,1}). \end{aligned}$$

In a similar fashion, let

$$I_{\text{rows}}(r) = \left\{ i \in \{0, \dots, q-1\} : \lambda_r^{i+1} > \lambda_r^i \right\}.$$

If there exists r such that $\mu_r > \lambda_r + 1$, then $|I_{\text{rows}}(r)| \geq 2$ and we take $j \in I_{\text{rows}}(r)$ to be the minimum and $i \in I_{\text{rows}}(r)$ to be another index. We have $\lambda_{r_{i,2}}^i > \lambda_{r_{i,2}}^j = \lambda_{r_{i,2}}$, so we get

$$\lambda_{r_{i,1}}^i - \lambda_{r_{i,2}}^i < \lambda_{r_{i,1}} - \lambda_{r_{i,2}} \leq s(r_{i,2} - r_{i,1}).$$

In both cases (13) holds strictly for at least an i . Therefore (12) holds strictly for this i , and (11) must hold strictly as a consequence. Thus, the Lemma is proven. \square

Example 6 Let

$$\lambda = (6, 4, 2), \quad \mu = (5, 3, 2, 1, 1), \quad \nu = (5, 3, 3, 1)$$

be three partitions of $n = 12$. We have that $\mu < \lambda$ and $\nu < \lambda$; moreover, in the pair (λ, μ) the first column differs by 2 boxes, while no column in (λ, ν) differs more than 1 box.

Since λ satisfies the 2-step condition, we can compute each term involved in (11). For (λ, μ) we have: $2 \cdot 6 \geq 8 + 2$ (which holds strictly), while for (λ, ν) we have: $2 \cdot 4 \geq 6 + 2$.

6 Inequalities on the Dimensions

In this section we will use the tools introduced in Section 5 to effectively compute the dimensions of the strata in Z .

Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a partition of $n := |\lambda|$ with $t := \lambda_1$ columns. We want to compare the dimension of Z_{τ^0} with the dimension of each other stratum Z_{τ} with $\tau \in \Lambda^{(\lambda)}$. Let $\mu = \pi(\tau)$, so that $\mu \leq \lambda$. The aim of this section is to give some sufficient combinatorial conditions on the pair of partitions (λ, μ) in order to secure a bound on the difference $\dim Z_{\tau^0} - \dim Z_{\tau}$.

The aim of the current section is to prove proposition 10 and Proposition 14. In order to introduce the stronger inequality in Proposition 14, we need an in-depth study of the combinatorics of Λ , so we prefer to introduce it later.

Proposition 10 *Let $\lambda \geq \mu$ be two partitions and let $Z = Z^{(\lambda)}$ be the variety built from λ . Let Z_{τ^0} be the only stratum of Z with $\pi(\tau_1^0) = \lambda$ and let Z_{τ} be a stratum with $\tau = (\tau_1, \dots, \tau_t)$ such that $\pi(\tau_1) = \mu$.*

For each real number x , we have that:

$$2r(\lambda, \mu) - c(\lambda, \mu) - q(\lambda, \mu) \geq 4x \implies \dim Z_{\tau^0} - \dim Z_{\tau} \geq x.$$

The proof will proceed as follows: we take the dimension formula of the strata of Z given by (4) then we will estimate each term on the difference. Each of the following lemmas deals with one of the terms.

Lemma 11 *Let $\lambda \geq \mu$ be two partitions of n . Let $t = \lambda_1$ be the number of columns of λ . Then*

$$\sum_{i=1}^t (\widehat{\mu}_i^2 - \widehat{\lambda}_i^2) = 2r(\lambda, \mu).$$

(We allow $\widehat{\mu}_i = 0$ for each $i > \mu_1$.)

Proof For every integer m we have: $m^2 = 2\binom{m+1}{2} - m$. So, by (5), we get:

$$\sum_{i=1}^t (\widehat{\mu}_i^2 - \widehat{\lambda}_i^2) = \sum_{i=1}^t \left(2\binom{\widehat{\mu}_i + 1}{2} - \widehat{\mu}_i - 2\binom{\widehat{\lambda}_i + 1}{2} + \widehat{\lambda}_i \right) = 2r(\lambda, \mu) - n + n.$$

□

In order to carry on the computation, we examine carefully all the strata Z_τ with $\tau \neq \tau^0$. We do so by grouping together the $\tau \in \Lambda$ which can appear in $\Theta^{-1}(C_\mu)$, for fixed μ . We already know that for every $\tau = (\tau_1, \dots, \tau_t)$ such that $\Theta(Z_\tau) = C_\mu$, we have that $\pi(\tau_1) = \mu$.

The conditions $\tau \in \Lambda$ and $\pi(\tau_1) = \mu$ place important combinatorial restrictions in the choice of the ab -diagrams τ_1, \dots, τ_t . We are going to describe them in the following paragraphs.

Let $Z^{(\mu)}$ be the variety Z built from the partition μ (rather than the partition λ). We will denote by σ^0 the unique stratum in $\Lambda^{(\mu)}$ such that $\Theta(Z_{\sigma^0}^{(\mu)}) = C_\mu$. As in the case of τ^0 , every ab -diagram σ_i^0 has only rows starting and ending by a , in particular they all have odd length, so:

$$\sum_{i=1}^t o(\sigma_i^0) = n = \sum_{i=1}^t o(\tau_i^0). \tag{14}$$

The sequence of ab -diagrams σ^0 does not belong to $\Lambda^{(\lambda)}$ because the integers n_i computed from μ are different from those computed from λ .

Nonetheless, for every $\tau \in \Lambda^{(\lambda)}$ such that $\pi(\tau_1) = \mu$, the condition $\rho(\tau_{i-1}) = \pi(\tau_i)$ implies that each τ_i can be obtained from σ_i^0 by adding an adequate amount of a 's and b 's letters. For example, if $i = 0$ we need some b 's, possibly.

For each $i = 1, \dots, t$, let d_i be the list of a 's and b 's we need to add to σ_i^0 in order to obtain τ_i . Let d_i^a (resp. d_i^b) be the number of a 's (resp. b 's) in d_i . We remark that d_i^a and d_i^b depend only on (λ, μ) and not on the particular τ . In fact, we can compute d_i^a by taking the differences

$$d_i^a = \sum_{j=i}^t (\widehat{\lambda}_j - \widehat{\mu}_j); \tag{15}$$

and therefore we have $d_i^b = d_{i+1}^a$.

Example 7 Let $\tau \in \Lambda^{(\lambda)}$ and let $\sigma^0 \in \Lambda^{(\mu)}$. Suppose that the i th diagrams are the following:

$$\sigma_i^0 = \begin{matrix} aba \\ aba \\ b \end{matrix}; \quad \tau_i = \begin{matrix} ababa \\ aba \\ ba \\ ab \end{matrix}.$$

Then, we have $d_i = (a, a, a, b, b)$, $d_i^a = 3$, $d_i^b = 2$.

Given an ab -diagram δ and a list d of a 's and b 's, we define $\text{aug}_\delta(d)$ to be the set of ortho-symmetric ab -diagrams obtainable from δ by adding the letters in d .

Example 8 Let $d = (a, b)$ and

$$\delta = \begin{matrix} aba \\ aba \\ b \end{matrix}.$$

Then $\text{aug}_\delta(d)$ has three elements:

$$\text{aug}_\delta(d) = \left\{ \begin{matrix} ababa & aba & aba \\ aba & aba & a \\ b & bab & b \\ & & b \end{matrix} \right\}.$$

Lemma 12 Let δ^0 be an ab -diagram with rows starting and ending only with a . Let d be a list of d^a a 's and d^b b 's. Let $\delta \in \text{aug}_{\delta^0}(d)$. Then the following holds:

$$o(\delta) - 2\Delta(\delta) - o(\delta^0) \leq \max\{d^a, d^b\}.$$

Proof The ab -diagram δ is obtained from δ^0 by adding d^a a 's and d^b b 's.

Let L (resp. S) be the set of rows of δ longer than 1 (resp. of length 1) built using only letters in d . As $o(\delta)$ counts some of the rows in δ , we compare the difference $o(\delta) - o(\delta^0)$ with the number of rows built only with letters in d and we get $o(\delta) - o(\delta^0) \leq |L| + |S|$.

Let S_a (resp. S_b) the rows of S starting with a (resp. b), so that $S = S_a \sqcup S_b$. Recall that a_i (resp. b_i) is the number of rows of length i starting with a (resp. b); in this particular ab -diagram δ we have $|S_b| = b_1$ and $|S_a| \leq a_1$.

$$\begin{aligned} o(\delta) - o(\delta^0) - 2\Delta(\delta) &\leq |L| + |S| - 2 \sum_{i \text{ odd}} a_i b_i \\ &\leq |L| + |S| - 2a_1 b_1 \\ &\leq |L| + |S_a| + |S_b| - 2|S_a||S_b| \\ &\leq |L| + \max\{|S_a|, |S_b|\}. \end{aligned}$$

If $|S_a| \geq |S_b|$, then $|L| + |S_a| \leq d_i^a$ because each row in L or in S_a contains at least one a . If the opposite holds true, then $|L| + |S_b| \leq d_i^b$ for the same reason. Thus the Lemma is proven. □

We can introduce a small, but very important, improvement on Lemma 12.

Lemma 13 *In the same setting of Lemma 12, if we furthermore assume that $d = (b)$, that is $d^a = 0$ and $d^b = 1$, and that δ^0 has l rows of length 1, we have the stronger equality:*

$$o(\delta) - 2\Delta(\delta) - o(\delta^0) = \max\{d^a, d^b\} - 2l = 1 - 2l.$$

Proof As all the rows in δ^0 have odd length, we cannot extend exactly one row with letter b , otherwise the resulting ab -diagram would not be ortho-symmetric. So we must place b on a new row.

In this case, $o(\delta) - o(\delta^0) = 1$ and $\Delta(\delta) = a_1b_1 = l \cdot 1$. □

Using this lemma we will be able to introduce the previously announced stronger version of proposition 10, namely:

Proposition 14 *Let $\lambda \geq \mu$ be two partitions and let $Z = Z^{(\lambda)}$ be the variety built from λ . Let Z_{τ^0} be the only stratum of Z with $\pi(\tau_1^0) = \lambda$ and let Z_{τ} be a stratum with $\tau = (\tau_1, \dots, \tau_t)$ such that $\pi(\tau_1) = \mu$. As in proposition 10, we suppose that there exists a real number x such that:*

$$2r(\lambda, \mu) - c(\lambda, \mu) - q(\lambda, \mu) \geq 4x.$$

If there exists an index i such that $d_i = (b)$ and σ_i^0 has l rows of length one, then:

$$\dim Z_{\tau^0} \geq \dim Z_{\tau} + x + l/2.$$

The proof of proposition 14 will be given together with the proof of proposition 10 at the end of this section.

We want an estimate of the sum of the terms $\max\{d_i^a, d_i^b\}$ depending only on λ, μ .

Lemma 15 *Let $\lambda \geq \mu$ be partitions, so that we can define d_i^a, d_i^b for all $i = 1, \dots, t$. Then*

$$\sum_{i=1}^t \max\{d_i^a, d_i^b\} \leq c(\lambda, \mu) + q(\lambda, \mu). \tag{16}$$

Proof First we prove the claim for $q(\lambda, \mu) = 1$.

In this case, μ is obtained from λ by moving down a single box B . Let us call c_1 (resp. c_2) the column where B lies in μ (resp. λ); so $c_1 < c_2$. By (15), one gets $d_i^a = 1$ for each $c_1 < i \leq c_2$, and 0 otherwise. Moreover one also gets that $d_i^b = 1$ for each $c_1 \leq i < c_2$ and 0 otherwise. Therefore $\max\{d_i^a, d_i^b\} = 1$ for all $c_1 \leq i \leq c_2$ and 0 otherwise. Therefore

$$\sum_{i=1}^t \max\{d_i^a, d_i^b\} = c_2 - c_1 + 1,$$

while

$$c(\lambda, \mu) = c_2 - c_1,$$

so

$$\sum_{i=1}^t \max\{d_i^a, d_i^b\} = c(\lambda, \mu) + 1 \tag{17}$$

and the conclusion holds in this case.

In the general case, let $q := q(\lambda, \mu)$ so, by Lemma 8, we get a sequence of partitions

$$\lambda = \lambda^0 > \lambda^1 > \dots > \lambda^q = \mu$$

such that $q(\lambda^j, \lambda^{j+1}) = 1$ for each $j = 0, \dots, q - 1$. By (17), for every j , we have

$$\sum_{i=1}^t \max\{d_i^a, d_i^b\}(\lambda^j, \lambda^{j+1}) = c(\lambda^j, \lambda^{j+1}) + 1$$

and, summing up over j ,

$$\sum_{i=1}^t \sum_{j=0}^{q-1} \max\{d_i^a, d_i^b\}(\lambda^j, \lambda^{j+1}) = \sum_{j=0}^{q-1} c(\lambda^j, \lambda^{j+1}) + q.$$

By (8), the function c is additive on the pairs $(\lambda^j, \lambda^{j+1})$, therefore the right hand side is equal to $c(\lambda, \mu) + q$. Also d_i^a and d_i^b are additive functions on the pairs $(\lambda^j, \lambda^{j+1})$ (as it follows from (15)), therefore

$$\begin{aligned} \sum_{j=0}^{q-1} \max\{d_i^a, d_i^b\}(\lambda^j, \lambda^{j+1}) &= \sum_{j=0}^{q-1} \max\left\{d_i^a(\lambda^j, \lambda^{j+1}), d_i^b(\lambda^j, \lambda^{j+1})\right\} \\ &\geq \max\left\{\sum_{j=0}^{q-1} d_i^a(\lambda^j, \lambda^{j+1}), \sum_{j=0}^{q-1} d_i^b(\lambda^j, \lambda^{j+1})\right\} \\ &= \max\left\{d_i^a(\lambda, \mu), d_i^b(\lambda, \mu)\right\} = \max\{d_i^a, d_i^b\} \end{aligned}$$

and the conclusion follows. □

We are finally able to prove both Proposition 10 and Proposition 14.

Proof of Proposition 10 and Proposition 14 We start from the dimension formula of the strata (4). We recall that $\Delta(\tau^0) = 0$ as no ab -diagram of τ^0 has rows starting with b . Therefore

$$\begin{aligned} 4(\dim Z_{\tau^0} - \dim Z_{\tau} - x) &= 2(\dim C_{\pi(\tau_1^0)} - \dim C_{\pi(\tau_1)}) + \\ &\quad + \left(o(\tau^0) - 2\Delta(\tau^0)\right) - \left(o(\tau) - 2\Delta(\tau)\right) - 4x \\ &= 2(\dim C_{\lambda} - \dim C_{\mu}) + o(\tau^0) - o(\tau) + 2\Delta(\tau) - 4x \\ &= \sum_{i=1}^t \left(\widehat{\mu}_i^2 - \widehat{\lambda}_i^2\right) + \sum_{i=1}^t \left(o(\tau_i^0) - o(\tau_i) + 2\Delta(\tau_i)\right) - 4x \\ &\text{(by Lemma 11)} = 2r(\lambda, \mu) + \sum_{i=1}^t \left(o(\tau_i^0) - o(\tau_i) + 2\Delta(\tau_i)\right) - 4x \end{aligned}$$

$$\begin{aligned}
 \text{(by (14))} &= 2r(\lambda, \mu) + \sum_{i=1}^t \left(o(\sigma_i^0) - o(\tau_i) + 2\Delta(\tau_i) \right) - 4x \\
 &\geq 2r(\lambda, \mu) - 4x + \\
 &\quad - \sum_{i=1}^t \max_{\tau_i \in \text{aug}_{\sigma_i^0}(d_i)} \left(o(\tau_i) - 2\Delta(\tau_i) - o(\sigma_i^0) \right) \\
 \text{(by Lemma 12)} &\geq 2r(\lambda, \mu) - \sum_{i=1}^t \max\{d_i^a, d_i^b\} - 4x \\
 \text{(by Lemma 15)} &\geq 2r(\lambda, \mu) - c(\lambda, \mu) - q(\lambda, \mu) - 4x.
 \end{aligned}$$

Therefore $\dim Z_{\tau^0} - \dim Z_{\tau} \geq x$ is guaranteed as soon as $2r(\lambda, \mu) - c(\lambda, \mu) - q(\lambda, \mu) \geq 4x$.

In order to obtain the sharper result of proposition 14, it is enough to use Lemma 13 in place of Lemma 12 in the second to last step. \square

7 Complete Intersection Conditions

In this section we want to give condition under which we can make sure that Z is a complete intersection variety.

Proposition 16 *Let λ be a partition and recall that $Z = \Phi^{-1}(0) \subseteq M$. We have*

$$\text{codim}_M(Z_{\tau^0}) = \dim N.$$

Proof In order to prove the equality of the required dimensions, we will work with the dimension formula given by (4). We are going to prove that

$$\dim Z_{\tau^0} = \dim M - \dim N. \tag{18}$$

We immediately have:

$$\dim M = \sum_{i=1}^t \dim L(V_{i-1}, V_i) = \sum_{i=1}^t n_{i-1}n_i$$

and

$$\dim N = \sum_{i=1}^{t-1} \dim \mathfrak{p}(V_i) = \sum_{i=1}^{t-1} \frac{1}{2}n_i^2 + \sum_{i=1}^{t-1} \frac{1}{2}n_i,$$

so

$$\dim M - \dim N = \sum_{i=1}^t n_{i-1}n_i - \frac{1}{2} \sum_{i=1}^{t-1} n_i^2 - \frac{1}{2} \sum_{i=1}^{t-1} n_i.$$

On the other hand, we have:

$$\begin{aligned}
 \dim Z_{\tau^0} &= \frac{1}{2} \dim C_{\pi(\tau^0)} + \sum_{i=0}^{t-1} \left(\frac{1}{2}n_i n_{i+1} - \frac{1}{4}(n_i + n_{i+1}) \right) + \\
 &+ \sum_{i=1}^t \left(\frac{1}{4}o(\tau_i^0) - \frac{1}{2}\Delta(\tau_i^0) \right).
 \end{aligned}$$

We have that $\pi(\tau_1^0) = \lambda$; we then use formula (2).

We also have that $o(\tau_i^0)$ is precisely the number of rows of τ_i^0 (as they all have odd length), the number of rows of τ_i^0 is equal to the number of rows of $\lambda^i := \pi(\tau_i^0)$, the partition λ^i has clearly $\widehat{\lambda}_1^i$ rows and $\widehat{\lambda}_1^i = \widehat{\lambda}_i$, that is the i th column of λ .

Finally, we have that $\Delta(\tau_i^0) = 0$, because there are no rows in τ_i^0 starting with b . Therefore:

$$\dim Z_{\tau^0} = \frac{1}{4} \left(n_0^2 - \sum_{i=1}^t \widehat{\lambda}_i^2 \right) + \sum_{i=0}^{t-1} \left(\frac{1}{2} n_i n_{i+1} - \frac{1}{4} (n_i + n_{i+1}) \right) + \frac{1}{4} \sum_{i=1}^t \widehat{\lambda}_i;$$

we recall that $\widehat{\lambda}_i = n_{i-1} - n_i$ and that $n_0 = n = |\lambda|, n_t = 0$; so

$$\begin{aligned} \dim Z_{\tau^0} &= \frac{1}{4} \left(n_0^2 - \sum_{i=1}^t (n_{i-1} - n_i)^2 \right) + \\ &\quad + \frac{1}{2} \sum_{i=0}^{t-1} n_i n_{i+1} - \frac{1}{4} n_0 - \frac{1}{2} \sum_{i=1}^{t-1} n_i + \frac{1}{4} \sum_{i=1}^t n_{i-1} - n_i \\ &= -\frac{1}{2} \sum_{i=1}^{t-1} n_i^2 + \frac{1}{2} \sum_{i=1}^{t-1} n_{i-1} n_i + \frac{1}{2} \sum_{i=0}^{t-1} n_i n_{i+1} - \frac{1}{2} \sum_{i=1}^{t-1} n_i \\ &= \dim M - \dim N. \end{aligned}$$

□

We can proceed to the main result of this section.

Proposition 17 *If λ satisfies the 2-step condition, then Z is a complete intersection variety.*

The argument is basically the same used in [5, Theorem in 3.3] and [7, Theorem in 5.3]. Here we merely put the pieces together. One of these pieces is the following technical lemma.

Lemma 18 *If the partition λ satisfies the 2-step condition and $\tau \in \Lambda$ is a string of ab -diagrams different from τ^0 , then*

$$\dim Z_{\tau^0} > \dim Z_{\tau}.$$

Proof of Proposition 17 We start recalling Proposition 6 and Proposition 16.

As $Z \setminus Z_{\tau^0}$ consists of finitely many strata Z_{τ} and, by Lemma 18, $\text{codim}_Z(Z_{\tau}) \geq 1$, we deduce that $Z = \overline{Z_{\tau^0}}$. That implies that Z is a complete intersection variety smooth in codimension 0 and Z is reduced as a scheme. □

We can finally use Lemma 9 and Propositions 10 and 14 to prove Lemma 18.

Proof of Lemma 18 Let $\mu = \pi(\tau)$, so in particular $\mu < \lambda$, as $\tau \neq \tau^0$. Therefore Lemma 9 implies

$$2r(\lambda, \mu) \geq c(\lambda, \mu) + q(\lambda, \mu) \tag{19}$$

and, by combining it with proposition 10 we immediately get

$$\dim Z_{\tau^0} \geq \dim Z_{\tau}.$$

So our concern is reduced to obtain a strict inequality. We will gain the strict inequality either by proving that (19) holds strictly for the pair (λ, μ) or by using proposition 14 with $l > 0$.

Let c be the first column such that $\widehat{\mu}_c > \widehat{\lambda}_c$ and let $r = \widehat{\mu}_c$. This means that the first $c - 1$ columns of λ and μ are the same and $\mu_r > \lambda_r$. On the basis of c and r , we distinguish three cases.

$\widehat{\mu}_c - \widehat{\lambda}_c \geq 2$: in this case we can use Lemma 9 with the column c to obtain that

$$2r(\lambda, \mu) > c(\lambda, \mu) + q(\lambda, \mu).$$

$\widehat{\mu}_c - \widehat{\lambda}_c = 1, \mu_r - \lambda_r \geq 2$: we can still use Lemma 9, this time with the row r , to obtain that

$$2r(\lambda, \mu) > c(\lambda, \mu) + q(\lambda, \mu).$$

$\widehat{\mu}_c - \widehat{\lambda}_c = 1, \mu_r - \lambda_r = 1$: in this case we can use Proposition 14 with $i = c$ and $l \geq 1$. Indeed, the following two facts are immediately seen: by (15), $d_c = (b)$; the ab -diagram σ_c^0 has at least one row equal to a single a , namely the $\widehat{\mu}_c$ th row.

□

8 Normality Conditions

In this section we prove Theorem 1. We start by proving the necessary condition as a consequence of the works by Ohta and Sekiguchi.

Theorem 19 (the *only if* part of Theorem 1) *Let λ be a partition of n and suppose that λ does not satisfy the 1-step condition. Then $\overline{C_\lambda} \subseteq \mathfrak{p}$ is not normal.*

Proof If λ does not satisfy the 1-step condition, then there is an i such that $\lambda_i \geq \lambda_{i+1} + 2$. Let

$$\mu = (\lambda_1, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1} + 1, \dots, \lambda_h)$$

be another partition of n , so that $\mu < \lambda$ and μ is a minimal degeneration of λ (in the same sense of [6, 10]). Let $m = \lambda_i - \lambda_{i+1} \geq 2$ and let us define the following two partitions of m :

$$\begin{aligned} \lambda' &= (m), \\ \mu' &= (m - 1, 1). \end{aligned}$$

The degeneration $\mu' < \lambda'$ is obtained from $\mu < \lambda$ by removing the common rows and columns (of μ and λ). By Theorem [10, Theorem 2], the singularity C_μ of $\overline{C_\lambda}$ is normal if and only if the singularity $C_{\mu'}$ of $\overline{C_{\lambda'}}$ is normal.

Moreover, by construction, $C_{\lambda'}$ is the variety of the principal (or regular) nilpotent elements of \mathfrak{p} (in dimension m); therefore we also have that $\overline{C_{\lambda'}} = \mathcal{N}(\mathfrak{p})$, that is the cone of the nilpotent elements in \mathfrak{p} . As μ' is the minimal degeneration of λ' , $C_{\mu'}$ is the variety of subregular nilpotent elements of \mathfrak{p} . The Theorem [11, §3.1, Theorem 7]. proves that the singularity $C_{\mu'}$ is not normal in $\mathcal{N}(\mathfrak{p})$. So this concludes the *only if* part. \square

We turn to the sufficient condition. We will make use of the construction of the variety Z .

Theorem 20 (the *if* part of Theorem 1) *If the partition λ satisfies the 1-step condition, then Z is a normal variety. In particular $\overline{C_{\lambda}}$ is a normal variety.*

As λ satisfies the 1-step condition, also λ satisfies the 2-step condition, so, by proposition 17, Z is a complete intersection variety. At this point the proof of Theorem 20 is a direct consequence of the following Lemma, as in [5, Section 3.7].

Lemma 21 *If the partition λ satisfies the 1-step condition, then $\dim(Z \setminus Z_{\tau^0}) \leq \dim Z - 2$.*

Proof Let $\tau \in \Lambda$ such that $\tau \neq \tau^0$. Thus $\mu := \pi(\tau_1) < \lambda$. We need to prove that

$$\dim Z_{\tau^0} - \dim Z_{\tau} \geq 2.$$

By Lemma 9, $r(\lambda, \mu) \geq c(\lambda, \mu) + q(\lambda, \mu)$ and hence

$$2r(\lambda, \mu) - c(\lambda, \mu) - q(\lambda, \mu) \geq 4x$$

where $x = \frac{1}{4}(c(\lambda, \mu) + q(\lambda, \mu))$. By proposition 10, we get

$$\dim Z_{\tau^0} - \dim Z_{\tau} \geq x,$$

so we are done if $x > 1$.

Let us assume that $x \leq 1$. For simplicity of notation, we put $c := c(\lambda, \mu)$ and $q := q(\lambda, \mu)$. We claim that there are only four cases left: $q = c = 2$ and $q = 1, c \leq 3$. Indeed, by (6), $\lambda > \mu$ if and only if both $c \geq 1$ and $q \geq 1$. By (10), we also have $c \geq q$. Since $4x = c + q \leq 4$, then $q < 3$. Indeed, $q \geq 3 \Rightarrow c \geq 3 \Rightarrow c + q \geq 6$. If $q = 2$, then $c \leq 2$ by the same reasoning. If $q = 1$, then $c = 4x - q \leq 4 - 1$.

We analyze the four cases separately and, in each case, we find an index i such that $d_i = (b)$, the ab -diagram σ_i^0 has l rows of length one and l is big enough so that we can conclude the proof by proposition 14 as follows:

$$\dim Z_{\tau^0} - \dim Z_{\tau} \geq x + \frac{l}{2} = \frac{1}{4}(c + q) + \frac{l}{2} > 1.$$

$q = 2, c = 2$ There are two boxes B, B' which are lowered from λ in order to obtain μ . Let i' be the column of B in λ and i be the column of B in μ . As $c = 2, i = i' - 1$. By (15), the list $d_i = (b)$. In σ_i^0 the rows with indices $\widehat{\mu}_i = \widehat{\lambda}_i + 1, \widehat{\lambda}_i$ and $\widehat{\lambda}_i - 1$ have length one. Indeed, since λ is 1-step, λ does not have two different columns with the same height. Thus, σ_i^0 has $l \geq 3$ rows of length one. We found that $\frac{1}{4}(c + q) + \frac{l}{2} \geq \frac{5}{2} > 1$.

$q = 1, c = 3$ There is a single box B which is lowered from λ in order to obtain μ .

Let i' be the column of B in λ and i be the column of B in μ . As $c = 3, i = i' - 3$.

By (15), the list $d_i = (b)$. In σ_i^0 the rows with indices $\widehat{\mu}_i, \widehat{\mu}_i - 1$ have length one.

Thus, σ_i^0 has $l \geq 2$ rows of length one. We found that $\frac{1}{4}(c + q) + \frac{1}{2} \geq 2 > 1$.

$q = 1, c = 2$ Similar to the previous case, there is a single box B moving from column $i' = i + 2$ to column i ; $d_i = (b)$; the rows with indices $\widehat{\mu}_i, \widehat{\mu}_i - 1$ have length one; so σ_i^0 has $l \geq 2$ rows of length one. We found that $\frac{1}{4}(c + q) + \frac{1}{2} \geq \frac{7}{4} > 1$.

$q = 1, c = 1$ There is a single box B moving from column $i' = i + 1$ to column i .

Similar to the case $q = 2, c = 2, d_i = (b)$, the rows with indices $\widehat{\mu}_i = \widehat{\lambda}_i + 1, \widehat{\lambda}_i$ and $\widehat{\lambda}_i - 1$ have length one and σ_i^0 has $l \geq 3$ rows of length one. We found that

$\frac{1}{4}(c + q) + \frac{1}{2} \geq 2 > 1$. □

Declarations

Conflict of Interest The author declares no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Beckenbach, E.F., Bellman, R.: Inequalities, vol. 30. Springer Science & Business Media, Berlin (2012)
2. Broer, B.: Normality of some nilpotent varieties and cohomology of line bundles on the cotangent bundle of the flag variety. In: Lie theory and geometry, pp. 1–19. Springer, Berlin (1994)
3. Hesselink, W.: Singularities in the nilpotent scheme of a classical group. Transactions of the American Mathematical Society **222**, 1–32 (1976)
4. Kostant, B.: Lie group representations on polynomial rings. American Journal of Mathematics **85**(3), 327–404 (1963)
5. Kraft, H., Procesi, C.: Closures of conjugacy classes of matrices are normal. Inventiones mathematicae **53**(3), 227–247 (1979)
6. Kraft, H., Procesi, C.: Minimal singularities in GL_n . Inventiones mathematicae **62**(3), 503–515 (1980)
7. Kraft, H., Procesi, C.: On the geometry of conjugacy classes in classical groups. Commentarii Mathematici Helvetici **57**(1), 539–602 (1982)
8. Kostant, B., Rallis, S.: Orbits and representations associated with symmetric spaces. American Journal of Mathematics **93**(3), 753–809 (1971)
9. Magyar, P., Weyman, J., Zelevinsky, A.: Symplectic multiple flag varieties of finite type I. Journal of Algebra **230**, 245–265 (2000)
10. Ohta, T.: The singularities of the closures of nilpotent orbits in certain symmetric pairs. Tohoku Mathematical Journal, Second Series **38**(3), 441–468 (1986)
11. Sekiguchi, J.: The nilpotent subvariety of the vector space associated to a symmetric pair. Publications of the Research Institute for Mathematical Sciences **20**(1), 155–212 (1984)
12. Sommers, E.: Normality of nilpotent varieties in E_6 . Journal of Algebra **270**(1), 288–306 (2003)

13. Sommers, E.: Normality of very even nilpotent varieties in D_{2l} . *Bulletin of the London Mathematical Society* **37**(3), 351–360 (2005)
14. Vinberg, E.B.: The Weyl group of a graded Lie algebra. *Mathematics of the USSR-Izvestiya* **10**(3), 463 (1976)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.