

The L^2 –Atiyah-Bott-Lefschetz theorem on manifolds with conical singularities. A heat kernel approach.

Francesco Bei

Dipartimento di Matematica, La Sapienza Università di Roma
E-mail addresses: bei@mat.uniroma1.it, francescobei27@gmail.com

Abstract

Using an approach based on the heat kernel we prove an Atiyah-Bott-Lefschetz theorem for the L^2 –Lefschetz numbers associated to an elliptic complex of cone differential operators over a compact manifold with conical singularities. We then apply our results to the case of the de Rham complex.

Keywords: Atiyah-Bott-Lefschetz theorem, elliptic complexes, differential cone operators, heat kernel, geometric endomorphisms, manifolds with conical singularities.

Mathematics subject classification: 58J10, 14F40, 14F43.

Introduction

The Atiyah-Bott-Lefschetz theorem for elliptic complexes, see [2], is a landmark of elliptic theory on closed manifold. After its publication in 1969, several papers have been devoted to this theorem, to explore its applications, to investigate new approaches to its proof and to find some generalizations. For example in [3] the authors use their first paper to explore applications to the classical elliptic complexes arising in differential geometry; in [7], [18], [24], [25] and [31] the heat kernel approach is developed, while in [6] an approach using probabilistic methods is employed. In [8], [29], [30] [34],[35], [37] and [38] the Atiyah-Bott-Lefschetz theorem is extended to some kind of manifolds that are not closed: for example [29] is devoted to the case of elliptic conic operators on manifold with conical singularities, in [34] the case of a manifold with cylindrical ends is studied and in [35] the case of a complex of Hecke operators over an arithmetic variety is studied. In particular the use of the heat kernel turned out to be a powerful tool in order to get alternative proofs and extensions of the theorem. Since the heat kernel associated to a conic operator has been intensively studied in the last thirty years, e.g. [10], [11] [12], [13], [15],[26] and [28], it is interesting to explore its applications in this context as well, that is to prove an Atiyah-Bott-Lefschetz theorem over a manifold with conical singularities using the heat kernel. This is precisely the goal of this paper.

Our geometric framework is the following: given a compact and orientable manifold with isolated conical singularities X , we consider over its regular part, $reg(X)$ (usually labeled M), a complex of elliptic conic differential operators:

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \xrightarrow{P_n} 0 \quad (1)$$

and a geometric endomorphism $T = (T_0, \dots, T_n)$ of the complex, that is for each $i = 0, \dots, n$, $T_i = \phi_i \circ f^*$ where $f : X \rightarrow X$ is an isomorphism and $\phi_i : f^*E_i \rightarrow E_i$ is a bundle homomorphism.

Using a conic metric over M we associate to (1) two Hilbert complexes $(L^2(M, E_i), P_{max/min,i})$ and then, as first step, we recall the following important property:

- The cohomology groups of $(L^2(M, E_i), P_{max/min,i})$ are finite dimensional.

This result follows directly from the Fredholm property of elliptic cone operators, see [26] Prop. 1.3.16 or [17] Prop. 3.14.

Afterwards assuming that f satisfies the following condition:

$$f : \overline{M} \rightarrow \overline{M} \text{ is a diffeomorphism}$$

(where \overline{M} is a manifold with boundary which desingularizes X , see Prop. 6, and f is supposed to admit an extension on the whole \overline{M}), we prove that:

- Each T_i extends to a bounded map acting on $L^2(M, E_i)$ such that $(T_{i+1} \circ P_{max/min,i})(s) = (P_{max/min,i} \circ T_i)(s)$ for each $s \in \mathcal{D}(P_{max/min,i})$.

In this way we can associate to T and (1) two L^2 -Lefschetz numbers $L_{2,max/min}(T)$ defined as

$$L_{2,max/min}(T) := \sum_{i=0}^n (-1)^i \text{Tr}(T_i^* : H_{2,max/min}^i(M, E_i) \rightarrow H_{2,max/min}^i(M, E_i)) \quad (2)$$

Subsequently, using the operators $\mathcal{P}_i := P_i^t \circ P_i + P_{i-1} \circ P_{i-1}^t$, its absolute and relative extension and the fact that respective heat operators $e^{-t\mathcal{P}_{abs/rel,i}} : L^2(M, E_i) \rightarrow L^2(M, E_i)$ are trace-class operators we prove the following results:

- $L_{2,max/min}(T) = \sum_{i=0}^n (-1)^i \text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}})$ for every $t > 0$.

After this, to improve the above formula, we require some particular properties about f ; more precisely we require that:

- f fixes each singular point of X .
- $Fix(f)$, the fixed points of f , is made only by *simple fixed points*.

The second requirement means that if $f(q) = q$ and $q \in M$ then the diagonal of $M \times M$ is transverse to the graph on f in (q, q) while if $f(q) = q$ and $q \in \text{sing}(X)$ then it means the following: over a neighborhood U_q of q , $U_q \cong C_2(L_q)$ the cone over L_q , f takes clearly the form

$$f(r, p) = (rA(r, p), B(r, p)). \quad (3)$$

(We make the additional assumption that $A(r, p) : [0, 1] \times L_q \rightarrow [0, 1]$ and $B(r, p) : [0, 1] \times L_q \rightarrow L_q$ are smooth up to zero). Then we will say that the fixed point is a **simple** fixed point if for each $p \in L_q$ at least one of the following conditions is satisfied (for more details see Definition 15):

1. $A(0, p) \neq 1$.
2. $B(0, p) \neq p$.

Under this conditions, we prove the formula below:

$$L_{2,max/min}(T) = \lim_{t \rightarrow 0} \left(\sum_{q \in Fix(f)} \sum_{i=0}^n (-1)^i \int_{U_q} \text{tr}(\phi_i \circ k_{abs/rel,i}(t, f(x), x)) d\text{vol}_q \right)$$

where $\phi_i \circ k_{abs/rel,i}(t, f(x), x)$ is the smooth kernel of $T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}$ and U_q is a neighborhood of q (obviously when $q \in \text{sing}(X)$ then we mean the regular part of U_q). Moreover under some additional hypothesis, in particular that (3) modifies in the following way:

$$f(r, p) = (rA(p), B(p)) \quad (4)$$

we prove the following formulas, (see Theorem 7), which are the main result of the paper :

$$L_{2,max/min}(T) = \sum_{q \in Fix(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\phi_i)}{|\det(\text{Id} - d_q(f))|} + \sum_{q \in \text{sing}(X)} \sum_{i=0}^n (-1)^i \zeta_{T_i, q}(\mathcal{P}_{abs/rel,i})(0) \quad (5)$$

$$\zeta_{T_i,q}(\mathcal{P}_{abs/rel,i})(0) = \frac{1}{2\nu} \int_0^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{abs/rel,i}}(A(p), B(p), 1, p)) d\text{vol}_h. \quad (6)$$

Finally, in the last part of the paper, we apply the previous results to the de Rham complex. We get an analytic construction of the Lefschetz numbers arising in intersection cohomology and a topological interpretation of the contributions given by the singular points to the L^2 -Lefschetz numbers. In particular, under suitable conditions, we prove the following formula:

$$\begin{aligned} I^m L(f) = L_{2,max}(T) &= \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn } \det(\text{Id} - d_q f) + \\ &+ \sum_{q \in \text{sing}(X)} \sum_{i < \frac{m+1}{2}} (-1)^i \text{Tr}(B^* : H^i(L_q) \rightarrow H^i(L_q)) \end{aligned} \quad (7)$$

where $I^m L(f)$ is the intersection Lefschetz number arising in intersection cohomology, T is the endomorphism of $(L^2\Omega^i(M, g), d_{max,i})$ induced by f and B is the diffeomorphism of the link L_q such that, in a neighborhood of q , f satisfies (4). In particular from (7) we get:

$$\sum_{i=0}^{m+1} (-1)^i \zeta_{T_i,q}(\Delta_{abs,i})(0) = \sum_{i < \frac{m+1}{2}} (-1)^i \text{Tr}(B^* : H^i(L_q) \rightarrow H^i(L_q)). \quad (8)$$

As recalled at the beginning of the introduction also [29] is devoted to the Atiyah-Bott-Lefschetz theorem on manifold with conical singularities. Anyway there are some substantial differences between our paper and [29]: the notion of ellipticity used there, which is taken from [33], is stronger than that one used in this paper; in particular the de Rham complex is not elliptic for the definition given in [33]. Moreover the complexes considered in [29] are complexes of weighted Sobolev space while our complexes are Hilbert complexes of unbounded operator defined on some natural extensions of their core domain; finally also the techniques used are different because we use the heat kernel while in [29] the existence of a parametrix of an elliptic cone operator is used. Some results of this paper are also close to results proved in [26]: indeed in [26] the heat kernel is studied in an equivariant situation and an equivariant index theorem is proved (see Corollary 2.4.7). Also in this case there are some relevant differences: the Lie group G acting in [26] is a *compact* Lie group of isometry, while in our work we just require that the map f is a diffeomorphism. Moreover the non degeneracy conditions that we require on the fixed point of f led us to different formulas to those stated in [26]. On the other hand, for the geometric endomorphisms considered in [26], that is those induced by isometries g lying in a compact Lie group G , the formula obtained by Lesch applies to a more general case than the ours because in his work there are not assumptions on the fixed points set while in our work there are.

Moreover, as recalled above, the last part of this paper contains several applications to the de Rham complex which are not mentioned in the other papers.

Acknowledgment. I wish to thank Paolo Piazza for having suggested this subject, for his help and for many helpful discussions. I wish also to thank Pierre Albin for having invited me to spend the months of March and April 2012 at the University of Illinois at Urbana-Champaign and for many interesting discussions.

1 Background

1.1 Hilbert complexes

In this first subsection we recall briefly the notion of Hilbert complex and how it appears in riemannian geometry. We refer to [9] for a thorough discussion about this subject.

Definition 1. A Hilbert complex is a complex, (H_*, D_*) of the form:

$$0 \rightarrow H_0 \xrightarrow{D_0} H_1 \xrightarrow{D_1} H_2 \xrightarrow{D_2} \dots \xrightarrow{D_{n-1}} H_n \rightarrow 0, \quad (9)$$

where each H_i is a separable Hilbert space and each map D_i is a closed operator called the differential such that:

1. $\mathcal{D}(D_i)$, the domain of D_i , is dense in H_i .
2. $\text{ran}(D_i) \subset \mathcal{D}(D_{i+1})$.
3. $D_{i+1} \circ D_i = 0$ for all i .

The cohomology groups of the complex are $H^i(H_*, D_*) := \text{Ker}(D_i)/\text{ran}(D_{i-1})$. If the groups $H^i(H_*, D_*)$ are all finite dimensional we say that it is a *Fredholm complex*.

Given a Hilbert complex there is a dual Hilbert complex

$$0 \leftarrow H_0 \xleftarrow{D_0^*} H_1 \xleftarrow{D_1^*} H_2 \xleftarrow{D_2^*} \dots \xleftarrow{D_{n-1}^*} H_n \leftarrow 0, \quad (10)$$

defined using $D_i^* : H_{i+1} \rightarrow H_i$, the Hilbert space adjoints of the differentials $D_i : H_i \rightarrow H_{i+1}$. The cohomology groups of $(H_j, (D_j)^*)$, the dual Hilbert complex, are

$$H^i(H_j, (D_j)^*) := \text{Ker}(D_{n-i-1}^*)/\text{ran}(D_{n-i}^*).$$

For all i there is also a laplacian $\Delta_i = D_i^* D_i + D_{i-1} D_{i-1}^*$ which is a self-adjoint operator on H_i with domain

$$\mathcal{D}(\Delta_i) = \{v \in \mathcal{D}(D_i) \cap \mathcal{D}(D_{i-1}^*) : D_i v \in \mathcal{D}(D_i^*), D_{i-1}^* v \in \mathcal{D}(D_{i-1})\} \quad (11)$$

and nullspace:

$$\mathcal{H}^i(H_*, D_*) := \ker(\Delta_i) = \text{Ker}(D_i) \cap \text{Ker}(D_{i-1}^*). \quad (12)$$

The following propositions are standard results for these complexes. The first result is a weak Kodaira decomposition:

Proposition 1. [[9], Lemma 2.1] Let (H_i, D_i) be a Hilbert complex and $(H_i, (D_i)^*)$ its dual complex, then:

$$H_i = \mathcal{H}^i \oplus \overline{\text{ran}(D_{i-1})} \oplus \overline{\text{ran}(D_i^*)}.$$

The reduced cohomology groups of the complex are:

$$\overline{H}^i(H_*, D_*) := \text{Ker}(D_i)/\overline{(\text{ran}(D_{i-1}))}.$$

By the above proposition there is a pair of weak de Rham isomorphism theorems:

$$\begin{cases} \mathcal{H}^i(H_*, D_*) \cong \overline{H}^i(H_*, D_*) \\ \mathcal{H}^i(H_*, D_*) \cong \overline{H}^{n-i}(H_*, (D_*)^*) \end{cases} \quad (13)$$

where in the second case we mean the cohomology of the dual Hilbert complex.

The complex (H_*, D_*) is said *weak Fredholm* if $\mathcal{H}_i(H_*, D_*)$ is finite dimensional for each i . By the next propositions it follows immediately that each Fredholm complex is a weak Fredholm complex.

Proposition 2. [[9], Corollary 2.5] If the cohomology of a Hilbert complex (H_*, D_*) is finite dimensional then, for all i , $\text{ran}(D_{i-1})$ is closed and $H^i(H_*, D_*) \cong \mathcal{H}^i(H_*, D_*)$.

Proposition 3 ([9], Corollary 2.6). A Hilbert complex (H_j, D_j) , $j = 0, \dots, n$ is a Fredholm complex (weak Fredholm) if and only if its dual complex, (H_j, D_j^*) , is Fredholm (weak Fredholm). If it is Fredholm then

$$\mathcal{H}_i(H_j, D_j) \cong H_i(H_j, D_j) \cong H_{n-i}(H_j, (D_j)^*) \cong \mathcal{H}_{n-i}(H_j, (D_j)^*). \quad (14)$$

Analogously in the the weak Fredholm case we have:

$$\mathcal{H}_i(H_j, D_j) \cong \overline{H}_i(H_j, D_j) \cong \overline{H}_{n-i}(H_j, (D_j)^*) \cong \mathcal{H}_{n-i}(H_j, (D_j)^*). \quad (15)$$

Proposition 4. *A Hilbert complex (H_j, D_j) , $j = 0, \dots, n$ is a Fredholm complex if and only if for each i the operator Δ_i defined in (11) is a Fredholm operator on its domain endowed with the graph norm.*

Proof. See [33], Lemma 1 pag 203. □

Now we recall another result which shows that it is possible to compute the cohomology groups of an Hilbert complex using a core subcomplex

$$\mathcal{D}^\infty(H_i) \subset H_i.$$

For all i we define $\mathcal{D}^\infty(H_i)$ as consisting of all elements η that are in the domain of Δ_i^l for all $l \geq 0$.

Proposition 5 ([9], Theorem 2.12). *The complex $(\mathcal{D}^\infty(H_i), D_i)$ is a subcomplex quasi-isomorphic to the complex (H_i, D_i)*

As it is well known, riemannian geometry offers a framework in which Hilbert and (sometimes) Fredholm complexes can be built in a natural way. The rest of this subsection is devoted to recall these constructions.

Let (M, g) be an open and oriented riemannian manifold of dimension m and let E_0, \dots, E_n be vector bundles over M . For each $i = 0, \dots, n$ let $C_c^\infty(M, E_i)$ be the space of smooth section with compact support. If we put on each vector bundle a metric h_i $i = 0, \dots, n$ the we can construct in a natural way a sequences of Hilbert space $L^2(M, E_i)$, $i = 0, \dots, n$ as the completion of $C_c^\infty(M, E_i)$. Now suppose that we have a complex of differential operators :

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} C_c^\infty(M, E_2) \xrightarrow{P_2} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \rightarrow 0, \quad (16)$$

To turn this complex into a Hilbert complex we must specify a closed extension of P_* that is an operator between $L^2(M, E_*)$ and $L^2(M, E_{*+1})$ with closed graph which is an extension of P_* . We start recalling the two canonical closed extensions of P .

Definition 2. *The maximal extension P_{max} ; this is the operator acting on the domain:*

$$\mathcal{D}(P_{max,i}) = \{\omega \in L^2(M, E_i) : \exists \eta \in L^2(M, E_{i+1})\} \quad (17)$$

$$s.t. \langle \omega, P_i^t \zeta \rangle_{L^2(M, E_i)} = \langle \eta, \zeta \rangle_{L^2(M, E_{i+1})} \quad \forall \zeta \in C_0^\infty(M, E_{i+1})\}$$

where P_i^t is the formal adjoint of P_i .

In this case $P_{max,i}\omega = \eta$. In other words $\mathcal{D}(P_{max,i})$ is the largest set of forms $\omega \in L^2(M, E_i)$ such that $P_i\omega$, computed distributionally, is also in $L^2(M, E_{i+1})$.

Definition 3. *The minimal extension $P_{min,i}$; this is given by the graph closure of P_i on $C_0^\infty(M, E_i)$ respect to the norm of $L^2(M, E_i)$, that is,*

$$\mathcal{D}(P_{min,i}) = \{\omega \in L^2(M, E_i) : \exists \{\omega_j\}_{j \in J} \subset C_0^\infty(M, E_i), \omega_j \rightarrow \omega, P_i\omega_j \rightarrow \eta \in L^2(M, E_{i+1})\} \quad (18)$$

and in this case $P_{min,i}\omega = \eta$

Obviously $\mathcal{D}(P_{min,i}) \subset \mathcal{D}(P_{max,i})$. Furthermore, from these definitions, it follows immediately that

$$P_{min,i}(\mathcal{D}(P_{min,i})) \subset \mathcal{D}(P_{min,i+1}), \quad P_{min,i+1} \circ P_{min,i} = 0$$

and that

$$P_{max,i}(\mathcal{D}(P_{max,i})) \subset \mathcal{D}(P_{max,i+1}), \quad P_{max,i+1} \circ P_{max,i} = 0.$$

Therefore $(L^2(M, E_*), P_{max/min,*})$ are both Hilbert complexes and their cohomology groups, respectively reduced cohomology groups, are denoted respectively by $H_{2,max/min}^i(M, E_*)$ and $\overline{H}_{2,max/min}^i(M, E_*)$.

Another straightforward but important fact is that the Hilbert complex adjoint of $(L^2(M, E_*), P_{max/min,*})$ is $(L^2(M, E_*), P_{min/max,*}^t)$, that is

$$(P_{max,i})^* = P_{min,i}^t, \quad (P_{min,i})^* = P_{max,i}^t. \quad (19)$$

Using Proposition 1 we obtain two weak Kodaira decompositions:

$$L^2(M, E_i) = \mathcal{H}_{abs/rel}^i(M, E_i) \oplus \overline{\text{ran}(P_{max/min,i-1})} \oplus \overline{\text{ran}(P_{min/max,i}^t)} \quad (20)$$

with summands mutually orthogonal in each case. For the first summand on the right, called the absolute or relative Hodge cohomology, we have by (12):

$$\mathcal{H}_{abs/rel}^i(M, E_*) = \text{Ker}(P_{max/min,i}) \cap \text{Ker}(P_{min/max,i-1}^t). \quad (21)$$

We can also consider the two natural laplacians associated to these Hilbert complexes, that is for each i

$$\mathcal{P}_{abs,i} := P_{min,i}^t \circ P_{max,i} + P_{max,i-1} \circ P_{min,i-1}^t \quad (22)$$

and

$$\mathcal{P}_{rel,i} := P_{max,i}^t \circ P_{min,i} + P_{min,i-1} \circ P_{max,i-1}^t \quad (23)$$

with domain described in (11). Using (12) and (13) it follows that the nullspace of (22) is isomorphic to the absolute Hodge cohomology which is in turn isomorphic to the reduced cohomology of the Hilbert complex $(L^2(M, E_*), P_{max,*})$. Analogously, using again (12) and (13), it follows that the nullspace of (23) is isomorphic to the relative Hodge cohomology which is in turn isomorphic to the reduced cohomology of the Hilbert complex $(L^2(M, E_*), P_{min,*})$. Finally we recall that we can define other two Hodge cohomology groups $\mathcal{H}_{max/min}^i(M, E_*)$ defined as

$$\mathcal{H}_{max/min}^i(M, E_*) = \text{Ker}(P_{max/min,i}) \cap \text{Ker}(P_{max/min,i-1}^t). \quad (24)$$

1.2 Manifolds with conical singularities and differential cone operators

Definition 4. Let L an open manifold. The cone over L , usually labeled $C(L)$, is the topological space defined as

$$L \times [0, \infty) / (\{0\} \times L). \quad (25)$$

The truncated cone, usually labeled $C_a(L)$, is defined as

$$L \times [0, a] / (\{0\} \times L). \quad (26)$$

Finally with $\overline{C_a(L)}$ we mean

$$L \times [0, a] / (\{0\} \times L). \quad (27)$$

In both the above cases, with v , we will label the vertex of the cone or the truncated cone, that is $C(L) - (L \times (0, \infty))$, $C_a(L) - (L \times (0, a))$ and $\overline{C_a(L)} - (L \times (0, a])$ respectively.

Definition 5. A manifold with conical singularities X is a metrizable, locally compact, Hausdorff space such that there exists a sequence of points $\{p_1, \dots, p_n, \dots\} \subset X$ which satisfies the following properties:

1. $X - \{p_1, \dots, p_n, \dots\}$ is a smooth open manifold.
2. For each p_i there exist an open neighborhood U_{p_i} , a closed manifold L_{p_i} and a map $\chi_{p_i} : U_{p_i} \rightarrow C_2(L_{p_i})$ such that $\chi_{p_i}(p_i) = v$ and $\chi_{p_i}|_{U_{p_i} - \{p_i\}} : U_{p_i} - \{p_i\} \rightarrow L_{p_i} \times (0, 2)$ is a diffeomorphism.

The regular and the singular part of X are defined as

$$\text{sing}(X) = \{p_1, \dots, p_n, \dots\}, \quad \text{reg}(X) := X - \text{sing}(X) = X - \{p_1, \dots, p_n, \dots\}.$$

The singular points p_i are usually called *conical points* and the smooth closed manifold L_{p_i} is usually called the *link* relative to the point p_i . If X is compact then it is clear, from the above definition, that the sequences of conical points $\{p_1, \dots, p_n, \dots\}$ is made of isolated points and

therefore on X there are just a finite number of conical points.

A manifold with conical singularities is a particular case of a compact smoothly stratified pseudomanifold; more precisely it is a compact smoothly stratified pseudomanifold with depth 1 and with the singular set made of a sequence of isolated points. Since in this paper we will work exclusively with compact manifolds with conical singularities we prefer to omit the definition of smoothly compact stratified pseudomanifold and the notions related to it and refer to [1] for a thorough discussion on this subject.

Remark 1. *Let X be a compact manifold with one conical singularity p and let L_p its link; it follows from Definition 5 that we can decompose X as*

$$X \cong \bar{Y} \cup_{L_p} \overline{C_1(L_p)}$$

where \bar{Y} is a compact manifold with boundary defined as $X - \chi_p^{-1}(C_1(L_p))$. Obviously this decomposition generalizes in a natural way when X has several conical points. As we will see in one of the following sections this decomposition is the starting point to study the heat kernel on X and we will use it to calculate the contribution given by the conical points to the Lefschetz number of some geometric endomorphisms.

Now we recall from [1] a particular case, which is suitable for our purpose, of an important result which describe a blowup process to resolve the singularities of a compact smoothly stratified pseudomanifold.

Proposition 6. *Let X be a compact manifold with conical singularities. Then there exists a manifold with boundary \bar{M} and a blow-down map $\beta : \bar{M} \rightarrow X$ which has the following properties:*

1. $\beta|_M : M \rightarrow \text{reg}(X)$, where M is the interior of \bar{M} , is a diffeomorphism.
2. If N is a connected component of $\partial\bar{M}$ and if $U \cong N \times [0, 1)$ is a collar neighborhood of N then $\beta(U) = N \times [0, 1)/(N \times \{0\})$. In particular $\beta(N) = p$ where p is a conical point of X and N becomes one of the connected components of the link of p .
3. If for each conical point p_i the relative link L_{p_i} is connected, then there is a bijection between the conical points of X and the connected components of $\partial\bar{M}$.

Proof. See [1], Proposition 2.5. □

Now we introduce a class of natural riemannian metrics on these spaces.

Definition 6. *Let X be a manifold with conical singularities. A conic metric g on $\text{reg}(X)$ is riemannian metric with the following property: for each conical point p_i there exists a map χ_{p_i} , as defined in Definition 5, such that*

$$(\phi_{p_i}^{-1})^*(g|_{U_{p_i}}) = dr^2 + r^2 h_{L_{p_i}}(r) \quad (28)$$

where $h_{L_{p_i}}(r)$ depends smoothly on r up to 0 and for each fixed $r \in [0, 1)$ it is a riemannian metric on L_{p_i} . Analogously, if \bar{M} is manifold with boundary and M is its interior part, then g is a conic metric on M if it is a smooth, symmetric section of $T^*\bar{M} \otimes T^*\bar{M}$, degenerate over the boundary, such that over a collar neighborhood U of $\partial\bar{M}$, g satisfies (28) with respect to some diffeomorphism $\chi : U \rightarrow [0, 1) \times \partial\bar{M}$.

The next step is to recall the notion of differential cone operator and its main properties. Before to proceed we introduce some notations that we will use steadily through the paper. Given an open manifold M and two vector bundles E, F over it, with $\text{Diff}^n(M, E, F)$, $n \in \mathbb{N}$, we will label the space of differential operator $P : C_c^\infty(M, E) \rightarrow C_c^\infty(M, F)$ of order n . Given \bar{M} , a manifold with boundary, we will label with N the boundary of \bar{M} and with M the interior part of \bar{M} . Given a vector bundle E over \bar{M} , with E_N we mean the restriction of E on N . Finally each metric ρ over E (riemannian if E is real or hermitian if E is complex) is assumed to be a non degenerate metric up to the boundary. The next definition is taken from [26]:

Definition 7. Let \overline{M} be a manifold with boundary $N = \partial\overline{M}$. Let E, F be two vector bundles on \overline{M} . Let \overline{U}_N be a collar neighborhood of N , $\overline{U}_N \cong [0, \epsilon) \times N$ and let $U_N = \overline{U}_N - N$. A differential cone operator of order $\mu \in \mathbb{N}$ and weight $\nu > 0$ is a differential operator $P : C_c^\infty(M, E) \rightarrow C_c^\infty(M, F)$ such that on U_N it takes the form:

$$P|_{U_N} = x^{-\nu} \sum_{i=0}^{\mu} A_k(-x \frac{\partial}{\partial x})^k \quad (29)$$

where $A_k \in C^\infty([0, \epsilon), \text{Diff}^{\mu-k}(N, E_N, F_N))$ and x is the coordinate on $[0, \epsilon)$. As in [26] we will label with $\text{Diff}_0^{\mu, \nu}(M, E, F)$ the space of differential cone operators between the bundles E and F .

Now we explain what we mean by *differential cone operator* on a manifold X with conical singularities. In the previous definition we recalled the notion of differential cone operator acting on the smooth sections with compact support of two vector bundles E, F defined on a manifold \overline{M} with boundary. In Proposition 6, given a manifold with conical singularities X , we stated the existence of a manifold with boundary \overline{M} endowed with a blow down map $\beta : \overline{M} \rightarrow X$ which desingularizes X . Therefore given two vector bundles E, F on $\text{reg}(X)$ and $P \in \text{Diff}(\text{reg}(X), E, F)$ we will say that P is a differential cone operators if the following properties are satisfied:

1. $\beta^*(E), \beta^*(F)$ that are vector bundles on M , the interior of \overline{M} , extend as smooth vector bundles over the whole \overline{M} . In the same way, if E and F are endowed with metrics ρ_1 and ρ_2 then $\beta^*\rho_1$ and $\beta^*\rho_2$ extend as non degenerate metric up to the boundary of \overline{M} .
2. The differential operator induced by P through β between $C_c^\infty(M, \beta^*E, \beta^*F)$ is a differential cone operator in the sense of Definition 7.

In the rest of the paper, with a slight abuse of notation, we will identify M with $\text{reg}(X)$, E with β^*E , F with β^*F and P with the operator that it induces through β between $C_c^\infty(M, \beta^*E, \beta^*F)$.

Remark 2. We can reformulate Definition 7 in the following way: P is differential cone operator of order μ and weight ν if and only if $x^\nu P$ is a b -differential operator of order μ in the sense of Melrose. For the definition of b -operator and the full development of this subject we refer to the monograph [27]. Using this approach we have $\text{Diff}_0^{\mu, \nu}(M, E, F) = x^{-\nu} \text{Diff}_b^\mu(M, E, F)$. This last point of view is used for example in [17].

Now we introduce the notion of ellipticity:

Definition 8. Let \overline{M} be a manifold with boundary and let E, F be two vector bundles over \overline{M} . Let $P \in \text{Diff}_0^{\mu, \nu}(\overline{M}, E, F)$ and let $\sigma^\mu(P)$ be its principal symbol. Then P is called elliptic if it is elliptic on M in the usual sense and if

$$x^\nu \sigma^\mu(P)(x, p, x^{-1}\tau, \xi) \quad (30)$$

is invertible for $(x, p) \in [0, \epsilon) \times N$ and $(\tau, \xi) \in T^*\overline{M} - \{0\}, \xi \in T_p^*(N)$.

In the above definition there is implicit the natural identification of $T^*\overline{M}|_{[0, \epsilon) \times N}$ with $\mathbb{R} \times T^*N$.

Definition 9. Let \overline{M}, E, F and P be as in the previous definition. The conormal symbol of P , as defined in [26], is the family of differential operators, acting between $C^\infty(N, E_N, F_N)$, defined as

$$\sigma_M^{\mu, \nu}(P)(z) := \sum_{k=0}^{\mu} A_k(0) z^k \quad (31)$$

Now we make some further comments about the notion of ellipticity introduced in Definition 8. The requirement (30) in Definition 8 means that

$$\sum_{k=0}^{\mu} \sigma^{\mu-k}(A_k(x))((p, \xi)) \sigma^k((-x \frac{\partial}{\partial x})^k)(x, x^{-1}\tau) = \sum_{k=0}^{\mu} \sigma^{\mu-k}(A_k(x))((p, \xi)) (-i\tau)^k$$

is invertible. On M this is covered by classical ellipticity and for $x = 0$ it is equivalent to require that (31) is a parameter dependent elliptic family of differential operators with parameters in $i\mathbb{R}$.

Using again the b framework of Melrose, Definition 8 is equivalent to say that the b -principal symbol of $P' := x^\nu P$, that is $\sigma_b^\mu(P') := \sigma^\mu(P')(x, p, x^{-1}\tau, \xi)$, as an object lying in $C^\infty(T_b^*\overline{M}, \text{Hom}(\pi_b^*E, \pi_b^*F))$, where $\pi_b : T_b^*\overline{M} \rightarrow \overline{M}$ is the b -cotangent bundle of \overline{M} , is an isomorphism on $T_b^*\overline{M} - \{0\}$. For further details on these approach see [17] and the relative bibliography.

Finally we remark that in Definition 8 we followed [26] and [17]. This is slightly different from those given, for example, in [29], [30] and [33]. The definition given in these papers, in fact, requires the invertibility of the conormal symbol on a certain weight line (for more details see the above papers). By the fact that we are interested to study the operators on their natural domains, that is the maximal and the minimal one, we can waive this requirement (see [26] pag. 13 for more comments about this).

Finally we conclude this subsection stating an important proposition on the theory of differential cone operators:

Theorem 1. *Let (\overline{M}, g) be a compact and oriented manifold of dimension m with boundary where g is a conic metric over M ; let E, F be two hermitian vector bundles over \overline{M} and let $P \in \text{Diff}_0^{\mu, \nu}(M, E, F)$ be an elliptic differential cone operator.*

1. *Each closed extension $\overline{P} : L^2(M, E) \rightarrow L^2(M, F)$ of P is a Fredholm operator on its domain, $\mathcal{D}(\overline{P})$, endowed with the graph norm.*
2. *Suppose that $E = F$ and that P is positive. Suppose, in addition, that on a collar neighborhood of $\partial\overline{M}$ the metric ρ on E does not depend on r and the conic metric g satisfies $g = dr^2 + r^2h$ where h is any riemannian metric over $\partial\overline{M}$ which does not depend on r . Then, for each positive self-adjoint extension \overline{P} of P , the heat operator $e^{-t\overline{P}} : L^2(M, E) \rightarrow L^2(M, E)$ is a trace-class operator. Moreover \overline{P} is discrete and the sequences of eigenvalues of \overline{P} satisfies $\lambda_j \sim Cj^{\frac{\mu}{m}}$.*

Proof. For the first statement see [26] Prop. 1.3.16 or [17] Prop. 3.14. For the second one see [26] Theorem 2.4.1 and Corollary 2.4.3. \square

1.3 Elliptic complex on manifolds with conical singularities

The aim of this subsection is to define the notion of elliptic complex on a manifold with conical singularities. As for the notion of ellipticity, the definition of elliptic complex on a manifold with conical singularities was introduced in [33], pag. 205, but our definition is slightly different because we waive some requirements about the sequence of conormal symbols on a certain weight line. The reason is still given by the fact that we are interested on the minimal and maximal extension of a complex differential cone operators.

Let \overline{M} be a manifold with boundary, E_0, \dots, E_n a sequence of vector bundle over \overline{M} and consider $P_i \in \text{Diff}_0^{\mu, \nu}(M, E_i, E_{i+1})$ such that

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \xrightarrow{P_n} 0 \quad (32)$$

is a complex. We have the following definition:

Definition 10. *The complex (32) is an elliptic complex if it is an elliptic complex in the usual sense on M and if the sequence*

$$0 \rightarrow \pi^*E_0 \rightarrow \pi^*E_1 \rightarrow \dots \rightarrow \pi^*E_n \rightarrow 0 \quad (33)$$

where the maps are given by $x^\nu \sigma^\mu(P_i)(x, p, x^{-1}\tau, \xi) : \pi_i^*E_i \rightarrow \pi_{i+1}^*E_{i+1}$ is an exact sequence up to $x = 0$ over $T^*\overline{M} - \{0\}$.

With the help of Melrose's b framework we can reformulate the previous definition in the following way: (32) is an elliptic complex if and only if the following sequence is exact over $T_b^*(\overline{M}) - \{0\}$:

$$0 \rightarrow \pi_b^* E_0 \xrightarrow{\sigma_b^\mu(P'_0)} \pi_b^* E_1 \xrightarrow{\sigma_b^\mu(P'_1)} \dots \xrightarrow{\sigma_b^\mu(P'_{n-1})} \pi_b^* E_n \xrightarrow{\sigma_b^\mu(P'_n)} 0 \quad (34)$$

where $P' = x^\nu P$, that is the b -operator naturally associated to P , $\pi_b : T_b^* \overline{M} \rightarrow \overline{M}$ is the b -cotangent bundle and $\sigma_b^\mu(P'_i) \in C^\infty(\overline{M}, \text{Hom}(\pi_b^* E_i, \pi_b^* E_{i+1}))$ is the b -principal symbol of P'_i .

We have the following proposition:

Proposition 7. *Consider a complex of differential cone operators as in (32). Suppose moreover that M is endowed with a conic metric g . Then the complex is an elliptic complex if and only if for each $i = 0, \dots, n$*

$$P_i^t \circ P_i + P_{i-1} \circ P_{i-1}^t : C_c^\infty(M, E_i) \rightarrow C_c^\infty(M, E_i)$$

is an elliptic differential cone operator.

Proof. It is clear that if $P \in \text{Diff}_0^{\mu, \nu}(M, E_i, E_{i+1})$ then also $P^t \in \text{Diff}_0^{\mu, \nu}(M, E_{i+1}, E_i)$ where $P_t : C_c^\infty(M, E_{i+1}) \rightarrow C_c^\infty(M, E_i)$ is the formal adjoint of P . Now, as in the previous comment, let $P'_i = x^\nu P$ be the b -operator that is naturally associated to P . It is well known that $\sigma_b^\mu(P'_{i+1} \circ P'_i) = \sigma_b^\mu(P'_{i+1}) \circ \sigma_b^\mu(P'_i)$ and that $\sigma_b^\mu((P'_i)^t) = (\sigma_b^\mu(P'_i))^t$. The proof follows now by standard arguments of linear algebra, in complete analogy with the case of an elliptic complex on a closed manifold. \square

From the above proposition it follows the following useful corollary:

Corollary 1. *In the same hypothesis of the previous proposition. The Hilbert complexes $(L^2(M, E_*), P_{\max/\min,*})$ are both Fredholm complexes. Moreover each Hilbert complex that extends $(L^2(M, E_*), P_{\min,*})$ and that is extended by $(L^2(M, E_*), P_{\max,*})$ is still an Fredholm complex.*

Proof. From Theorem 1 it follows that $P_{\min,i}^t \circ P_{\max,i} + P_{\max,i-1} \circ P_{\min,i-1}^t$ and $P_{\max,i}^t \circ P_{\min,i} + P_{\min,i-1} \circ P_{\max,i-1}^t$ are both Fredholm operators on their natural domain endowed with the graph norm. Now the statement follows from Prop. 4 \square

We remark the fact that we gave the definition of an elliptic complex of differential cone operators on a manifold with boundary \overline{M} . Following the remark after Definition 7 the notion of elliptic complex of differential cone operators is naturally extended on a manifold X with conical singularities.

1.4 A brief reminder on the heat kernel

The aim of this subsection is to recall briefly the main local properties of the heat kernel on an open and oriented riemannian manifold (M, g) .

Let (M, g) be an open and oriented riemannian manifold, E a vector bundle over M , $P_0 : C_c^\infty(M, E) \rightarrow C_c^\infty(M, E)$ a non-negative symmetric differential operator and $P : \mathcal{D}(P) \subset L^2(M, E) \rightarrow L^2(M, E)$ a non-negative, self-adjoint extension of P_0 . It is well known that, using the spectral theorem for unbounded self-adjoint operators and its associated functional calculus (see [16], chap. XXII), it is possible to construct the operator e^{-tP} . The next result we are going to recall summarizes the main local properties of e^{-tP} that we will use in the rest of the paper. We start with the following definitions:

Definition 11. *A cut-off function is a smooth function $\eta : [0, \infty) \rightarrow [0, 1]$ which admits a $\epsilon > 0$ such that $\eta(x) = 1$ for $x \leq \frac{\epsilon}{4}$ and $\eta = 0$ for $x \geq \epsilon$.*

Definition 12. *Let (M, g) be an open manifold, E a vector bundle over M and $P_0 : C_c^\infty(M, E) \rightarrow C_c^\infty(M, E)$ a differential operator of second order. Then P_0 is a generalized Laplacian if its principal symbol satisfies:*

$$\sigma^2(P_0)(x, \xi) = \|\xi\|^2.$$

An operator of this type is clearly elliptic. We refer to [5] for a comprehensive discussion on this class of operators.

Theorem 2. Let (M, g) be an open and oriented riemannian manifold, E a vector bundle over M , $P_0 : C_c^\infty(M, E) \rightarrow C_c^\infty(M, E)$ a non-negative symmetric differential operator of order d and $P : \mathcal{D}(P) \subset L^2(M, E) \rightarrow L^2(M, E)$ a non-negative, self-adjoint extension of P_0 . Then e^{-tP} satisfies the following properties:

- e^{-tP} has a C^∞ -kernel, that is usually labeled $e^{-tP}(s, q)$ or $k_P(t, s, q)$, which lies in $C^\infty((0, \infty) \times M \times M, E \boxtimes E^*)$.
- If K_1, K_2 are compact subset of M such that $K_1 \cap K_2 = \emptyset$ then

$$\|k_P(t, s, q)\|_{C^k(K_1 \times K_2, E \boxtimes E^*)} = O(t^n), \quad t \rightarrow 0$$

for all $k, n \in \mathbb{N}$.

- Let $\phi, \chi \in C_c^\infty(M)$; then the operator $\phi e^{-tP} \chi$ is a trace-class operator and we have, on $C^l(K_1 \times K_2, E \boxtimes E^*|_{K_1 \times K_2})$ for each $l \in \mathbb{N}$,

$$(\phi e^{-tP} \chi)(q, q) \sim_{t \rightarrow 0} \sum_{n=0}^{\infty} \phi(q) \chi(q) \Phi_n(q) t^{\frac{n-m}{d}}$$

and

$$\text{Tr}((\phi e^{-tP} \chi)(q, q)) \sim_{t \rightarrow 0} \sum_{n=0}^{\infty} \left(\int_M \phi(q) \chi(q) \text{tr}(\Phi_n(q)) d\text{vol}_g \right) t^{\frac{n-m}{d}}$$

where $q \in M$, $\{\Phi_1, \dots, \Phi_n, \dots\}$ is a suitable sequence of sections in $C^\infty(M, \text{End}(E))$, $K_1 = \text{supp}(\phi)$ and $K_2 = \text{supp}(\chi)$.

Finally if P_0 is a generalized Laplacian then the last property above can be sharpened in the following way:

- Let $\phi, \chi \in C_c^\infty(M)$; then the operator $\phi e^{-tP} \chi$ is a trace-class operator and we have

$$\phi(s) e^{-tP}(s, q) \chi(q) \sim_{t \rightarrow 0} h_t(s, q) \sum_{n=0}^{\infty} \phi(s) \chi(q) \Phi_n(s, q) t^n$$

where $(s, q) \in M \times M$, $\{\Phi_1, \dots, \Phi_n, \dots\}$ is a suitable sequence of sections in $C^\infty(M \times M, E \boxtimes E^*)$ and $h_t(s, q) = (4\pi t)^{\frac{-n}{2}} e^{\frac{-d(s, q)^2}{4t}} \eta(d(s, q)^2)$ with η a cut-off function. As in the previous case the above expansion holds in $C^l(K_1 \times K_2, E \boxtimes E^*|_{K_1 \times K_2})$ for each $l \in \mathbb{N}$, where $K_1 = \text{supp}(\phi)$ and $K_2 = \text{supp}(\chi)$.

Proof. For the first three properties we refer to [26], Theorem 1.1.18. As explained there these properties are proved globally, for example in [19], when M is a closed manifold. A careful examination of those proofs shows that the same properties remain true locally when M is an open manifold. The same argumentation applies to the last property which is proved globally, on a closed manifold, in [5] Prop. 2.46 or in [32] Theorem 7.15. \square

The rest of the subsection is a brief reminder about the heat kernel of a differential cone operator. For more details and for the proof we refer to [26]. As already recalled in Theorem 1 we know that, if \overline{M} is a compact and oriented manifold with boundary, M its interior part, $P_0 \in \text{Diff}_0(M, E; E)$ is a positive operator and g is a conic metric over M , then for each positive self-adjoint extension P of P_0 , $e^{-tP} : L^2(M, g) \rightarrow L^2(M, g)$ is a trace-class operator. Now we want to recall an important property named **scaling property**. Before doing this we need to introduce some notations:

Let N be a compact manifold; consider $C(N)$ and endow it with a product metric $g = dr^2 + h$ where h is a riemannian metric over N . Finally let E be a vector bundle over $\text{reg}(C(N))$.

Define $U_t : L^2(\text{reg}(C(N)), E) \rightarrow L^2(\text{reg}(C(N)), E)$ as $s(r, p) \mapsto t^{\frac{1}{2}} s(tr, p)$. It is immediate to show that $U_t : L^2(\text{reg}(C(N)), E) \rightarrow L^2(\text{reg}(C(N)), E)$ is an isometry and that $U_{t_1} \circ U_{t_2} = U_{t_1 t_2}$.

Proposition 8. Let N be a compact manifold, E a vector bundle over $\text{reg}(C(N))$, let $P_0 \in \text{Diff}_0^{\mu, \nu}(\text{reg}(C(N)), E, E)$ be a symmetric differential cone operator and let P be a self-adjoint extension of P_0 . Endow $\text{reg}(C(N))$ with a product metric g , that is $g = dr^2 + h$ where h is a

riemannian metric over N . Finally let $P_t = t^\nu U_t P U_t^*$ and let $f : \mathbb{R} \rightarrow \mathbb{R}$ a function such that $f(P)$ has a measurable kernel. Then for each $\lambda > 0$

$$f(P)(r, p, s, q) = \frac{1}{\lambda} f(\lambda^{-\nu} P_\lambda)\left(\frac{r}{\lambda}, p, \frac{s}{\lambda}, q\right), \quad \lambda > 0 \quad (35)$$

As particular case, given $P_0 \in \text{Diff}_0^{\mu, \nu}(\text{reg}(C(N)), E, E)$ positive and P a positive self-adjoint extension then

$$e^{-tP}(r, p, r, q) = \frac{1}{r} e^{-tr^{-\nu} P_r}(1, p, 1, q) \quad (36)$$

Proof. See [26] Lemma 2.2.3. \square

Now we modify the above proposition for the heat operator in the case that g is a conic metric over M . As we will see, we are interested to the study of the L^2 -Lefschetz numbers where the L^2 space are built using a conic metric. The reason is that when the considered complex is the L^2 de Rham complex (built using a conic metric) then its L^2 -cohomology has a topological meaning. More precisely, as showed by Cheeger in [14], we have the following theorem:

Theorem 3. *Let (F, h) be a compact and oriented riemannian manifold of dimension f . Consider the cone $C_b(F)$ with b a positive real number and endow $C_b(F)$ with the conic metric $g = dr^2 + r^2h$. Then*

$$H_{2, \max}^i(C_b(F), g) \cong \begin{cases} H^i(F) & i < \frac{f}{2} + \frac{1}{2} \\ 0 & i \geq \frac{f}{2} + \frac{1}{2} \end{cases} \quad (37)$$

If X is a compact and oriented manifold with conical singularities and if g is a conic metric over $\text{reg}(X)$ then

$$H_{2, \max}^i(\text{reg}(X), g) \cong I^m H^i(X), \quad H_{2, \min}^i(\text{reg}(X), g) \cong I^{\overline{m}} H^i(X). \quad (38)$$

Proof. See [14]. \square

For the definition and the main properties of intersection cohomology we refer to [20] and [21]

Lemma 1. *Let N be a compact manifold of dimension n , E a vector bundle over $\text{reg}(C(N))$, let $P_0 \in \text{Diff}_0^{\mu, \nu}(\text{reg}(C(N)), E, E)$ be a positive differential cone operator and let P be a positive self-adjoint extension of P_0 . Endow $\text{reg}(C(N))$ with a conic metric g , that is $g = dr^2 + r^2h$ where h is a riemannian metric over N . Then for each $\lambda > 0$*

$$e^{-tP}(r, p, s, q) = \frac{1}{\lambda^{n+1}} e^{-t\lambda^{-\nu} P_\lambda}\left(\frac{r}{\lambda}, p, \frac{s}{\lambda}, q\right), \quad \lambda > 0 \quad (39)$$

In particular we have

$$e^{-tP}(r, p, r, q) = \frac{1}{r^{n+1}} e^{-tr^{-\nu} P_r}(1, p, 1, q). \quad (40)$$

Proof. The proof is completely analogous to the proof of Proposition 8. We have just to add the natural modifications caused by the fact that now the Hilbert space $L^2(\text{reg}(C(N)), E)$ is built using the conic metric $g = dr^2 + r^2h$ and this means that given $\gamma \in L^2(\text{reg}(C(N)), E)$ we have $\|\gamma\|_{L^2(\text{reg}(C(N)), E)} = \int_{\text{reg}(C(N))} \langle \gamma, \gamma \rangle r^n dr d\text{vol}_h$ where $\langle \gamma, \gamma \rangle$ is the pointwise inner product induced by the metric on E (which is a riemannian metric if E is a real vector bundle and is a Hermitian metric if E is complex.). This implies that now the isometry U_t , introduced above Proposition 8, is defined as $U_t : L^2(\text{reg}(C(N)), E) \rightarrow L^2(\text{reg}(C(N)), E)$, $U_t(\gamma) = t^{\frac{n+1}{2}} \gamma(tr, p)$. The proof follows now in completely analogy to that one of Proposition 8. Moreover, in the case that P is a positive self-adjoint extension of $\Delta_i : \Omega_c^i(\text{reg}(C(N))) \rightarrow \Omega_c^i(\text{reg}(C(N)))$, the Laplacian constructed using a conic metric and acting on the space of smooth i -forms with compact support, the proof is given in [15], pag. 582. \square

Finally we conclude the section with the following proposition; before to state it we introduce some notations. Given $\lambda \in \mathbb{R}$ we define

$$p^+(\lambda) := \left| \lambda + \frac{1}{2} \right| \text{ and}$$

$$p^-(\lambda) := \begin{cases} \left| \lambda - \frac{1}{2} \right| & |\lambda| \geq \frac{1}{2} \\ \lambda - \frac{1}{2} & |\lambda| < \frac{1}{2} \end{cases} \quad (41)$$

Moreover we recall that $I_a(x)$ is the modified Bessel function of order a . For the definition see [26] pag. 67.

Proposition 9. *Let (N, h) be a compact and oriented riemannian manifold of dimension n . Consider $C(N)$ and let E be a vector bundle over $\text{reg}(C(N))$ endowed with a metric ρ (hermitian if it is complex or riemannian if it is real). Suppose that E admits an extension over all $[0, \infty) \times N$ that we denote \bar{E} . Let $E_N = \bar{E}|_N$ and suppose that (E, ρ) is isometric to $\pi^*(E_N, \rho|_N)$ where $\pi : (0, \infty) \times N \rightarrow N$ is the natural projection. Finally let $P : C_c^\infty(E) \rightarrow C_c^\infty(E)$ be an elliptic differential cone operator of order one. Then:*

1. *On $L^2(\text{reg}(C_2(N)), E)$ built with the product metric $g_p = dr^2 + h$, if P satisfies $P = \frac{\partial}{\partial r} + \frac{1}{r}S$, where $S \in \text{Diff}^1(N, E_N)$ is elliptic, we have*

$$e^{-tP_{max}^t \circ P_{min}}(r, p, s, q) = \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (rs)^{\frac{1}{2}} I_{p^+(\lambda)}\left(\frac{rs}{2t}\right) e^{-\frac{r^2+s^2}{4t}} \Phi_\lambda(p, q) \quad (42)$$

and

$$e^{-tP_{min} \circ P_{max}^t}(r, p, s, q) = \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (rs)^{\frac{1}{2}} I_{p^-(\lambda)}\left(\frac{rs}{2t}\right) e^{-\frac{r^2+s^2}{4t}} \Phi_\lambda(p, q)$$

where $\Phi_\lambda(p, q)$ is the smooth kernel of $\Phi_\lambda : L^2(N, E_N) \rightarrow V_\lambda$, the orthogonal projection on the eigenspace V_λ .

2. *On $L^2(\text{reg}(C_2(N)), E)$ built with the conic metric $g_c = dr^2 + r^2h$, if P satisfies $P = \frac{n}{2r} + \frac{\partial}{\partial r} + \frac{1}{r}S$, where $S \in \text{Diff}^1(N, E_N)$ is elliptic, we have*

$$e^{-tP_{max}^t \circ P_{min}}(r, p, s, q) = \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (rs)^{\frac{1-n}{2}} I_{p^+(\lambda)}\left(\frac{rs}{2t}\right) e^{-\frac{r^2+s^2}{4t}} \Phi_\lambda(p, q) \quad (43)$$

and

$$e^{-tP_{min} \circ P_{max}^t}(r, p, s, q) = \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (rs)^{\frac{1-n}{2}} I_{p^-(\lambda)}\left(\frac{rs}{2t}\right) e^{-\frac{r^2+s^2}{4t}} \Phi_\lambda(p, q)$$

where $\Phi_\lambda(p, q)$ is the smooth kernel of $\Phi_\lambda : L^2(N, E_N) \rightarrow V_\lambda$, the orthogonal projection on the eigenspace V_λ .

Proof. The first assertion is proved in [26], see Proposition 2.3.11 and pag. 68. The second statement follows using the following argument. Only for the remaining part of this proof let us label $L^2(\text{reg}(C_2(N)), E, g_p)$ the L^2 space of sections built using the product metric $g_p = dr^2 + h$ and $L^2(\text{reg}(C_2(N)), E, g_c)$ the L^2 space of sections built using the conic metric $g_c = dr^2 + r^2h$. The measure induced by g_p is $drd\text{vol}_h$ while the measure induce by g_c is $r^n drd\text{vol}_h$. Therefore it is clear that the map $\tau : L^2(\text{reg}(C(N)), E, g_c) \rightarrow L^2(\text{reg}(C_2(N)), E, g_p)$, $\tau(\gamma) = r^{\frac{n}{2}}\gamma$ is an isometry with inverse given by $\tau^{-1}(\gamma) = r^{-\frac{n}{2}}\gamma$. A simple calculation shows that $\tilde{P} := \tau^{-1} \circ P \circ \tau$ satisfies $\tilde{P} = \frac{\partial}{\partial r} + \frac{1}{r}S$. Therefore $\tilde{P}_{max}^t \circ \tilde{P}_{min} = r^{\frac{n}{2}} P_{max}^t \circ P_{min} r^{-\frac{n}{2}}$ and this implies that

$$e^{-t\tilde{P}_{max}^t \circ \tilde{P}_{min}} = r^{\frac{n}{2}} e^{-tP_{max}^t \circ P_{min}} r^{-\frac{n}{2}}.$$

Therefore if we call $\tilde{k}(t, r, p, s, q)$ the heat kernel relative to $e^{-t\tilde{P}_{max}^t \circ \tilde{P}_{min}}$ and analogously $k(t, r, p, s, q)$ the heat kernel relative to $e^{-tP_{max}^t \circ P_{min}}$ we have, for each $\gamma \in L^2(\text{reg}(C_2(N)), E, g_p)$,

$$\int_{\text{reg}(C_2(N))} \tilde{k}(t, r, p, s, q) \gamma(s) ds d\text{vol}_h = \int_{\text{reg}(C_2(N))} r^{\frac{n}{2}} k(t, r, p, s, q) s^{-\frac{n}{2}} \gamma(s) s^n ds d\text{vol}_h$$

and therefore $\tilde{k}(t, r, p, s, q) = r^{\frac{n}{2}} k(t, r, p, s, q) s^{\frac{n}{2}}$. Finally, applying this last equality to (42), we get (43). For the heat kernel of $e^{-tP_{min} \circ P_{max}^t}$ the proof is completely analogous to the previous one. \square

2 Geometric endomorphisms

The goal of this section is to introduce and study the notion of **geometric endomorphism** of an elliptic complex of differential cone operators.

Let X be a compact manifold with conical singularities and let M be its regular part that, as explained after Definition 7, we identify with the interior part of \overline{M} the manifold with boundary which desingularizes X , see Prop. 6. Finally consider an elliptic complex of differential cone operators as described in Definition 10:

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \xrightarrow{P_n} 0 \quad (44)$$

Definition 13. A *geometric endomorphism* T of (44) is given by a n -tuple of maps $T = (T_1, \dots, T_n)$, where each T_i maps $C^\infty(M, E_i)$ to itself, constructed in the following way: there exists a smooth map $f : \overline{M} \rightarrow \overline{M}$ and a n -tuples of morphisms of bundles $\phi_i : f^*E_i \rightarrow E_i$ such that the following properties hold:

1. $f : \overline{M} \rightarrow \overline{M}$ is a diffeomorphism.
2. If, with a little abuse of notation, we still label with $f : X \rightarrow X$ the isomorphism that $f : \overline{M} \rightarrow \overline{M}$ induces on X then we require that $f(q) = q$ for each $q \in \text{sing}(X)$.
3. $T_i = \phi_i \circ f^*$ where f^* acts naturally between $C^\infty(M, E_i)$ and $C^\infty(M, f^*E_i)$.
4. $P_i \circ T_i = T_{i+1} \circ P_i$.

We make a little comment on the above definition. The third and the fourth property are exactly the definition of geometric endomorphism of an elliptic complex over a closed manifold given in [2]. However our definition is not a complete extension of that one given by Atiyah and Bott in [2]. The reason is that in the closed case any smooth map is allowed. For our purposes we need that T_i induce a bounded map from $L^2(M, E_i)$ to itself and clearly this prevents us to allow every smooth map in Definition 13. As we will see in the following lemma, the property that $f : \overline{M} \rightarrow \overline{M}$ is a diffeomorphism is a reasonable sufficient condition in order to get a bounded extension of T_i on $L^2(M, E_i)$.

Lemma 2. *In the same hypothesis of the above definition the endomorphism T satisfies that the following properties:*

1. For each i and for each $\psi \in C_c^\infty(M, E_i)$ we have $T_i(\psi) \in C_c^\infty(M, E_i)$.
2. For each i T_i extends as a bounded operator from $L^2(M, E_i)$ to itself; with a small abuse of notation, we denote this again by T_i .
3. Let $T_i^* : L^2(M, E_i) \rightarrow L^2(M, E_i)$ be the adjoint of T_i . Then for each $\psi \in C_c^\infty(M, E_i)$ we have $T_i^*(\psi) \in C_c^\infty(M, E_i)$.

Proof. The first two properties follow immediately by the fact that $f : \overline{M} \rightarrow \overline{M}$ is a diffeomorphism and that \overline{M} is compact. For the third properties, we observe first of all that T_i admits an adjoint because it is densely defined and that T_i^* is bounded and defined over the whole $L^2(M, E_i)$ because T_i is bounded. Now consider the bundle f^*E_i . The metric over E_i induces in a natural way through f a metric over f^*E_i . Therefore it make sense consider the bundle homomorphism $\phi_i^* : E_i \rightarrow f^*E_i$ defined in each fiber as the adjoint of ϕ_i . Now consider the pull-back under f of the volume form $dvol_g$. Then there exists a smooth function τ such that $\tau dvol_g = f^*dvol_g$ and $\tau > 0$ if f preserves the orientation of M , $\tau < 0$ if f reverses the orientation of M . Finally define $S : C_c^\infty(M, E_i) \rightarrow C_c^\infty(M, E_i)$ as

$$S_i(\psi) := \begin{cases} \tau(\phi_i^* \circ (f^{-1})^*)(\psi) & \text{if } f \text{ preserves the orientation} \\ -\tau(\phi_i^* \circ (f^{-1})^*)(\psi) & \text{if } f \text{ reserves the orientation} \end{cases} \quad (45)$$

It is immediate to check that for each $\psi_1, \psi_2 \in C_c^\infty(M, E_i)$ we have

$$\langle T_i(\psi_1), \psi_2 \rangle_{L^2(M, E_i)} = \langle \psi_1, S_i(\psi_2) \rangle_{L^2(M, E_i)}.$$

Therefore, over $C_c^\infty(M, E_i)$, T_i^* coincides with S and so from this the third property follows immediately. \square

Now we state the following property :

Proposition 10. *Let M be an open and oriented riemannian manifold and let g be an incomplete riemannian metric on M . Let E_0, \dots, E_n be a sequence of vector bundles over M and consider a complex of differential operators:*

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \xrightarrow{P_n} 0 \quad (46)$$

Let $T = (T_0, \dots, T_n)$ be an endomorphism of (46) that satisfies the second and the third properties of Lemma 2. Then we have the following properties:

1. For each $i = 0, \dots, n$, for each $s \in \mathcal{D}(P_{min,i})$ we have $T_i(s) \in \mathcal{D}(P_{min,i})$ and $P_{min,i} \circ T_i = T_{i+1} \circ P_{min,i}$.
2. For each $i = 0, \dots, n$, for each $s \in \mathcal{D}(P_{max,i})$ we have $T_i(s) \in \mathcal{D}(P_{max,i})$ and $P_{max,i} \circ T_i = T_{i+1} \circ P_{max,i}$.

Proof. Let $i \in \{0, \dots, n\}$ and let $s \in \mathcal{D}(P_{min,i})$. Then there exists a sequence $\{s_j\}_{j \in \mathbb{N}}$ such that $s_j \rightarrow s$ in $L^2(M, E_i)$ and $P_i(s_j) \rightarrow P_i(s)$ in $L^2(M, E_{i+1})$. By the assumptions, we know that $\{T_i(s_j)\}_{j \in \mathbb{N}}$ is a sequence of smooth sections with compact support contained in $C_c^\infty(M, E_i)$ such that $T_i(s_j) \rightarrow T_i(s)$ in $L^2(M, E_i)$ and $T_{i+1}(P_i(s_j)) \rightarrow T_{i+1}(P_i(s))$ in $L^2(M, E_{i+1})$. But $T_{i+1}(P_i(s_j)) = P_i(T_i(s_j))$. Therefore $P_i(T_i(s_j))$ converges in $L^2(M, E_{i+1})$ and this implies that $T_i(s) \in \mathcal{D}(P_{min,i})$ and that $P_{min,i} \circ T_i = T_{i+1} \circ P_{min,i}$.

Now we give the proof of the second statement. From the first part of the proof it follows that, if we look at $T_{i+1} \circ P_{min,i}$, $P_{min,i} \circ T_i$ as unbounded operator with domain $\mathcal{D}(P_{min,i})$ then $T_{i+1} \circ P_{min,i} = P_{min,i} \circ T_i$ and therefore $(T_{i+1} \circ P_{min,i})^* = (P_{min,i} \circ T_i)^*$. Moreover, by the fact that T_{i+1} is bounded, it follows that $(T_{i+1} \circ P_{min,i})^* = P_{min,i}^* \circ T_{i+1}^*$ with domain given by $(T_{i+1}^*)^{-1}(\mathcal{D}(P_{min,i}^*))$. Now let $s \in \mathcal{D}(P_{max,i})$ and let $\phi \in C_c^\infty(M, E_{i+1})$. Then

$$\begin{aligned} \langle T_i(s), P_i^t(\phi) \rangle_{L^2(M, E_i)} &= \langle s, T_i^*(P_i^t(\phi)) \rangle_{L^2(M, E_i)} = \langle s, (P_{min,i} \circ T_i)^*(\phi) \rangle_{L^2(M, E_i)} = \\ &= \langle s, P_{min,i}^*(T_{i+1}^*(\phi)) \rangle_{L^2(M, E_i)} = (\text{because } T_{i+1}^*(\phi) \in C_c^\infty(M, E_{i+1})) \\ &= \langle s, P_{max,i}^*(T_{i+1}^*(\phi)) \rangle_{L^2(M, E_i)} = \langle P_{max,i}(s), (T_{i+1}^*(\phi)) \rangle_{L^2(M, E_i)} \\ &= \langle T_{i+1}(P_{max,i}(s)), \phi \rangle_{L^2(M, E_i)}. \end{aligned}$$

So we can conclude that $T_i(s) \in \mathcal{D}(P_{max,i})$ and that $T_{i+1} \circ P_{max,i} = P_{max,i} \circ T_i$. \square

In the rest of this section we describe the notion of **non degeneracy condition** for a fixed point of a map $f : X \rightarrow X$. As we will see, over the regular part of X , this is the same of the one used in [2].

Let X be a compact manifold with conical singularities and let $f : X \rightarrow X$ a continuous map such that $f(\text{sing}(X)) \subset \text{sing}(X)$, $f(\text{reg}(X)) \subset \text{reg}(X)$ and $f|_{\text{reg}(X)}$ is a smooth map. Define

$$\text{Fix}(f) := \{p \in X : f(p) = p\} \quad (47)$$

Definition 14. *A point $p \in \text{reg}(X) \cap \text{Fix}(f)$ is said to be simple if $\det(\text{Id} - d_p f) \neq 0$.*

Obviously this definition make sense because, being p a fixed point, it follows that $d_p f$ is an endomorphism of $T_p(\text{reg}(X))$. Moreover it is easy to show that Definition 14 is equivalent to require that, on $\text{reg}(X) \times \text{reg}(X)$, $\mathcal{G}(f)$ meets transversely $\Delta_{\text{reg}(X)}$ on (p, p) , where $\mathcal{G}(f)$ is the graph of $f|_{\text{reg}(X)}$ and $\Delta_{\text{reg}(X)}$ is the diagonal of $\text{reg}(X)$. In this way we get the following useful corollary:

Corollary 2. *Each simple fixed point in $\text{reg}(X) \cap \text{Fix}(f)$ is an isolated fixed point.*

Now, following [29], [30] but with little modifications, we recall what is a simple fixed point $p \in \text{Fix}(f) \cap \text{sing}(X)$. As we said above, we assumed that $f(\text{sing}(X)) \subset \text{sing}(X)$ and that $f(\text{reg}(X)) \subset \text{reg}(X)$. Therefore if $q \in \text{sing}(X) \cap \text{Fix}(f)$ is a fixed conical point it follows that, on a neighborhood $U_q \cong C_2(L_q)$ of q , f takes the form:

$$f(r, p) = (rA(r, p), B(r, p)) \quad (48)$$

We make the additional assumption that $A(r, p)$ and $B(r, p)$ are smooth up to zero, that is

$$A(r, p) : [0, 2) \times L_q \rightarrow [0, 2)$$

is smooth up to 0 and analogously

$$B(r, p) : [0, 2) \times L_q \rightarrow L_q$$

is smooth up to 0. Moreover, by the fact that $f(\text{sing}(X)) \subset \text{sing}(X)$ and that $f(\text{reg}(X)) \subset \text{reg}(X)$ it follows that $A(r, p) \neq 0$ for $r > 0$. Obviously if our starting point is a diffeomorphism $\bar{f} : \bar{M} \rightarrow \bar{M}$ as in Definition 13, then these requirements are automatically satisfied.

Definition 15. *A point $q \in \text{Fix}(f) \cap \text{sing}(X)$ is a **simple fixed point** if for each $p \in L_q$ at least one of the following conditions is satisfied:*

1. $A(0, p) \neq 1$.
2. $B(0, p) \neq p$.

A natural question follows from Definition 15: what is the meaning of these requirements? The answer is that if f satisfies one of the two requirements above then a sequence of fixed points converging to q cannot exist and therefore q is an isolated fixed point. We can show this last property in the following way: suppose that $\{(r_j, p_j)\}$ is a sequence of fixed point of f contained in $U_q \cong C_2(L_q)$ such that $r_j \rightarrow 0$ when $j \rightarrow \infty$. Then $\{p_j\}$ is a sequence of point in L_q which is compact and therefore there exists a subsequence, that with a little abuse of notations we still label $\{p_j\}$, such that p_j converges to some $p \in L_q$. By the assumptions, for each j , $(r_j, p_j) = (r_j A(r_j, p_j), B(r_j, p_j))$. Therefore $1 = \lim_{j \rightarrow \infty} A(r_j, p_j) = A(0, p)$, $B(r_j, p_j) = p_j$ for each j and this implies that f does not satisfy both the properties of Definition 15.

So we can state the following useful corollary:

Corollary 3. *Let X be a compact manifold with conical singularities and let $f : X \rightarrow X$ a map such that $f(\text{sing}(X)) \subset \text{sing}(X)$, $f(\text{reg}(X)) \subset \text{reg}(X)$, $f|_{\text{reg}(X)} : \text{reg}(X) \rightarrow \text{reg}(X)$ is smooth and, on a neighborhood of a conical point, $A(r, p)$ and $B(r, p)$ are smooth up to 0. Then, if f has only simple fixed point, $\text{Fix}(f)$ is made of a finite number of points.*

Proof. If f has only simple fixed points then we already know that each of this fixed points is an isolated fixed point and this implies that $\text{Fix}(f)$ is a sequence without accumulation points. Therefore, by the compactness of X , it follows that $\text{Fix}(f)$ is made of a finite number of points. \square

Now we state the following definition:

Definition 16. *Let f be as in the previous corollary. Let $q \in \text{Fix}(f) \cap \text{sing}(X)$ a simple fixed point for f such that f satisfies the first requirement of Definition 15. Then if for each $p \in L_q$*

$$A(0, p) < 1 \tag{49}$$

*q is called **attractive simple fixed point** while if*

$$A(0, p) > 1 \tag{50}$$

*then q is called **repulsive simple fixed point**.*

Clearly if for each $q \in \text{sing}(X)$ the relative link L_q is connected then each simple fixed point $q \in \text{sing}(X)$ satisfying the first property of Definition 15 is necessarily attractive or repulsive.

Finally we conclude the section observing that in [22], pag. 384, Goresky and MacPherson introduced the notion of contracting fixed point. An elementary check shows that (49) is equivalent to the definition given by Goresky and MacPherson.

3 L^2 –Lefschetz numbers of a geometric endomorphism

Let X be a compact manifold with conical singularities of dimension $m+1$. Consider an elliptic complex of cone differential operators as defined in Definition 10:

$$0 \rightarrow C_c^\infty(M, E_0) \xrightarrow{P_0} C_c^\infty(M, E_1) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(M, E_n) \xrightarrow{P_n} 0 \quad (51)$$

where $P_i \in \text{Diff}_0^{\mu, \nu}(M, E_i, E_{i+1})$ and let $T = \phi \circ f$ be a geometric endomorphism of (51) as in Definition 13. Obviously, with a small abuse of notation, we are using the same notation for the diffeomorphism $f : \overline{M} \rightarrow \overline{M}$ and for the isomorphism that it induces on X . We recall that the isomorphism $f : X \rightarrow X$ satisfies:

1. $f|_{\text{reg}(X)} : \text{reg}(X) \rightarrow \text{reg}(X)$ is a diffeomorphism
2. For each $p \in \text{sing}(X)$ we have $f(p) = p$
3. $A(r, p)$ and $B(r, p)$ (see (48)) are smooth up to 0.

Using Corollary 1 we know that both the complexes $(L^2(M, E_i), P_{\max/\min, i})$ are Fredholm complexes, that is the cohomology groups $H_{2, \max/\min}^i(M, E_i)$ are finite dimensional.

Moreover by Proposition 10 we know that T is a morphism of both complexes $(L^2(M, E_i), P_{\max/\min, i})$. Therefore, for each $i = 0, \dots, n$, it induces an endomorphism

$$T_i^* : H_{2, \max}^i(M, E_i) \rightarrow H_{2, \max}^i(M, E_i) \text{ and analogously } T_i^* : H_{2, \min}^i(M, E_i) \rightarrow H_{2, \min}^i(M, E_i).$$

So we are in position to give the following definition:

Definition 17. *The L^2 –Lefschetz numbers of T are defined in the following way:*

$$L_{2, \max}(T) = \sum_{i=0}^n (-1)^i \text{tr}(T_i^* : H_{2, \max}^i(M, E_i) \rightarrow H_{2, \max}^i(M, E_i)) \quad (52)$$

and analogously

$$L_{2, \min}(T) = \sum_{i=0}^n (-1)^i \text{tr}(T_i^* : H_{2, \min}^i(M, E_i) \rightarrow H_{2, \min}^i(M, E_i)) \quad (53)$$

The L^2 –Lefschetz numbers satisfy the following property:

Proposition 11. *$L_{2, \max/\min}(T)$ do not depend on the conic metric g we fix on M and on the metrics ρ_0, \dots, ρ_n that we fix on E_0, \dots, E_n*

Proof. By the fact that \overline{M} is compact and that, as explained above Definition 7, (E_i, ρ_i) are defined over all \overline{M} and ρ_i is non degenerate up to the boundary, it follows that all the metrics we consider on E_i are quasi-isometric. Moreover, using [4] Proposition 9, it follows that if g and g' are two conic metric over M then they are quasi-isometric, that is there exists a positive real number c such that $g' \leq g \leq cg'$. Therefore, for each $i = 0, \dots, n$, $L^2(M, E_i)$ doesn't depend on the metric that we fix on E_i and on the conic metric that we fix over M . This in turn implies that same conclusion holds for $H_{2, \max}^i(M, E_i)$ and for $H_{2, \min}^i(M, E_i)$, that is they do not depend on the metric that we fix on E_i and on the conic metric that we fix over M . In this way we can conclude that also the traces of $T_i^* : H_{2, \max}^i(M, E_*) \rightarrow H_{2, \max}^i(M, E_*)$ and $T_i^* : H_{2, \min}^i(M, E_*) \rightarrow H_{2, \min}^i(M, E_*)$ satisfy the same property and so the proposition is proved. \square

- From the above proposition it follows that in order to calculate $L_{2, \max/\min}(T)$ we can use any conic metric g on M and any metrics ρ_0, \dots, ρ_n over E_0, \dots, E_n . Therefore, in the remaining part of this section, we make the following assumptions: for each singular point q there exists U_q , an open neighborhood of q satisfying $U_q \cong C_2(L_q)$, such that on $\text{reg}(C_2(L_q))$ the conic metric g satisfies $g = dr^2 + r^2h$ where h is any riemannian metric over L_q that does not depend on r . Moreover we assume that each metric ρ_i on E_i does not depend on r in a collar neighborhood of $\partial\overline{M}$.

Consider, for each $i = 0, \dots, n$, the operator

$$\mathcal{P}_i := P_i^t \circ P_i + P_{i-1} \circ P_{i-1}^t : C_c^\infty(M, E_i) \rightarrow C_c^\infty(M, E_i).$$

It is clearly a positive operator. As stated in Proposition 7, we know that \mathcal{P}_i is an elliptic differential cone operator. Therefore, by Theorem 1, we know that for each positive self-adjoint extension of \mathcal{P}_i , the relative heat operator is a trace-class operator. In particular this is true for $\mathcal{P}_{abs,i}$ that we recall it is defined as $P_{min,i}^t \circ P_{max,i} + P_{max,i-1} \circ P_{min,i-1}^t$ and for $\mathcal{P}_{rel,i}$ that it is defined as $P_{max,i}^t \circ P_{min,i} + P_{min,i-1} \circ P_{max,i-1}^t$. A well known and basic result of operators theory (see [32], Prop. 8.8) says that, given an Hilbert space H , the space of trace-class operators is a two sided ideal of $\mathcal{B}(H)$, the space of bounded operators of H , and that the trace doesn't depend on the order of composition. In this way we know that for each $i = 0, \dots, n$

$$T_i \circ e^{-t\mathcal{P}_{abs/rel,i}} : L^2(M, E_i) \rightarrow L^2(M, E_i)$$

are trace-class operator and that $\text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) = \text{Tr}(e^{-t\mathcal{P}_{abs/rel,i}} \circ T_i)$ ¹. Moreover it is clear that $T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}$ are operators with smooth kernel given by

$$\phi_i \circ k_{abs,i}(t, f(x), y) \text{ for } T_i \circ e^{-t\mathcal{P}_{abs,i}} \quad (54)$$

and analogously

$$\phi_i \circ k_{rel,i}(t, f(x), y) \text{ for } T_i \circ e^{-t\mathcal{P}_{rel,i}} \quad (55)$$

where $k_{abs/rel,i}(t, x, y)$ are respectively the smooth kernel of $e^{-t\mathcal{P}_{abs/rel,i}}$. In both the expressions above ϕ_i acts on the x variable of $k_{abs/rel,i}(t, f(x), y)$ because $k_{abs/rel,i}(t, f(x), y)$ is a section of $f^*E_i \boxtimes E_i^*$ and $\phi_i : f^*E_i \rightarrow E_i$ is a morphism of bundle. So the kernels $\phi_i \circ k_{abs/rel,i}(t, f(x), y)$ are well defined and they are smooth sections of $E \boxtimes E^*$.

Now we are in position to state the following theorem which is one of the main results of this section:

Theorem 4. *Consider an elliptic complex of differential cone operators as in (51) and let T be a geometric endomorphism as in Definition 13. Then for each t :*

$$L_{2,max}(T) = \sum_{i=0}^n (-1)^i \text{Tr}(T_i e^{-t\mathcal{P}_{abs,i}}) \quad (56)$$

and analogously

$$L_{2,min}(T) = \sum_{i=0}^n (-1)^i \text{Tr}(T_i e^{-t\mathcal{P}_{rel,i}}) \quad (57)$$

In particular, in both the equalities, the member on the right hand side does not depend on t .

We need to state some propositions in order to prove the above theorem. We give the proof only for the complex $(L^2(M, E_i), P_{max,i})$. The other one is completely analogous.

Lemma 3. *Consider an abstract Fredholm complex as in (9) and let T be an endomorphism of this complex, that is $T = (T_0, \dots, T_n)$, for each $i = 0, \dots, n$ $T_i : H_i \rightarrow H_i$ is bounded and $D_i \circ T_i = T_{i+1} \circ D_i$ on $\mathcal{D}(D_i)$. Let $\pi_i : H_i \rightarrow \mathcal{H}_i(H_*, D_*)$ be the orthogonal projection induced by the Kodaira decomposition of Proposition 1. Then for each $i = 0, \dots, n$ we have*

$$\text{Tr}(\pi_i \circ T_i : \mathcal{H}^i(H_*, D_*) \rightarrow \mathcal{H}^i(H_*, D_*)) = \text{Tr}(T_i^* : H^i(H_*, D_*) \rightarrow H^i(H_*, D_*))$$

Proof. Let $\gamma : \mathcal{H}^i(H_*, D_*) \rightarrow H^i(H_*, D_*)$ the isomorphism of (14). Then it is clear that T_i^* , that is the endomorphism of $H^i(H_*, D_*)$ induced by T_i , satisfies $T_i^* = \gamma \circ \pi_i \circ T_i \circ \gamma^{-1}$. Now from this it follows immediately that $\text{Tr}(\pi_i \circ T_i : \mathcal{H}^i(H_*, D_*) \rightarrow \mathcal{H}^i(H_*, D_*)) = \text{Tr}(T_i^* : H^i(H_*, D_*) \rightarrow H^i(H_*, D_*))$. \square

Lemma 4. *We have the following properties.*

1. *Let $E_i(\lambda)$ be the eigenspace relative to $\mathcal{P}_{abs,i}$ and the eigenvalue λ . Then $E_i(\lambda)$ is finite dimensional and made of eigensections which are smooth in the interior.*

¹This is the reason because we need to require that $f : \overline{M} \rightarrow \overline{M}$ is a diffeomorphism. In this way each $T_i : L^2(M, E_i) \rightarrow L^2(M, E_i)$ is bounded and so we can conclude that $T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}$ is a trace-class operator

2. For each $\lambda \neq 0$ consider the following complex:

$$\dots \xrightarrow{P_{max,i-1}^\lambda} E_i(\lambda) \xrightarrow{P_{max,i}^\lambda} E_{i+1}(\lambda) \xrightarrow{P_{max,i+1}^\lambda} E_{i+2}(\lambda) \xrightarrow{P_{max,i+2}^\lambda} \dots \quad (58)$$

where $P_{max,i}^\lambda := P_{max,i}|_{E_i(\lambda)}$. Then it is an acyclic complex.

Proof. Consider the eigenspaces $E_i(\lambda)$. That is finite dimensional for each $\lambda \neq 0$ follows by the fact that $e^{-t\mathcal{P}_{abs,i}}$ is a trace-class operator while that it is finite dimensional for $\lambda = 0$ follows by the fact that $\mathcal{P}_{abs,i}$ is a Fredholm operator on its domain endowed with the graph norm. Moreover elliptic regularity tells us that $E_i(\lambda)$ is made of eigensections which are smooth in the interior. Finally, given $\lambda > 0$, consider

$$\dots \xrightarrow{P_{max,i-1}^\lambda} E_i(\lambda) \xrightarrow{P_{max,i}^\lambda} E_{i+1}(\lambda) \xrightarrow{P_{max,i+1}^\lambda} E_{i+2}(\lambda) \xrightarrow{P_{max,i+2}^\lambda} \dots \quad (59)$$

where $P_{max,i}^\lambda := P_{max,i}|_{E_i(\lambda)}$.

Let $s \in \text{Ker}(P_{max,i})$. Then $\mathcal{P}_{abs,i}(s) = \lambda s = P_{max,i-1}(P_{min}^t(s))$. Therefore $s \in \text{ran}(P_{max,i-1})$ and this implies that (59) is a long exact sequences, or in other words, it is an acyclic complex. \square

Now we state the last result we need to prove Theorem 4. We take it from [2].

Lemma 5. Consider a complex of finite dimensional vector space

$$0 \rightarrow V_0 \xrightarrow{f_0} \dots \xrightarrow{f_{i-1}} V_i \xrightarrow{f_i} V_{i+1} \xrightarrow{f_{i+1}} V_{i+2} \xrightarrow{f_{i+2}} \dots \xrightarrow{f_{n-1}} V_n \xrightarrow{f_n} 0. \quad (60)$$

and for each i let $G_i : V_i \rightarrow V_i$ an endomorphism such that $f_i \circ G_i = G_{i+1} \circ f_i$. Then

$$\sum_{i=0}^n (-1)^i \text{Tr}(G_i) = \sum_{i=0}^n (-1)^i \text{Tr}(G_i^*)$$

where G_i^* is the endomorphism of the i -th cohomology group of the complex (60) induced by G_i .

Proof. See [2]. \square

Proof. (of Theorem 4). As said above we give the proof only for (56). The proof for (57) is completely analogous. Consider the heat operator $e^{-t\mathcal{P}_{abs,i}} : L^2(M, E_i) \rightarrow L^2(M, E_i)$. By the third point of Theorem 1 it follows that there exists an Hilbert base of $L^2(M, E_i)$, $\{\phi_j\}_{j \in \mathbb{N}}$, made of smooth eigensections of $\mathcal{P}_{abs,i}$, in such way the smooth kernel of $e^{-t\mathcal{P}_{abs,i}}$ satisfies $k(t, x, y) = \sum_j e^{-t\lambda_j} \phi_j(x) \boxtimes \phi_j^*(y)$. Moreover, by the fact that $T_i : L^2(M, E_i) \rightarrow L^2(M, E_i)$ is bounded, we know that $T_i \circ e^{-t\mathcal{P}_{abs,i}}$ and $e^{-t\mathcal{P}_{abs,i}} \circ T_i$ are trace class and that $\text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs,i}}) = \text{Tr}(e^{-t\mathcal{P}_{abs,i}} \circ T_i)$. Now, if we label $\pi(i, \lambda_j)$ the orthogonal projection $\pi(i, \lambda_j) : L^2(M, E_i) \rightarrow E_i(\lambda_j)$, then we can write $e^{-t\mathcal{P}_{abs,i}} = \sum_j e^{-t\lambda_j} \pi(i, \lambda_j)$ and therefore $e^{-t\mathcal{P}_{abs,i}} \circ T_i = (\sum_j e^{-t\lambda_j} \pi(i, \lambda_j)) \circ T_i = \sum_j e^{-t\lambda_j} (\pi(i, \lambda_j) \circ T_i)$. In this way we get

$$\text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs,i}}) = \text{Tr}(e^{-t\mathcal{P}_{abs,i}} \circ T_i) = \sum_j e^{-t\lambda_j} \text{Tr}((\pi(i, \lambda_j) \circ T_i)). \quad (61)$$

Consider $\sum_{i=0}^n (-1)^i \text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs,i}})$. Then $\sum_{i=0}^n (-1)^i \text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs,i}}) =$

$$= \sum_{i=0}^n (-1)^i \sum_j e^{-t\lambda_j} \text{Tr}((\pi(i, \lambda_j) \circ T_i)) = \sum_j e^{-t\lambda_j} \sum_{i=0}^n (-1)^i \text{Tr}((\pi(i, \lambda_j) \circ T_i)). \quad (62)$$

Now examine carefully this last expression. Both $\pi(i, \lambda_j) \circ T_i : L^2(M, E_i) \rightarrow E_i(\lambda_j)$ and $\pi(i, \lambda_j) : L^2(M, E_i) \rightarrow E_i(\lambda_j)$ are trace-class operators. This implies that $\text{Tr}(\pi(i, \lambda_j) \circ T_i) = \text{Tr}(\pi(i, \lambda_j) \circ \pi(i, \lambda_j) \circ T_i) = \text{Tr}(\pi(i, \lambda_j) \circ T_i \circ \pi(i, \lambda_j))$ and this last one is equal to the trace of $\pi(i, \lambda_j) \circ T_i : E_i(\lambda_j) \rightarrow E_i(\lambda_j)$. But if we take the following complex for $\lambda_j \neq 0$

$$\dots \xrightarrow{P_{max,i-1}^\lambda} E_i(\lambda_j) \xrightarrow{P_{max,i}^\lambda} E_{i+1}(\lambda_j) \xrightarrow{P_{max,i+1}^\lambda} E_{i+2}(\lambda_j) \xrightarrow{P_{max,i+2}^\lambda} \dots \quad (63)$$

we know that (63) is an acyclic complex. Moreover it is immediate to check that $\pi(i, \lambda_j) \circ T_i$ is an endomorphism of (63) and therefore, applying Lemma 60, we can conclude that $\sum_{i=0}^n (-1)^i \text{Tr}(\pi(i, \lambda_j) \circ T_i) = 0$ for $\lambda_j \neq 0$. This leads to a relevant simplification of (62):

$$\sum_{i=0}^n (-1)^i \text{Tr}(T_i e^{-t\mathcal{P}_{abs,i}}) = \sum_j e^{-t\lambda_j} \sum_{i=0}^n (-1)^i \text{Tr}(\pi(i, \lambda_j) \circ T_i) = \sum_{i=0}^n (-1)^i \text{Tr}(\pi(i, 0) \circ T_i). \quad (64)$$

Finally, using Lemma 3, it follows that $\text{Tr}(\pi(i, 0) \circ T_i) = \text{Tr}(T_i^*)$ and therefore the theorem is proved. \square

As an immediate consequence of Theorem 4 we have the following corollary

Corollary 4. *In the same assumptions of Theorem 4 then*

$$L_{2,max}(T) = \lim_{t \rightarrow 0} \sum_{i=0}^n (-1)^i \text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs,i}}) \quad (65)$$

and analogously

$$L_{2,min}(T) = \lim_{t \rightarrow 0} \sum_{i=0}^n (-1)^i \text{Tr}(T_i \circ e^{-t\mathcal{P}_{rel,i}}) \quad (66)$$

Before to go ahead we add some comments to Theorem 4.

Remark 3. *In the statement of Theorem 4 we assume that the endomorphism T satisfies Definition 13. But from the proof it is clear that the particular structure of the endomorphism, that is $T_i = \phi_i \circ f^*$ doesn't play any role. It is just a sufficient condition to assure that each T_i induces a bounded map acting on $L^2(M, E_i)$ and that T is an endomorphism of $(L^2(M, E_i), P_{max/min,i})$. Therefore if we have a n -tuple of map $T = (T_1, \dots, T_n)$ such that, for each $i = 0, \dots, n$, $T_i : L^2(M, E_i) \rightarrow L^2(M, E_i)$ is bounded and $T_{i+1} \circ P_{max/min,i} = P_{max/min,i} \circ T_i$ on $\mathcal{D}(P_{max/min,i})$ then we can state and prove Theorem 4 in the same way.*

Remark 4. *We stated Theorem 4 in the case of an elliptic complex of differential cone operators over a compact manifold with conical singularities. This is because, using the result coming from the theory of elliptic differential cone operators, we know that $(L^2(M, E_i), P_{max/min,i})$ are Fredholm complexes and that $e^{-t\mathcal{P}_{abs/rel,i}}$ are trace-class operators. Therefore it is possible to define maximal and minimal L^2 -Lefschetz numbers and to prove Theorem 4. A priori it is not possible to do the same for an arbitrary elliptic complex of differential operators over a (possibly incomplete) riemannian manifold (M, g) . But it is clear that if we know that the maximal and the minimal extension of our complex are Fredholm complexes and that for each i the heat operator constructed from the i -th laplacian associated to the maximal/minimal complex is a trace-class operator, then it is possible to state and prove in the same way formulas (56) and (57) for the L^2 -Lefschetz numbers associated to the maximal and minimal extension of our complex.*

We conclude the section with the following theorems:

Theorem 5. *Let X be a compact manifold with conical singularities of dimension $m + 1$ and let g be a conic metric on $\text{reg}(X) = M$. Consider an elliptic complex of differential cone operators as in (51) and let $T = \phi \circ f^*$ be a geometric endomorphism of (51) as in Definition 13. Finally suppose that f has only simple fixed points. Then we have:*

$$L_{2,max/min}(T) = \lim_{t \rightarrow 0} \left(\sum_{q \in \text{Fix}(f)} \sum_{i=0}^n (-1)^i \int_{U_q} \text{tr}(T \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g \right) \quad (67)$$

where U_q is an open neighborhood of $q \in \text{Fix}(f)$ (clearly, when $q \in \text{sing}(X) \cap \text{Fix}(f)$ then we mean $U_q - \{q\}$).

Proof. We know, by the assumptions, that f has only simple fixed points. For each of these points, that we label q , let U_q be an open neighborhood of q . Then, using again Corollary 4, we

know that $L_{2,max/min}(T) = \lim_{t \rightarrow 0} \sum_i (-1)^i \int_M \text{tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}})$. Obviously we can break the member on the right as

$$\sum_{q \in \text{Fix}(f)} \sum_{i=0}^n (-1)^i \int_{U_q} \text{tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g + \sum_{i=0}^n (-1)^i \int_V \text{tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g$$

where $V = M - \cup_{q \in \text{Fix}(f)} U_q$. Now, as remarked previously, we know that $f(q) = q$ for each $q \in \text{sing}(X)$. This implies $\{(f(q), q) : q \in V\}$ is a compact subset of $M \times M$ disjoint from Δ_M . So we can use the second property of Theorem 2 to conclude that

$$\lim_{t \rightarrow 0} \int_V \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel,i}}(f(q), q)) d\text{vol}_g = \int_V \lim_{t \rightarrow 0} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel,i}}(f(q), q)) d\text{vol}_g = 0.$$

This complete the proof. \square

The second point in the above theorem suggests to break the Lefschetz numbers as a contribution of two terms, that is

$$L_{2,max/min}(T) = \mathcal{L}_{max/min}(T, \mathcal{R}) + \mathcal{L}_{max/min}(T, \mathcal{S}) \quad (68)$$

where $\mathcal{L}_{max/min}(T, \mathcal{R})$ is the contribution given by the simple fixed point lying in $\text{reg}(X)$, that is

$$\mathcal{L}_{max/min}(T, \mathcal{R}) = \lim_{t \rightarrow 0} \left(\sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \sum_{i=0}^n (-1)^i \int_{U_q} \text{tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g \right)$$

and analogously $\mathcal{L}_{max/min}(T, \mathcal{S})$ is the contribution given by the simple fixed point lying in $\text{Fix}(f) \cap \text{sing}(X)$, that is

$$\mathcal{L}_{max/min}(T, \mathcal{S}) = \lim_{t \rightarrow 0} \left(\sum_{q \in \text{Fix}(f) \cap \text{sing}(X)} \sum_{i=0}^n (-1)^i \int_{U_q - \{q\}} \text{tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g \right).$$

Theorem 6. *In the hypothesis of the previous theorem, suppose furthermore that for each $i = 0, \dots, n$*

$$P_i^t \circ P_i + P_{i-1} \circ P_{i-1}^t : C_c^\infty(M, E_i) \rightarrow C_c^\infty(M, E_i)$$

is a generalized Laplacian (see Definition 12). Then we get :

$$L_{2,max}(T) = \sum_{q \in \text{Fix}(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\phi_i)}{|\det(\text{Id} - d_q f)|} + \mathcal{L}_{2,max}(T, \mathcal{S}).$$

Analogously for $L_{2,min}(T)$ we have

$$L_{2,min}(T) = \sum_{q \in \text{Fix}(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\phi_i)}{|\det(\text{Id} - d_q f)|} + \mathcal{L}_{2,min}(T, \mathcal{S}).$$

Proof. By Theorem 5, we know that the L^2 -Lefschetz numbers depend only on the simple fixed point of f and that we can localize their contribution, that is,

$$L_{2,max/min}(T) = \lim_{t \rightarrow 0} \left(\sum_{q \in \text{Fix}(f)} \sum_{i=0}^n (-1)^i \int_{U_q} \text{tr}(T \circ e^{-t\mathcal{P}_{abs/rel,i}}) d\text{vol}_g \right)$$

where U_q is an arbitrary open neighborhood of q (and clearly when $q \in \text{sing}(X)$ then we mean the regular part of U_q). If $q \in \text{reg}(X) \cap \text{Fix}(f)$, by the assumptions, we can use the local asymptotic expansion recalled in the last point of Theorem 2. Now, to get the conclusion, the proof is exactly the same as in the closed case; see for example [5] Theorem 6.6 or [32] Theorem 10.12. \square

We have the following immediate corollary:

Corollary 5. *In the same hypothesis of Theorem 6; Then:*

1. $\mathcal{L}_{max}(T, \mathcal{R}) = \mathcal{L}_{min}(T, \mathcal{R})$ that is, the simple fixed points in M give the same contributions for both the Lefschetz numbers $L_{2,max/min}(T)$.
2. $\mathcal{L}_{max/min}(T, \mathcal{S})$ do not depend on the particular conic metric fixed on M and on the metrics ρ_0, \dots, ρ_n respectively fixed on E_0, \dots, E_n .

Proof. The first assertion is an immediate consequence of the second point of Theorem 6. For the second statement, by Proposition 11, we know that $L_{2,max/min}(T)$ are independent on the conic metric we put over M and on the metric ρ_0, \dots, ρ_n respectively on E_0, \dots, E_n . Again, by the second point of Theorem 6, we know that also $\mathcal{L}_{max/min}(T, \mathcal{R})$ are independent from the conic metrics and on the metric ρ_0, \dots, ρ_n respectively on E_0, \dots, E_n . Therefore the same conclusion holds for $\mathcal{L}_{max/min}(T, \mathcal{S})$. The corollary is proved. \square

4 The contribution of the singular points

The aim of this section is to give, in some particular cases, an explicit formula for $\mathcal{L}_{max/min}(T, \mathcal{S})$, that is for the contribution given by the singular points to the Lefschetz numbers $L_{2,max/min}(T)$. Consider the same situation described in Theorem 5. Suppose moreover that the following properties hold:

1. For each $q \in \text{sing}(X)$ there exists an isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$ such that on $[0, 2) \times L_q$, using (29), each operator A_k is **constant** in x and, using the decomposition (48), the map f takes the form:

$$f = (rA(p), B(p)). \quad (69)$$

2. On $\text{reg}(C_2(L_q))$, using again the isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$, the conic metric g satisfies $g = dr^2 + r^2h$ with h that does not depend on r and each metric ρ_i on E_i does not depend on r in a collar neighborhood of ∂M .

Before stating the next theorem we recall a definition from [26].

Definition 18. Consider the isometry $U_t : L^2(\text{reg}(C(N)), E) \rightarrow L^2(\text{reg}(C(N)), E)$ as defined in the proof of Lemma (1), that is $U_t(\gamma) = t^{\frac{n+1}{2}} \gamma(tr, p)$. Consider an operator $P_0 \in \text{Diff}_0^{\mu, \nu}(\text{reg}(C(N)))$ such that, using the expression (29), each A_k is constant in x . Then a closed extension P of P_0 is said *scalable* if $U_t^* P U_t = t^\nu P$.

Lemma 6. Given $P_0 \in \text{Diff}_0^{\mu, \nu}(\text{reg}(C(N)))$ as in Definition 18 then $P_{0,max}$ and $P_{0,min}$ are always scalable. If we take P_0^t , the formal adjoint of P_0 , then also $P_{0,min}^t \circ P_{0,max}$, $P_{0,max}^t \circ P_{0,min}$, $P_{0,min} \circ P_{0,max}^t$ and $P_{0,max} \circ P_{0,min}^t$ are scalable extensions of $P_0^t \circ P_0$ and $P_0 \circ P_0^t$ respectively. Finally, if in a complex we consider $\mathcal{P}_i := P_i^t \circ P_i + P_{i-1} \circ P_{i-1}^t$ (see the statement of Theorem 6) then also the closed extension $\mathcal{P}_{abs,i}$ and $\mathcal{P}_{rel,i}$ (see (22) and (23)) are scalable extensions.

Proof. For the first assertion see [26] pag. 58. The others assertions are an immediate consequence of the previous one and of the definition of scalable extension. \square

Now we are ready to state the following theorem:

Theorem 7. In the same hypothesis of Theorem 5. Suppose moreover that the two properties described above Definition 18 hold. Then we have:

$$\mathcal{L}_{max/min}(T, \mathcal{S}) = \sum_{q \in \text{sing}(X)} \sum_{i=0}^n (-1)^i \frac{1}{2\nu} \int_0^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{abs/rel,i}}(A(p), B(p), 1, p)) d\text{vol}_h. \quad (70)$$

Proof. Let $q \in \text{sing}(X)$. By the hypothesis we know that there exists an open neighborhood U_q and an isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$ such that, on $C_2(L_q)$, f takes the form (69) and each A_k is constant in x . Using the properties stated in [26] pag. 42-43, we get that the limit

$$\lim_{t \rightarrow 0} \int_{\text{reg}(U_q)} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel,i}}(rA(p), B(p), r, p)) d\text{vol}_g$$

is equal to

$$\lim_{t \rightarrow 0} \int_{\text{reg}(C_2(L_q))} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{\text{abs}/\text{rel},i}}(rA(p), B(p), r, p)) r^m \text{dvol}_h \text{d}r$$

where, with a little abuse of notation, in the second expression we mean the heat kernel associated to the absolute and relative extension of the operator, induced by $\mathcal{P}_i|_{U_q}$ through χ_q , acting on $C_c^\infty(\text{reg}(C_2(L_q)), (\chi_q^{-1})^* E_i)$. So, for each $i = 0, \dots, n$, we have to calculate

$$\lim_{t \rightarrow 0} \int_{\text{reg}(C_2(L_q))} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{\text{abs}/\text{rel},i}}(rA(p), B(p), r, p)) r^m \text{d}r \text{dvol}_h.$$

Moreover, we assumed that, on $\text{reg}(C_2(L_q))$, the conic metric g satisfies $g = dr^2 + r^2h$ with h that does not depend on r and that each metric ρ_i on E_i does not depend on r in a neighborhood of $\partial\bar{M}$. This implies that, for each $i = 0, \dots, n$, the operator \mathcal{P}_i satisfies the assumption at the beginning of the subsection, that is each A_k does not depend on x . Therefore, using Lemma 6, we get that $\mathcal{P}_{\text{abs}/\text{rel},i}$ are scalable extensions of \mathcal{P}_i . Now, after these observations, we can go on to calculate

$$\lim_{t \rightarrow 0} \int_{\text{reg}(C_2(L_q))} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{\text{abs}/\text{rel},i}}(rA(p), B(p), r, p)) \text{dvol}_g.$$

Using Lemma 1 and the fact that $\mathcal{P}_{\text{abs}/\text{rel},i}$ are scalable extensions of \mathcal{P}_i we get

$$\begin{aligned} & \int_{\text{reg}(C_2(L_q))} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{\text{abs}/\text{rel},i}}(rA(p), B(p), r, p)) r^m \text{d}r \text{dvol}_h = \\ & = \int_0^2 \int_{L_q} \frac{1}{r} \text{tr}(\phi_i \circ e^{-tr^{-2\nu}\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h \text{d}r. \end{aligned}$$

Now if we put $\frac{t}{r^{2\nu}} = x$ we get $\frac{-2\nu t \text{d}r}{r^{2\nu+1}} = dx$ which implies that

$$\frac{dx}{x} = \frac{-2\nu t \text{d}r}{r^{2\nu+1}} \frac{r^{2\nu}}{t} = -2\nu \frac{\text{d}r}{r}$$

Moreover when r goes to 0 then x goes to ∞ and when r goes to 2 then x goes to $\frac{t}{4}$. So we get

$$\begin{aligned} & \int_0^2 \int_{L_q} \frac{1}{r} \text{tr}(\phi_i \circ e^{-tr^{-2\nu}\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h \text{d}r = \\ & = \frac{1}{2\nu} \int_{t/4}^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h. \end{aligned} \quad (71)$$

Therefore to conclude we have to evaluate the limit

$$\lim_{t \rightarrow 0} \frac{1}{2\nu} \int_{t/4}^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h \quad (72)$$

To do this consider the term $\int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h$. We know, by the hypothesis, that f has only simple fixed points. In particular each $q \in \text{sing}(X)$ is a simple fixed point. The conditions described in Definition 15 together with (69) implies that either $A(p) \neq 1$ for all $p \in L_q$ or $B : L_q \rightarrow L_q$ has not fixed points. Anyway each of these conditions implies that when p runs over L_q then $\{(A(p), B(p), 1, p)\}$ is a compact subset of $\text{reg}(C_2(L_q)) \times \text{reg}(C_2(L_q))$ that doesn't intersect the diagonal. Therefore we can use the second property stated in Theorem 2 to conclude that, when $x \rightarrow 0$,

$$\int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h = O(x^N) \text{ for each } N > 0. \quad (73)$$

In this way we can conclude that the limit (72) exists and we have

$$\begin{aligned} & \lim_{t \rightarrow 0} \frac{1}{2\nu} \int_{t/4}^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h = \\ & = \frac{1}{2\nu} \int_0^\infty \frac{dx}{x} \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{\text{abs}/\text{rel},i}}(A(p), B(p), 1, p)) \text{dvol}_h. \end{aligned} \quad (74)$$

□

Now, for each $i = 0, \dots, n$, using again the hypothesis and the notations of Theorem 7, and assuming still that q is a simple fixed point for f , define the following "modified version" of the classical ζ -function:

$$\zeta_{T_i, q}(\mathcal{P}_{abs/rel, i})(s) := \frac{1}{2\nu} \int_0^\infty x^{s-1} dx \int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{abs/rel, i}}(A(p), B(p), 1, p)) dvol_h. \quad (75)$$

The definition makes sense for each $s \in \mathbb{C}$ because, as observed in the proof of Theorem 7, $\{(A(p), B(p), 1, p)\}$ is a compact subset of $reg(X) \times reg(X)$ that is disjoint from the diagonal $\Delta_{reg(X)}$. Therefore we can apply the second point of Theorem 2 to conclude that, when $x \rightarrow 0$,

$$\int_{L_q} \text{tr}(\phi_i \circ e^{-x\mathcal{P}_{abs/rel, i}}(A(p), B(p), 1, p)) dvol_h = O(x^N) \text{ for each } N > 0. \quad (76)$$

and this implies that $\zeta_{T_i, q}(\mathcal{P}_{abs/rel, i})(s)$ is a holomorphic function over the whole complex plane. The reason behind (74) is that if we compare (74) with the definitions of the zeta functions for a generalized Laplacian, see for example [5] pag. 295, then it natural to think at (74) as a sort of zeta function for the operators $\mathcal{P}_{abs/rel, i}$ valued in 0, which takes account of the action of T_i in its definition. In this way, using (75), we can reformulate Theorem 7 in a more concise way:

$$\mathcal{L}_{max/min}(T, \mathcal{S}) = \sum_{q \in \text{sing}(X)} \sum_{i=0}^n (-1)^i \zeta_{T_i, q}(\mathcal{P}_{abs/rel, i})(0). \quad (77)$$

Before to conclude the section we make the following remarks.

In the same hypothesis of Theorem 5 consider a point $q \in \text{sing}(X)$ such that q is an attractive simple fixed point. We recall that over a neighborhood $U_q \cong [0, 2) \times L_q$ of q we can look at f as a map given by $(rA(r, p), B(r, p)) : [0, 2) \times L_q \rightarrow [0, 2) \times L_q$ with A and B smooth up to 0. From Definition 16 we know that q is attractive if $A(0, p) < 1$ for each fixed $p \in L_q$. Clearly this implies that $f(U_q) \subset U_q$. Therefore it follows that, if we consider the complex

$$0 \rightarrow C_c^\infty(U_q, E_0|_{U_q}) \xrightarrow{P_0} C_c^\infty(U_q, E_1|_{U_q}) \xrightarrow{P_1} \dots \xrightarrow{P_{n-1}} C_c^\infty(U_q, E_n|_{U_q}) \xrightarrow{P_n} 0 \quad (78)$$

then T is also a geometric endomorphism of (78) and, using Proposition 10, we get that T extends as a bounded endomorphism of the complexes $(L^2(U_q, E_i|_{U_q}), (P|_{U_q})_{max/min, i})$. Moreover, by the results proved in the first and the second chapter of [26], it follows that $(L^2(U_q, E_i|_{U_q}), (P|_{U_q})_{max/min, i})$ are both Fredholm complexes and that, the respective heat operators, $e^{-t(\mathcal{P}|_{U_q})_{abs/rel, i}} : L^2(U_q, E_i|_{U_q}) \rightarrow L^2(U_q, E_i|_{U_q})$, are trace-class operators. Using again the properties stated in [26] at pag. 42-43, it follows that for each open neighborhood V_q of q , such that \bar{V}_q is a subset of U_q , we have

$$\begin{aligned} & \lim_{t \rightarrow 0} \int_{V_q} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel, i}}(rA(r, p), B(r, p), r, p)) dvol_g = \\ & = \lim_{t \rightarrow 0} \int_{V_q} \text{tr}(\phi_i \circ e^{-t(\mathcal{P}|_{U_q - \{q\}})_{abs/rel, i}}(rA(r, p), B(r, p), r, p)) dvol_g. \end{aligned}$$

Suppose now that we are in the hypothesis of Theorem 7. By the proof of the same theorem, it follows that for each $0 < b \leq 2$

$$\begin{aligned} & \lim_{t \rightarrow 0} \int_0^b \int_{L_q} \text{tr}(\phi_i \circ e^{-t(\mathcal{P}|_{U_q - \{q\}})_{abs/rel, i}}(rA(p), B(p), r, p)) r^m dvol_h dr = \\ & \int_0^\infty x^{-1} dx \int_{L_q} \text{tr}(\phi_i \circ e^{-x(\mathcal{P}|_{U_q - \{q\}})_{abs/rel, i}}(A(p), B(p), 1, p)) dvol_h \end{aligned}$$

that is it does not depend on the particular b we fixed. Therefore we can conclude that

$$\begin{aligned} & \lim_{t \rightarrow 0} \int_{U_q - \{q\}} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel, i}}(rA(p), B(p), r, p)) dvol_g = \\ & = \lim_{t \rightarrow 0} \int_{U_q - \{q\}} \text{tr}(\phi_i \circ e^{-t(\mathcal{P}|_{U_q - \{q\}})_{abs/rel, i}}(rA(p), B(p), r, p)) dvol_g. \end{aligned} \quad (79)$$

Summarizing we obtained that it makes sense to define, for an attractive simple fixed point, $L_{2,max/min}(T|_{U_q})$ as the L^2 -Lefschetz numbers of T acting on the maximal/minimal extension of (78) and that, under the hypothesis of Theorem 7, it satisfies

$$L_{2,max/min}(T|_{U_q}) = \lim_{t \rightarrow 0} \sum_{i=0}^n (-1)^i \int_{U_q - \{q\}} \text{tr}(\phi_i \circ e^{-t\mathcal{P}_{abs/rel,i}}(rA(p), B(p), r, p)) dvol_g. \quad (80)$$

Now we proceed making another remark before the conclusion.

As showed in the second section, T_i^* , the adjoint of T_i , has the following form:

$$T_i^* = \theta_i \circ (f^{-1})^* \quad (81)$$

where $\theta_i = \tau\phi_i^*$ with τ positive or negative function respectively if f preserves or reverses the orientation. Moreover, a simple computation, shows that T^* is an endomorphism of the following Fredholm complexes: $(L^2(M, E_i), P_{max/min,i}^t)$. By the fact that, if $Q : H \rightarrow H$ is a trace-class operator acting on the Hilbert space H then also Q^* is trace-class and $\text{Tr}(Q) = \text{Tr}(Q^*)$, it follows that

$$\text{Tr}(T_i \circ e^{-t\mathcal{P}_{abs/rel,i}}) = \text{Tr}(e^{-t\mathcal{P}_{abs/rel,i}} \circ T_i^*) = \text{Tr}(T_i^* \circ e^{-t\mathcal{P}_{abs/rel,i}}). \quad (82)$$

In particular, from (82), it follows that:

$$L_{2,max/min}(T) = L_{2,min/max}(T^*) \quad (83)$$

where T acts on $(L^2(M, E_i), P_{max/min,i})$ and T^* acts on $(L^2(M, E_i), P_{min/max,i}^t)$.

A second consequence is the following: consider a point $q \in \text{sing}(X)$ such that q is a repulsive simple fixed point. Clearly, by the fact that f on $U_q \cong C_2(L_q)$ takes the form $f = (rA(p), B(p))$ it follows that $f^{-1} = (rG(p), B^{-1}(p))$ where $G = \frac{1}{A \circ B^{-1}}$. The fact that q is repulsive means that $A > 1$. Therefore it follows that q is an **attractive** simple fixed point for T^* .

Finally we are in position to conclude with the following results:

Corollary 6. *In the same hypothesis of Theorem 7; Suppose moreover that $q \in \text{sing}(X)$ is an attractive fixed point. Then*

$$\sum_{i=0}^n (-1)^i \zeta_{T_i,q}(\mathcal{P}_{abs/rel,i})(0) = L_{2,max/min}(T|_{U_q}).$$

In particular this tells us that $\sum_{i=0}^n (-1)^i \zeta_{T_i,q}(\mathcal{P}_{abs/rel,i})(0)$ has a geometric meaning itself.

Proof. It follows immediately from Theorem 7 and (80). \square

Theorem 8. *In the same hypothesis of Theorem 6. Suppose moreover that the first property stated at the beginning of the section holds. Then we have:*

$$L_{2,max/min}(T) = \sum_{p \in \text{Fix}(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\phi_i)}{|\det(Id - d_q f)|} + \sum_{q \in \text{sing}(X)} \sum_{i=0}^n (-1)^i \zeta_{T_i,q}(\mathcal{P}_{abs/rel,i})(0) \quad (84)$$

where in (84) the contribution given by the singular points is calculated fixing any conic metric g on $\text{reg}(X)$ and any metrics ρ_0, \dots, ρ_n on E_0, \dots, E_n which satisfy the hypothesis of Theorem 7.

Moreover if each point $q \in \text{sing}(X)$ is an attractive fixed point we have:

$$L_{2,max/min}(T) = \sum_{p \in \text{Fix}(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\phi_i)}{|\det(Id - d_q f)|} + \sum_{q \in \text{sing}(X)} L_{2,max/min}(T|_{U_q}). \quad (85)$$

while if each $q \in \text{sing}(X)$ is a repulsive fixed point then we have :

$$L_{2,max/min}(T) = \sum_{p \in \text{Fix}(f) \cap M} \sum_{i=0}^n \frac{(-1)^i \text{Tr}(\theta_i)}{|\det(Id - d_q (f^{-1}))|} + \sum_{q \in \text{sing}(X)} L_{2,min/max}(T^*|_{U_q}). \quad (86)$$

Finally we remark again that, when \mathcal{P}_i is a generalized Laplacian, the contribution given by the singular simplex fixed points, that is

$$\mathcal{L}_{max/min}(T, \mathcal{S}) = \sum_{q \in \text{sing}(X)} \sum_{i=0}^n (-1)^i \zeta_{T_i, q}(\mathcal{P}_{abs/rel, i})(0)$$

does not depend on the particular conic metric that we fix on $\text{reg}(X)$ and on the metrics ρ_0, \dots, ρ_n that we fix on E_0, \dots, E_n .

Proof. As showed in Corollary 5, when each \mathcal{P}_i is a generalized Laplacian, then $L_{2, max/min}(T)$, $\mathcal{L}(T, \mathcal{R})$ and $\mathcal{L}_{max/min}(T, \mathcal{S})$ do not depend on the conic metric we fix on $\text{reg}(X)$ and do not depend on the metrics we fix ρ_0, \dots, ρ_n on E_0, \dots, E_n . Therefore, without loss of generality, we can assume that for each $q \in \text{sing}(X)$, using the isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$ of (69), the conic metric g satisfies $g = dr^2 + r^2h$ with h that does not depend on r and that each metric ρ_i on E_i does not depend on r in a neighborhood of $\partial\bar{M}$. In this way we are in position to apply Theorem 7 and so (84) follows combining the theorems 6 and 7. Moreover this tell us that, in (84), the contribution of the singular points is well defined and does not depend on the metrics g, ρ_0, \dots, ρ_n (satisfying the assumptions of Theorem 7) used to calculate it. The second assertion follows from Corollary 6 while the last assertion follows from (81) and (83). \square

Remark 5. We stress on the fact that, unlike Theorem 7, in Theorem 8 there are not assumptions about the conic metric g on $\text{reg}(X)$ and about the metrics ρ_0, \dots, ρ_n on E_0, \dots, E_n respectively.

Finally we conclude the section with the following comment.

The condition that we required at the beginning of the subsection for each operator P_i , that each A_k does not depend on x , might appear as to be too strong at first right. Obviously this is indeed a strong assumption but it is at the same time quite natural because the most natural complex arising in differential geometry, the de Rham complex, satisfies this assumption.

The requirement (69), about the behavior of f near the point p , is justified by the idea to evaluate $\mathcal{L}_{max/min}(T, \mathcal{S})$ using the scaling invariance of the heat kernel, see Lemma 1. In fact if $f = (rA(r, p), B(r, p))$ then, after the scaling invariance is used, we get in our expression the term $\text{tr}(\phi_i \circ e^{-tr^{-2\nu}\mathcal{P}_{abs/rel, i}}(A(r, p), B(r, p), 1, p))$. To have that this last expression make sense we need that $(A(r, p), B(r, p), 1, p) \in \mathcal{G}(f)$, where $\mathcal{G}(f) \subset X \times X$ is the graph of f , and therefore this leads us to assume (69).

4.1 The case of a short complex

The aim of this subsection is to give a formula for the L^2 -Lefschetz numbers in the particular case of a short complex, that is is an elliptic conic operator $P : C_c^\infty(M, E) \rightarrow C_c^\infty(M, E)$, using the result stated in Proposition 9. To do this we start describing our geometric situation which is the same of the previous results with some additional requirements: let X be a compact and oriented manifold with conical singularities of dimension $m + 1$. Let M be its regular part and let \bar{M} be the compact manifold with boundary which desingularize X . Endow M with a conic metric g . Let (E, ρ) be a vector bundle endowed with a metric (riemannian or hermitian) according if E is complex or real. Let (\bar{E}, ρ) be the extension of (E, ρ) over \bar{M} . Let $T = (T_1, T_2)$ be a geometric endomorphism where, as we already know, $T_i = \phi_i \circ f^*$ with $f : \bar{M} \rightarrow \bar{M}$ is a diffeomorphism as described in Definition 13 and $\phi : f^*E \rightarrow E$ a bundle homomorphism. Suppose that $\text{Fix}(f)$ is made only by simple fixed points. Finally, suppose that in each neighborhood $U_q \cong C_2(L_q)$ of $q \in \text{sing}(X)$ the operator P take the form

$$P = \frac{n}{2r} + \frac{\partial}{\partial r} + \frac{1}{r}S \quad (87)$$

where $S \in \text{Diff}^1(N, E_N)$ is an elliptic operator and the map f take the form

$$f = (rc, B(p)), \quad c \neq 1 \quad (88)$$

where $c > 0$ and depends only on q .

Theorem 9. *In the same hypothesis of Theorem 7; suppose moreover that the properties described above hold. Then for each $q \in \text{sing}(X)$ we have:*

$$\zeta_{T_0,q}(P_{max}^t \circ P_{min})(0) = \frac{c^{\frac{1-n}{2}}}{4} \int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^+(\lambda)}\left(\frac{uc}{2}\right) du \text{Tr}(\tilde{\Phi}_{0,\lambda,q}) \quad (89)$$

and analogously

$$\zeta_{T_1,q}(P_{min} \circ P_{max}^t)(0) = \frac{c^{\frac{1-n}{2}}}{4} \int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^-(\lambda)}\left(\frac{uc}{2}\right) du \text{Tr}(\tilde{\Phi}_{1,\lambda,q}) \quad (90)$$

where

$$\text{Tr}(\tilde{\Phi}_{j,\lambda,q}) = \int_{L_q} \text{tr}(\phi_j \Phi_{\lambda,q}(B(p), p)) d\text{vol}_h, \quad j = 0, 1.$$

Proof. We give the proof only for (89) because for (90) is completely analogous. To prove the assertion we have to calculate

$$\lim_{t \rightarrow 0} \int_{\text{reg}(C_2(L_q))} \text{tr}(T_0 \circ e^{-P_{max}^t \circ P_{min}}) d\text{vol}_g.$$

By the assumptions we are in position to use the second statement of Proposition 9 and therefore it is clear that the smooth kernel of $T_0 \circ e^{-P_{max}^t \circ P_{min}}$ is

$$\sum_{\lambda \in \text{spec } S} \frac{1}{2t} (crs)^{\frac{1-n}{2}} I_{p^+(\lambda)}\left(\frac{crs}{2t}\right) e^{-\frac{c^2 r^2 + s^2}{4t}} \phi_0 \Phi_\lambda(B(p), q) \quad (91)$$

In this way we have to calculate

$$\lim_{t \rightarrow 0} \int_0^2 \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (cr^2)^{\frac{1-n}{2}} I_{p^+(\lambda)}\left(\frac{cr^2}{2t}\right) e^{-\frac{r^2(c^2+1)}{4t}} r^m dr \int_{L_q} \text{tr}(\phi_0 \Phi_\lambda(B(p), q)) d\text{vol}_h.$$

Clearly $\int_{L_q} \text{tr}(\phi_0 \Phi_\lambda(B(p), q)) d\text{vol}_h$ does not depend on t and so, if we label it $\text{Tr}(\tilde{\Phi}_{0,\lambda,q})$, our task now is to calculate

$$\lim_{t \rightarrow 0} \int_0^2 \sum_{\lambda \in \text{spec } S} \frac{1}{2t} (cr^2)^{\frac{1-n}{2}} I_{p^+(\lambda)}\left(\frac{cr^2}{2t}\right) e^{-\frac{r^2(c^2+1)}{4t}} r^m dr.$$

To do this put $\frac{r^2}{t} = u$. Then $r dr = \frac{t du}{2}$. Moreover when r goes to 2 u goes to $\frac{4}{t}$ while when r goes to 0 u goes to zero. So, applying this change of variable, we get

$$\lim_{t \rightarrow 0} \frac{c^{\frac{1-n}{2}}}{4} \int_0^{\frac{4}{t}} e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^+(\lambda)}\left(\frac{uc}{2}\right) du.$$

Now, by the asymptotic behavior of the integrand, we know that this limit exists and is equal to

$$\frac{c^{\frac{1-n}{2}}}{4} \int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^+(\lambda)}\left(\frac{uc}{2}\right) du.$$

So we proved the statement. \square

From Theorem 9 we have the following immediate corollary:

Corollary 7. *In the same hypothesis of Theorem 9 but without any assumptions about the conic metric g on $\text{reg}(X)$ and the metric ρ on E . Suppose moreover that $P^t \circ P : C_c^\infty(M, E) \rightarrow C_c^\infty(M, E)$ is a generalized Laplacian. Then we have the following formula:*

$$L_{2,min}(T) = \sum_{q \in M \cap \text{Fix}(f)} \sum_{j=0}^1 \frac{(-1)^j \text{Tr}(\phi_j)}{|\det(\text{Id} - d_q f)|} + \quad (92)$$

$$\begin{aligned}
& + \sum_{q \in \text{sing}(X)} \frac{c^{\frac{1-n}{2}}}{4} \int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^+(\lambda)}\left(\frac{uc}{2}\right) du \text{Tr}(\tilde{\Phi}_{0,\lambda,q}) + \\
& - \sum_{q \in \text{sing}(X)} \int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_{\lambda \in \text{spec } S} I_{p^-(\lambda)}\left(\frac{uc}{2}\right) du \text{Tr}(\tilde{\Phi}_{1,\lambda,q})
\end{aligned}$$

where the contribution of the singular points is calculated fixing any conic metric g on $\text{reg}(X)$ and any metric ρ on E which satisfy the assumptions of Theorem 9.

Proof. As observed in the proof of Theorem 8, by the fact that $P^t \circ P$ is a generalized Laplacian, it follows that $\mathcal{L}(T, \mathcal{S})$ does not depend on the conic metric we fix on $\text{reg}(X)$ and does not depend on the metric ρ we fix on E . Therefore, without loss of generality, we can assume that for each $q \in \text{sing}(X)$, using the isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$ of (69), the conic metric g satisfies $g = dr^2 + r^2h$ with h that does not depend on r and that each metric ρ_i on E_i does not depend on r in a neighborhood of $\partial\bar{M}$. In this way we are in position to apply Theorem 9 and therefore (92) follows. \square

5 A thorough analysis of the de Rham case

5.1 Applications of the previous results

As remarked previously, Theorems 6 and 8, Corollary 6 and in particular (84) hold for the Hilbert complexes $(L^2\Omega^i(M, g), d_{\max/\min, i})$. More explicitly, we have the following result:

Theorem 10. *Let X be a compact and oriented manifold with isolated conical singularities and of dimension $m+1$. Let g be a conic metric over its regular part $\text{reg}(X)$. Let $f : X \rightarrow X$ be a map induced by a diffeomorphism $f : \bar{M} \rightarrow \bar{M}$ such that $f : X \rightarrow X$ fixes each singular point of X . Consider $T := (df)^* \circ f^*$, the natural endomorphism of the de Rham complex induced by f . Finally suppose that f has only simple fixed points. Then we have:*

$$L_{2,\max/\min}(T) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_q f) + \mathcal{L}_{\max/\min}(T, \mathcal{S}). \quad (93)$$

If in a neighborhood of each simple fixed point q f satisfies the condition described in (69), then we have: $L_{2,\max/\min}(T) =$

$$= \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_q f) + \sum_{q \in \text{sing}(X)} \sum_{i=0}^{m+1} (-1)^i \zeta_{T_i, q}(\Delta_{\text{abs/rel}, i})(0) \quad (94)$$

where in (94) the contribution of the singular points is calculated using any conic metric g on $\text{reg}(X)$ such that, again through the isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$ of (69), g takes the form $dr^2 + r^2h$ and h does not depend on r .

In particular if each $q \in \text{sing}(X)$ is an attractive simple fixed point then we have:

$$L_{2,\max/\min}(T) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_q f) + \sum_{q \in \text{sing}(X)} L_{2,\max/\min}(T|_{U_q}). \quad (95)$$

while if each $q \in \text{sing}(X)$ is a repulsive simple fixed point then we have:

$$L_{2,\max/\min}(T) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_q(f^{-1})) + \sum_{q \in \text{sing}(X)} L_{2,\min/\max}(T^*|_{U_q}). \quad (96)$$

Moreover in (94) the member on the right, that is $\mathcal{L}_{\max/\min}(T, \mathcal{S})$, does not depend on the particular conic metric that we fix on $\text{reg}(X)$.

Proof. (93) follows immediately from Theorem 6. In particular the expression for $\mathcal{L}_{\max/\min}(T, \mathcal{R})$ follows by a standard argument of linear algebra; see for example [3] or [32]. (94) follows as in the proof of Theorem (8); in particular, as remarked in the proof of Lemma 1, the scaling invariance property for the heat operator associated to positive self-adjoint extension of Δ_i , was proved by Cheeger in [15]. Finally (95) and (96) follows again from Theorem 8. \square

By the assumptions on f it follows that $f(\text{sing}(X)) = \text{sing}(X)$ and $f(\text{reg}(X)) = \text{reg}(X)$. This implies, see for example [20], that if we fix a perversity p then f induces a well defined map, f^* , between the intersection cohomology groups respect to the perversity p . In particular we have $f^* : I^{\overline{m}}H^i(X) \rightarrow I^{\overline{m}}H^i(X)$ and $f^* : I^mH^i(X) \rightarrow I^mH^i(X)$. Therefore it is natural to define in this context, as it is showed in [22], the **intersection Lefschetz number** respects to a given perversity p as

$$I^pL(f) = \sum_{i=0}^n \text{tr}(f^* : I^pH^i(X) \rightarrow I^pH^i(X)). \quad (97)$$

$I^pL(f)$ is deeply studied, from a topological point of view, in [22] and [23] in the more general context of a stratified pseudomanifold; our goal in the next corollaries is to give an analytic description of $I^{\overline{m}}L(f)$ and $I^mL(f)$ when X is a compact manifold with conical singularities. In particular in (102) we will give an analytic proof of a formula already proved in [22]. So, using Theorem 93 and Theorem 37, we get the following results:

Proposition 12. *In the same hypothesis of Theorem 10; let $q \in \text{sing}(X)$ be an attractive fixed point. Let U_q be an open neighborhood of q isomorphic to $C_2(L_q)$ and suppose that f satisfies (69) and g takes the form $g = dr^2 + r^2h$ where h does not depend on r . Then, for $i < \frac{m+1}{2}$, we have:*

$$\text{Tr}((f|_{U_q})^* : H_{2,\max}^i(U_q, g|_{U_q}) \rightarrow H_{2,\max}^i(U_q, g|_{U_q})) = \text{Tr}(B^* : H^i(L_q) \rightarrow H^i(L_q)) \quad (98)$$

Proof. As it is showed in [14], in (37) the isomorphism between $H_{2,\max}^i(\text{reg}(C_2(L_q)), g)$ and $H^i(L_q)$, for $i < \frac{m}{2} + \frac{1}{2}$, is given by the pull-back π^* where $\pi : (0, b) \times F \rightarrow F$ is the projection on the second factor and inverse is given by v_a , the evaluation map in a , where a is any point $(0, 2)$. Now by the hypothesis, over U_q f can be written as $(rA(p), B(p))$. An immediate check shows that $\pi^* \circ B^* = B^* \circ \pi^*$ and therefore $\text{Tr}((f|_{U_q})^*) = \text{Tr}(B^*)$. \square

Corollary 8. *In the same hypothesis of Theorem 10, suppose moreover that near each point $q \in \text{sing}(X)$ f satisfies (69). Then we have:*

$$I^mL(f) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_qf) + \sum_{q \in \text{sing}(X)} \sum_{i=0}^{m+1} (-1)^i \zeta_{T_i, q}(\Delta_{\text{abs}, i})(0) \quad (99)$$

and analogously

$$I^{\overline{m}}L(f) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_qf) + \sum_{q \in \text{sing}(X)} \sum_{i=0}^{m+1} (-1)^i \zeta_{T_i, q}(\Delta_{\text{rel}, i})(0) \quad (100)$$

Finally, if $q \in \text{sing}(X)$ is an attractive fixed point, then we have

$$\sum_{i=0}^{m+1} (-1)^i \zeta_{T_i, q}(\Delta_{\text{abs}, i})(0) = \sum_{i < \frac{m}{2} + \frac{1}{2}} (-1)^i \text{tr}(B^* : H^i(L_q) \rightarrow H^i(L_q)) \quad (101)$$

and therefore from (99) we get:

$$I^mL(f) = L_{2,\max}(T) = \sum_{q \in \text{Fix}(f) \cap \text{reg}(X)} \text{sgn det}(Id - d_qf) + \sum_{q \in \text{sing}(X)} \sum_{i < \frac{m+1}{2}} (-1)^i \text{Tr}(B^* : H^i(L_q) \rightarrow H^i(L_q)). \quad (102)$$

Proof. As in Theorem 10, to get the Lefschetz numbers, we can use a conic metric g such that, in each neighborhood U_q of $q \in \text{sing}(X)$, using the isomorphism $\chi_q : U_q \rightarrow C_2(L_q)$, g takes the form $g = dr^2 + r^2h$ where h does not depend on r . Now (99) and (100) follow immediately by the previously theorems. Finally (101) and (102) follow immediately from Proposition 12. \square

Finally we have this last corollary; before stating it we recall that a manifold with conical singularities of dimension $m+1$ is a **Witt** space if $m+1$ is even or, when it is odd, if $H^{\frac{m}{2}}(L_q) = 0$ for each link L_q . For more details see, for example, [20].

Corollary 9. *In the same hypothesis of Corollary 8. Suppose moreover that X is a Witt space. Then we get:*

$$L_{2,max}(T) = L_{2,min}(T), \quad \mathcal{L}_{max}(T, \mathcal{S}) = \mathcal{L}_{min}(T, \mathcal{S}) \quad (103)$$

and, if each $q \in \text{sing}(X)$ is an attractive fixed point then

$$\begin{aligned} \mathcal{L}_{max}(T, \mathcal{S}) &= \mathcal{L}_{min}(T, \mathcal{S}) = \sum_{q \in \text{sing}(X)} L_{2,max}(T|_{U_q}) = \sum_{q \in \text{sing}(X)} L_{2,min}(T|_{U_q}) = \\ &= \sum_{q \in \text{sing}(X)} \sum_{i < \frac{m+1}{2}} (-1)^i \text{Tr}(B_a^* : H^i(L_q) \rightarrow H^i(L_q)). \end{aligned} \quad (104)$$

Finally if each $q \in \text{sing}(X)$ is repulsive then we have:

$$\mathcal{L}_{max}(T, \mathcal{S}) = \mathcal{L}_{min}(T, \mathcal{S}) = \sum_{q \in \text{sing}(X)} L_{2,max}(T^*|_{U_q}) = \sum_{q \in \text{sing}(X)} L_{2,min}(T^*|_{U_q}). \quad (105)$$

Proof. (103) follows by the fact that, as it is showed in [14], if X is a Witt space then for each i , $\Delta_i : \Omega_c^i(\text{reg}(X)) \rightarrow \Omega_c^i(\text{reg}(X))$ is essentially self-adjoint as unbounded operator acting on $L^2\Omega(\text{reg}(X), g)$ and this implies that $d_{max,i} = d_{min,i}$ for $i = 0, \dots, m+1$. (104) follows by (103) combined with (95) and (102). Finally (105) follows from the fact that X is Witt and from Theorem 8. \square

5.2 Some further results arising from Cheeger's work on the heat kernel

The aim of this section is to approach the L^2 -Lefschetz numbers of the L^2 -de Rham complex using the results of Cheeger stated in [14] and in [15]. For simplicity assume that X is a Witt space. As recalled previously, if X is a Witt space and if over $\text{reg}(X)$ we put a conic metric, then $\Delta_i : L^2\Omega^*(\text{reg}(X), g) \rightarrow L^2\Omega^*(\text{reg}(X), g)$ is essentially self-adjoint for each $i = 0, \dots, m+1$, with core domain given by the smooth compactly supported forms. In particular this implies that, if $\dim X = m+1$, then for each $i = 0, \dots, m+1$, $d_{max,i} = d_{min,i}$. Therefore, for each map $f : X \rightarrow X$ that induces a geometric endomorphism T as in Theorem 10, we have just one L^2 -Lefschetz number that we label $L_2(T)$.

Now we recall briefly the results we need and we refer to [14] and in particular to [15], section 3, for the complete details and for the proofs. Let N be an oriented closed manifold of dimension m and let $C(N)$ be the cone over N . Endow $\text{reg}(C(N))$ with a conic metric $g = dr^2 + r^2h$ where h is a riemannian metric over N . In the mentioned papers Cheeger introduce four types of differential forms over $\text{reg}(C(N))$, called forms of type 1, 2, 3 and 4, such that each eigenform of Δ_i , the Laplacian acting on the i -forms over $\text{reg}(C(N))$, can be expressed as convergent sum of these forms. For the definition of these forms see [15] pag. 586-588.

The main reason to introduce these four types of forms is that now we can break the heat operator in four pieces, see [15] pag. 90-92:

$$e^{-t\Delta_i} = {}_1e^{-t\Delta_i} + {}_2e^{-t\Delta_i} + {}_3e^{-t\Delta_i} + {}_4e^{-t\Delta_i}$$

where, for each $l = 1, \dots, 4$, ${}_l e^{-t\Delta_i}$ is the heat operator built using the i -forms of type l . As it is showed in [15], pag. 590-592, it is possible to give an explicit expression for ${}_l e^{-t\Delta_i}$. In particular for type 1 forms we have:

$${}_1e^{-t\Delta_i} = (r_1 r_2)^{a(i)} \sum_j \int_0^\infty e^{-t\lambda^2} J_{\nu_j(i)}(\lambda r_1) J_{\nu_j(i)}(\lambda r_2) \lambda d\lambda \phi_j^i(p_1) \otimes \phi_j^i(p_2) = \quad (106)$$

$$= (r_1 r_2)^{a(i)} \sum_j \frac{1}{2t} e^{-\frac{r_1^2 + r_2^2}{4t}} I_{\nu_j(i)}\left(\frac{r_1 r_2}{2t}\right) \phi_j^i(p_1) \otimes \phi_j^i(p_2) \quad (107)$$

where $I_{\nu_j(i)}$ is the modified Bessel function (see [26] pag. 67), $a(i) = \frac{1}{2}(1 + 2i - m)$, $\nu_j(i) = (\mu_j + a^{\pm}(i))^{\frac{1}{2}}$ and $a^{\pm}(i) = a(i) \pm \nu_j(i)$. The corresponding expression for type 2 forms is

$${}_2e^{-t\Delta_i} = \sum_j d_1 d_2 ((r_1 r_2)^{a(i-1)}) \int_0^\infty e^{-t\lambda^2} J_{\nu_j(i-1)}(\lambda r_1) J_{\nu_j(i-1)}(\lambda r_2) \lambda^{-1} d\lambda \phi_j^{i-1}(p_1) \otimes \phi_j^{i-1}(p_2) \quad (108)$$

The expression for forms of type 3 is:

$$\begin{aligned}
{}_3e^{-t\Delta_i} &= \sum_j \int_0^\infty e^{-t\lambda^2} ((-a(i-1)r_1^{a(i-1)})J_{\nu_j(i-1)}(\lambda r_1) + r_1^{a(i-1)+1}J'_{\nu_j(i-1)}(\lambda r_1)\lambda) \frac{d\phi_j^{i-1}(p_1)}{\sqrt{\mu_j}} \\
&\quad + r_1^{a(i-1)-1}J_{\nu_j(i-1)}(\lambda r_1)dr_1 \wedge \sqrt{\mu_j}\phi_j^{i-1}(p_1) \otimes ((-a(i-1)r_2^{a(i-1)})J_{\nu_j}(\lambda r_2) + \\
&\quad + r_2^{a(i-1)+1}J'_{\nu_j(i-1)}(\lambda r_2)\lambda) \frac{d\phi_j^{i-1}(p_2)}{\sqrt{\mu_j}} + r_2^{a(i-1)-1}J_{\nu_j}(\lambda r_2)dr_2 \wedge \sqrt{\mu_j}\phi_j^{i-1}(p_2))\lambda^{-1}d\lambda
\end{aligned} \tag{109}$$

Finally for forms of type 4 we have:

$$\begin{aligned}
{}_4e^{-t\Delta_i} &= (r_1r_2)^{a(i-1)} \sum_j \int_0^\infty e^{-t\lambda^2} J_{\nu_j(i-2)}(\lambda r_1)J_{\nu_j(i-2)}(\lambda r_2)\lambda d\lambda dr_1 \wedge \frac{d\phi_j^{i-2}(p_1)}{\sqrt{\mu_j}} \otimes dr_2 \wedge \frac{d\phi_j^{i-2}(p_2)}{\sqrt{\mu_j}} = \\
&= (r_1r_2)^{a(i-2)} \sum_j \frac{1}{2t} e^{-\frac{r_1^2+r_2^2}{4t}} I_{\nu_j(i-2)}\left(\frac{r_1r_2}{2t}\right) dr_1 \wedge \frac{d\phi_j^{i-2}(p_1)}{\sqrt{\mu_j}} \otimes dr_2 \wedge \frac{d\phi_j^{i-2}(p_2)}{\sqrt{\mu_j}}
\end{aligned} \tag{110}$$

Now suppose that for each point $q \in \text{sing}(X)$, over a neighborhood $U_q \cong C_2(L_q)$, f satisfies (88). Using Cheeger's results recalled above, it make sense to break $T \circ e^{-t\Delta_i}$, over $C_2(L_q)$, as a sum of four pieces such that:

$$\lim_{t \rightarrow 0} \text{Tr}(T \circ e^{-t\Delta_i}) = \lim_{t \rightarrow 0} \text{Tr}(T \circ {}_1e^{-t\Delta_i} + T \circ {}_2e^{-t\Delta_i} + T \circ {}_3e^{-t\Delta_i} + T \circ {}_4e^{-t\Delta_i}). \tag{111}$$

Moreover, using (54), (88), (107) and (110) it is clear that on $\text{reg}(C_2(L_q))$ we have:

$$\text{tr}(T \circ {}_1e^{-t\Delta_i})(r, p) = (cr^2)^{a(i)} \sum_j \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} I_{\nu_j(i)}\left(\frac{cr^2}{2t}\right) \text{tr}(B^*\phi_j^i \otimes B^*\phi_j^i) \tag{112}$$

and analogously

$$\text{tr}(T \circ {}_4e^{-t\Delta_i})(r, p) = (cr^2)^{a(i-2)} \sum_j \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} I_{\nu_j(i-2)}\left(\frac{cr^2}{2t}\right) \text{tr}\left(dr \wedge \frac{d(B^*\phi_j^{i-2})}{\sqrt{\mu_j}} \otimes dr \wedge \frac{d(B^*\phi_j^{i-2})}{\sqrt{\mu_j}}\right) \tag{113}$$

Now we are in position to state the following result:

Theorem 11. *Let X , g and f be as in Theorem 10 such that $\dim X = m + 1$. Suppose moreover that X is a Witt space and that, on each neighborhood $U_q \cong C_2(L_q)$ of each point $q \in \text{sing}(X)$, f satisfies (88) and g takes the form $g = dr^2 + r^2h$ where h does not depend on r . Then, for each $q \in \text{sing}(X)$, we have:*

1. *The forms of type 1 give a contribution only in degree 0.*
2. *The contribution given by q in degree zero depends only on the forms of type 1 and we have*

$$\zeta_{T_0, q}(\Delta_0)(0) = \frac{c^{\frac{1-m}{2}}}{4} \left(\int_0^\infty e^{-u(c^2+1)} \sum_j I_{\nu_j(0)}\left(\frac{cu}{2}\right) du \right) (\text{Tr}(B^*\phi_j^i \otimes B^*\phi_j^i)) \tag{114}$$

3. *The forms of type 4 give a contribution only in degree 2 and this contribution is*

$$\text{Tr}(T_2 \circ {}_4e^{-t\Delta_2}) = \frac{c^{\frac{1-m}{2}}}{4} \left(\int_0^\infty e^{-\frac{u(c^2+1)}{4}} \sum_j I_{\nu_j(0)}\left(\frac{cu}{2}\right) du \right) (\text{Tr}(dr \wedge \frac{d(B^*\phi_j^{i-2})}{\sqrt{\mu_j}} \otimes dr \wedge \frac{d(B^*\phi_j^{i-2})}{\sqrt{\mu_j}})) \tag{115}$$

where $\text{Tr}(T_2 \circ {}_4e^{-t\Delta_2})$ is taken over $\text{reg}(C_2(L_q))$.

4. *The contribution given by q in the others degrees, that is $i \neq 0, 2$, depends only on the forms of type 2 and 3.*

Proof. First of all we note that from (107), (108), (109) and (110) it follows that ${}_1e^{-t\Delta_i} = e^{-t\Delta_i}$ for $i = 0$ and that ${}_4e^{-t\Delta_i}$ occurs only for $i \geq 2$. Now, using (107) and (112) we know that, over $\text{reg}(C_2(L_q))$,

$$\lim_{t \rightarrow 0} \text{Tr}(T_i \circ {}_1e^{-t\Delta_i}) = \lim_{t \rightarrow 0} \int_0^2 \int_{L_q} (cr^2)^{a(i)} \sum_j \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} I_{\nu_j(i)}\left(\frac{cr^2}{2t}\right) \text{tr}(B^* \phi_j^i \otimes B^* \phi_j^i) r^m dr d\text{vol}_h.$$

Clearly this last term it is in turn equal to

$$\lim_{t \rightarrow 0} \left(\int_0^2 (cr^2)^{a(i)} \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} \sum_j I_{\nu_j(i)}\left(\frac{cr^2}{2t}\right) r^m dr \right) (\text{Tr}(B^* \phi_j^i \otimes B^* \phi_j^i)) \quad (116)$$

and therefore, to get the first two points we have to calculate

$$\lim_{t \rightarrow 0} \int_0^2 (cr^2)^{a(i)} \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} \sum_j I_{\nu_j(i)}\left(\frac{cr^2}{2t}\right) r^m dr \quad (117)$$

First of all remember that $a(i) = \frac{1}{2}(1 - m + 2i)$; therefore $r^{2a(i)} r^m = r^{2i+1}$. Now put $\frac{r^2}{t} = u$. It follows immediately that $dr = \frac{t du}{2r}$. Now, by the fact that $r^2 = tu$ it follows that $r^{2i+1} = t^i u^i r$ and therefore we also get $r^{2i+1} dr = \frac{t^{i+1} u^i du}{2}$. Moreover when r goes to 2 then u goes to $\frac{2}{t}$ and when r goes to 0 then u goes to 0. In this way we have

$$\lim_{t \rightarrow 0} \frac{c^{a(i)}}{4} t^i \int_0^{\frac{2}{t}} e^{-u(c^2+1)} \sum_j I_{\nu_j(i)}\left(\frac{c^2 u}{2}\right) u^i du \quad (118)$$

Now, by the asymptotic behavior of the integrand it follows that

$$\lim_{t \rightarrow 0} \frac{c^{a(i)}}{4} \int_0^{\frac{2}{t}} e^{-u(c^2+1)} \sum_j I_{\nu_j(i)}\left(\frac{c^2 u}{2}\right) u^i du = \frac{c^{a(i)}}{4} \int_0^\infty e^{-u(c^2+1)} \sum_j I_{\nu_j(i)}\left(\frac{c^2 u}{2}\right) u^i du.$$

Therefore we can conclude that

$$(118) = \begin{cases} \frac{c^{\frac{1-m}{2}}}{4} \int_0^\infty e^{-u(c^2+1)} \sum_j I_{\nu_j(0)}\left(\frac{c^2 u}{2}\right) du & i = 0 \\ 0 & i > 0 \end{cases} \quad (119)$$

In this way we proved the first and the second assertion. For the third statement the proof is completely analogous to the previous one. Also in this case it is clear that in order to establish the assertion we have to calculate:

$$\lim_{t \rightarrow 0} c^{a(i-2)+1} \int_0^2 \frac{1}{2t} e^{-\frac{r^2(c^2+1)}{4t}} \sum_j I_{\nu_j(i-2)}\left(\frac{cr^2}{2t}\right) r^{2i-3} dr.$$

Now if we put again $\frac{r^2}{t} = u$ the remaining part of the proof is completely analogous to that one of the first two points.

Finally the last point follows from the first three points. \square

References

- [1] P. Albin, E. Leichtnam, R. Mazzeo, P. Piazza, The signature package on Witt spaces, *Annales Scientifiques de l'École Normale Supérieure*, volume 45, 2012.
- [2] M. F. Atiyah, R. Bott, A Lefschetz fixed point formula for elliptic complexes, I. *Ann. Math.*, 86, 374-407, (1967).
- [3] M. F. Atiyah, R. Bott, A Lefschetz fixed point formula for elliptic complexes, II. *Ann. Math.*, 88, 451-491, (1968).
- [4] F. Bei, General perversities and L^2 de Rham and Hodge theorems on stratified pseudo-manifolds, to appear on *Bulletin the Sciences Mathématiques*

- [5] N. Berline, E. Getzler, M. Vergne., Heat kernels and Dirac operators, Springer-Verlag, New York 1992.
- [6] J. M. Bismut, The Atiyah-Singer theorems, a probabilistic approach. Part II: The Lefschetz fixed point formulas, *J. Funct. Anal.* 57(3), 329-348, 1984.
- [7] J. M. Bismut, Infinitesimal Lefschetz formulas: A heat equation proof, *J. Funct. Anal.* 62, 437-457, 1985.
- [8] V. A. Brenner, M. A. Shubin, The Atiyah-Bott-Lefschetz formula for elliptic complexes on manifold with boundary, *J. Soviet Math.* 64 (1993), 1069-1111.
- [9] J. Brüning, M. Lesch, Hilbert complexes, *J. Func. Anal.* 108 1992 88-132.
- [10] J. Brüning, R. Seeley, Regular Singular Asymptotics, *Adv. Math.* 58 (1985), 133-148.
- [11] J. Brüning, R. Seeley, The resolvent expansion for second order regular singular operators, *J. Func. Anal.*, 73 (1987), 369-429.
- [12] J. Brüning, R. Seeley, An index theorem for first order regular singular operators, *Amer. J. Math.* 110 (1988), 659-714.
- [13] J. Brüning, R. Seeley, The expansion of the resolvent near a singular stratum of conical type. *J. Funct. Anal.* 95 (1991), 255-290.
- [14] J. Cheeger, On the spectral geometry of spaces with cone-like singularities, *Proc. Nat. Acad. Sci. USA* 76 (1979), no. 5, 2103-2106.
- [15] J. Cheeger, Spectral geometry of singular riemannian spaces, *J. Differential Geometry*, 18, 575-6657, (1983).
- [16] N. T. Dunford, J. T. Schwartz, Linear operator Part II: Spectral theory, Wiley, 1963.
- [17] J. B. Gil, G. A. Mendoza, Adjoints of elliptic cone operators. *Amer. J. Math.* 125, 357-408, (2003).
- [18] P. B. Gilkey, Lefschetz fixed point formulas and the heat equations, *Comm. Pure Appl. Math.*, 48, 91-147, (1979).
- [19] P. B. Gilkey, Invariance theory, the heat equations and the Atiyah-Singer index theorem. *Mathematics Lecture Series*, 11. Publish or Perish, Wilmington, DE (1984).
- [20] M. Goresky, R. MacPherson, Intersection homology theory, *Topology* 19 (1980) 135-162.
- [21] M. Goresky, R. MacPherson, Intersection homology II, *Invent. Math.* 72 (1983), 77-129.
- [22] M. Goresky, R. MacPherson, Lefschetz fixed point theorem for intersection homology, *Comment. Math. Helvetici*, 60 (1985), 366-391.
- [23] M. Goresky, R. MacPherson, Local contribution to the Lefschetz fixed point formula, *Inv. Math.*, 111, 1-33 (1993).
- [24] T. Kotake, The fixed point theorem of Atiyah-Bott via parabolic operator, *Comm. Pure and Appl. Math.*, 22 (1969), 789-806.
- [25] J. Lafferty, Y. Yu, W. Zhang, A direct geometric proof of the Lefschetz fixed point formulas, *Trans. Am. Math. Soc.*, 329 (1992), 571-573.
- [26] M. Lesch, Operators of Fuchs type, conic singularities and asymptotic methods, *Teubner Texte zur Mathematik Vol. 136*, Teubner-Verlag, Leipzig, 1997.
- [27] R. B. Melrose, The Atiyah-Patodi-Singer index theorem, *Research Notes in Mathematics*, 4. A K Peters, Ltd., Wellesley, MA, 1993.
- [28] E. Mooers, Heat kernel asymptotics on manifolds with conical singularities, *J. Anal. Math.* 78 (1999), 1-36.

- [29] V. Nazaikinskii, B.-W. Schulze, B. Sternin, V. Shatalov, The Atiyah-Bott-Lefschetz theorem on manifolds with conical singularities. *Ann. Global Anal. Geom.*, 17, 409-439, (1998).
- [30] V. Nazaikinskii, B. Sternin, Lefschetz theory on manifolds with singularities, C^* -algebras and Elliptic Theory, *Trend in Mathematics*, 157-186, Birkäuser Verlag Basel/Switzerland, (2006).
- [31] V. K. Patodi, Holomorphic Lefschetz fixed point formula, *Bull. A.M.S.* 81, no. 6, 1971, 1133-1134.
- [32] J. Roe, Elliptic operators, topology and asymptotic methods. Second edition. Pitman Research Notes in Mathematics Series, 395. Longman, Harlow (1998).
- [33] B.-W. Schulze, Elliptic complexes on manifolds with conical singularities, *Teubner Texte zur Mathematik Vol. 106*, pages 170-223, Teubner-Verlag, Leipzig, (1988).
- [34] S. Seyfarth, M. A. Shubin, Lefschetz fixed point formula for manifold with cylindrical ends, *Ann. of Global Anal Geom.* 9 (1991), 99-108.
- [35] M. A. Stern, Lefschetz formulae for arithmetic varieties, *Inventiones Math.* 115, 241-296, 1994.
- [36] M. A. Stern, Fixed point theorems from a de Rham perspective, *Asian journal of Mathematics* 13, 65-88, 2009.
- [37] D. Toledo, On the Atiyah-Bott formula, *J. Diff. Geom.*, 8 (1973), 401-436.
- [38] D. Toledo, Y. Tong, Duality and intersection theory on complex manifolds, II, the holomorphic Lefschetz formula, *Ann. of Math.*, 108 (1978), 519-538.